

Washington University School of Medicine Digital Commons@Becker

Presentations

2005: Alcoholism and Comorbidity

2005

Genetic epidemiologic approaches to understanding of comorbidity of substance abuse and psychiatric disorders

Kathleen Ries Merikangas

Follow this and additional works at: <http://digitalcommons.wustl.edu/guzepresentation2005>

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Merikangas, Kathleen Ries, "Genetic epidemiologic approaches to understanding of comorbidity of substance abuse and psychiatric disorders" (2005). *Presentations*. Paper 1 Samuel B. Guze Symposium on Alcoholism.
<http://digitalcommons.wustl.edu/guzepresentation2005/1>

This Presentation is brought to you for free and open access by the 2005: Alcoholism and Comorbidity at Digital Commons@Becker. It has been accepted for inclusion in Presentations by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Genetic Epidemiologic Approaches to Understanding of Comorbidity of Substance Abuse and Psychiatric Disorders

Kathleen Ries Merikangas, Ph.D.

Senior Investigator

Section on Developmental Genetic Epidemiology



In recognition of the scientific contributions of Samuel B. Guze

- **Emphasis on empiricism in psychiatry**
- **Validation of criteria for psychiatric disorders**
(Robins E. & Guze, S. Am J Psychiatry, 1970)
- **Application of family studies to investigate subtypes and overlap between syndromes**
- **Integration of clinical work and research**




Goals

- **Population-based data on comorbidity**
- **Familial patterns of co-aggregation to illustrate genetic epidemiologic approach**
- **Patterns of co-occurrence and order of onset among high risk youth**
- **Clinical and research implications**



Diseases with Greatest Global Burden : % Total Disability Adjusted Life Years



World Health Organization, 2002

Guze SB *Psychiatr Clin North Am.* 1990

Dec;13(4):651-9.

Secondary depression: observations in alcoholism, Briquet's syndrome, anxiety disorder, schizophrenia, and antisocial personality. A form of **comorbidity**?

Patterns of Comorbidity in the Community

12-month Prevalence of Substance Use Disorders in National Surveys of United States

Site	NCS-R	NES
Alc Abuse	3.8	4.7
Alc Dep	1.9	3.8
Drug Abuse	1.8	1.4
Drug Dep	0.7	0.6
Any Sub Abuse	9.4	9.4

NCS = National Comorbidity Survey Replication, Kessler et al; 1st 5000 cases

NES = National Epidemiologic Survey on Alcohol and Related Conditions, Grant et al

Comorbidity of Alcoholism and Mood Disorders in Community Studies

Author (yr)	Subtype	Alc Abuse	Alc Dep
Brady (92)	BPI	3.0	5.5
	BPII	3.9	3.1
	MDD	0.9	1.6
Kessler (90)	Mania	0.3	9.7
	MDE	1.0	2.7
Grant (04)	Mania	1.4	5.7
	Hypomania	1.7	5.2
	MDE	1.2	3.7

Comorbidity of Alcoholism and Anxiety Disorders in Community Studies

Author (yr)	Subtype	Alc Abuse	Alc Dep
Brady (92)	BPI	3.0	5.5
	BPII	3.9	3.1
	MDD	0.9	1.6
Kessler (90)	Mania	0.3	9.7
	MDE	1.0	2.7
Grant (04)	Mania	1.4	5.7
	Hypom	1.7	5.2
	MDE	1.2	3.7

International Consortium in Psychiatric Epidemiology:
**Comorbidity of Drug and Psychiatric
Disorders across Sites**

(Median Odds Ratio)

	Use	Drug Problems	Dependence
Mood	2.2	3.1	3.5
Anxiety	1.9	2.5	4.0
Behavior	3.3	5.7	5.6

Substance Abuse/Dependence Lifetime Comorbidity in Puerto Rican in San Juan and New Haven

	Affective Dx	Anxiety Dx
Alcohol		
New Haven	40 %	38 %
San Juan	42 %	42 %
Drug		
New Haven	48 %	49 %
San Juan	40 %	44 %


Guze SB. Semin Psychiatry. 1970 Nov;2(4):392-402.
The role of **follow-up studies**: their contribution to
diagnostic classification as applied to hysteria.

Prospective Studies

Zurich Cohort Study of Young Adults

- **Sample from general community of Zurich, Switzerland**
- **Methods:**
 - **Diagnostic interview for psychiatric and somatic disorders**
 - **Comprehensive assessment of risk factors and correlates**
 - **Evaluation of spectrum of expression of mental disorders in the community**

Zurich Cohort Study (1978-1999)



Statistical Methods

- Regression models were fit using generalized ordinal logistic models that yield an odds ratio for each cut-point in the ordinal outcome (Stata).
- The odds ratios represent the relative odds of being above the cut-point (*e.g., alcohol abuse or dependence vs. none or use; and alcohol dependence vs. none, use, or abuse*).
- These cut-points may be regarded as diagnostic thresholds.

Mood Disorders as Predictors of Alcohol Abuse/Dependence

	ALCOHOL	
	Abuse	Dependence
	<i>Odds Ratio (CI)</i>	
Major Dep	1.3 (0.6,2.9)	2.2 (0.7, 7.2)
Manic Sx	2.4 (1.2, 4.8)	4.4 (1.5,12.7)
Bipolar II	9.1 (2.7,31.2)	21.0 (6.6,67.5)

Smoking as Predictor of Alcohol and Cannabis Abuse or Dependence

	Alcohol Abuse	Dependence
	<i>Odds Ratio (CI)</i>	
	ALCOHOL	
	6.3 (2.9,13.6)	7.6 (2.7, 21.7)
	CANNABIS	
Smoking	8.6 (4.7,15.9)	40.4 (11.2,144.9)

Alcohol Use Disorders as Predictors of Cannabis Use/ Abuse/ Dependence

	CANNABIS	
	Use	Abuse/Dep
Alcohol	<i>Odds Ratio (CI)</i>	
Abuse	1.7 (1.2, 4.1)	2.2 (0.7,6.9)
Dependence	3.8 (0.8, 4.1)	4.1 (1.0,15.8)



Results of Community Surveys

- **Mood disorders are more strongly associated with alcohol dependence than with alcohol abuse**
- **The bipolar subtype has a significantly larger association with alcoholism than major depression**

What are the potential explanations for comorbidity?

- Comorbidity is a marker of severity of the index disorder
- Different syndromes are developmentally different manifestations of the same underlying pathogenesis
- Comorbid disorder is a consequence of another index disorder, or vice versa
- Comorbid disorders are alternate manifestations of the same underlying familial liability

Sources of Comorbidity



Guze SB, Cloninger CR, Martin RL, Clayton PJ.
Br J Psychiatry. 1986 Jul;149:17-23.
A follow-up and family study of Briquet's syndrome.

**Family Study of Explanations for
Comorbidity**

Mechanisms for Comorbidity: Family Studies

Common Etiology

Increased risk of “comorbid” disorder alone among relatives of probands with index disorder

Causal (Precursor of Consequence)

Increased risk of “comorbid” disorder only in combination with same index disorder



Investigators: Yale Family Study of Comorbidity of Substance Disorders & Psychopathology

K.R. Merikangas

K. Conway

B. Fenton

J. Merikangas

N. Risch

D. Stevens

P. Szatmari

S. Avenevoli

L. Dierker

C. Grillon

M. Preisig

B. Rounsaville

M. Stolar

H. Zhang

Yale Comorbidity Family Study : Sample


Probands (N=262)

Drug Alcohol Anxiety Control

N of Probands	87	89	76	61
Sex (Males %)	58	71	26	43
Age (Mean Yrs.)	36	40	40	41
N of Relatives	604	408	359	255

Relatives (N = 1626)

What disorder runs in this family?



Orange square: BIPOLAR

Red square: ALCOHOLISM

Green square: DRUG ABUSE


Yellow square: ANXIETY

Blue square: DEPRESSION

Substance Disorders in Relatives

Probands

Comorbidity in Relatives



Mechanisms for Comorbidity: Yale Family Study

Common Etiology

Panic, Major Depression & Substance Use Disorders have shared underlying etiologic factors.

Causal (Precursor of Consequence)

The familial associations between Social Phobia, Bipolar Disorder & Behavior Disorders with Substance Use Disorders are independent, despite the high magnitude of comorbidity between them.

Common familial liability for cannabis and alcohol dependence


Variable	OR	95% CI
Latent Familial	8.4	1.2- 58.5
PROBANDS		
Anxiety	0.7	0.4- 1.5
Depression	0.9	0.5- 1.7
Antisocial	0.4	0.1- 1.4
Alcohol Dep	2.6	1.2- 5.5
RELATIVES		
Anxiety	1.5	0.9- 2.9
Alcohol Dep	4.3	2.0- 9.2
Antisocial	3.7	1.3-10.6

Yale High Risk Study: Age and Sex of Sample at Wave I

Proband Group

	Substance +Anxiety N=39	Anxiety N=58	Substance N= 38	Normal N = 57
Age (% \geq12)	51.3	46.6	50.0	49.1
Sex (% male)	53.8	55.2	50.0	45.6
			Total N = 203	



Pathways to Substance Disorders through Psychopathology (Relative Risk)



Overlap in Use and Abuse/Dependence on Specific Substances among Offspring

Use

Abuse/Dependence



Prospective Association between Psychiatric and Substance Use Disorders by Ages 13-23 (N=203)

Pre-existing Dx	Use	Adjusted Risk Ratio	
		Substance Abuse	Dependence
Affective	0.6	1.7	3.2
Conduct	4.2	6.0	6.0
Oppositional	4.2	3.3	4.1
ADHD	0.9	2.0	3.6
Anxiety	0.9	1.9	5.5
ANY DX	1.3	3.0	5.7

Impact of Parental History and Premorbid Psychopathology on Substance Use and Disorders

	<i>Attributable Risk</i>	
	Use	Disorder
Family History	20%	12%
Premorbid Disorder	20%	18%
BOTH	32%	20%

Summary

- **Mood and anxiety disorders co-occur with alcohol and drug dependence in both clinical and community surveys**
- **Alcoholism is largely transmitted independently of most other comorbid conditions with the exception of panic and cannabis use disorder**
- **The onset of bipolar syndromes and social anxiety tend to precede that of alcohol problems.**



Implications

- **Etiology:** Identification of pathways and risk factors for the development of substance use disorders
- **Treatment:** Integration of psychiatric symptoms/syndromes in defining treatment strategies; Family-based approaches
- **Prevention:** Intervention in psychiatric syndromes may reduce incidence of substance use disorders; Offspring of substance abusers are important target for prevention

Clinic-based Prevention Opportunities

- **Offspring of parents in treatment for mental illness**
- **Incorporation of potential sequelae of primary disorders in treatment**
(e.g., Geller, et al, 1998)