

Energetic Contributions to Channel Gating of Residues in the Muscle Nicotinic Receptor $\beta 1$ Subunit

Gustav Akk, Megan Eaton, Ping Li, Steven Zheng, Joshua Lo, Joe Henry Steinbach*

Department of Anesthesiology and the Taylor Family Institute for Innovative Psychiatric Research, Washington University School of Medicine, Saint Louis, Missouri, United States of America

Abstract

In the pentameric ligand-gated ion channel family, transmitter binds in the extracellular domain and conformational changes result in channel opening in the transmembrane domain. In the muscle nicotinic receptor and other heteromeric members of the family one subunit does not contribute to the canonical agonist binding site for transmitter. A fundamental question is whether conformational changes occur in this subunit. We used records of single channel activity and rate-equilibrium free energy relationships to examine the $\beta 1$ (non-ACh-binding) subunit of the muscle nicotinic receptor. Mutations to residues in the extracellular domain have minimal effects on the gating equilibrium constant. Positions in the channel lining (M2 transmembrane) domain contribute strongly and relatively late during gating. Positions thought to be important in other subunits in coupling the transmitter-binding to the channel domains have minimal effects on gating. We conclude that the conformational changes involved in channel gating propagate from the binding-site to the channel in the ACh-binding subunits and subsequently spread to the non-binding subunit.

Citation: Akk G, Eaton M, Li P, Zheng S, Lo J, et al. (2013) Energetic Contributions to Channel Gating of Residues in the Muscle Nicotinic Receptor $\beta 1$ Subunit. *PLoS ONE* 8(10): e78539. doi:10.1371/journal.pone.0078539

Editor: Israel Silman, Weizmann Institute of Science, Israel

Received: May 14, 2013; **Accepted:** September 19, 2013; **Published:** October 23, 2013

Copyright: © 2013 Akk et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Research reported in this publication was supported by the National Institute of Neurological Disorders and Stroke of the National Institutes of Health under award numbers R21NS72770 and R01NS22356. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: jhs@morpheus.wustl.edu

Introduction

The pentameric ligand-gated ion channel (pLGIC) family includes the vertebrate nicotinic, GABA_A, serotonin-type A and glycine receptors, as well as prokaryotic and invertebrate receptors [1–3]. Each receptor comprises a pentamer of related subunits; the transmitter-binding sites are located at the interface between 2 subunits. The canonical acetylcholine (ACh) binding sites occur between a subunit that contributes the "principal" side and a second subunit that contributes the "complementary" side. In the case of heteromeric pLGIC receptors the result is that 2 pairs of the subunits (4 subunits) contribute to such a transmitter binding site whereas the 5th subunit does not. However, it is clear that the "non-binding subunit" can have profound effects on the activation of the receptor by transmitter [4,5]. Mutations to residues in the channel-lining region of the non-binding subunit affect gating with energetic contributions approximately equal to the effects of homologous mutations in transmitter-binding subunits [6,7]. To date, few studies have been made of the effects on receptor activation of mutations to residues outside the channel-lining region in the non-binding subunit. In the muscle nicotinic

receptor the canonical ACh-binding sites are located between the $\alpha 1$ subunit (principal face) and the δ and ϵ subunits (complementary face). We examined the effects of mutations in the $\beta 1$ subunit of the muscle nicotinic receptor (that does not bind acetylcholine) to determine the energetic consequences and, when possible, the inferred timing of energetic contributions to gating [8–10]. Our results indicate that the amino-terminal extracellular region of the $\beta 1$ subunit and the regions proposed to be involved in coupling between extracellular and transmembrane domains have few residues that make significant energetic contributions to the overall receptor gating equilibrium. In contrast, residues in the channel-lining region of the $\beta 1$ subunit do make energetic contributions and the timing indicates that the change occurs later in the gating process than for homologous residues in the $\alpha 1$ subunit. These findings indicate that the transduction of binding energy to gating flows from the binding regions of the transmitter-binding subunits to the channel and only subsequently is transmitted to the non-binding subunit.

Methods

Constructs and expression

Mouse muscle nicotinic subunits ($\alpha 1$, $\beta 1$, δ , ϵ) were expressed in HEK293 cells, using the pcDNA3 vector (Invitrogen, San Diego, CA). HEK293 cells were obtained from ATTC (Manassas VA). Mutations were introduced by QuikChange (Stratagene, San Diego, CA) mutagenesis, and the entire subunit was sequenced to verify that no additional mutations were introduced. Cells were transfected using the calcium-phosphate precipitation method [11]. The aligned sequences for the mouse $\alpha 1$, $\beta 1$, δ and ϵ subunits are shown in Figure S1, with positions studied indicated.

Physiological recordings

One to 3 days after transfection recordings were made in the cell-attached mode, and single channel events were recorded and analyzed [12]. Cells were bathed in recording bath solution (140 mM NaCl, 5 mM KCl, 1 mM $MgCl_2$, 2 mM $CaCl_2$, 10 mM glucose, and 10 mM HEPES, pH 7.4). The pipette solution contained (in mM): 142 KCl, 1.8 $CaCl_2$, 1.7 $MgCl_2$, 5.4 NaCl, and 10 HEPES, pH 7.4 with added choline. Recordings were made at a membrane potential of -50 mV (determined assuming that the reversal potential is at 0 mV) and room temperature (20-24 °C) using an Axopatch 200B amplifier (Molecular Devices, Union City, CA). Signals were low-pass filtered at 10 kHz, digitized with a Digidata 1320 series interface at 50 kHz using pClamp software (Molecular Devices) and analyzed using the QuB Suite (<http://www.qub.buffalo.edu>). Events were idealized using the SKM routine in QuB.

Choline was used as agonist in all cases. We used choline because the apparent channel opening rate constant is low. Recordings were made at a low concentration (50-100 μM) to determine the apparent channel closing rate constant (k_c). The apparent opening rate (k_o) was determined from recordings using a high concentration (20 mM) of choline. Because k_o is low, at low concentration apparent openings contain only a single opening, so the apparent closing rate (k_c) can be readily measured. At high concentration the closed periods reflecting channel opening are clearly resolvable in the experimental data. For wild-type receptors and 9 mutations k_o was estimated at both 10 mM and 20 mM choline, to verify that the k_o was estimated at a saturating concentration. In no case did the estimates differ significantly between the two concentrations (data not shown), so it is likely that our estimates do not significantly underestimate the actual k_o .

We adopted the methods introduced by the Auerbach Laboratory [9] to examine the consequences of mutations on channel gating. The di-liganded equilibrium gating constant (E_2) was computed as the ratio k_o / k_c for each mutation at a position. The "range energy" for that position was estimated from the range of values for E_2 (including wild-type) as $0.59 (\ln(E_{2,max}) - \ln(E_{2,min}))$ kcal/mol [8]. The range energy is defined only on the basis of the tested mutations, and so is a minimal estimate of the possible value. The obtained values for k_o , k_c and E_2 are shown in Table S3. The mutations made had a range of divergence from the original residue, as assessed

from the BLOSUM62 [13] value, but the estimated range energy did not depend on the most negative BLOSUM62 score at a location (Figure S3).

The parameter ϕ was calculated from the slope of a logarithmic plot of k_o on E_2 (a "rate-equilibrium plot"). The value of ϕ has been interpreted to reflect the position along the reaction coordinate from closed to open at which a residue has an effect on the overall energetic change in channel opening. A value near 1 indicates that at the transition state between closed and open the environment at that position is similar to the open state, and a value near 0 that the environment is closed-like [10,14,15]. Rate-equilibrium plots are shown for all the positions examined in Figure S2.

Molecular model

The homology model for the mouse muscle receptor was made by threading the mouse sequences ($\alpha 1$: NP_031415.2, $\beta 1$: NP_033731.3, δ : NP_067611.2, ϵ : NP_033733.1) on the crystal structure of the glutamate-activated chloride channel from *C. elegans* (GluCl; 3RHW; [16]) using the SWISS-MODEL web tool (<http://swissmodel.expasy.org/>). Structures were visualized and displays generated using Chimera 1.6.2 (<http://www.cgl.ucsf.edu/chimera>). The GluCl structure was chosen as the template based on reports [17–19] that it is a better representation of the muscle receptor structure than the *Torpedo marmorata* cryoelectron microscopic structure [20]. The positions of the transmembrane regions differ in the two structures (see Figure S1). We have chosen to call residues that are in the second transmembrane region (TM2) in both structures "TM2," while those that are placed in TM2, TM3 or the linker between them in the different structures as "TM2-link-TM3."

Results

The adult muscle nicotinic receptor (AChR) is a pentamer containing 2 copies of the $\alpha 1$ subunit and 1 each of the $\beta 1$, δ and ϵ subunits (Figure 1). The 2 canonical binding sites for acetylcholine (ACh) and other cholinergic agonists are located at the interfaces between the $\alpha 1$ and δ subunits and the $\alpha 1$ and ϵ subunits. The $\alpha 1$ subunits contribute the "principal" side of the binding site, with regions named the A, B and C loops. The δ or ϵ subunits contribute the complementary side, with regions named the D, E and F loops. The $\beta 1$ subunit does not contribute to a canonical binding site. After binding of transmitter, a series of conformational changes takes place between the transmitter-binding site (located in the extracellular portion of the receptor) and the channel (located in the membrane-spanning portion) that results in channel opening. Several regions of the $\alpha 1$ subunit have been proposed to be essential for establishing the connection between the extracellular domain (ECD) and the transmembrane domain (TMD), including the "PreM1" and "TM2-link-TM3" regions [21,22].

We wished to obtain a picture of the $\beta 1$ subunit in terms of its energetic contributions to overall channel gating, to compare to those of subunits that bind transmitter. The $\beta 1$ subunit has not been extensively studied: at least 113 positions have been




Figure 1. Schematic summaries of AChR structure. Panel A shows the locations of relevant regions in the primary sequences. There are 4 transmembrane regions (TM1-TM4). The channel is formed by the TM2 regions (highlighted in red) from the 5 subunits. The ACh-binding loops are in the N-terminal extracellular region (loops A-C form the "principal" side, D-F the "complementary" side). Panel B shows then membrane topology of a subunit (ACh-binding site: green circle, TM2 helix red cylinder). Panel C shows the arrangement of subunits in the pentamer (viewed from the extracellular side; green circles - ACh-binding interfaces). The channel is located in the center of the rosette of subunits. Panel D shows a homology model of the mouse $\beta 1$ subunit, threaded on the *C. elegans* GluCl E subunit [16]. The main chain is shown in blue, and the extracellular domain (ECD) and transmembrane domain (TMD) are indicated by brackets. The region coupling the ECD to the TMD is indicated by "coupling," and includes residues in the PreM1, loop 9, the TM2-link-TM3 region and the "principal pathway" (see text). In the ECD mutated residues in "loop 9" are shown in cyan, "Pre M1" in green, K46 in orange, V132 in yellow and other residues in light gray. In the TMD mutated residues in "TM2" are in blue and in the TM2-link-TM3 region in red. Many of the residues studied are located at or near the interface between the ECD and TMD regions of the subunit. Panel E shows the kinetic scheme used to interpret the data. A receptor with a closed channel (C) binds 2 molecules of agonist (A) then the channel opens (O) with an apparent opening rate k_o and apparent closing rate k_c .

doi: 10.1371/journal.pone.0078539.g001

studied in the $\alpha 1$ subunit, 52 in the ϵ , 29 in δ but only 15 in $\beta 1$ (see references to Table S1). We examined positions in the $\beta 1$ subunit that are homologous to positions that had already been studied in other subunits of the AChR. We made mutations to 27 positions, 13 in the ECD and 14 in transmembrane regions (summarized in Table 1; sequences shown in Figure S1). In general, we chose positions for which relatively large energetic changes had been reported for one or more subunits. In the ECD some positions corresponding to ACh-binding loops in other subunits were studied. We also examined positions proposed to be involved in coupling binding to gating in the $\alpha 1$ subunit. In the TMD we extended the analysis of residues in the transmembrane regions.

At each position we made 2 to 4 mutations, and determined the apparent rates for channel opening (k_o) and closing (k_c) (see Figure 1E for the kinetic model for activation; there is further consideration of kinetic models and rates in the Discussion). We used the partial agonist choline for all studies. Representative data are shown in Figure 2.

"Range energy"

We initially characterized the effects of mutations in terms of the ratio of channel opening to closing rate constants (E_2), to determine the consequences on the overall gating equilibrium. A parameter derived from values of E_2 is the "range energy" for a given position, defined as the observed maximal range of

Table 1. Mutations created and analyzed.

Residue	Structure	mutations	$\phi \pm SE$	Range energy (kcal/mol)
K46	PP	Q, M, R	0.29 \pm 0.58	0.43
Y55	D loop	F, Q, S	0.56 \pm 0.38	0.24
L93	A loop	A, T, Y	0.52 \pm 0.32	0.47
N96	Col	S, V*, W	0.07 \pm 0.24	0.52
V132		A, Q, L	-0.12 \pm 0.10	0.77
Y149	B loop & Col	A, D, Q, S	0.73 \pm 0.26	0.48
N182		D, E, S	0.15 \pm 0.29	0.18
G183		F, W, Y	0.69 \pm 0.15	0.55
Q184		T, W	-0.37 \pm 0.46	0.40
I218	Pre-M1	T, V	0.69 \pm 0.66	0.25
R219	Pre-M1	I, K, Q	0.27 \pm 0.07	0.76
R220	Pre-M1 & PP	I*, Q, K	6.88 \pm 7.88	0.02
K221	Pre-M1	I, Q, R	0.52 \pm 0.28	0.54
S257	TM2	C, G, I	-0.05 \pm 0.32	0.67
A260	TM2	C, G, V	0.03 \pm 0.25	0.80
T265	TM2	P, S, Y	0.42 \pm 0.31	0.93
V266	TM2 & Col	A, F, T	0.48 \pm 0.07	2.01
L270	TM2	A, T, Y	0.53 \pm 0.18	2.80
V275	TM2-link-TM3	A, L, M	0.24 \pm 0.27	1.19
P276	TM2-link-TM3 & Col	G, K, T	0.24 \pm 0.27	1.26
L280	TM2-link-TM3 & PP	A, T, Y	-0.30 \pm 0.27	0.50
A281	TM2-link-TM3	F, T, W	0.34 \pm 0.24	0.45
V282	TM2-link-TM3	A, Q, L	0.31 \pm 0.08	0.67
P283	TM2-link-TM3 & PP & Col	A, G, S	0.16 \pm 0.28	0.62
I284	TM2-link-TM3	F, L, T	0.52 \pm 0.12	0.78
I285	TM2-link-TM3	F, L, T	0.24 \pm 0.15	1.44
I286	TM2-link-TM3	F, L, T	0.48 \pm 0.04	1.03

The first column shows the residue studied. The second shows structural features, where PP indicates a residue in the Principal Pathway and "Col" a residue in the high energy column (see Text). The 3rd column shows the mutations made (an asterisk indicates that no functional channels were recorded from receptors containing that mutated subunit). The 4th column gives $\phi \pm$ the estimated SE returned by the fitting program. The final column gives range energy (kcal/mol). Positions at which the range energy is greater than 0.7 kcal/mol are shown in **bold**. doi: 10.1371/journal.pone.0078539.t001

energetic effects on E_2 . A large value for range energy indicates that the gating equilibrium is highly sensitive to the interactions between the amino-acid residue and its local environment. A small value indicates that that particular position is unlikely to undergo a large change in interactions during opening, at least for the tested substitutions. As can be seen in Figure 3, the range energy is low for positions in the ECD of the $\beta 1$ subunit, and higher in the TMD.

A low value of the range energy might be expected for residues homologous to positions in the transmitter-binding site (the A through F loops), because the $\beta 1$ subunit does not participate in the canonical ACh-binding site. However, the range energy is also low at other positions in the ECD, including regions proposed to be involved in coupling the extracellular binding domain to the transmembrane channel domain (Table 1, Figures 1 & 3). The locations we studied include some that show high range energies for the δ or ϵ ACh-

binding subunits, including the "loop 9" [23] and "pre-M1" [24] regions. In the $\beta 1$ subunit, these locations appear to make little contribution to the energetics of gating (Figure 3). Two proposals have been made for residues that are particularly important for coupling binding to gating in the $\alpha 1$ subunit. In one, a complex between the extracellular domain and the transmembrane domain containing $\alpha 1E45$, V46, R209, S269 and P272 was proposed (a "principal pathway" [25]. In the second, a "column" of residues with large range energies was proposed [8], including $\alpha 1E45$, A96, Y127, W149, P265, S268 and P272. Several of these positions were tested in the $\beta 1$ subunit, and none showed large range energies. We note that previous work has also shown that mutations to some of these positions in non- α subunits have only small effects on gating [25–27].

In the TMD the energetic contributions made by residues in $\beta 1$ are more comparable to those of other subunits. However, a comparison of our estimates of the range energies for positions in $\beta 1$ to estimates for homologous positions in other subunits indicates that 11 of 14 positions in $\beta 1$ have a lower value than for $\alpha 1$. This observation suggests that, overall the energetic contributions from the $\beta 1$ subunit are lower than those from homologous positions of the $\alpha 1$ subunit. We compared values by simply tabulating the numbers of positions that had lower as opposed to equal or greater values for range energy, and then compared positions in $\beta 1$ to homologous positions in $\alpha 1$. Based on the binomial distribution with equal probability of being greater or lesser, 11 or more positions would have lower energy with $P = 0.03$. Similar comparisons of the data from the $\beta 1$ subunit to data from the δ subunit show 4 of 7 with lower range energy ($P = 0.5$) and to the ϵ subunit 8 of 11 have lower range energy ($P = 0.11$).

Timing

When the range energy is large enough it is possible to accurately calculate the parameter ϕ , an estimate of the time at which the interaction between a residue and its local environment changes during the channel gating reaction. In practice, ϕ is calculated as the slope of a plot of the logarithm of k_0 to the logarithm of E_2 . A value near 1 indicates an early change event, while a value near 0 indicates a later event [10,14,15]. The reliability of the estimate for ϕ depends on the spread in values for E_2 . To allow for consistency with previously reported values, we adopted the criterion that the range energy needed to be greater than 0.7 kcal/mol (corresponding to a 3.3-fold ratio of values of E_2) for the estimate to be considered reliable.

Only two residues in the ECD ($\beta 1V132$ and $\beta 1R219$) have range energy greater than 0.7 kcal/mol, so little timing information is available in that region (Figure 4; Table 1). Both these residues have lower values for ϕ than agonist-binding subunits, suggesting late participation.

In contrast, 10 of 14 residues examined in the TMD had range energy > 0.7 kcal/mol (Figure 3). The comparison of ϕ values between subunits shows that positions in $\beta 1$ TMD have lower values (i.e. later timing) in 7 of 8 comparisons to $\alpha 1$ ($P = 0.04$). However, in comparison to δ no positions in $\beta 1$ have lower ϕ (0 of 4, $P = 0.06$), while in comparison to ϵ 3 of 6 do (P




Figure 2. Rate equilibrium free energy relationships. Panel A shows sample traces of data for wild-type receptors and receptors containing $\beta 1$ (L270A). On the left data at a low concentration (100 μM) of choline are shown, with histograms of the open durations shown below. The fits to the histograms were used to estimate the channel closing rate. The openings are clearly prolonged by the mutation, and the estimates for k_o are 2128 s^{-1} for wild type and 360 s^{-1} for the mutant. The number of events in the histogram are 506 for $\beta 1$ (L270A) and 695 for wild type. On the right data at a high concentration (20 mM) are shown, with histograms of the closed durations below. The estimates for k_o are 44 s^{-1} for wild type and 153 s^{-1} for the mutant (estimated from the major, slower component). The number of events in the histogram are 2390 for $\beta 1$ (L270A) and 2259 for wild type. Note that open channel block reduces the channel current amplitude at this high concentration. Panel B shows logarithmic plots of k_o on E_2 for mutations at 3 positions in the $\beta 1$ subunit. The lines show the linear regression of $\log(k_o)$ on $\log(E_2)$. The range energy is calculated from the range of E_2 values and ϕ is the slope of the linear regression (given as regression value \pm SE of fit value). $\beta 1$ (Y149) illustrates a position at which the range energy is small, $\beta 1$ (V266) a position with a linear relationship, and $\beta 1$ (V275) a position at which the slope of the line is poorly defined. The hollow symbol shows data for wild-type receptors. Data points are identified with the residue at the position, and are mean \pm SE. Rate-equilibrium plots for all positions are shown in Figure S2.

doi: 10.1371/journal.pone.0078539.g002

= 0.5). These comparisons suggest that on the whole the contributions of $\beta 1$ occur after those of $\alpha 1$, about the same time as those of ϵ and perhaps before those of δ .

It is possible that conformational changes at the extracellular end of the channel (that is, the TM2-link-TM3 region) might participate in coupling the ECD and TMD, and so might experience earlier movements during gating. In the $\alpha 1$ subunit




Figure 3. Range energy at homologous positions in the subunits. Panel A shows data for positions in the ECD, while panel B shows data for the TMD. The sites are identified by the residue and position in the $\beta 1$ subunit, and some structural features are shown below (see Text). Residues proposed to be part of the "principal pathway" for coupling binding to gating in the $\alpha 1$ subunit [25] are indicated by stars above the data, and residues lying in a "high range energy column" in the $\alpha 1$ subunit [8] are indicated by plus signs. Results for the $\beta 1$ subunit are shown by blue triangles (results from the present study shown as filled triangles, while results obtained by others as open triangles). Values for $\alpha 1$ are shown with filled red circles, for δ by yellow inverted triangles and ϵ by bold crosses. Values are given on a per subunit basis. The dashed line at 0.7 kcal/mol shows the discriminator chosen for the minimal range energy for considering an estimate of ϕ to be reliable. The values for the present data are shown in Table 1, and all values are shown in Tables S1 and S2.

doi: 10.1371/journal.pone.0078539.g003




Figure 4. Summary of results for timing. Values are shown for ϕ , in a similar format to that of Figure 3. The small symbols shown below the solid horizontal line indicate positions at which the range energy was less than 0.7 kcal/mol and for which ϕ could not be reliably calculated. The values for the present data are shown in Table 1, and all values are shown in Tables S1 and S2.

doi: 10.1371/journal.pone.0078539.g004

residues at this region show larger values for ϕ than in $\beta 1$ (4 of 4 comparisons), indicating an earlier movement. Only 2 comparisons could be made to the ϵ subunit, but in both cases ϕ for residues in ϵ was greater. It is interesting to note that the ϕ values for the $\beta 1$ subunit are largest and most similar to those of $\alpha 1$ in the middle of TM2 (Figure 4), which might suggest that the conformational change of gating is communicated from the $\alpha 1$ to the $\beta 1$ subunit at this level.

As shown in Table 1 several of the residues in the TMD of the $\beta 1$ subunit have a large uncertainty in our estimate of ϕ . It has been reported that some positions in other subunits may show non-linear relationships between $\log(k_o)$ and $\log(E_2)$ (for example 28). The reason for this lack of linearity is not known, but the interpretation of ϕ in terms of timing may be incorrect for these positions.




Figure 5. Parameters in subunits divided by structural regions. The data for positions examined in the present study are summarized in terms of the structural region in which the residue is located. The regions shown are ECD (from $\beta 1$ (V46) to Q184), Pre-M1 ($\beta 1$ (I218) to K221), TM2 ($\beta 1$ (S257) to L270) and TM2-link-TM3 ($\beta 1$ (V275) to I286). The upper panel shows data for range energy, with the median values for the individual subunits indicated by the horizontal bars, while the lower panel shows similar data for ϕ .

doi: 10.1371/journal.pone.0078539.g005

Comparisons across subunits

The results obtained are summarized in Figures 3 & 4 for different structural domains in Figure 5.

Previous studies have noted that mutations to homologous positions in different subunits may have very different energetic consequences for channel gating [25]. The present data confirm this observation, as shown in the scatter plots of range energies in Figure 6. The data are presented in terms of the region (ECD in panel A, TMD in panel B) and by whether the residue in $\alpha 1$ is part of the "principal pathway" or high energy column (panel C, "Pathways"). The regression coefficient does not differ significantly from 0 for any comparison (for ECD slope = -0.1, $P = 0.39$; for TMD slope = 0.55, $P = 0.10$; for pathway slope = 0.0, $P = 0.98$). Qualitatively speaking, it seems that positions that have a large range energy in the $\alpha 1$ subunit ECD tend to have low range energy in other subunits, and *vice versa*. In the TMD there is a large scatter of energies but more trend to similar energies in different subunits. This might reflect unique roles of the ECD in the different subunits, but more common roles for homologous residues in the TMD. Overall the proposed activation pathways appear to be unique to the $\alpha 1$ subunit, although both δ and ϵ show high energetic contributions at the positions homologous to $\alpha 1$ R209 ("Pre-

M1") and $\alpha 1$ V255 (TM2). In contrast, the range energies are lower for $\beta 1$ at these positions.

Discussion

The results indicate that residues in the extracellular domain of the $\beta 1$ subunit show relatively weak changes in interactions with the local environment during channel gating. The results are not consistent with the idea of circumferential transmission of energy in the extracellular domain; that is, little indication of conformational changes spreading from the transmitter-binding subunits to the $\beta 1$ subunit through interactions of residues in the ECD. Similarly, residues proposed to be involved in coupling between the ECD and TMD in other subunits do not appear to undergo significant changes in interactions during gating. In contrast, residues in the TMD of the $\beta 1$ subunit show larger energetic contributions to the gating equilibrium than residues in the ECD. Overall, the results indicate that the $\beta 1$ subunit makes significant energetic contributions to the gating process only when gating has progressed to the actual channel region. This contribution could reflect conformational changes in the $\beta 1$ subunit, altered interactions with adjacent subunits, or both.

We interpreted our observations in terms of a simple kinetic scheme ([29], Figures 1E, 7A) that has been used extensively in studies of the muscle nicotinic receptor. However, a number of recent studies have demonstrated that this model is not adequate, and that there are transitions between closed states that can be kinetically apparent [8,30,31]. A strongly supported version is the "Flip" model, in which there is a closed-closed transition (Flip) immediately before channel opening (Figure 7B) [30]. The data indicate that the actual channel opening rate constant is quite similar between choline and ACh, but that the transition from A_2C to A_2F (where F denotes a receptor with a closed channel that is in the "flipped" state) is slower for choline [32]. In the elegant analysis performed by Lape et al. [32] the component in the closed time distribution that we analyzed to obtain an estimate of k_o actually reflects dwells of the receptor in all the states preceding A_2F , and so does not give an estimate of the true channel opening rate constant. This result raises the question of whether the overall interpretation of the observations is appropriate. The goal of our work was to compare the energetic contributions of the $\beta 1$ subunit to the overall process of receptor activation - the equilibrium between closed channel states and the open state. As such, the kinetic details determining this equilibrium are of lesser importance than the global equilibrium. However, if mutations had different effects when receptor function was probed using different agonists then the comparison might well be between apples and oranges. This seems unlikely to be generally true. In several studies multiple mutations were made a single location, and the effects determined using either choline or ACh as agonist. The data fall along the same line on a logarithmic rate-equilibrium plot, suggesting that the data with the two agonists reflect the same basic energetic consequences of the mutations (for studies with many mutations at a given position see 26,33). However, we note that in two instances mutations do appear to have different




Figure 6. Range energies at homologous positions. Scatter plots for range energy are shown for positions in the $\beta 1$, δ or ϵ subunits (ordinate) plotted against the value for the homologous position in $\alpha 1$ (abscissa). All data shown in Figure 3 are plotted. The regions shown are ECD ($\beta 1$ (V46) to K221), TMD ($\beta 1$ (S257) to I286) and Pathways (see Text, $\beta 1$ (E45), K46, N96, S127, Y149, R220, V266, P276, S279, L280 and P283). Two positions are identified by their locations in the $\alpha 1$ subunit, that have large range energies in the complementary face subunits (δ and ϵ) but lower values in the $\beta 1$ subunit. Positions at which all 4 subunits have the same amino acid are shown by triangles, while positions at which one or more differ are shown by circles. Data for the $\beta 1$ subunit are shown as filled red symbols. The solid lines show the line of equality, the dashed lines show the regression lines (no slope differed significantly from 0).

doi: 10.1371/journal.pone.0078539.g006

consequences for receptors activated by choline than by other agonists: $\beta 1$ (R220Q) (Table S2) and $\alpha 1$ (E45R) [34]. We used choline for all constructs tested, and the majority of studies of homologous positions in other subunits used choline for activation of some or all constructs (about 60%, see Table S1). In sum, we feel that our studies of the effects of mutations in the $\beta 1$ subunit on both the range energy and the timing of energetic contributions can be compared to effects of mutations to homologous residues in other subunits.

There were only 2 residues in the ECD of the $\beta 1$ subunit for which the value of ϕ could be reliably calculated, but these values indicate late events. In the TMD, for residues at homologous positions $\alpha 1$ appears to have the earliest effects, δ the latest, with $\beta 1$ and ϵ intermediate. These observations suggest that the conformational changes involved in gating occur initially in the $\alpha 1$ subunit, then propagate to the $\beta 1$ and ϵ subunits and finally to the δ subunit. This difference in timing supports the idea that there is a conformational change in the $\beta 1$ TMD during channel opening, rather than solely conformational changes in the adjacent subunits that alter the environment around static residues in $\beta 1$.

Earlier studies have noted that mutations to homologous residues in the ECD do not have identical effects when made in different subunits [25,35,36]. This may indicate unique roles or dissimilar movements in different subunits. Our observations confirm this observation, and indicate that the ECD in the $\beta 1$ subunit makes the lowest overall contribution to the gating energy, at least for the locations tested.

Although this picture shows the $\beta 1$ subunit in the muscle nicotinic receptor as a relatively passive element in the overall




Figure 7. A kinetic model with additional closed states. Panel A shows the kinetic scheme used in interpreting our data. Panel B shows an extended scheme incorporating a closed-closed conformational change ("flip") preceding the channel opening step. In this scheme, A receptor with a closed channel (C) binds 2 molecules of agonist (A) then the receptor enters the flipped state (F) while keeping a closed channel. The channel opens (O) from the flipped state. The scheme is discussed further in the Text.

doi: 10.1371/journal.pone.0078539.g007

function of the receptor, it is clear that the nature of the subunit that does not contribute to a canonical transmitter-binding site (the "non-binding" subunit) can have a significant effect on the pharmacology and biophysics of a heteropentameric receptor. It can contribute to a binding site for allosteric drugs - most notably for the benzodiazepines acting on the GABA_A receptor [37], but also in neuronal nicotinic receptors [38,39] - and in the

nicotinic $\alpha 4\beta 2$ receptor there is evidence for a non-canonical ACh-binding site at an $\alpha 4$ - $\alpha 4$ interface [39,40]. In addition, the non-binding subunit can alter the gating by transmitters and other agonists [4,5,41]. Similarly, the non-binding subunit can affect the potency of competitive inhibitors [4,42], in which case it is less likely that a change in conformational equilibrium occurs. It has been suggested that the nature of the non-binding subunit may affect the conformation of the pentameric receptor extracellular domain and thereby change the fine structure of the transmitter-binding site [42]. These results suggest a picture in which the non-binding subunit affects the structure of the ECD of the resting receptor. If our observations on the $\beta 1$ subunit can be generalized to these related receptors, they would generally support this picture with the extension that there is relatively little change in the structure of the ECD for the non-binding subunit during channel gating. However, this then begs the question of the precise intersubunit interactions by which the change in the agonist-binding site is mediated.

A previous study examined coupling between the ECD and TMD in a homopentameric receptor, utilizing mutated chimeric subunits formed from the ECD of the nicotinic $\alpha 7$ subunit and the TMD and intracellular domain of the 5HT 3_A subunit [43]. The "coupling region" studied was a 7 amino acid sequence at the start (extracellular end) of the first transmembrane domain; when these residues were from the $\alpha 7$ subunit channel opening did not occur, while when they were from the 5HT 3_A subunit it did [44]. The data indicated that the coupling region of a subunit mutated so that it did not contribute to an agonist-binding site did contribute to the overall stability of the open-channel state. This observation is not consistent with our finding that there is no indication that coupling between the ECD and TMD in the $\beta 1$ subunit affects channel gating of the muscle nicotinic receptor. There are, however, some significant differences between the experiments. First, the studies conducted by Anderson et al [43] were conducted with a basically homopentameric receptor, rather than a heteropentameric receptor. It is possible that the symmetry of the subunits alters some of the structural changes in gating. The second difference is that 6 of the 7 residues swapped in the TM1 region were altered simultaneously. It is possible that this larger perturbation affected gating more generally, rather than coupling specifically. In our experiments we did not mutate any of the residues altered by Anderson et al.

There are two major caveats to the interpretation of our observations. The first is the possibility that we examined "incorrect" residues in the ECD. Perhaps there is a unique, as yet unidentified, constellation of positions in the $\beta 1$ subunit which does make a significant contribution to channel gating. The second caveat is the possibility that a change in the $\beta 1$ subunit takes place as a rigid body motion that may remain undetected by our studies, as they likely focus largely on side-chain interactions. A recent study by Unwin and Fujiyoshi [45], examined cryo-electron microscopic images of AChR from *Torpedo marmorata* that were obtained from specimens frozen within milliseconds after exposure to agonist and were interpreted to reflect the open channel state. The images were compared to previously obtained images, interpreted to reflect

the closed channel state [20]. The major change seen in a non- α subunit was a tilt in the $\beta 1$ ECD (an outward movement of about 0.1 nm). It was proposed that the ECD rocks on the TMD at the interface between the two domains and that this motion is central to the mechanism by which binding is communicated to the gate, indicating a major role of the $\beta 1$ subunit in channel gating. We mutated many residues at this interface in the $\beta 1$ subunit (Figure 1), and our data do not indicate a major energetic contribution. Furthermore, the data about timing of energetic contributions in the TMD are not consistent with the idea that conformational changes occur first in the $\beta 1$ subunit. However, one concern about our data is that it might not have detected changes in position of the backbone polypeptide chain, for example if a tilt "pulled" on the top of TM1 with little change in side-chain interactions.

Overall, our results indicate that the $\beta 1$ subunit is passive in the transfer of the energy of binding from the ECD to the channel gate. However, our data indicate that the non-binding subunit participates in the conformational changes in the channel during opening, and results of others suggest that it may influence the conformation of the transmitter-binding site of the resting receptor.

Supporting Information

Table S1. Data for homologous positions in all subunits.
(DOCX)

Table S2. Comparative values for positions in $\beta 1$.
(DOCX)

Table S3. Measured rates.
(DOCX)

Figure S1. Aligned primary sequences of the subunits.
(DOCX)

Figure S2. REFER plots for the 27 locations.
(DOCX)

Figure S3. Range energy versus BLOSUM62.
(DOCX)

File S1. The file "Mouse beta1 only on GluCl E Fig 1D.py" is a Python file for Chimera 1.6.2 (<http://www.cgl.ucsf.edu/chimera>) that contains the homology model shown in Figure 1.
(PY)

Acknowledgements

We thank Chris Lingle for comments on the manuscript. JHS is the Russell and Mary Professor of anesthesiology.

Author Contributions

Conceived and designed the experiments: GA JHS. Performed the experiments: GA ME PL SZ JL. Analyzed the data: GA JHS. Wrote the manuscript: GA JHS.

References

- Ortells MO, Lunt GG (1995) Evolutionary history of the ligand-gated ion-channel superfamily of receptors. *Trends Neurosci* 18: 121-127. doi:10.1016/0166-2236(95)93887-4. PubMed: 7754520.
- Tasneem A, Iyer LM, Jakobsson E, Aravind L (2005) Identification of the prokaryotic ligand-gated ion channels and their implications for the mechanisms and origins of animal Cys-loop ion channels. *Genome Biol* 6: R4. doi:10.1186/gb-2005-6-4-p4. PubMed: 15642096.
- Dent JA (2006) Evidence for a diverse Cys-loop ligand-gated ion channel superfamily in early bilateria. *J Mol Evol* 62: 523-535. doi:10.1007/s00239-005-0018-2. PubMed: 16586016.
- Moroni M, Zwart R, Sher E, Cassels BK, Bermudez I (2006) $\alpha 4\beta 2$ nicotinic receptors with high and low acetylcholine sensitivity: pharmacology, stoichiometry, and sensitivity to long-term exposure to nicotine. *Mol Pharmacol* 70: 755-768. doi:10.1124/mol.106.023044. PubMed: 16720757.
- Akk G, Bracamontes J, Steinbach JH (2004) Activation of GABA(A) receptors containing the $\alpha 4$ subunit by GABA and pentobarbital. *J Physiol* 556: 387-399. doi:10.1113/jphysiol.2003.058230. PubMed: 14966300.
- Chang Y, Weiss DS (1999) Allosteric activation mechanism of the $\alpha 1\beta 2\gamma 2$ gamma-aminobutyric acid type A receptor revealed by mutation of the conserved M2 leucine. *Biophys J* 77: 2542-2551. doi:10.1016/S0006-3495(99)77089-X. PubMed: 10545355.
- Labarca C, Nowak MW, Zhang H, Tang L, Deshpande P et al. (1995) Channel gating governed symmetrically by conserved leucine residues in the M2 domain of nicotinic receptors. *Nature* 376: 514-516. doi:10.1038/376514a0. PubMed: 7637783.
- Auerbach A (2010) The gating isomerization of neuromuscular acetylcholine receptors. *J Physiol* 588: 573-586. doi:10.1113/jphysiol.2009.182774. PubMed: 19933754.
- Grosman C, Zhou M, Auerbach A (2000) Mapping the conformational wave of acetylcholine receptor channel gating. *Nature* 403: 773-776. doi:10.1038/35001586. PubMed: 10693806.
- Auerbach A (2007) How to turn the reaction coordinate into time. *J Gen Physiol* 130: 543-546. doi:10.1085/jgp.200709898. PubMed: 18040056.
- Akk G (2002) Contributions of the non- α subunit residues (loop D) to agonist binding and channel gating in the muscle nicotinic acetylcholine receptor. *J Physiol* 544: 695-705. doi:10.1113/jphysiol.2002.029413. PubMed: 12411516.
- Akk G, Steinbach JH (2005) Galantamine activates muscle-type nicotinic acetylcholine receptors without binding to the acetylcholine-binding site. *J Neurosci* 25: 1992-2001. doi:10.1523/JNEUROSCI.4985-04.2005. PubMed: 15728839.
- Henikoff S, Henikoff JG (1992) Amino acid substitution matrices from protein blocks. *Proc Natl Acad Sci U S A* 89: 10915-10919. doi:10.1073/pnas.89.22.10915. PubMed: 1438297.
- Zhou Y, Pearson JE, Auerbach A (2005) Phi-value analysis of a linear, sequential reaction mechanism: theory and application to ion channel gating. *Biophys J* 89: 3680-3685. doi:10.1529/biophysj.105.067215. PubMed: 16183877.
- Colquhoun D (2005) From shut to open: what can we learn from linear free energy relationships? *Biophys J* 89: 3673-3675. doi:10.1529/biophysj.105.071563. PubMed: 16319077.
- Hibbs RE, Gouaux E (2011) Principles of activation and permeation in an anion-selective Cys-loop receptor. *Nature* 474: 54-60. doi:10.1038/nature10139. PubMed: 21572436.
- Cymes GD, Ni Y, Grosman C (2005) Probing ion-channel pores one proton at a time. *Nature* 438: 975-980. doi:10.1038/nature04293. PubMed: 16355215.
- Bondarenko V, Mowrey D, Tillman T, Cui T, Liu LT et al. (2012) NMR structures of the transmembrane domains of the $\alpha 4\beta 2$ nAChR. *Biochim Biophys Acta* 1818: 1261-1268. doi:10.1016/j.bbame.2012.02.008. PubMed: 22361591.
- Mnatsakanyan N, Jansen M (2013) Experimental determination of the vertical alignment between the second and third transmembrane segments of muscle nicotinic acetylcholine receptors. *J Neurochem* (2013) doi:10.1111/jnc.12260. PubMed: 23565737.
- Unwin N (2005) Refined structure of the nicotinic acetylcholine receptor at 4 Å resolution. *J Mol Biol* 346: 967-989. doi:10.1016/j.jmb.2004.12.031. PubMed: 15701510.
- Sine SM (2012) End-plate acetylcholine receptor: structure, mechanism, pharmacology, and disease. *Physiol Rev* 92: 1189-1234. doi:10.1152/physrev.00015.2011. PubMed: 22811427.
- Corringer PJ, Poitevin F, Prevost MS, Sauguet L, Delarue M et al. (2012) Structure and pharmacology of pentameric receptor channels: from bacteria to brain. *Structure* 20: 941-956. doi:10.1016/j.str.2012.05.003. PubMed: 22681900.
- Jha A, Gupta S, Zucker SN, Auerbach A (2012) The energetic consequences of loop 9 gating motions in acetylcholine receptor-channels. *J Physiol* 590: 119-129. PubMed: 22025664.
- Bruhova I, Auerbach A (2010) Subunit symmetry at the extracellular domain-transmembrane domain interface in acetylcholine receptor channel gating. *J Biol Chem* 285: 38898-38904. doi:10.1074/jbc.M110.169110. PubMed: 20864527.
- Lee WY, Sine SM (2005) Principal pathway coupling agonist binding to channel gating in nicotinic receptors. *Nature* 438: 243-247. doi:10.1038/nature04156. PubMed: 16281039.
- Purohit P, Auerbach A (2007) Acetylcholine receptor gating at extracellular transmembrane domain interface: the "pre-M1" linker. *J Gen Physiol* 130: 559-568. doi:10.1085/jgp.200709857. PubMed: 18040058.
- Jha A, Purohit P, Auerbach A (2009) Energy and structure of the M2 helix in acetylcholine receptor-channel gating. *Biophys J* 96: 4075-4084. doi:10.1016/j.bpj.2009.02.030. PubMed: 19450479.
- Mitra A, Cymes GD, Auerbach A (2005) Dynamics of the acetylcholine receptor pore at the gating transition state. *Proc Natl Acad Sci U S A* 102: 15069-15074. doi:10.1073/pnas.0505090102. PubMed: 16217024.
- del Castillo J, Katz B (1957) Interaction at end-plate receptors between different choline derivatives. *Proc R Soc Lond B Biol Sci* 146: 369-381. doi:10.1098/rspb.1957.0018. PubMed: 13431862.
- Lape R, Colquhoun D, Sivilotti LG (2008) On the nature of partial agonism in the nicotinic receptor superfamily. *Nature* 454: 722-727. PubMed: 18633353.
- Mukhtasimova N, Lee WY, Wang HL, Sine SM (2009) Detection and trapping of intermediate states priming nicotinic receptor channel opening. *Nature* 459: 451-454. doi:10.1038/nature07923. PubMed: 19339970.
- Lape R, Krashia P, Colquhoun D, Sivilotti LG (2009) Agonist and blocking actions of choline and tetramethylammonium on human muscle acetylcholine receptors. *J Physiol* 587: 5045-5072. doi:10.1113/jphysiol.2009.176305. PubMed: 19752108.
- Purohit P, Auerbach A (2007) Acetylcholine receptor gating: movement in the alpha-subunit extracellular domain. *J Gen Physiol* 130: 569-579. doi:10.1085/jgp.200709858. PubMed: 18040059.
- Mukhtasimova N, Sine SM (2013) Nicotinic receptor transduction zone: invariant arginine couples to multiple electron-rich residues. *Biophys J* 104: 355-367. doi:10.1016/j.bpj.2012.11.1968. PubMed: 23442857.
- Lee WY, Free CR, Sine SM (2009) Binding to gating transduction in nicotinic receptors: Cys-loop energetically couples to pre-M1 and M2-M3 regions. *J Neurosci* 29: 3189-3199. doi:10.1523/JNEUROSCI.6185-08.2009. PubMed: 19279256.
- Shen XM, Ohno K, Tsujino A, Brengman JM, Gingold M et al. (2003) Mutation causing severe myasthenia reveals functional asymmetry of AChR signature cysteine loops in agonist binding and gating. *J Clin Invest* 111: 497-505. doi:10.1172/JCI16997. PubMed: 12588888.
- Sigel E, Buhr A (1997) The benzodiazepine binding site of GABA receptors. *Trends Pharmacol Sci* 18: 425-429. doi:10.1016/S0165-6147(97)90675-1. PubMed: 9426470.
- Seo S, Henry JT, Lewis AH, Wang N, Levandoski MM (2009) The positive allosteric modulator morantel binds at noncanonical subunit interfaces of neuronal nicotinic acetylcholine receptors. *J Neurosci* 29: 8734-8742. doi:10.1523/JNEUROSCI.1859-09.2009. PubMed: 19587280.

39. Mazzaferro S, Benallegue N, Carbone A, Gasparri F, Vijayan R et al. (2011) Additional acetylcholine (ACh) binding site at alpha4/alpha4 interface of (alpha4beta2)2alpha4 nicotinic receptor influences agonist sensitivity. *J Biol Chem* 286: 31043-31054. doi:10.1074/jbc.M111.262014. PubMed: 21757735.
40. Harpsøe K, Ahring PK, Christensen JK, Jensen ML, Peters D et al. (2011) Unraveling the high- and low-sensitivity agonist responses of nicotinic acetylcholine receptors. *J Neurosci* 31: 10759-10766. doi: 10.1523/JNEUROSCI.1509-11.2011. PubMed: 21795528.
41. Fisher JL, Macdonald RL (1997) Single channel properties of recombinant GABAA receptors containing gamma 2 or delta subtypes expressed with alpha 1 and beta 3 subtypes in mouse L929 cells. *J Physiol Lond* 505: 283-297. doi:10.1111/j.1469-7793.1997.283bb.x. PubMed: 9423172.
42. Hogg RC, Hopping G, Alewood PF, Adams DJ, Bertrand D (2003) Alpha-conotoxins Pn1A and [A10L]Pn1A stabilize different states of the alpha7-L247T nicotinic acetylcholine receptor. *J Biol Chem* 278: 26908-26914. doi:10.1074/jbc.M212628200. PubMed: 12746432.
43. Andersen N, Corradi J, Bartos M, Sine SM, Bouzat C (2011) Functional relationships between agonist binding sites and coupling regions of homomeric Cys-loop receptors. *J Neurosci* 31: 3662-3669. doi:10.1523/JNEUROSCI.5940-10.2011. PubMed: 21389221.
44. Bouzat C, Bartos M, Corradi J, Sine SM (2008) The interface between extracellular and transmembrane domains of homomeric Cys-loop receptors governs open-channel lifetime and rate of desensitization. *J Neurosci* 28: 7808-7819. doi:10.1523/JNEUROSCI.0448-08.2008. PubMed: 18667613.
45. Unwin N, Fujiyoshi Y (2012) Gating movement of acetylcholine receptor caught by plunge-freezing. *J Mol Biol* 422: 617-634. doi:10.1016/j.jmb.2012.07.010. PubMed: 22841691.