

Washington University School of Medicine Digital Commons@Becker

Open Access Publications

2012

The new nucleoporin: Regulator of transcriptional repression and beyond

Nayan J. Sarma

Washington University School of Medicine in St. Louis

Kristine Willis

Washington University School of Medicine in St. Louis

Follow this and additional works at: http://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Sarma, Nayan J. and Willis, Kristine, "The new nucleoporin: Regulator of transcriptional repression and beyond." *Nucleus*.3,6. 1-8. (2012).

http://digitalcommons.wustl.edu/open_access_pubs/2512

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

The new nucleoporin

Regulator of transcriptional repression and beyond

Nayan J. Sarma¹ and Kristine Willis^{2,*}

¹Department of Surgery; Washington University School of Medicine; St. Louis, MO USA; ²Department of Biology; Georgetown University; Washington, DC USA

Transcriptional regulation is a complex process that requires the integrated action of many multi-protein complexes. The way in which a living cell coordinates the action of these complexes in time and space is still poorly understood. Recent work has shown that nuclear pores, well known for their role in 3' processing and export of transcripts, also participate in the control of transcriptional initiation. We have recently begun to explore how nuclear pores interface with the well-described machinery that regulates initiation. This work led to the discovery that specific nucleoporins are required for binding of the repressor protein Mig1 to its site in target promoters. Nuclear pores are therefore involved in repressing, as well as activating, transcription. Here we discuss in detail the main models explaining our result and consider what each implies about the roles that nuclear pores play in the regulation of gene expression.

Introduction

Cells respond to environmental cues to grow and differentiate by turning discrete sets of genes on and off. Although no one model appears to apply to all genes,^{1,2} work done over the past 40 y has given us a general understanding of how a eukaryotic cell accomplishes this on/off switch.²⁻⁹ The process is complex, requiring the interaction of nucleosomes, chromatin remodelers, chromatin modifiers, sequence-specific repressors and activators, co-activators, basal transcription factors, positive and negative regulators

of elongation, termination factors and of course the DNA itself. Control of transcriptional initiation, though historically appreciated as important, is only one way in which information in the genome is managed. Once the signal to express a certain gene has been received and the necessary factors are all associated with the DNA, the polymerase must escape from the promoter and transition to productive elongation, travel down the body of the gene and stop correctly once the termination signal has been reached. At the same time, the mRNA must be capped, spliced and polyadenylated, then exported to the cytoplasm for translation when finished. Each step in this process is highly interdependent and subject to multiple levels of regulation.^{2,10-12} Developing a precise understanding of the mechanisms that enforce these multiple levels of regulation and the ways in which they interact with each other is now the central challenge in the field of gene expression.

Recently, it has been shown that several different subunits of the nuclear pore, a 60 MDa complex best known for its role in nucleocytoplasmic transport, also play a role in upstream transcriptional regulation. Nuclear pore proteins (nucleoporins) have been found in physical association with both active and repressed regions of the budding yeast, fruit fly, rat and human genomes.¹³⁻²¹ Importantly, a subset of nucleoporins interact with DNA in the nuclear interior, independently of the presence of RNA polymerase II. This finding strongly suggests that proteins of the nuclear pore play a role in transcription

Keywords: transcriptional regulation, RNA polymerase II, nuclear pore complex, NPC, Nup120, Nup133, acute myeloid leukemia, glucose, glucose-regulated gene expression, glucose metabolism, AMPK

Abbreviations: NPC, nuclear pore complex; TBP, TATA binding protein; GFP, green fluorescent protein; TSS, transcription start site; HDAC, histone deacetylase

Submitted: 07/30/12

Revised: 09/26/12

Accepted: 10/02/12

<http://dx.doi.org/10.4161/nucleus.22427>

*Correspondence to: Kristine Willis; Email: kaw85@georgetown.edu

Table 1. Nuclear pore proteins in physical interaction with genomic loci

	nucleoporin	yeast homolog	references
<i>Saccharomyces cerevisiae</i>			
active	Nup60		16
	Nup116		16
	Nic96		16
	Mlp1		16
	Mlp2		16
active and repressed	Nup2		16, 17, 41
	Nup145C		13, 16
	Nup53		13
	Nup133		13
	Pom152		13
<i>Drosophila melanogaster</i>			
active	mTOR	Mlp1	19, 22
	Nup153	Nup60	19
	Nup50	none	20
	Nup62	Nsp1	20
	Nup98	Nup145N	14, 20
	Sec13	Sec13	14
repressed	Nup88	Nup82	14
<i>Rattus norvegicus</i>			
active and repressed	Nup155	Nup170	15
<i>Homo sapiens</i>			
active and repressed	Nup93	Nic96	18

that is independent of mRNA transport.^{19,20} Consistent with such a role, experiments in both flies and rats have shown that knock-down or overexpression of nucleoporins reciprocally alters mRNA levels of the genes with which they interact.^{14,15,19,20,22} However, almost nothing is known about the nature of the nuclear pore-genome interaction, or the mechanics of how nucleoporins contribute to transcriptional regulation.

Recently, though, we have identified a specific mechanism through which nuclear pore proteins can influence transcription. Deletion of either of two particular subunits of the nuclear pore, *NUPI20* or *NUPI33*, results in a loss of repression of *SUC2*, one of the canonical model genes in the *Saccharomyces cerevisiae* system on which our current understanding of transcriptional regulation has been built. This loss of repression is not attributable to defects in the

nucleocytoplasmic shuttling of Mig1, the DNA-binding factor primarily responsible for inhibition of *SUC2* transcription. Instead, nuclear Mig1 must physically associate with intact nuclear pore complexes (NPCs) in order to bind DNA and repress transcription. In other words, the ability of the repressor to find its site in the DNA is dependent on its interaction with NPCs; both interaction with NPCs and DNA binding are lost if either *NUPI20* or *NUPI33* is deleted. Viewed in the context of earlier work, this result suggests that nuclear pore proteins function as transcriptional regulators by exerting a global influence on chromatin biology, nuclear organization, or both.

Different Biological Functions for Nuclear Pores

Nuclear pores were first identified as electron-dense structures visible in

microscopic analyses of the nuclear membrane performed during the 1950s.²³⁻²⁸ Over the next 25 y, their universality in eukaryotic cells, as well as a consensus regarding their overall shape and size, became well established. By 1975 two distinct but mutually compatible functions of nuclear pores had been proposed: nucleocytoplasmic transport and chromatin organization.²⁹ NPCs are now known to be the sole conduit through which macromolecules are exchanged between the nucleus and cytoplasm (reviewed in refs. 30 and 31). The second postulated function, which was predicated on numerous observations of the physical proximity of chromatin to nuclear pores,³²⁻³⁹ has received less attention. Recent work, though, has reintroduced the idea that this physical juxtaposition may indicate a functional link.

ChIP,^{13-15,40} ChIP-chip¹⁶⁻¹⁹ and chromatin cleavage^{20,41} analyses have shown that subunits of the nuclear pore, or nucleoporins, interact with both active and inactive regions of the genomes of budding yeast, fruit flies, rats and humans (Table 1). Several lines of evidence suggest that these interactions serve a purpose other than facilitating the export of newly synthesized transcripts. First and simplest, some nucleoporins preferentially associate with regions of the genome that are not transcribed.^{14,18} The next argument for a transport-independent function stems from the observation that some nucleoporins localize to both the nuclear periphery and the nuclear interior,⁴²⁻⁴⁶ also called the nucleoplasm; nucleoplasmic pools of these nucleoporins have been shown to interact with loci that are also found in the nuclear interior and to influence steady-state levels of mRNA transcribed from these genes.^{14,19,20} Finally, work done in fruit flies has directly demonstrated that the association between nucleoporins and active genes is stable to treatment with RNase A and thus not tethered by RNA molecules in the process of being exported.¹⁴

Since RNA-only models are ruled out, the way in which nucleoporins interact with the genome is unclear. With one possible exception,⁴⁷ nucleoporins do not contain recognizable domains for binding DNA or chromatin directly. It therefore seems likely that they recognize

specific loci through another protein or proteins, but the identity of these is currently unknown. However, examining the function of nucleoporin-DNA interactions may provide a hint about mechanism of interaction. In budding yeast, nucleoporins interact specifically with intergenic regions, places where the transcriptional machinery and the DNA first come into contact; this contact occurs independently of active transcription.^{13,41} Two *Drosophila* nucleoporins, Nup98 and Sec13, have been shown to make contact with their target loci prior to RNA polymerase.¹⁴ Consistent with these data, the yeast nucleoporin Nup2 has been shown to contact the promoter of the *GALI,10* locus prior to TBP.⁴¹ Sequence specific transcription factors, chromatin remodelers and chromatin modifying complexes are therefore the most logical candidates for proteins that serve as intermediates between nucleoporins and DNA, since only they arrive at the promoter prior to TBP.

Regulation of *SUC2* Transcription is Linked to Nuclear Pores

Our understanding of glucose-regulated gene expression in the model eukaryote *Saccharomyces cerevisiae* has been established largely through study of the *SUC2* promoter. Work completed over the past 25 y has thoroughly identified the set of proteins that control its activity (Fig. 1), although the way in which they work together is still not fully understood. Two factors central to this regulatory system are the Snf1 kinase and the DNA-binding repressor protein Mig1. Together they control transcription of not only *SUC2*, but approximately 300 other genes.⁴⁸

Recent work has shown that the promoters of three Mig1 targets—*SUC2*, *HXX1* and *GALI, 10*^{13,41}—are also bound by nuclear pore proteins; furthermore, these genes very likely spend some time interacting with intact NPCs, since GFP tagging shows they periodically visit the nuclear periphery when repressed and localize there for prolonged intervals when active.^{13,49,50} We therefore wondered if the Snf1 and Mig1 proteins themselves might also interact with NPCs. We found that Snf1 co-fractionates with intact NPCs only in the absence of glucose,




Figure 1. Regulation of glucose-repressed gene expression in *Saccharomyces cerevisiae*. Regulation of *SUC2* expression is well characterized, and the proteins that carry out this regulation have been studied extensively. The *SUC2* system therefore provides a good model for understanding how other genes controlled by the same factors are regulated. **(A)** A kinase-transcription factor pair work together to control expression of *SUC2*. Left panel. In the presence of glucose (repressed), the C2H2 zinc finger protein Mig1 binds DNA and inhibits transcription of *SUC2*, plus approximately 300 other genes. Complete inhibition also requires the hexokinase Hxk2. Snf1 kinase, a structural and functional homolog of mammalian AMPK, is found in the cytoplasm. Right panel. When glucose is depleted or withdrawn (derepressed), Snf1 is phosphorylated by upstream kinases, enters the nucleus and phosphorylates Mig1, which is then exported together with Hxk2. For detailed review, see reference 48. **(B)** Chromatin structure of the *SUC2* promoter. Left panel. In the presence of glucose (repressed), four nucleosomes (numbered 1 through 4) cover the *SUC2* promoter; a gradient of histone H3 acetylation decreases from 5' to 3'.⁵⁵ Two GC-rich sequences are required for repression of transcription (red t-bars) from the TSS (dashed black arrow).⁸⁸ These sites are interchangeably bound in vivo by Mig1 and its homolog Mig2; function of the former, but not the latter, is regulated by the Snf1 kinase.⁴⁸ The average location of the first Mig1/2 site (left-most red rectangle) is between nucleosomes 1 and 2. The second Mig1/2 site (right-most red rectangle) is close to the end of the DNA covered by nucleosome 2;⁸⁹⁻⁹¹ nucleosome and repressor may compete for occupancy of this site. The TATA box (green rectangle) is covered by nucleosome 4.⁸⁹⁻⁹³ Right panel. Full induction of transcription requires the Swi/Snf chromatin remodeling complex.^{57,90,91,94-98} Swi/Snf associates more stably with promoter nucleosomes that have been acetylated by SAGA and NuA4 complexes;⁵⁷ in the absence of glucose, acetylation of H3 and H4 tails increases for all nucleosomes. The DNA formerly covered by two nucleosomes (nucleosomes 2 and 3) is more frequently covered by only one. Nucleosome 4, covering the TATA box, is hyperacetylated and becomes unstable.⁵⁵ Initiation occurs at the TSS (black arrow). Additionally required for full induction of *SUC2* transcription are the nucleosome remodeling protein Spt6^{99,100} and the transcriptional activator Gcr1 (blue oval, Gcr1 binding site).^{90,101-104} Nuclear pore proteins also interact with the *SUC2* promoter;¹³ it is important to learn how they collaborate with the well-characterized transcription factors described here.

when *SUC2*, *HXX1* and *GALI,10* are actively transcribed and stably associated with nuclear pores. In contrast, Mig1 co-fractionates with NPCs only in the presence of glucose, when these three target genes are repressed and appear to interact with nuclear pores transiently.¹³ However, Snf1 and Mig1 undergo

nucleocytoplasmic transport in response to changes in the availability of glucose and must associate with the nuclear pores for this purpose. Both proteins cross the pore when glucose appears as well as when it disappears; therefore it is not straightforward to explain the differential physical association we observe by invoking a

transport-only function, but we still considered it a possibility. Accordingly, we looked for a way to test whether the association of these regulators with nuclear pores was only a reflection of their transport. It has long been known that deletion of *HXX2*, which encodes the predominant form of hexokinase in budding yeast, causes defects in Mig1-dependent glucose repression.⁵¹⁻⁵³ We found that in an *hxx2Δ* mutant, Mig1 is properly localized to the nucleus in response to the appearance of glucose; however, despite being localized to the nucleus in this mutant, Mig1 no longer co-fractionates with nuclear pores.¹³ In other words, our analysis of an *hxx2Δ* mutant shows that repressor function of Mig1 requires its physical association with NPCs in a way that is independent of the protein's nucleocytoplasmic transport.

Specific subunits of the NPC Mediate Binding of the Mig1 Repressor to Target Promoters

We next asked whether mutation of nuclear pores themselves might cause defects in the transcriptional regulation of *SUC2*. Of particular interest were cells deleted for either *NUP120* or *NUP133*, in which we observed a loss of repression of *SUC2* approximately equal to that seen on deleting *MIG1* itself.⁵⁴ Since a reduction in export of *SUC2* mRNA would not be expected to lead to an increase in levels of the translated protein product, we reasoned that these nucleoporins likely impact regulation of *SUC2* expression at the level of transcription. Visualization of Mig1-GFP shows that import of Mig1 into the nucleus does not require Nup120 or Nup133.⁵⁴ Given our above-described data on Hxk2, though, we wondered if Nup120 and Nup133 might be required for the interaction of Mig1 with NPCs. We found that this is true; although Mig1 is correctly transported into and out of the nucleus in the absence of these nucleoporins, it is no longer able to co-fractionate with nuclear pores. In the absence of Nup120 or Nup133, nuclear Mig1 is also unable to bind its site in DNA.⁵⁴ This striking result provides a clear explanation of how these nucleoporins influence *SUC2* repression.

How Nuclear Pores Might Influence Gene Expression: Models and Possible Mechanisms

Why does Mig1 require Nup120 and Nup133 to bind DNA? We have shown that Mig1 co-purifies with intact NPCs in glucose-grown cells, in which the repressor binds its target loci.^{13,54} We have also shown that both Nup133 and Pom152 interact with the *SUC2* promoter in glucose-grown cells.⁵⁴ Nup133 is a structural component of the NPC, and Pom152 is an integral nuclear membrane protein. We therefore interpret this result to mean that in the presence of glucose, the repressed *SUC2* promoter makes transient contact with intact NPCs embedded in the nuclear membrane. This interpretation is consistent with the observation that the *SUC2* locus can be seen to periodically “visit” the nuclear periphery in glucose grown cells.¹³ An obvious possibility, therefore, is that during these visits, the NPC facilitates binding of the repressor protein to the DNA by simultaneously interacting with both. Bringing together a gene and its transcriptional regulator in a particular location, at the nuclear periphery, might not only facilitate DNA binding, but also reduce the time it takes that transcription factor to search the genome for its site.

Drawing the simplest possible analogy to cooperative DNA binding by two transcription factors (Fig. 2A), we can infer two significant corollaries to this model. First, *SUC2* is required to contact the NPC independently of Mig1. As noted above, this second, Mig1-independent contact may be direct, but more likely occurs through some other, currently unidentified factor. Second, deletion of any one of the components shown in Figure 2A has the potential to reduce the affinity with which the others interact. Consistent with these requirements, we observe that in glucose-grown *mig1Δ* cells, Nup133 still binds the *SUC2* promoter, although the strength of the interaction appears to be reduced (Fig. 2B).

Although interesting, this finding is far from conclusive, and further work must be done to test the accuracy of the idea. If it is correct, classical biochemical theory predicts that adding either individual

nucleoporin(s) or intact nuclear pores to in vitro DNA-binding reactions should increase the affinity of Mig1 for its site. However, if in vivo nuclear context is important—perhaps for reducing search time by compressing three dimensions into two—then in vitro analyses such as this may not be informative. In which case, quantitative measurement of in vivo DNA binding at different nuclear locations, either by fluorescent resonance energy transfer (FRET) or bimolecular fluorescence complementation (BiFC), may provide the best test of this model. It will also be critical to definitively address the question of where in the yeast nucleus promoter-nucleoporin interactions truly occur. Circumstantial evidence—the periodic localization of the DNA to the nuclear periphery, together with the relative immobility of the nucleoporins we observe to interact with it—strongly suggests that intact, membrane-embedded NPCs influence Mig1 DNA binding. However, at this time, we cannot rule out a role for soluble nucleoplasmic pools of nucleoporins. Determining whether an interaction occurs in one or the other of two specific locations—the nuclear periphery or the nucleoplasm—may seem straightforward at first. In reality, the need for precise localization in a small nucleus, together with the likelihood that no more than a very few molecules of nucleoporin are localized to the nuclear interior, means that a combination of super-resolution microscopy and single molecule detection techniques will be necessary to answer this question in a satisfactory way.

A second model, not mutually exclusive with the one described above, is that NPCs collaborate with chromatin modifying complexes and chromatin remodelers to fine-tune the post-translational modification and/or position of nucleosomes over many Mig1 target promoters (Fig. 2C). In repressed wild type cells, nucleosomes 1 and 2 of the *SUC2* promoter are at least 2- to 5-fold more acetylated than nucleosome 4;⁵⁵ 1 and 2 flank the first Mig1 binding site, 2 covers the second Mig1 binding site and 4 covers the TATA box (Fig. 1B). The effect of histone acetylation on Mig1 binding has not been studied directly, but modeling of

nucleosome positioning data suggests that these are less stably associated with the DNA in the absence of glucose.⁵⁶ A change in the targeting or activity of histone acetyltransferases and/or in acetylation-dependent Swi/Snf remodeling, might occur in *nup120Δ* and *nup133Δ* mutants, thus altering the position or occupancy of promoter nucleosomes so that binding sites for Mig1 become obscured. In this model, Mig1 cannot repress transcription because it is unable to outcompete these nucleosomes.

Some data exist to support a model in which nucleoporins influence transcriptional regulation by co-operating with chromatin modifying complexes, particularly those that control histone acetylation. This is significant, since acetylation state is believed to be an important determinant of the affinity between nucleosomes and DNA. In *Drosophila*, the histone acetyltransferase MOF has been found in complex with both Nup153 and the NPC associated protein mTOR.²² Nup153 has also been found to preferentially interact with genomic loci where H4K16Ac-containing nucleosomes are enriched.¹⁹ In *Saccharomyces*, the MOF homolog Esa1, catalytic subunit of NuA4, constitutively associates with the *SUC2* promoter.⁵⁷ Deletion of *ESAI* is synthetically defective with deletion of *NUP120*, suggesting the products of these genes share some common function;⁵⁸ this interesting possibility has not yet been explored. Yeast Gcn5, the catalytic subunit of SAGA, has been shown to associate directly with Mlp1 and 2, homologs of *Drosophila* mTOR.⁵⁹ Most recently, the *Rattus norvegicus* nucleoporin Nup155 has been shown to interact with the histone deacetylase HDAC4 and to regulate the association of specific genomic loci with that nucleoporin.¹⁵

An intriguing but currently even less-well elaborated connection may exist between nuclear pores and chromatin remodeling complexes. The RSC complex is essential both for the localization of NPCs to the nuclear envelope and for normal nuclear morphology, leading to the suggestion that RSC might be required for establishing contacts between chromatin and the NPC.⁶⁰ *SUC2* is not known to be a target of RSC, but the promoters of other Mig1-regulated genes,




Figure 2. How nucleoporins might influence access to a specific site in DNA. **(A)** Model for NPC facilitated Mig1 binding to DNA. (a) In the presence of glucose, Mig1 associates with NPCs; the *SUC2* locus contacts NPCs transiently. (b) Increased local concentration of both Mig1 and its consensus site facilitates DNA binding by the repressor. (c) The repressor-bound promoter can dissociate from the NPC. **(B)** Interaction of Nup133 with the *SUC2* promoter is reduced in *mig1Δ* cells. TAP-tagged Nup133 was immunoprecipitated from *mig1Δ* or wild type cells grown in media containing glucose as the carbon source, then fixed with formaldehyde as described previously.¹³ Crosslinks were reversed and PCR was used to amplify the promoters of *SUC2* and *ACT1* (negative control) from recovered material. Amplified target is expressed as a ratio of *SUC2*/*ACT1*, normalized to the amount of product amplified whole cell extracts (input). Adding increasing amounts of input DNA shows that amplification of the product is linear. A representative experiment is shown; similar results were obtained for Nup145C (not shown). **(C)** A model for collaborative nucleosome positioning. Top panel. Mig1 associates with nuclear pores as in (A) above. *SUC2* is not associated with nuclear pores, and the nucleosomes across its promoter are poorly positioned (three yellow circles, one distinct and two indistinct, represent the average position of one nucleosome, over time and across a population). Bottom panel. One or more subunits of the nuclear pore associate with the *SUC2* locus, serving as a marker or barrier against which chromatin remodelers can position a single nucleosome, thus seeding an ordered array. Alternatively, nuclear pore proteins may direct the activity of chromatin modifiers such as SAGA and NuA4, which in turn direct the activity of remodelers such as Swi/Snf. Models shown in A and C are not mutually exclusive; for example, nuclear pores contribute to nucleosome positioning at step (b) of model A.

including *HXX1* and *FBP1*, are bound at a low level by RSC in glucose-grown cells.⁶¹ Again, further work must be done to determine whether changes in the position or acetylation state of promoter nucleosomes are responsible for the failure of Mig1 to bind DNA in *nup120Δ* and *nup133Δ* cells.

Relevance to Leukemia: Nucleoporins as Transcription Factors in Human Cells

Genes encoding components of the NPC are the target of chromosomal translocations in several different types of cancer;⁶²⁻⁶⁶ possibly the best studied of these

translocations are found in hematological malignancies, including Myelodysplastic Syndrome (MDS), Acute Myeloid Leukemia (AML), Chronic Myeloid Leukemia (CML) and T-cell acute lymphocytic leukemia (T-ALL).⁶⁷⁻⁶⁹ Different mechanisms have been proposed to explain how nucleoporin fusions contribute to the development of disease. One hypothesis is that interaction of nucleoporin chimeras with endogenous nucleoporins or RNA export factors may contribute to their leukemogenic properties.⁷⁰⁻⁷² Another is that nucleoporins are involved in spindle formation and this function is disrupted in oncogenic fusions, leading to aneuploidy.⁶⁹ However, in many cases the translocation partner of the nucleoporin is a transcription factor. NUP98-HOXA9, probably the best-characterized of these leukemogenic nucleoporin fusion proteins,⁷³⁻⁷⁷ has long been known to have the ability to bind to DNA and interact with components of the basal transcription machinery, leading to aberrant gene expression.^{14,20,78-82} The recent work described above strongly suggests that nucleoporins play a role in normal transcriptional regulation; it seems likely, then, that conscription of this function is a primary mechanism of oncogenesis.⁶⁹ This conclusion is supported by studies of the fusion proteins themselves.

Of particular relevance to our work on Mig1 is a recent study linking NUP98 with the transcription factor AES. AES is a homolog of the Mig1-interacting protein Tup1; both AES and Tup1 are Groucho family proteins that act as co-repressors of transcription. Importantly, AES interacts with both wild type NUP98 and NUP98-HOXA9, consistent with the idea of a conserved, nucleoporin-dependent mechanism of transcriptional regulation that is hijacked in leukemogenesis. Together, AES and NUP98-HOXA9 cooperate to regulate transcription, cell transformation and the impairment of hematopoietic differentiation.^{79,80}

Functional and physical links have also been described between NUP98 fusions and chromatin modifiers. The FG repeats in the NUP98 portion of a NUP98-HOXA9 fusion physically interact with CREB-binding protein

(CBP)/p300; transcriptional activation of a reporter gene by NUP98-HOXA9 is dependent on the histone acetyltransferase activity of CBP/p300.⁸¹ Chromosomal translocations also fuse NUP98 to the lysine methyltransferases NSD1 and MLL, to the lysine demethylase JARID1A and to the poorly-characterized protein PHF23, which recognizes methylated H3K4.⁶⁹ NUP98-NSD1 promotes the methylation of histone H3-Lys36 (H3K36) at loci encoding several proto-oncogenes, including HoxA7, HoxA9, HoxA10, Meis1; enforced activation of these transcription factors then blocks differentiation.⁸³ NUP98-JARID1A and NUP98-PHF23 fusion proteins inhibit demethylation of histone H3-Lys 4 (H3K4me3) at loci encoding lineage-specific transcription factors, again blocking differentiation.^{84,85} It is not clear how the deregulation of lysine methylation by these NUP98 chimeras relates to nucleoporin-mediated regulation of gene expression described above. However, in *Drosophila* cells, the methylation state of H3K36 is linked to levels of H4K16 acetylation,^{86,87} and nucleoporins preferentially bind loci enriched for H4K16Ac.^{19,20}

Conclusions and Future Directions

These observations bring up an important question: to what extent is transcriptional regulation by nucleoporins conserved in healthy and cancerous cells? The work summarized above suggests that it is, to at least some extent. Where this is true, with what partner proteins do they work, and how do these differ at different stages of transcript production? How can their contribution be integrated into existing models of gene regulation? Addressing these questions should help us to understand how multiple steps of transcript synthesis are coupled, bringing us closer to a full understanding of how gene expression is regulated in a nuclear context.

Acknowledgments

We thank George Santangelo for important theoretical contributions; Dan Larson and Vytas Bankaitis for comments on the manuscript and Patrick Varga-Weisz for

comments on the manuscript, as well as for a generous amount of time spent in helpful discussions of the text. This work was supported by National Institutes of Health grant GM083309 to K.A.W.

References

1. Cosma MP. Ordered recruitment: gene-specific mechanism of transcription activation. *Mol Cell* 2002; 10:227-36; PMID:12191469; [http://dx.doi.org/10.1016/S1097-2765\(02\)00604-4](http://dx.doi.org/10.1016/S1097-2765(02)00604-4).
2. Orphanides G, Reinberg D. A unified theory of gene expression. *Cell* 2002; 108:439-51; PMID:11909516; [http://dx.doi.org/10.1016/S0092-8674\(02\)00655-4](http://dx.doi.org/10.1016/S0092-8674(02)00655-4).
3. Krishnamurthy S, Hampsey M. Eukaryotic transcription initiation. *Curr Biol* 2009; 19:R153-6; PMID:19243687; <http://dx.doi.org/10.1016/j.cub.2008.11.052>.
4. Weake VM, Workman JL. Inducible gene expression: diverse regulatory mechanisms. *Nat Rev Genet* 2010; 11:426-37; PMID:20421872; <http://dx.doi.org/10.1038/nrg2781>.
5. Hahn S, Young ET. Transcriptional regulation in *Saccharomyces cerevisiae*: transcription factor regulation and function, mechanisms of initiation, and roles of activators and coactivators. *Genetics* 2011; 189:705-36; PMID:22084422; <http://dx.doi.org/10.1534/genetics.111.127019>.
6. Malik S, Roeder RG. The metazoan Mediator coactivator complex as an integrative hub for transcriptional regulation. *Nat Rev Genet* 2010; 11:761-72; PMID:20940737; <http://dx.doi.org/10.1038/nrg2901>.
7. Saunders A, Core LJ, Lis JT. Breaking barriers to transcription elongation. *Nat Rev Mol Cell Biol* 2006; 7:557-67; PMID:16936696; <http://dx.doi.org/10.1038/nrm1981>.
8. Bryant GO, Ptashne M. Independent recruitment in vivo by Gal4 of two complexes required for transcription. *Mol Cell* 2003; 11:1301-9; PMID:12769853; [http://dx.doi.org/10.1016/S1097-2765\(03\)00144-8](http://dx.doi.org/10.1016/S1097-2765(03)00144-8).
9. Carey MF, Peterson CL, Smale ST. *Transcriptional Regulation in Eukaryotes: Concepts, Strategies, & Techniques*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2009.
10. Maniatis T, Reed R. An extensive network of coupling among gene expression machines. *Nature* 2002; 416:499-506; PMID:11932736; <http://dx.doi.org/10.1038/416499a>.
11. Pandit S, Wang D, Fu XD. Functional integration of transcriptional and RNA processing machineries. *Curr Opin Cell Biol* 2008; 20:260-5; PMID:18436438; <http://dx.doi.org/10.1016/j.ceb.2008.03.001>.
12. Proudfoot NJ, Furger A, Dye MJ. Integrating mRNA processing with transcription. *Cell* 2002; 108:501-12; PMID:11909521; [http://dx.doi.org/10.1016/S0092-8674\(02\)00617-7](http://dx.doi.org/10.1016/S0092-8674(02)00617-7).
13. Sarma NJ, Haley TM, Barbara KE, Buford TD, Willis KA, Santangelo GM. Glucose-responsive regulators of gene expression in *Saccharomyces cerevisiae* function at the nuclear periphery via a reverse recruitment mechanism. *Genetics* 2007; 175:1127-35; PMID:17237508; <http://dx.doi.org/10.1534/genetics.106.068932>.
14. Capelson M, Liang Y, Schulte R, Mair W, Wagner U, Hetzer MW. Chromatin-bound nuclear pore components regulate gene expression in higher eukaryotes. *Cell* 2010; 140:372-83; PMID:20144761; <http://dx.doi.org/10.1016/j.cell.2009.12.054>.
15. Kehat I, Accornero F, Aronow BJ, Molkenin JD. Modulation of chromatin position and gene expression by HDAC4 interaction with nucleoporins. *J Cell Biol* 2011; 193:21-9; PMID:21464227; <http://dx.doi.org/10.1083/jcb.201101046>.

16. Casolari JM, Brown CR, Komili S, West J, Hieronymus H, Silver PA. Genome-wide localization of the nuclear transport machinery couples transcriptional status and nuclear organization. *Cell* 2004; 117:427-39; PMID:15137937; [http://dx.doi.org/10.1016/S0092-8674\(04\)00448-9](http://dx.doi.org/10.1016/S0092-8674(04)00448-9).
17. Dilworth DJ, Tackett AJ, Rogers RS, Yi EC, Christmas RH, Smith JJ, et al. The mobile nucleoporin Nup2p and chromatin-bound Prp20p function in endogenous NPC-mediated transcriptional control. *J Cell Biol* 2005; 171:955-65; PMID:16365162; <http://dx.doi.org/10.1083/jcb.200509061>.
18. Brown CR, Kennedy CJ, Delmar VA, Forbes DJ, Silver PA. Global histone acetylation induces functional genomic reorganization at mammalian nuclear pore complexes. *Genes Dev* 2008; 22:627-39; PMID:18316479; <http://dx.doi.org/10.1101/gad.1632708>.
19. Vaquerizas JM, Suyama R, Kind J, Miura K, Luscombe NM, Akhtar A. Nuclear pore proteins nup153 and megator define transcriptionally active regions in the *Drosophila* genome. *PLoS Genet* 2010; 6:e1000846; PMID:20174442; <http://dx.doi.org/10.1371/journal.pgen.1000846>.
20. Kalverda B, Pickersgill H, Shloma VV, Fornerod M. Nucleoporins directly stimulate expression of developmental and cell-cycle genes inside the nucleoplasm. *Cell* 2010; 140:360-71; PMID:20144760; <http://dx.doi.org/10.1016/j.cell.2010.01.011>.
21. Finlan LE, Sproul D, Thomson I, Boyle S, Kerr E, Perry P, et al. Recruitment to the nuclear periphery can alter expression of genes in human cells. *PLoS Genet* 2008; 4:e1000039; PMID:18369458; <http://dx.doi.org/10.1371/journal.pgen.1000039>.
22. Mendjan S, Taipale M, Kind J, Holz H, Gebhardt P, Schelder M, et al. Nuclear pore components are involved in the transcriptional regulation of dosage compensation in *Drosophila*. *Mol Cell* 2006; 21:811-23; PMID:16543150; <http://dx.doi.org/10.1016/j.molcel.2006.02.007>.
23. Watson ML. Further observations on the nuclear envelope of the animal cell. *J Biophys Biochem Cytol* 1959; 6:147-56; PMID:13843146; <http://dx.doi.org/10.1083/jcb.6.2.147>.
24. Callan HG, Randall JT, Tomlin SG. An electron microscope study of the nuclear membrane. *Nature* 1949; 163:280; PMID:18112254; <http://dx.doi.org/10.1038/163280a0>.
25. Callan HG, Tomlin SG. Experimental studies on amphibian oocyte nuclei. I. Investigation of the structure of the nuclear membrane by means of the electron microscope. *Proc R Soc Lond B Biol Sci* 1950; 137:367-78; PMID:14786306; <http://dx.doi.org/10.1098/rspb.1950.0047>.
26. Gall JG. Observations on the nuclear membrane with the electron microscope. *Exp Cell Res* 1954; 7:197-200; PMID:13200519; [http://dx.doi.org/10.1016/0014-4827\(54\)90054-3](http://dx.doi.org/10.1016/0014-4827(54)90054-3).
27. Anderson E, Beams HW. Evidence from electron micrographs for the passage of material through pores of the nuclear membrane. *J Biophys Biochem Cytol* 1956; 2(Suppl):439-44; PMID:13357583; <http://dx.doi.org/10.1083/jcb.2.4.439>.
28. Beams HW, Tahmisiyan TN, Anderson E, Devine RL. The structure of nuclear membrane in larval gonads of *Heliothis obsoleta*. *Proc Soc Exp Biol Med* 1956; 91:473-5; PMID:13322969.
29. Aaronson RP, Blobel G. Isolation of nuclear pore complexes in association with a lamina. *Proc Natl Acad Sci U S A* 1975; 72:1007-11; PMID:1055359; <http://dx.doi.org/10.1073/pnas.72.3.1007>.
30. Grünwald D, Singer RH, Rout M. Nuclear export dynamics of RNA-protein complexes. *Nature* 2011; 475:333-41; PMID:21776079; <http://dx.doi.org/10.1038/nature10318>.
31. Hoelz A, Debler EW, Blobel G. The structure of the nuclear pore complex. *Annu Rev Biochem* 2011; 80:613-43; PMID:21495847; <http://dx.doi.org/10.1146/annurev-biochem-060109-151030>.
32. DuPraw E. *Cell and Molecular Biology*. New York: Academic Press, 1958:424.
33. Dupraw EJ. *Cell and Molecular Biology*. NY: Academic Press, 1954.
34. Comings DE, Okada TA. Association of nuclear membrane fragments with metaphase and anaphase chromosomes as observed by whole mount electron microscopy. *Exp Cell Res* 1970; 63:62-8; PMID:5531486; [http://dx.doi.org/10.1016/0014-4827\(70\)90331-9](http://dx.doi.org/10.1016/0014-4827(70)90331-9).
35. Kirschner RH, Rusli M, Martin TE. Characterization of the nuclear envelope, pore complexes, and dense lamina of mouse liver nuclei by high resolution scanning electron microscopy. *J Cell Biol* 1977; 72:118-32; PMID:556616; <http://dx.doi.org/10.1083/jcb.72.1.118>.
36. Chan JK, Park PC, De Boni U. Association of DNase sensitive chromatin domains with the nuclear periphery in 3T3 cells in vitro. *Biochem Cell Biol* 2000; 78:67-78; PMID:10874467; <http://dx.doi.org/10.1139/o99-074>.
37. Hutchison N, Weintraub H. Localization of DNAase I-sensitive sequences to specific regions of interphase nuclei. *Cell* 1985; 43:471-82; PMID:4075401; [http://dx.doi.org/10.1016/0092-8674\(85\)90177-1](http://dx.doi.org/10.1016/0092-8674(85)90177-1).
38. Markovics J, Glass L, Maul GG. Pore patterns on nuclear membranes. *Exp Cell Res* 1974; 85:443-51; PMID:4827878; [http://dx.doi.org/10.1016/0014-4827\(74\)90148-7](http://dx.doi.org/10.1016/0014-4827(74)90148-7).
39. Arlucca J, Andrade R, Alonso R, Aréchaga J. The nuclear basket of the nuclear pore complex is part of a higher-order filamentous network that is related to chromatin. *J Struct Biol* 1998; 124:51-8; PMID:9931273; <http://dx.doi.org/10.1006/jbsi.1998.4054>.
40. Light WH, Brickner DG, Brand VR, Brickner JH. Interaction of a DNA zip code with the nuclear pore complex promotes H2A.Z incorporation and INO1 transcriptional memory. *Mol Cell* 2010; 40:112-25; PMID:20932479; <http://dx.doi.org/10.1016/j.molcel.2010.09.007>.
41. Schmid M, Arib G, Laemmler C, Nishikawa J, Durussel T, Laemmler UK. Nup-Pl: the nucleopore-promoter interaction of genes in yeast. *Mol Cell* 2006; 21:379-91; PMID:16455493; <http://dx.doi.org/10.1016/j.molcel.2005.12.012>.
42. Griffis ER, Altan N, Lippincott-Schwartz J, Powers MA. Nup98 is a mobile nucleoporin with transcription-dependent dynamics. *Mol Biol Cell* 2002; 13:1282-97; PMID:11950939; <http://dx.doi.org/10.1091/mbc.01-11-0538>.
43. Griffis ER, Craige B, Dimaano C, Ullman KS, Powers MA. Distinct functional domains within nucleoporins Nup153 and Nup98 mediate transcription-dependent mobility. *Mol Biol Cell* 2004; 15:1991-2002; PMID:14718558; <http://dx.doi.org/10.1091/mbc.E03-10-0743>.
44. Dilworth DJ, Suprapto A, Padovan JC, Chait BT, Wozniak RW, Rout MP, et al. Nup2p dynamically associates with the distal regions of the yeast nuclear pore complex. *J Cell Biol* 2001; 153:1465-78; PMID:11425876; <http://dx.doi.org/10.1083/jcb.153.7.1465>.
45. Nakielny S, Shaikh S, Burke B, Dreyfuss G. Nup153 is an M9-containing mobile nucleoporin with a novel Ran-binding domain. *EMBO J* 1999; 18:1982-95; PMID:10202161; <http://dx.doi.org/10.1093/emboj/18.7.1982>.
46. Rabut G, Doye V, Ellenberg J. Mapping the dynamic organization of the nuclear pore complex inside single living cells. *Nat Cell Biol* 2004; 6:1114-21; PMID:15502822; <http://dx.doi.org/10.1038/ncb1184>.
47. Rasala BA, Orjalo AV, Shen Z, Briggs S, Forbes DJ. ELYS is a dual nucleoporin/kinetochore protein required for nuclear pore assembly and proper cell division. *Proc Natl Acad Sci U S A* 2006; 103:17801-6; PMID:17098863; <http://dx.doi.org/10.1073/pnas.0608484103>.
48. Santangelo GM. Glucose signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* 2006; 70:253-82; PMID:16524925; <http://dx.doi.org/10.1128/MMBR.70.1.253-282.2006>.
49. Cabal GG, Genovesio A, Rodriguez-Navarro S, Zimmer C, Gadal O, Lesne A, et al. SAGA interacting factors confine sub-diffusion of transcribed genes to the nuclear envelope. *Nature* 2006; 441:770-3; PMID:16760982; <http://dx.doi.org/10.1038/nature04752>.
50. Taddei A, Van Houwe G, Hediger F, Kalck V, Cubizolles F, Schober H, et al. Nuclear pore association confers optimal expression levels for an inducible yeast gene. *Nature* 2006; 441:774-8; PMID:16760983; <http://dx.doi.org/10.1038/nature04845>.
51. Entian KD. Genetic and biochemical evidence for hexokinase PII as a key enzyme involved in carbon catabolite repression in yeast. *Mol Gen Genet* 1980; 178:633-7; PMID:6993859; <http://dx.doi.org/10.1007/BF00337871>.
52. Neigeborn L, Carlson M. Genes affecting the regulation of SUC2 gene expression by glucose repression in *Saccharomyces cerevisiae*. *Genetics* 1984; 108:845-58; PMID:6392017.
53. Neigeborn L, Carlson M. Mutations causing constitutive invertase synthesis in yeast: genetic interactions with snf mutations. *Genetics* 1987; 115:247-53; PMID:3549450.
54. Sarma NJ, Buford TD, Haley T, Barbara-Haley K, Santangelo GM, Willis KA. The nuclear pore complex mediates binding of the Mig1 repressor to target promoters. *PLoS One* 2011; 6:e27117; PMID:22110603; <http://dx.doi.org/10.1371/journal.pone.0027117>.
55. Boukaba A, Georgieva EI, Myers FA, Thorne AW, López-Rodas G, Crane-Robinson C, et al. A short-range gradient of histone H3 acetylation and Tup1p redistribution at the promoter of the *Saccharomyces cerevisiae* SUC2 gene. *J Biol Chem* 2004; 279:7678-84; PMID:14670975; <http://dx.doi.org/10.1074/jbc.M310849200>.
56. Zawadzki KA, Morozov AV, Broach JR. Chromatin-dependent transcription factor accessibility rather than nucleosome remodeling predominates during global transcriptional restructuring in *Saccharomyces cerevisiae*. *Mol Biol Cell* 2009; 20:3503-13; PMID:19494041; <http://dx.doi.org/10.1091/mbc.E09-02-0111>.
57. Geng F, Laurent BC. Roles of SWI/SNF and HATs throughout the dynamic transcription of a yeast glucose-repressible gene. *EMBO J* 2004; 23:127-37; PMID:14685262; <http://dx.doi.org/10.1038/sj.emboj.7600035>.
58. Mitchell L, Lambert JP, Gerdes M, Al-Madhoun AS, Skerjanc IS, Figeys D, et al. Functional dissection of the NuA4 histone acetyltransferase reveals its role as a genetic hub and that Eaf1 is essential for complex integrity. *Mol Cell Biol* 2008; 28:2244-56; PMID:18212056; <http://dx.doi.org/10.1128/MCB.01653-07>.
59. Luthra R, Kerr SC, Harreman MT, Apponi LH, Fasken MB, Ramineni S, et al. Actively transcribed GAL genes can be physically linked to the nuclear pore by the SAGA chromatin modifying complex. *J Biol Chem* 2007; 282:3042-9; PMID:17158105; <http://dx.doi.org/10.1074/jbc.M608741200>.
60. Titus LC, Dawson TR, Rexer DJ, Ryan KJ, Wente SR. Members of the RSC chromatin-remodeling complex are required for maintaining proper nuclear envelope structure and pore complex localization. *Mol Biol Cell* 2010; 21:1072-87; PMID:20110349; <http://dx.doi.org/10.1091/mbc.E09-07-0615>.
61. Ng HH, Robert F, Young RA, Struhl K. Genome-wide location and regulated recruitment of the RSC nucleosome-remodeling complex. *Genes Dev* 2002; 16:806-19; PMID:11937489; <http://dx.doi.org/10.1101/gad.978902>.

62. Gould VE, Orucevic A, Zentgraf H, Gattuso P, Martinez N, Alonso A. Nup88 (karyoporin) in human malignant neoplasms and dysplasias: correlations of immunostaining of tissue sections, cytologic smears, and immunoblot analysis. *Hum Pathol* 2002; 33:536-44; PMID:12094380; <http://dx.doi.org/10.1053/hupa.2002.124785>.
63. Graux C, Cools J, Melotte C, Quentmeier H, Ferrando A, Levine R, et al. Fusion of NUP214 to ABL1 on amplified episomes in T-cell acute lymphoblastic leukemia. *Nat Genet* 2004; 36:1084-9; PMID:15361874; <http://dx.doi.org/10.1038/ng1425>.
64. Soman NR, Wogan GN, Rhim JS. TPR-MET oncogene rearrangement: detection by polymerase chain reaction amplification of the transcript and expression in human tumor cell lines. *Proc Natl Acad Sci U S A* 1990; 87:738-42; PMID:2300559; <http://dx.doi.org/10.1073/pnas.87.2.738>.
65. Ma Z, Hill DA, Collins MH, Morris SW, Sumegi J, Zhou M, et al. Fusion of ALK to the Ran-binding protein 2 (RANBP2) gene in inflammatory myofibroblastic tumor. *Genes Chromosomes Cancer* 2003; 37:98-105; PMID:12661011; <http://dx.doi.org/10.1002/gcc.10177>.
66. Köhler A, Hurt E. Gene regulation by nucleoporins and links to cancer. *Mol Cell* 2010; 38:6-15; PMID:20385085; <http://dx.doi.org/10.1016/j.molcel.2010.01.040>.
67. Xu S, Powers MA. Nuclear pore proteins and cancer. *Semin Cell Dev Biol* 2009; 20:620-30; PMID:19577736; <http://dx.doi.org/10.1016/j.semcdb.2009.03.003>.
68. Lam DH, Aplan PD. NUP98 gene fusions in hematologic malignancies. *Leukemia* 2001; 15:1689-95; PMID:11681408; <http://dx.doi.org/10.1038/sj.leu.2402269>.
69. Gough SM, Slape CI, Aplan PD. NUP98 gene fusions and hematopoietic malignancies: common themes and new biologic insights. *Blood* 2011; 118:6247-57; PMID:21948299; <http://dx.doi.org/10.1182/blood-2011-07-328880>.
70. Funasaka T, Nakano H, Wu Y, Hashizume C, Gu L, Nakamura T, et al. RNA export factor RAE1 contributes to NUP98-HOXA9-mediated leukemogenesis. *Cell Cycle* 2011; 10:1456-67; PMID:21467841; <http://dx.doi.org/10.4161/cc.10.9.15494>.
71. Ren Y, Seo HS, Blobel G, Hoelz A. Structural and functional analysis of the interaction between the nucleoporin Nup98 and the mRNA export factor Rael. *Proc Natl Acad Sci U S A* 2010; 107:10406-11; PMID:20498086; <http://dx.doi.org/10.1073/pnas.1005389107>.
72. Takeda A, Sarma NJ, Abdul-Nabi AM, Yaseen NR. Inhibition of CRM1-mediated nuclear export of transcription factors by leukemogenic NUP98 fusion proteins. *J Biol Chem* 2010; 285:16248-57; PMID:20233715; <http://dx.doi.org/10.1074/jbc.M109.048785>.
73. Nakamura T, Largaespada DA, Lee MP, Johnson LA, Ohyashiki K, Toyama K, et al. Fusion of the nucleoporin gene NUP98 to HOXA9 by the chromosome translocation t(7;11)(p15;p15) in human myeloid leukemia. *Nat Genet* 1996; 12:154-8; PMID:8563753; <http://dx.doi.org/10.1038/ng0296-154>.
74. Raza-Egilmez SZ, Jani-Sait SN, Grossi M, Higgins MJ, Shows TB, Aplan PD. NUP98-HOXD13 gene fusion in therapy-related acute myelogenous leukemia. *Cancer Res* 1998; 58:4269-73; PMID:9766650.
75. Nakamura T, Yamazaki Y, Hatano Y, Miura I. NUP98 is fused to PMX1 homeobox gene in human acute myelogenous leukemia with chromosome translocation t(1;11)(q23;p15). *Blood* 1999; 94:741-7; PMID:10397741.
76. Panagopoulos I, Isaksson M, Billström R, Strömbeck B, Mittelman F, Johansson B. Fusion of the NUP98 gene and the homeobox gene HOXC13 in acute myeloid leukemia with t(11;12)(p15;q13). *Genes Chromosomes Cancer* 2003; 36:107-12; PMID:12461755; <http://dx.doi.org/10.1002/gcc.10139>.
77. Jankovic D, Gorello P, Liu T, Ehret S, La Starza R, Desjoberg C, et al. Leukemogenic mechanisms and targets of a NUP98/HHEX fusion in acute myeloid leukemia. *Blood* 2008; 111:5672-82; PMID:18388181; <http://dx.doi.org/10.1182/blood-2007-09-108175>.
78. Ghannam G, Takeda A, Camarata T, Moore MA, Viale A, Yaseen NR. The oncogene Nup98-HOXA9 induces gene transcription in myeloid cells. *J Biol Chem* 2004; 279:866-75; PMID:14561764; <http://dx.doi.org/10.1074/jbc.M307280200>.
79. Yassin ER, Sarma NJ, Abdul-Nabi AM, Dombrowski J, Han Y, Takeda A, et al. Dissection of the transformation of primary human hematopoietic cells by the oncogene NUP98-HOXA9. *PLoS One* 2009; 4:e6719; PMID:19696924; <http://dx.doi.org/10.1371/journal.pone.0006719>.
80. Sarma NJ, Yaseen NR. Amino-terminal enhancer of split (AES) interacts with the oncoprotein NUP98-HOXA9 and enhances its transforming ability. *J Biol Chem* 2011; 286:38989-9001; PMID:21937451; <http://dx.doi.org/10.1074/jbc.M111.297952>.
81. Kasper LH, Brindle PK, Schnabel CA, Pritchard CE, Cleary ML, van Deursen JM. CREB binding protein interacts with nucleoporin-specific FG repeats that activate transcription and mediate NUP98-HOXA9 oncogenicity. *Mol Cell Biol* 1999; 19:764-76; PMID:9858599.
82. Menon BB, Sarma NJ, Pasula S, Deminoff SJ, Willis KA, Barbara KE, et al. Reverse recruitment: the Nup84 nuclear pore subcomplex mediates Rap1/Gcr1/Gcr2 transcriptional activation. *Proc Natl Acad Sci U S A* 2005; 102:5749-54; PMID:15817685; <http://dx.doi.org/10.1073/pnas.0501768102>.
83. Wang GG, Cai L, Pasillas MP, Kamps MP. NUP98-NSD1 links H3K36 methylation to Hox-A gene activation and leukemogenesis. *Nat Cell Biol* 2007; 9:804-12; PMID:17589499; <http://dx.doi.org/10.1038/ncb1608>.
84. Wang GG, Song J, Wang Z, Dormann HL, Casadio F, Li H, et al. Haematopoietic malignancies caused by dysregulation of a chromatin-binding PHD finger. *Nature* 2009; 459:847-51; PMID:19430464; <http://dx.doi.org/10.1038/nature08036>.
85. Reader JC, Meekins JS, Gojo I, Ning Y. A novel NUP98-PHF23 fusion resulting from a cryptic translocation t(11;17)(p15;p13) in acute myeloid leukemia. *Leukemia* 2007; 21:842-4; PMID:17287853.
86. Wu SF, Zhang H, Cairns BR. Genes for embryo development are packaged in blocks of multivalent chromatin in zebrafish sperm. *Genome Res* 2011; 21:578-89; PMID:21383318; <http://dx.doi.org/10.1101/gr.113167.110>.
87. Bell O, Wirbelauer C, Hild M, Scharf AN, Schwaiger M, MacAlpine DM, et al. Localized H3K36 methylation states define histone H4K16 acetylation during transcriptional elongation in *Drosophila*. *EMBO J* 2007; 26:4974-84; PMID:18007591; <http://dx.doi.org/10.1038/sj.emboj.7601926>.
88. Lundin M, Nehlin JO, Ronne H. Importance of a flanking AT-rich region in target site recognition by the GC box-binding zinc finger protein MIG1. *Mol Cell Biol* 1994; 14:1979-85; PMID:8114729.
89. Pérez-Ortín JE, Estruch F, Matallana E, Franco L. Fine analysis of the chromatin structure of the yeast SUC2 gene and of its changes upon derepression. Comparison between the chromosomal and plasmid-inserted genes. *Nucleic Acids Res* 1987; 15:6937-56; PMID:2821486; <http://dx.doi.org/10.1093/nar/15.17.6937>.
90. Wu L, Winston F. Evidence that Snf-Swi controls chromatin structure over both the TATA and UAS regions of the SUC2 promoter in *Saccharomyces cerevisiae*. *Nucleic Acids Res* 1997; 25:4230-4; PMID:9336451; <http://dx.doi.org/10.1093/nar/25.21.4230>.
91. Gavin IM, Simpson RT. Interplay of yeast global transcriptional regulators Ssn6p-Tup1p and Swi-Snf and their effect on chromatin structure. *EMBO J* 1997; 16:6263-71; PMID:9321405; <http://dx.doi.org/10.1093/emboj/16.20.6263>.
92. Hirschhorn JN, Bortvin AL, Ricupero-Hovasse SL, Winston F. A new class of histone H2A mutations in *Saccharomyces cerevisiae* causes specific transcriptional defects in vivo. *Mol Cell Biol* 1995; 15:1999-2009; PMID:7891695.
93. Hirschhorn JN, Brown SA, Clark CD, Winston F. Evidence that SNF2/SWI2 and SNF5 activate transcription in yeast by altering chromatin structure. *Genes Dev* 1992; 6(12A):2288-98; PMID:1459453; <http://dx.doi.org/10.1101/gad.6.12a.2288>.
94. Schwabish MA, Struhl K. The Swi/Snf complex is important for histone eviction during transcriptional activation and RNA polymerase II elongation in vivo. *Mol Cell Biol* 2007; 27:6987-95; PMID:17709398; <http://dx.doi.org/10.1128/MCB.00717-07>.
95. Dror V, Winston F. The Swi/Snf chromatin remodeling complex is required for ribosomal DNA and telomeric silencing in *Saccharomyces cerevisiae*. *Mol Cell Biol* 2004; 24:8227-35; PMID:15340082; <http://dx.doi.org/10.1128/MCB.24.18.8227-8235.2004>.
96. Fleming AB, Pennings S. Tup1-Ssn6 and Swi-Snf remodelling activities influence long-range chromatin organization upstream of the yeast SUC2 gene. *Nucleic Acids Res* 2007; 35:5520-31; PMID:17704134; <http://dx.doi.org/10.1093/nar/gkm573>.
97. Pollard KJ, Peterson CL. Role for ADA/GCN5 products in antagonizing chromatin-mediated transcriptional repression. *Mol Cell Biol* 1997; 17:6212-22; PMID:9343382.
98. Sudarsanam P, Cao Y, Wu L, Laurent BC, Winston F. The nucleosome remodeling complex, Snf/Swi, is required for the maintenance of transcription in vivo and is partially redundant with the histone acetyltransferase, Gcn5. *EMBO J* 1999; 18:3101-6; PMID:10357821; <http://dx.doi.org/10.1093/emboj/18.11.3101>.
99. Neigeborn L, Celenza JL, Carlson M. Ssn20 is an essential gene with mutant alleles that suppress defects in SUC2 transcription in *Saccharomyces cerevisiae*. *Mol Cell Biol* 1987; 7:672-8; PMID:3547080.
100. Adkins MW, Tyler JK. Transcriptional activators are dispensable for transcription in the absence of Spt6-mediated chromatin reassembly of promoter regions. *Mol Cell* 2006; 21:405-16; PMID:16455495; <http://dx.doi.org/10.1016/j.molcel.2005.12.010>.
101. Türkel S, Turgut T, López MC, Uemura H, Baker HV. Mutations in GCR1 affect SUC2 gene expression in *Saccharomyces cerevisiae*. *Mol Genet Genomics* 2003; 268:825-31; PMID:12655409.
102. Sasaki H, Uemura H. Influence of low glycolytic activities in *ger1* and *ger2* mutants on the expression of other metabolic pathway genes in *Saccharomyces cerevisiae*. *Yeast* 2005; 22:111-27; PMID:15645478; <http://dx.doi.org/10.1002/yea.1198>.
103. Barbara KE, Haley TM, Willis KA, Santangelo GM. The transcription factor Gcr1 stimulates cell growth by participating in nutrient-responsive gene expression on a global level. *Mol Genet Genomics* 2007; 277:171-88; PMID:17124610; <http://dx.doi.org/10.1007/s00438-006-0182-0>.
104. López MC, Baker HV. Understanding the growth phenotype of the yeast *ger1* mutant in terms of global genomic expression patterns. *J Bacteriol* 2000; 182:4970-8; PMID:10940042; <http://dx.doi.org/10.1128/JB.182.17.4970-4978.2000>.