

Washington University School of Medicine Digital Commons@Becker

Open Access Publications

2010

Images in clinical medicine. Meckel's diverticulum

Brian T. Bucher

Washington University School of Medicine in St. Louis

Martin S. Keller

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation


Bucher, Brian T. and Keller, Martin S., "Images in clinical medicine. Meckel's diverticulum." *The New England Journal of Medicine*.363,21. 2045. (2010).

https://digitalcommons.wustl.edu/open_access_pubs/3145

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

IMAGES IN CLINICAL MEDICINE

Meckel's Diverticulum



A 10-YEAR-OLD BOY PRESENTED TO THE EMERGENCY DEPARTMENT WITH increasing periumbilical and epigastric pain after a 3-week period of intermittent abdominal discomfort. After clinical evaluation, computed tomography showed a proximal small-bowel intussusception. The results of a small-bowel series suggested the presence of a Meckel's diverticulum (Panel A, arrow), but no heterotopic gastric mucosa was identified on subsequent radionuclide scanning with the use of technetium-99m-labeled pertechnetate. Laparoscopy revealed a diverticulum on the antimesenteric border of the terminal ileum (Panel B, arrow), with its embryonic origin illustrated by the prominent vestige of the vitelline artery (arrowhead). No heterotopic gastric mucosal tissue was identified on pathological analysis of the resected surgical specimen. Intussusception that presents after infancy should raise suspicion regarding a pathologic lead point, including a Meckel's diverticulum. Symptomatic Meckel's diverticula frequently contain heterotopic tissue, predominantly heterotopic gastric mucosa. Gastric mucosal tissue is identifiable on radionuclide scanning, since it concentrates technetium-99m-labeled pertechnetate, but for diverticula without this tissue, the scan is normal. In this patient, the suggestion of a Meckel's diverticulum on the small-bowel series was confirmed. The child's abdominal symptoms resolved shortly after surgery and further episodes of intussusception have not been clinically evident during 1 year of follow-up.

Copyright © 2010 Massachusetts Medical Society.

Brian T. Bucher, M.D.
Martin S. Keller, M.D.

Washington University School
of Medicine
St. Louis, MO
kellerm@wudosis.wustl.edu