

## Washington University School of Medicine Digital Commons@Becker

---

### Open Access Publications

---

2005

# Potassium channels in *C. elegans*

Lawrence B. Salkoff

*Washington University School of Medicine in St. Louis*

Aguan Wei

*Washington University School of Medicine in St. Louis*

Beravan Baban

*Washington University School of Medicine in St. Louis*

Alice Butler

*Washington University School of Medicine in St. Louis*

Gloria L. Fawcett

*Washington University School of Medicine in St. Louis*

*See next page for additional authors*

Follow this and additional works at: [http://digitalcommons.wustl.edu/open\\_access\\_pubs](http://digitalcommons.wustl.edu/open_access_pubs)

---

### Recommended Citation

Salkoff, Lawrence B.; Wei, Aguan; Baban, Beravan; Butler, Alice; Fawcett, Gloria L.; Ferreira, Gonzalo; and Santi, Celia M., "Potassium channels in *C. elegans*." *WormBook*,. 1-15. (2005).

[http://digitalcommons.wustl.edu/open\\_access\\_pubs/3481](http://digitalcommons.wustl.edu/open_access_pubs/3481)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [engeszer@wustl.edu](mailto:engeszer@wustl.edu).

---

**Authors**

Lawrence B. Salkoff, Aguan Wei, Beravan Baban, Alice Butler, Gloria L. Fawcett, Gonzalo Ferreira, and Celia M. Santi

---

# Potassium channels in *C. elegans*\*

L. Salkoff<sup>1,2,§</sup>, A.D. Wei<sup>1</sup>, B. Baban<sup>1</sup>, A. Butler<sup>1</sup>, G. Fawcett<sup>1</sup>,  
G. Ferreira<sup>1</sup>, C.M. Santi<sup>1</sup>

<sup>1</sup>Department of Anatomy and Neurobiology and

<sup>2</sup>Department of Genetics, Washington University School of Medicine,  
Saint Louis MO, 63110 USA

## Table of Contents

1. Introduction .....	2
1.1. Many potassium channels in <i>C. elegans</i> have close mammalian orthologues .....	2
1.2. Genetic screens for abnormal behavioral phenotypes reveal potassium channel mutants .....	3
2. The 6TM gene families .....	4
2.1. Voltage-gated potassium channels .....	4
2.2. KQT potassium channels .....	4
2.3. <i>Eag</i> -like potassium channels .....	5
2.4. Calcium-activated <i>Slo</i> -like potassium channels .....	6
2.5. Cyclic nucleotide-gated cation channels .....	7
2.6. SK Small conductance, voltage insensitive calcium-activated potassium channels .....	7
3. 2TM potassium channels .....	7
4. 4TM potassium channels .....	7
4.1. Evolution may tune the excitability of individual neurons using 4TM channels .....	8
5. Usefulness of the information in <i>C. elegans</i> .....	9
5.1. <i>C. elegans</i> can be a useful system to dissect apart the components of cell excitability using its versatile genetic and molecular resources .....	9
5.2. Conserved genes represent natural structural variants .....	9
5.3. <i>C. elegans</i> could be a useful model system for studying genetic disease which involves ion channels .....	9
5.4. The study of ion channel functional properties in <i>C. elegans</i> has revealed unusual properties never before seen .....	10
5.5. Mutant analysis in <i>C. elegans</i> has revealed gain-of-function mutants with possible therapeutic uses .....	10
6. Acknowledgments .....	10
7. References .....	10

---

\*Edited by Erik M. Jorgensen and Joshua M. Kaplan. Last revised December 19, 2005. Published December 30, 2005. This chapter should be cited as: Salkoff, et al. Potassium channels in *C. elegans* (December 30, 2005), *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.42.1, <http://www.wormbook.org>.

**Copyright:** © 2005 Salkoff et al., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<sup>§</sup>To whom correspondence should be addressed. E-mail: [salkoffl@pcg.wustl.edu](mailto:salkoffl@pcg.wustl.edu)

## Abstract

Ion channels are the “transistors” (electronic switches) of the brain that generate and propagate electrical signals in the aqueous environment of the brain and nervous system. Potassium channels are particularly important because, not only do they shape dynamic electrical signaling, they also set the resting potentials of almost all animal cells. Without them, animal life as we know it would not exist, much less higher brain function.

Until the completion of the *C. elegans* genome sequencing project the size and diversity of the potassium channel extended gene family was not fully appreciated. Sequence data eventually revealed a total of approximately 70 genes encoding potassium channels out of the more than 19,000 genes in the genome. This seemed to be an unexpectedly high number of genes encoding potassium channels for an animal with a small nervous system of only 302 neurons. However, it became clear that potassium channels are expressed in all cell types, not only neurons, and that many cells express a complex palette of multiple potassium channels. All types of potassium channels found in *C. elegans* are conserved in mammals.

Clearly, *C. elegans* is “simple” only in having a limited number of cells dedicated to each organ system; it is certainly not simple with respect to its biochemistry and cell physiology.

## 1. Introduction


Potassium channels function in cellular tasks as ubiquitous as the setting of resting potentials (Katz, 1966), and as specific as defining the interspike intervals of endogenously beating neurons (Connor and Stevens, 1971). Not surprisingly, potassium channels are found in virtually all living cells including bacteria (Milkman, 1994; Saimi et al., 1999; Doyle et al., 1998; Jiang et al., 2002; Mackinnon and Doyle, 1997), yeast (Ketchum et al., 1995; Zhu et al., 1995), *Paramecia* (Jegla and Salkoff, 1995; Saimi et al., 1999), plants (Anderson et al., 1992; Sentenac et al., 1992 and Ketchum et al., 1996) and all animal cells, including electrically excitable tissues as well as inexcitable tissues, such as the liver (Pearson et al., 1999) and cells of the immune system (Cahalan and Chandy, 1997). Thus, potassium channels serve diverse functions such as shaping the dynamic electrical activity of neurons (Connor, 1980), modulating the immune response (Cahalan and Chandy, 1997) and controlling intestinal function (Koh et al., 1995). With the arrival of molecular biology and biochemical techniques, potassium currents in many cell types could be described not only electrophysiologically and pharmacologically, but also at the molecular level. Molecular techniques have also revealed a diversity far more extensive than was anticipated nearly two decades ago, when the first potassium channel gene was cloned from the *Drosophila Shaker* locus (Papazian et al., 1987; Kamb et al., 1987; Pongs et al., 1988).

### 1.1. Many potassium channels in *C. elegans* have close mammalian orthologues

Until the completion of the *C. elegans* genome sequencing project (*C. elegans* Sequencing Consortium, 1998) the size and diversity of the potassium channel extended gene family was not fully appreciated. Sequence data revealed a total of approximately 70 genes encoding potassium channels out of the more than 19,000 genes in the genome (Wei et al., 1996; Bargmann, 1998). This seemed to be an unexpectedly high number of genes encoding potassium channels for an animal with a small nervous system of only 302 neurons. However, it became clear that many potassium channels are expressed in all cell types, not only neurons (Salkoff et al., 1998) and that *C. elegans* is “simple” only in having a limited number of cells dedicated to each organ system; it is certainly not “simple” with respect to its biochemistry and physiology.

The genes encoding potassium channels can be divided into eight conserved families shown diagrammatically in Figure 1 (Wei et al., 1996). A universal feature of all potassium channels is a hyperconserved signature sequence (sometimes referred to as the “P” domain) that forms the potassium-selective filter of the pore (Heginbotham et al., 1992, 1994). These eight families fall into one of three major subunit classes as defined by the number of “P” domains and transmembrane domains per subunit. Those possessing two transmembrane domains per subunit (2TM) bracketing a single P domain are categorized as “inward rectifier” channels. Those possessing four transmembrane domains (4TM) contain two P domains per subunit and represented a novel class of channels. Those possessing six transmembrane domains per subunit (6TM) with one P domain, include the voltage-gated potassium channel family. All potassium channels are symmetrical or semi-symmetrical structures and require four P domains per channel. Thus, four 2TM or four 6TM subunits are required to form a channel, while only two of the 4TM subunits are required to form a single channel of that class. Families, such as voltage-gated channels, can be further divided into multiple **subfamilies** (e.g., Shaker, Shab, Shaw and Shal; Figure 1). Within a single species there are commonly

multiple members of each subfamily, such as Kv1.1 through Kv1.8, which represent eight mammalian homologues of the Shaker subfamily.



**Figure 1. Diagram of potassium channel families and subfamilies.** Three structural classes are shown defined by their subunit structure, having six (6TM), four (4TM), or two (2TM) transmembrane segments. The structural class of 6TM channels has six conserved gene families. See: <http://nt-salkoff.wustl.edu/> for information about mutant strains and cDNAs for expression studies.

## 1.2. Genetic screens for abnormal behavioral phenotypes reveal potassium channel mutants

An advantage of studying potassium channels in *C. elegans* is the ease with which the animal is subject to genetic manipulation. Because of the critical involvement of potassium channels in determining the excitability of neurons and muscles, it is perhaps not unexpected that genetic screens based on several *C. elegans* behavioral phenotypes have yielded examples of mutants linked to nearly every major class of potassium channel subunit. Mutant behavioral phenotypes have included the inability to lay eggs (*egl*, egg-laying abnormal), uncoordinated body movements (*unc*, uncoordinated), and abnormal defecation motor programs, in particular failure of the expulsion step (*exp*, expulsion defective; Brenner, 1974; Trent et al., 1983, Thomas, 1990). Surprisingly, nearly all of these potassium channel mutants were first isolated as dominant alleles that encode gain-of-function channel subunits. Loss-of-function mutants of these subunits, in general, largely resemble wild-type animals. These genetic results are consistent with an interpretation whereby a large degree of functional overlap or redundancy may be shared between wild-type potassium channels to determine normal excitability. The loss of an individual potassium channel may thus be compensated and not result in an overt phenotype. In contrast, any gain-of-function mutation which acts to hyper-activate a potassium channel may inappropriately suppress cellular excitability, causing a dramatic behavioral phenotype. Alternatively, specific functional roles for individual potassium channels may exist, but the deficits resulting from the loss of function of single potassium channel genes may be too subtle to be detected with coarse behavioral assays, and ideal laboratory culturing conditions. These genetic studies have provided insights into important regions of potassium channel subunits involved in gating, as well as potential auxiliary subunits required for activity. Among the 6TM potassium channel subunit genes, six examples are known

of mutants isolated by forward genetic screens (*egl-36*[Shaw], *egl-2*, *unc-103*[Eag], *exp-2* [Shab-like], *exp-3* [SK] and *slo-1*[Slo]). Currently, no mutants have been identified of the 2TM class of potassium channels but, as will be described, several of the 4TM class have been identified.

## 2. The 6TM gene families

### 2.1. Voltage-gated potassium channels

Voltage-gated potassium channels are the largest class of 6TM channels. These channels are activated by depolarization, but function with a wide diversity of activation and inactivation kinetics. In mammals, these channels are widely expressed in brain, muscle and heart where they control complex waveforms of electrical activity. They are also widely expressed in tissues outside the nervous system such as cells of the immune system (Cahalan and Chandy, 1997). Four conserved subfamilies, first cloned from *Drosophila*, comprise this family of channels: Shaker (Papazian et al., 1987; Kamb et al., 1987; Pongs et al., 1988), Shab, Shaw and Shal (Wei et al., 1990). Many vertebrate orthologues in each subfamily have been isolated (Chandy and Gutman, 1995). A total of eleven *C. elegans* genes were identified encoding voltage-gated potassium channel subunits that fall within the four subfamilies of Shaker (Kv1), Shab (Kv2), Shaw (Kv3) and Shal (Kv4). This group includes channels with biophysical properties conserved with other species (Bianchi et al., 2003) as well as those with unexpected biophysical properties (Davis et al., 1999; Fleischhauer et al., 2000). Examples of physiological and genetic studies undertaken with members of this family follow.

**Shaw Channels (Kv3 family)** *egl-36* mutants were isolated from screens for egg-laying deficient mutants. These mutants (*n728*, *n2332*) are also moderately defective in generating the expulsion step of the defecation motor program, and genetically behave as dominant alleles. Intragenic second mutations that result in null alleles, nearly completely revert the mutant phenotypes. *egl-36* encodes a voltage-gated potassium channel subunit which is an ortholog of vertebrate Kv3 (Shaw-like, KCNC) subunits (Elkes, et al., 1997, Johnstone et al., 1997). Heterologous expression studies of EGL-36 channels revealed voltage-gated potassium currents with functional properties similar to vertebrate Kv3 channels. The two dominant mutant alleles encode missense mutations which alter single residues within the amino-terminal T1 subunit tetramerization domain (*n728*, E142K) and the S6 transmembrane segment (*n2332*, P438S) of the predicted channel protein. Both mutations cause a hyperpolarizing shift in the voltage-dependence for activation, which could confer abnormal inhibition of cellular excitability. Promoter::GFP fusion experiments indicate expression of *egl-36* in uterine and vulva muscles, and in enteric motoneurons (AVL, DVB), consistent with the observed behavioral phenotype.


**Shab (Kv2)-like:** *Exp-2* encodes a potassium channel subunit related to the class of voltage-gated potassium channel subunits, but without a clear vertebrate ortholog (Davis et al., 1999). However, sequence comparisons clearly suggest that it is distantly related to the Shab (Kv2) family. Interestingly, homozygous *exp-2 gf* animals are inviable and die as young larvae, most likely because they are unable to feed. Heterozygous *+/exp-2(gf)* animals survive, but are severely expulsion defective, egg-laying defective, and exhibit abnormally brief pharyngeal contractions. *Exp-2* null animals grossly resemble wild-type and are viable, but these animals exhibit abnormally prolonged pharyngeal contractions. Promoter::GFP fusion experiments indicate prominent expression of *exp-2* in pharyngeal muscles, as well as enteric muscles and sensory neurons in the head and tail ganglia. Heterologous expression studies of EXP-2 channels revealed an unusual voltage-gated potassium channel, which gates by briefly conducting repolarizing current upon hyperpolarizations, but only when preceded by a prior depolarization (Davis et al., 1999; Fleischhauer et al., 2000). This unusual combination of gating properties functionally resembles the vertebrate *ether-a-go-go* related HERG (KCNH2) potassium channel (Trudeau et al., 1995; Curran et al., 1995; Sanguinetti et al., 1995), but because these two channels share low sequence similarity, this is likely to be an example of convergent evolution. Like HERG channels in vertebrate cardiomyocytes, EXP-2 functions to limit the length of plateau potentials generated by pharyngeal muscle cells, by providing a rapid repolarizing conductance at the end of the plateau potential. The *exp-2(sa26) gf* mutation alters a single residue in S6 (the sixth membrane spanning segment of the alpha subunit) (C480Y). This mutation shifts the voltage-dependence of activation towards negative potentials, creating a mutant channel that is constitutively active at normal resting potentials (Davis et al., 1999). These *exp-2* studies highlight the central importance of S6 structures in controlling the gating of potassium channels, even for functionally and structurally diverse types of channel subunits.

### 2.2. KQT potassium channels

KQT potassium channels are related to the human KvLQT (KCNQ) channel family (Wang et al., 1996), recently implicated in one form of long-QT syndrome, a hereditary cardiac syndrome. There are five of these genes

in humans (KCNQ1-5). Remarkably, four of the five human genes are associated with hereditary diseases. In addition to long-QT syndrome (associated with KCNQ1), KCNQ mutations have been linked with Benign Familial Neonatal Convulsions (KCNQ2 and KCNQ3; Biervert et al., 1998; Charlier et al., 1998; Singh et al., 1998 and 2003), and the Non-syndromic Autosomal Deafness, DFNA2 (KCNQ4; Kubisch et al., 1999).

*C. elegans* has three genes encoding channels of the KQT family. The *C. elegans* homologue of the human cardiac KvLQT1 channel (KQT-1) may have a conserved role that offers insight into its function in both species. As discussed previously, the orthologues in humans (called the KCNQ family) are the loci of many genetic diseases. Human KCNQ channels are expressed in both cardiac cells where they limit the length of the plateau phase of the cardiac action potential, and in the intestine where they may help provide the negative membrane potential necessary for apical chloride efflux mediating fluid secretion. In *C. elegans* genes of the KQT family are expressed both in the pharynx which has many cardiac-like electrical properties (Avery and Horvitz, 1989; Raizen and Avery, 1994) and also in the intestine (figure 2). Deletions of these genes have now been isolated (Wei et al., 2002) which may help in the further analysis of their physiological roles in *C. elegans*.



**Figure 2.** A. Functional expression of *C. elegans* KQT cDNA's in *Xenopus* oocytes using the two electrode voltage-clamp technique; test pulses were applied at 10 mV intervals from a holding potential of  $-80$  mV. The slow onset of the voltage-dependent current resembles that of human KCNQ counterparts (Wei et al., 2005). B. Tissue specific expression patterns in the cardiac-like pharynx and intestine as revealed by Promoter-GFP expression experiments. C. Cardiac-like action potentials in the *C. elegans* pharynx (Salkoff, unpublished; see also Raizen and Avery, 1994 and Shtonda and Avery, 2005).

### 2.3. *Eag*-like potassium channels

First discovered in *Drosophila* (Warmke et al., 1991), these channels form a diverse family (Warmke and Ganetzky, 1994) that may be divided into three distinct subfamilies (*Eag*, *Erg* and *Elk*). One human orthologue, *Herg*, with unusual gating properties, is also implicated in long-QT syndrome (Smith et al., 1996; Trudeau et al., 1995; Curran et al., 1995). *C. elegans* has two genes of this family, one which is an *eag* orthologue (*egl-2*) and one which is an *erg* orthologue (*unc-103*). Similar genetic and functional studies have been observed for mutants of both (Wienshenker et al., 1999; Reiner et al., 1999; Petersen et al., 2004). *Egl-2* *gf* mutants exhibit pleiotropic behavioral defects. In addition to egg-laying deficiency, these mutants show sluggish movement, defective expulsion-

defecation cycles, and more subtle defects in mechanosensation and chemosensation. Two independently isolated *gf* alleles of *egl-2* (*n693*, *n2656*) encode the same missense mutation, altering a single highly conserved residue (A533V) in the S6 transmembrane segment of the channel subunit. Heterologous expression studies confirmed that this mutation shifts the voltage-dependence of activation for this channel towards hyperpolarized potentials, allowing these mutant channels to abnormally inhibit excitability. Promoter::GFP fusion experiments indicate expression in vulva muscles, enteric muscles, mechanosensory and chemosensory neurons. Likewise *unc-103(e1597)* *gf* mutants exhibit behavioral defects resulting from a missense mutation altering a similar conserved residue in S6 (A334T), though functional analysis of the UNC-103 channel by electrophysiological studies has not been reported. Though *unc-103* null hermaphrodites are grossly wild-type, male worms carrying homozygous null mutations of *unc-103* show abnormal spicule protraction, consistent with hyperactive protractor muscle activity (Garcia and Sternberg, 2003). In addition, a genetic interaction between UNC-103 channels and calcium/calmodulin-dependent kinase II (CaMKII) (encoded by *unc-43*) is suggested, because the *unc-103* null mutation suppresses expulsion defects caused by an *unc-43(n498)* *gf* mutation (Reiner et al., 1999).

## 2.4. Calcium-activated Slo-like potassium channels

Channels of this family, were also first cloned in *Drosophila* (Elkins et al., 1986; Atkinson et al., 1991) and form large conductance ‘BK-type’ channels, gated by both voltage and intracellular calcium (Adelman et al., 1992; Butler et al., 1993). This class can be divided into three subfamilies, typified by the vertebrate genes *mSlo* (*Slo-1*), *Slack* (*Slo-2*) and *mSlo-3* (*Slo-3*). In mammals, *Slo-1* and *Slack* (*Slo-2*) are expressed in brain, various muscles, and most cell types that also express voltage-dependent calcium channels. *Slo-1* channels control the myogenic tone of arterial and airway smooth muscle, and are also key regulators of neurotransmitter release. *C. elegans* possess a single Slo-1 orthologue (*slo-1*) that plays an important regulatory role in synaptic transmission (Wang et al., 2001). Slo-2 is a somewhat more distantly related *C. elegans* Slo member; its vertebrate orthologue is the Slack gene (Joiner et al., 1998, Yuan et al., 2003). Interestingly, the *slo-2* gene expresses large conductance potassium channels with sensitivity to both intracellular calcium and chloride ions (Yuan et al., 2000).

In contrast to other potassium channel mutants, *slo-1* mutants were isolated as recessive loss-of-function mutations, from two independent forward genetic screens. Phenotypically, *slo-1 lof* mutants are grossly wild-type, though their movement is slightly jerky, and they exhibit decreased resistance to aldicarb paralysis compared to wild-type animals, consistent with increased cholinergic synaptic transmission.

Mutants of SLO-1 potassium channels were useful tools to show that the SLO-1 channel regulates the quantal content of neurotransmitter release. The absence of electrophysiological techniques to characterize synaptic physiology had been a handicap in studying synaptic function in *C. elegans* until the innovative studies by Janet Richmond (Richmond and Jorgensen, 1999). The techniques developed for those studies were further developed by Wang et al. (2001) to permit quantal analysis of neurotransmitter release. Large conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channels (Slo-1) have been implicated in the control of neurotransmitter release in vertebrates by pharmacological methods. Mutant analysis and physiological methods in *C. elegans* similarly showed that *slo-1* controls the quantal content of neurotransmitter release. Six *Slo-1* mutants were isolated in a screen for suppressors of a hypomorphic syntaxin mutant (*unc-64*), a mutation that confers behavioral lethargy. The selection of suppressor mutations was based on the reversal of lethargy. The quantal content of evoked excitatory postsynaptic currents (EPSCs) in *unc-64* mutants was ~5% of wild-type, but the addition of a *slo-1* mutation increased this fourfold. *Slo-1* mutations were also found to increase the quantal content of release even in a wild-type (*unc-64+*) background (Wang et al., 2001). Independently, multiple *slo-1 lof* mutants were also isolated from a screen for resistance to acute ethanol intoxication (Davies et al., 2003). This study argued that SLO-1 channels are specific targets for up-regulation by ethanol, which could mediate the suppression of movement caused by ethanol intoxication.

**Slo-2, a BK channel requiring both chloride and calcium** (Yuan et al., 2000; Jospin et al., 2002). Potassium channels have been reported to be modulated by many diverse factors, but modulation of gating by intracellular chloride ion has previously been unsuspected. However, when *slo-2*, was cloned in *C. elegans* and expressed in the heterologous *Xenopus* oocyte expression system, it was found to have a novel requirement for both chloride and calcium; no activity is detectable in the absence of either. The sequence divergence between *slo-2* and *slo-1* may underlie some of their functional differences, especially in the “tail” domain of the channel where calcium-sensing may occur. The unique property of chloride dependence has never been described for any other ion channel. Perhaps the SLO-2 channel is an intracellular Cl<sup>-</sup> sensor linking changes in Cl<sub>i</sub><sup>-</sup> with membrane electrical behavior. *C. elegans* transformed with a *slo-2* promoter::GFP transgene revealed that *slo-2* is widely expressed in both neurons and muscles and is highly abundant as suggested by the large number of ESTs. In muscle cells the SLO-2 current has been shown to be larger than the voltage-dependent potassium currents which are contributed by Shaker (Kv1)



and Shal (Kv4) channel types (Santi et al., 2003). Intriguingly, the *slo-2* tissue expression pattern overlaps with that of *slo-1* suggesting that the two subunits might form functional heteromeric channels *in vivo*. The rat homologue of SLO-2 has been reported to possibly form heteromultimers with the mammalian BK channel, mSLO-1 (Joiner et al., 1998). SLO-2's dependence on Cl<sup>-</sup> suggests that intracellular Cl<sup>-</sup> homeostasis may play an important role in regulating cellular excitability through this unusual K<sup>+</sup> channel. Recently, the mammalian orthologue of the SLO-2 channel was shown to be a sodium-activated potassium channel (which is also co-activated by chloride ions) (Yuan et al., 2003). The sodium-activated potassium channel was originally identified in cardiac myocytes and thought to confer resistance to hypoxia. Interestingly, *slo-2* mutants in *C. elegans* are especially susceptible to hypoxia, suggesting a conserved role for Slo-2 channels. Another point of interest is the fact that calcium and chloride ions co-activate SLO-2 channels in *C. elegans*, while sodium and chloride ions co-activate SLO-2 channels in mammals. This may be an evolutionary adaptation reflecting the absence of the classical voltage-dependent sodium channel in *C. elegans*.

## 2.5. Cyclic nucleotide-gated cation channels

These channels are gated by intracellular cyclic nucleotides (Kaupp et al., 1989), and are expressed in receptor neurons of vertebrate retina and olfactory epithelium, where they transduce primary sensory information into changes in membrane potential. This transduction process is mediated by the intracellular messengers cAMP (olfactory receptors; Goulding et al., 1992), or cGMP (visual photoreceptors; Kaupp et al., 1989). Though these channels are non-selective cation channels (Heginbotham et al., 1992), they are structurally similar to 6TM potassium channels and are considered here as a related family. Five *C. elegans* genes encode 6TM subunits with high sequence conservation to cyclic nucleotide-gated cation channels in other species. *C. elegans* with mutations in either of two cyclic nucleotide-gated channel genes (*tax-2* and *tax-4*) result in animals which are defective in chemotaxis and thermotaxis (Coburn and Bargmann 1996; Komatsu et al., 1996).

## 2.6. SK Small conductance, voltage insensitive calcium-activated potassium channels

Small-conductance calcium-activated potassium channels (SK channels) are fundamental regulators of excitability. Unlike high conductance calcium-activated potassium channels which are activated by both voltage and calcium ions, SK channels are activated solely by intracellular calcium ions and therefore may function at rest or at hyperpolarized potentials. SK channels underlie the slow after-hyperpolarization (sAHP) following an action potential. Regulation of the sAHP represents a powerful modulatory influence on excitability. In addition to calcium, many neurotransmitters act on the sAHP. When these channels are present, changes in intracellular calcium exert strong effects on neuronal excitability. SK channels are also found in cells of the immune system. There are four genes encoding SK channels in *C. elegans*, one of which appears to be the locus of the *exp-3* mutation. *exp-3* mutants exhibit a similar phenotype to other described dominant potassium channel mutants. Mutant *exp-3(n2372)* *gf* animals are severely defecation-expulsion and egg-laying defective (Thomas, 1990). We have recently mapped *exp-3* to B0399.1, which is the *C. elegans* ortholog of a vertebrate SK gene that encodes a small-conductance calcium activated potassium channel subunit (Salkoff et al., unpublished). We have generated a null deletion of this gene. Similar to other potassium channel mutants, homozygous null animals grossly resemble wild-type, though there may be a slight temperature-dependent larval lethality at higher temperatures (>20°C).

## 3. 2TM potassium channels

Three *C. elegans* genes, *nIRK1-3* have the same subunit structure as inward rectifier potassium channel subunits from other species. Vertebrate 2TM inward rectifier subunits can be grouped into at least six subfamilies termed Kir1 through Kir6 (Doupnik et al., 1995). However, the two nematode nIRK genes appear more closely related to each other than to any of the vertebrate genes. This raises the question of whether vertebrate 2TM subfamilies have specific homologues in *C. elegans* and other invertebrates, or if they differentiated in the vertebrate line of evolution. No mutants have yet been reported to be associated with any of the three genes encoding 2TM potassium channel subunits.

## 4. 4TM potassium channels

Surprisingly, the most frequently encountered type of potassium channel belongs to the 4TM class of channel. Forty six genes of this class are currently known. It has been suggested that this large number may reflect a special adaptation of *C. elegans* to customize the electrical properties of individual cell types (Salkoff et al., 2001). In *Drosophila* there are approximately eleven genes encoding channels of this type (Adams et al., 2000; Littleton and Ganetzky, 2000; and Rubin et al., 2000), while in humans there are approximately fifteen genes.

Four mutants are known of the 4TM “TWK” subunit class (*sup-9*, *twk-18*, *unc-58*, *egl-23*). *sup-9* encodes TWK-38 (de la Cruz et al., 2003), which is a clear structural orthologue of the human TASK (KCNK9) gene (Duprat, et al., 1997). SUP-9 likely functions in a native channel complex with at least two other gene products (SUP-10 and UNC-93), based on extensive genetic evidence (de la Cruz, et al., 2003; Levin and Horvitz, 1993; Levin and Horvitz, 1992). *unc-93* is a member of an extended gene family encoding multipass membrane proteins conserved among *C. elegans*, *Drosophila*, and humans. This family includes the sulfonyleurea receptor that forms a channel complex with the vertebrate K<sub>ATP</sub> channel (see below). SUP-10 is a unique putative membrane protein. Gain-of-function alleles of *sup-9*, *unc-93* or *sup-10* all cause a similar behavioral phenotype, characterized by sluggish body movements, egg-laying and defecation defects, and a “rubberband” uncoordinated response to prodding of the head, whereas loss-of-function mutants resemble wild-type. However, loss of function of any gene in this group suppresses the mutant phenotype caused by a gain of function of either of the remaining two genes. Genetic suppression of *sup-9(gf)* is observed by the loss of *unc-93* function, even when *sup-9(gf)* is overexpressed as an extrachromosomal array, suggesting an absolute requirement for both gene products (de la Cruz, et al., 2003). Taken together, these results argue that SUP-9, UNC-93, and SUP-10 combine to form a native channel complex, perhaps analogous to the vertebrate K<sub>ATP</sub> channel complex formed by 2TM Kir6.2 potassium channel subunits and the SUR sulfonyleurea receptor (Auilar-Bryan, et al., 1995; Inagaki, et al., 1996; Shyng and Nichols, 1997). The *sup-9(n1550)* gain-of-function allele encodes a missense mutation that substitutes threonine for alanine 236 in the fourth transmembrane segment of the subunit. This presumably acts to stabilize the open conformation of the channel. Electrophysiological analysis of the putative SUP-9 channel complex, however, has not been reported thus far by heterologous expression studies, and may require *in situ* recordings from *C. elegans* cells.

By contrast, combined genetic and electrophysiological analyses of *twk-18* mutants have been possible. Two separate dominant alleles of *twk-18* (also referred to as *unc-110* and *mah-2* in the literature; Reiner, et al., 1995 and Hosono, et al., 1985, respectively) were isolated from independent behavioral screens for uncoordinated movement (*e1913*) and temperature-sensitive paralysis (*cn110*). Homozygous *twk-18(e1913)* animals die as embryonic lethals, but heterozygous *twk-18(e1913)/+* animals display flaccid paralysis and uncoordinated movement. The second allele, *cn110*, is viable as a homozygous animal and moves well at lower temperatures (15°-22°C), but becomes uncoordinated and paralyzed at elevated temperatures (>22°C). *Twk-18* encodes a 4TM potassium channel subunit without a clear vertebrate ortholog (Kunkel, et al., 2000), similar to most of the ~ 46 *C. elegans twk* genes. Expression of *twk-18* is confined to bodywall muscles based on promoter::GFP reporter studies, consistent with its mutant behavioral phenotype. Electrophysiological analysis of TWK-18 by heterologous expression studies in *Xenopus* oocytes revealed non-inactivating potassium-selective currents with little voltage-dependence, but activated by elevated temperatures (>23°C), with a steep temperature-dependence. The effect of the gain-of-function mutations is a 4- to 5-fold increase in the level of currents measured in *Xenopus* oocytes without an obvious change in unitary conductance through single channels. However, the exact mechanism of this increase remains unresolved. The gain-of-function alleles encode missense mutations G165D (*e1913*) at the base of the second transmembrane segment, and M280I (*cn110*) at the top of the fourth transmembrane segment of the subunit, suggesting that these regions of the channel subunit may be critical for gating.

Further genetic analyses of two behavioral mutants have identified linkages to two additional 4TM TWK subunits, *unc-58* (encoding TWK-19) and *egl-23* (encoding TWK-41). As with *sup-9* and *twk-18*, these additional mutants were isolated as semidominant alleles which were reverted by intragenic second mutations that render the genes effectively null. These null mutants grossly resemble wild-type animals. The gain-of-function *unc-58(e665)* mutant displays rigid paralysis with a characteristic “shaking” phenotype, constitutive egg-laying, and short “dumpy” body length consistent with hypercontracted bodywall muscles. This mutant phenotype is suppressible by treatment with the pesticide endosulfan which may directly block the UNC-58 channel (Tzoneva and Thomas, personal communications). Promoter::GFP fusion studies indicate expression of *unc-58* in ventral cord motoneurons and interneurons (Salkoff et al., 2000). These results are consistent with an *unc-58 gf* mutation which alters ionic selectivity, perhaps analogous to the *weaver* mutation of GIRK2 (Silverman, et al., 1996), which converts an inhibitory potassium-selective channel to an excitatory non-selective cation channel. *Egl-23(n601) gf* mutant animals are defective in locomotion, egg-laying and enteric muscle contractions. The *egl-23(n601) gf* mutant phenotype can be reverted by intragenic second mutations, or by RNAi (Johnstone and Thomas, personal communications). Electrophysiological analyses of the channels encoded by either *unc-58* or *egl-23* have not yet been reported.

#### 4.1. Evolution may tune the excitability of individual neurons using 4TM channels

The relationship between the genome and the evolution of the nervous system may differ between an animal like *C. elegans* with 302 neurons, and mammals with tens of billions of neurons. It was found that the 4TM class of nonconserved potassium channels is highly expanded in *C. elegans* and may play a special role in the evolution of its nervous system. The *C. elegans* genome contains an extended gene family of potassium channels whose members fall into two evolutionary divergent classes. One class constitutes an ancient conserved “set” of K<sup>+</sup> channels with orthologues in both humans and *Drosophila*; a second larger class is made up of rapidly evolving genes unique to *C. elegans*. Chief among this second class are the novel “TWK” (4TM) type potassium channels that may function as regulated leak conductances to modulate cell electrical excitability. This inventory of TWK channels is far larger in *C. elegans* than in humans or *Drosophila*. Evidence for their rapid evolution is the observation that they tend to be less conserved between *C. elegans* and a sibling species, *C. briggsae*, than are voltage-dependent channels, like Shaker. Unlike conserved channel genes, many TWK genes are expressed in very few cells. DNA enhancer elements associated with these genes were also identified that direct gene expression to individual neurons. *C. elegans* may maintain an exceptionally large inventory of novel TWK channels (as well as ligand-gated channels) as an adaptive mechanism to “fine tune” the electrical properties of individual neurons, making the most of its limited circuitry (Salkoff et al., 2001).

### 5. Usefulness of the information in *C. elegans*

#### 5.1. *C. elegans* can be a useful system to dissect apart the components of cell excitability using its versatile genetic and molecular resources

GFP-promoter experiments showed that at least nine genes encoding potassium channel subunits are expressed in *C. elegans* muscle. By using genetic, RNAi and physiological techniques, the molecular identity of all major components of the outward K<sup>+</sup> currents in body wall muscle cells in culture and *in situ*, were revealed (Jospin et al. 2002; Santi et al. 2003). Under physiological conditions, outward current is dominated by the products of only two genes, *Shaker* (Kv1) and *Shal* (Kv4), both expressing voltage-dependent potassium channels. Both of these voltage-dependent currents can be removed either by the use of mutants or by treating cultured cells with appropriate RNAi's. GFP-promoter experiments also indicated that the high conductance Ca<sup>2+</sup>&Cl<sup>-</sup> activated SLO-2 channels are prominently expressed. A deletion mutant was generated to identify the SLO-2 current in native cells. It was shown that native SLO-2 channels were active only when intracellular Ca<sup>2+</sup> and Cl<sup>-</sup> were raised above normal physiological conditions, as occurs during hypoxia. However, under such conditions, SLO-2 is the largest outward current, contributing up to 87% of the total current. Other genes encoding potassium channels are expressed in muscle, but do not appear to contribute a large component of total outward current under physiological conditions. However they may contribute currents conditional on other factors such as temperature, or stringent conditions of salinity, osmolarity or pH. Hence, excitable membranes may contain a complex combination of household conductances functioning under normal circumstances, with additional conductances that are activated during unusual physiological circumstances.

#### 5.2. Conserved genes represent natural structural variants

The large number of conserved genes encoding ion channels in *C. elegans* represent structural variants useful to: 1. Biophysicists, protein structure analysts and X-ray crystallographers. Similarities and differences in biophysical properties between species can be mapped in structure-function studies to gain insight into mechanisms and structural domains responsible for particular properties. 2. Pharmacologists. In an analogous manner similarities and differences in pharmacological properties between the *C. elegans* and vertebrate channels can be mapped in structure-function studies to gain insight into binding sites and mechanisms of channel-drug interactions.

#### 5.3. *C. elegans* could be a useful model system for studying genetic disease which involves ion channels

Hereditary diseases involving ion channels cover various fields of medicine such as neurology (nocturnal frontal lobe epilepsy, benign neonatal convulsions, episodic ataxia, hemiplegic migraine, deafness, stationary night blindness), nephrology (X-linked recessive nephrolithiasis, Bartter), myology (hypokalemic and hyperkalemic periodic paralysis, myotonia congenita, paramyotonia, malignant hyperthermia), and cardiology (LQT syndrome). All types of voltage-gated ion channels for cations (sodium, calcium, and potassium channels) and anions (chloride channels) are involved (reviewed in Lehmann-Horn and Jurkat-Rott, 1999). It is difficult to predict all the ways in which *C. elegans* might be useful in contributing to a better understanding of hereditary disease. For example, the functions of the polycystins and the molecular basis of kidney cystogenesis are now better understood thanks to

research done in *C. elegans* (Barr and Sternberg, 1999). One polycystin was discovered to be an ion channel. Based on the role of polycystins in *C. elegans* sensory neuron function and the conservation of PKD pathways it is now proposed that polycystins act as sensors of the extracellular environment, mediating intracellular transduction events which are important in both physiological and morphogenetic processes (Kaletta et al., 2003).

#### **5.4. The study of ion channel functional properties in *C. elegans* has revealed unusual properties never before seen**

Potassium channels have many varied properties, some of which were seen for the first time in *C. elegans*. Two examples are a temperature-sensitive potassium channel (Kunkel et al., 2000), and a chloride-sensitive potassium channel (Yuan et al., 2000). These findings may encourage further studies in the perception and sensation of temperature-induced pain, or studies of how cells and tissues compensate for temperature-changes. The discovery of a chloride sensitive channel may encourage further studies of the role of chloride in controlling membrane potential and in repolarizing active responses. Since the mammalian orthologue of Slo-2 is expressed in the kidney, its role in kidney function *vis-à-vis* its possible chloride sensitivity may also be an important area of investigation.

#### **5.5. Mutant analysis in *C. elegans* has revealed gain-of-function mutants with possible therapeutic uses**

In studying several instances of mutant potassium channels in *C. elegans*, several examples of gain-of-function mutants were uncovered that suggest ways in which transgenic techniques may be useful in future clinical applications. These are mutants in which potassium channels are open inappropriately either because their voltage range of activation has been shifted negatively, or because their activity level is abnormally high for other reasons. These results could guide the construction of vectors designed to electrically silence cells, possibly without detrimental effects on cell viability. The introduction of such vectors into selected sets of sensory neurons in mammals, such as C-fibers, may offer an effective remedy for such conditions as chronic pain. Naturally such uses have to wait for the development of tissue-specific promoters and viral or “gene-gun” techniques for the introduction of vectors into cell populations, but *C. elegans* may also offer an effective system for “proof-of-principle” experiments.

## **6. Acknowledgments**

Supported by grants R24 RR017342-01 and R01 GM067154-01A1 from the National Institutes of Health to LS, and IBN-0117341 from the National Science Foundation to AW. Special thanks to Michael Nonet for many helpful suggestions.

## **7. References**

- Adams, M.D., et al. (2000). The genome sequence of *Drosophila melanogaster*. *Science* 287, 2185–2195. [Abstract Article](#)
- Adelman, J.P., Shen, K.Z., Kavanaugh, M.P., Warren, R.A., Wu, Y.N., Lagrutta, A., Bond, C.T., and North, R.A. (1992). Calcium-activated potassium channels expressed from cloned complementary DNAs. *Neuron* 9, 209–216. [Abstract Article](#)
- Aguilar-Bryan, L., Nichols, C.G., Wechsler, S.W., Clement, J.P.T., Boyd, A.E. III, Gonzalez, G., Herrera-Sosa, H., Nguy, K., Bryan, J., and Nelson, D.A. (1995). Cloning of the beta cell high-affinity sulfonylurea receptor: a regulator of insulin secretion. *Science* 268, 423–426. [Abstract](#)
- Anderson, J.A., Huprikar, S.S., Kochian, L.V., Lucas, W.J., and Gaber, R.F. (1992). Functional expression of a probable *Arabidopsis thaliana* potassium channel in *Saccharomyces cerevisiae*. *PNAS* 89, 3736–3740. [Abstract](#)
- Atkinson, N.S., Robertson, G.A., and Ganetzky, B. (1991). A component of calcium-activated potassium channels encoded by the *Drosophila Slo* locus. *Science* 253, 551–555. [Abstract](#)
- Bargmann, C.I. (1998). Neurobiology of the *Caenorhabditis elegans* genome. *Science* 282, 2028–2033. [Abstract Article](#)

- Barr, M.M., and Sternberg, P.W. (1999). A polycystic kidney-disease gene homologue required for male mating behavior in *C. elegans*. *Nature* 401, 386–389. [Abstract Article](#)
- Bianchi, L., Kwok, S.M., Driscoll, M., and Sesti, F. (2003). A potassium channel-MiRP complex controls neurosensory function in *Caenorhabditis elegans*. *J. Biol. Chem.* 278, 12415–12424. [Abstract Article](#)
- Biervert, C., Schroeder, B.C., Kubisch, C., Berkovic, S.F., Propping, P., Jentsch, T.J., and Steinlein, O.K. (1998). A potassium channel mutation in neonatal human epilepsy. *Science* 279, 403–406. [Abstract Article](#)
- Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71–94. [Abstract](#)
- Butler, A., Tsunoda, S., McCobb, D.P., Wei, A., and Salkoff, L. (1993). *mSlo*, a complex mouse gene encoding “maxi” calcium-activated potassium channels. *Science* 261, 221–224. [Abstract](#)
- Cahalan, M.D., and Chandy, K.G. (1997). Ion channels in the immune system as targets for immunosuppression. *Curr. Opin. Biotechnol.* 8, 749–756. [Abstract](#)
- C. elegans* Sequencing Consortium. (1998). Genome sequence of the nematode *C. elegans*: A platform for investigating biology. *Science* 282, 2012–2018. [Article](#)
- Chandy, K.G., and Gutman, G.A. (1995). Voltage-gated K<sup>+</sup> channel genes. In: *CRC Handbook of Receptors and Channels*, P.A. North, ed., CRC Press, Boca Raton, FL, pp. 1–71.
- Charlier, C., Singh, N.A., Ryan, S.G., Lewis, T.B., Reus, B.E., Leach, R.J., and Leppert, M. (1998). A pore mutation in a novel KQT-like potassium channel gene in an idiopathic epilepsy family. *Nat. Genet.* 18, 53–55. [Abstract Article](#)
- Coburn, C.M., and Bargmann, C.I. (1996). A putative cyclic nucleotide-gated channel is required for sensory development and function in *C. elegans*. *Neuron* 17, 695–706. [Abstract Article](#)
- Connor, J.A., and Stevens, C.F. (1971). Voltage-clamp studies of a transient outward membrane current in gastropod neural somata. *J. Physiol.* 213, 21–30. [Abstract](#)
- Connor, J.A. (1980). The fast K<sup>+</sup> Channel and Repetitive Firing. In: *Cold Spring Harbor Reports in the Neurosciences*, Vol. 1, J. Koester and J. Byrne, eds., CSHL, pp. 125–133.
- Curran, M.E., Splawski, I., Timothy, K.W., Vincent, G.M., Green, E.D., and Keating, M.T. (1995). A molecular basis for cardiac arrhythmia: HERG mutations cause long QT syndrome. *Cell* 80, 795–803. [Abstract Article](#)
- Davies, A.G., Pierce-Shimomura, J.T., Kim, H., VaHoven, M.K., Thiele, T.R., Bonci, A., Bargmann, C.I., and McIntire, S.L. (2003). A central role of the BK potassium channel in behavioral responses to ethanol in *C. elegans*. *Cell* 115, 655–666. [Abstract Article](#)
- Davis, M.W., Fleischhauer, R., Dent, J.A., Joho, R.H., and Avery, L. (1999). A mutation in the *C. elegans* EXP-2 potassium channel that alters feeding behavior. *Science* 286, 2501–2504. [Abstract Article](#)
- De la Cruz, I.P., Levin, J.Z., Cummins, C., Anderson, P., and Horvitz, H.R. (2003). *Sup-9*, *sup-10* and *unc-93* may encode components of a two-pore K<sup>+</sup> channel that coordinates muscle contraction in *Caenorhabditis elegans*. *J. Neurosci.* 23, 9133–9145. [Abstract](#)
- Doyle, D.A., Morais Cabral, J., Pfuetzner, R.A., Kuo, A., Gulbis, J.M., Cohen, S.L., Chait, B.T., and MacKinnon, R. (1998). The structure of the potassium channel: molecular basis of K<sup>+</sup> conduction and selectivity. *Science* 280, 69–77. [Abstract Article](#)
- Doupnik, C.A., Davidson, N., and Lester, H.A. (1995). The inward rectifier potassium channel family. *Curr. Opin. Neurobiol.* 5, 268–277. [Abstract Article](#)
- Duprat, F., Lesage, F., Fink, M., Ryeyes, R., Heurteaux, C., and Lazdunski, M. (1997). TASK, a human background K<sup>+</sup> channel to sense external pH variations near physiological pH. *EMBO J.* 16, 5464–5471. [Abstract Article](#)

Elkes, D.A., Cardozo, D.L., Madison, J., and Kaplan, J.M. (1997). EGL-36 Shaw channels regulate *C. elegans* egg-laying muscle activity. *Neuron* 19, 165–174. [Abstract Article](#)

Elkins, T., Ganetzky, B., and Wu, C.F. (1986). A *Drosophila* mutation that eliminates a calcium-dependent potassium current. *PNAS* 83, 8415–8419. [Abstract](#)

Fleischhauer, R., Davis, W., Dzhura, I., Neely, A., Avery, L., and Joho, R. (2000). Ultra-fast inactivation causes inward rectification in a voltage-gated K<sup>+</sup> channel from *C.elegans*. *J. Neurosci.*

Garcia, L.R., and Sternberg, P.W. (2003). *Caenorhabditis elegans* UNC-103 ERG-like potassium channel regulates contractile behaviors of sex muscles in males before and during mating. *J. Neurosci.* 23, 2696–2705. [Abstract](#)

Goulding, E.H., Nagi, J., Kramer, R.H., Colicos, S., Axel, R., Siegelbaum, S.A., and Chess, A. (1992). Molecular cloning and single-channel properties of the cyclic nucleotide-gated channel from catfish olfactory neurons. *Neuron* 8, 45–58. [Abstract Article](#)

Heginbotham, L., Abramson, T., and MacKinnon, R. (1992). A functional connection between the pores of distantly related ion channels as revealed by mutant K<sup>+</sup> channels. *Science* 258, 1152–1155. [Abstract](#)

Heginbotham, L., Lu, Z., Abramson, T., and MacKinnon, R. (1994). Mutations in the K<sup>+</sup> channel signature sequence. *Biophys. J.* 66, 1061–1067. [Abstract](#)

Hosono, R., Juno, S., and Midsukami, M. (1985). Temperature-sensitive mutations causing reversible paralysis in *Caenorhabditis elegans*. *J. Exp. Zool.* 235, 409–421. [Abstract Article](#)

Inagaki, N., Gono, T., Clement, J.P., Wang, C.Z., Aguilar-Bryan, L., Bryan, J., and Seino, S. (1996). A family of sulfonylurea receptors determines the pharmacological properties of ATP-sensitive K<sup>+</sup> channels. *Neuron* 16, 1011–1017. [Abstract Article](#)

Jegla, T., and Salkoff, L. (1995). A multigene family of novel K<sup>+</sup> channels from *Paramecium tetraurelia*. *Receptors Channels* 3, 51–60. [Abstract](#)

Jiang, Y., Lee, A., Chen, J., Cadene, M., Chait, B.T., MacKinnon, R. (2002). Crystal structure and mechanism of a calcium-gated potassium channel. *Nature* 417, 515–522. [Abstract Article](#)

Johnstone, D., Wei, A., Butler, A., Salkoff, L., and Thomas, J. (1997). Behavioral defects in *C. elegans egl-36* mutants result from potassium channels shifted in voltage-dependence of activation. *Neuron* 19, 151–164. [Abstract Article](#)

Joiner, W.J., Tang, M.D., Wang, L.Y., Dworetzky, S.I., Boissard, C.G., Gan, L., Gribkoff, V.K., and Kaczmarek, L.K. (1998). Formation of intermediate-conductance calcium-activated potassium channels by interaction of SLACK and SLO subunits. *Nature Neurosci.* 1, 462–469. [Abstract Article](#)

Jospin, M., Mariol, M.C., Segalat, L., and Allard, B. (2002). Characterization of K(+) currents using an *in situ* patch clamp technique in body wall muscle cells from *Caenorhabditis elegans* 1. *J. Physiol.* 544(Pt 2), 373–384. [Abstract Article](#)

Kaletta, T., Van der Craen, M., Van Gell, A., Dewulf, N., Bogaret, T., Branden, M., King, K.V., Buechner, M., Barstead, R., Hyink, D., Li, H.P., Geng, L., Burrow, C., and Wilson, P. (2003). Towards understanding the polycystins. *Nephron. Exp. Nephrol.* 93, e9–e17. [Abstract Article](#)

Kamb, A., Iverson, L.E., and Tanouye, M.A. (1987). Molecular characterization of *Shaker*, a *Drosophila* gene that encodes a potassium channel. *Cell* 50, 405–413. [Abstract Article](#)

Kaupp, B.U., Niidome, T., Tanabe, T., Terada, S., Bonigk, W., Stühmer, W., Cook, N., Kangawa, K., Matsuo, H., Hirose, T., and Numa, S. (1989). Primary structure and functional expression from complementary DNA of the rod photoreceptor cyclic GMP-gated channel. *Nature* 342, 762–766. [Abstract Article](#)

Katz, B. (1966). *Nerve, Muscle, and Synapse*. McGraw-Hill, New York.

- Ketchum, K.A., Joiner, W.J., Sellers, A.J., Kaczmarek, L.K., and Goldstein, S.A.N. (1995). A new family of outwardly rectifying potassium channel proteins with two pore domains in tandem. *Nature* 376, 690–695. [Abstract Article](#)
- Ketchum, K.A., and Slayman, C.W. (1996). Isolation of an ion channel gene from *Arabidopsis thaliana* using the H5 signature sequence from voltage-dependent K<sup>+</sup> channels. *FEBS Lett.* 378, 19–26. [Abstract Article](#)
- Koh, S.D., Campbell, J.D., Carl, A., and Sanders, K.M. (1995). Nitric oxide activates multiple potassium channels in canine colonic smooth muscle. *J. Physiol.* 489, 735–43. [Abstract](#)
- Kohler, M., Hirschberg, B., Bond, C.T., Kinzie, J.M., Marrion, N.V., Maylie, J., and Adelman, J.P. (1996). Small-conductance, calcium-activated potassium channels from mammalian brain. *Science* 273, 1709–1714. [Abstract](#)
- Komatsu, H., Mori, I., Rhee, J.S., Akaike, N., and Ohshima, Y. (1996). Mutations in a cyclic nucleotide-gated channel lead to abnormal thermosensation and chemosensation in *C. elegans*. *Neuron* 17, 707–718. [Abstract Article](#)
- Kubisch, C., Schroeder, B.C., Friedrich, T., Lutjohann, B., El-Amraoui, A., Marlin, S., Petit, C., and Jentsch, T.J. (1999). KCNQ4, a novel potassium channel expressed in sensory outer hair cells, is mutated in dominant deafness. *Cell* 96, 437–446. [Abstract Article](#)
- Kunkel, M.T., Johnstone, D., Thomas, J.H., and Salkoff, L. (2000). Mutants of a temperature-sensitive 2-P domain potassium channel. *J. Neurosci.* 20, 7517–7524. [Abstract](#)
- Lehmann-Horn, F., and Jurkat-Rott, K. (1999). Voltage-gated ion channels and hereditary disease. *Physiol. Rev.* 79, 1317–1372. [Abstract](#)
- Levin, J.Z., and Horvitz, H.R. (1992). The *Caenorhabditis elegans unc-93* gene encodes a putative transmembrane protein that regulates muscle contraction. *J. Cell Biol.* 117, 143–155. [Article](#)
- Levin and Horvitz (1993). Three New Classes of Mutations in the *Caenorhabditis elegans* Muscle Gene *sup-9*. *Genetics* 135, 53–70. [Abstract](#)
- Littleton, J.T., and Ganetzky, B. (2000). Ion channels and synaptic organization: analysis of the *Drosophila* genome. *Neuron* 26, 35–43. [Abstract Article](#)
- MacKinnon, R., and Doyle, D.A. (1997). Prokaryotes offer hope for potassium channel structural studies. *Nat. Struct. Biol.* 4, 877–879. [Abstract Article](#)
- Milkman, R. (1994). An *Escherichia coli* homologue of eukaryotic potassium channel proteins. *PNAS* 91, 3510–3514. [Abstract](#)
- Papazian, D.M., Schwarz, T.L., Tempel, B.L., Jan, Y.N., and Jan, L.Y. (1987). Cloning of genomic and complementary DNA from Shaker, a putative potassium channel gene from *Drosophila*. *Science* 237, 749–753. [Abstract](#)
- Pearson, W.L., Dourado, M., Schreiber, M., Salkoff, L., and Nichols, C.G. (1999). Expression of a functional Kir4 family inward rectifier K<sup>+</sup> channel from a gene cloned from mouse liver. *J. Physiol.* 514, 639–653. [Abstract Article](#)
- Petersen, C.I., McFarland, T.R., Stepanovic, S.Z., Yang, P., Reiner, D.J., Hayashi, K., George, A.L., Roden, D.M., Thomas, J.H., and Balsler, J.R. (2004). *In vivo* identification of genes that modify ether-a-go-go-related gene activity in *Caenorhabditis elegans* may also affect human cardiac arrhythmia. *Proc. Natl. Acad. Sci. USA* 101, 11773–11778. [Abstract Article](#)
- Pongs, O., Kecskemethy, N., Muller, R., Krah-Jentgens, I., Baumann, A., Kiltz, H.H., Canal, I., Llamazares, S., and Ferrus, A. (1988). Shaker encodes a family of putative potassium channel proteins in the nervous system of *Drosophila*. *EMBO J.* 7, 1087–1096. [Abstract](#)
- Raizen, D.M., and Avery, L. (1994). Electrical activity and behavior in the pharynx of *Caenorhabditis elegans*. *Neuron* 12, 483–495. [Abstract Article](#)

- Richmond, J.E., and Jorgensen, E.M. (1999). One GABA and two acetylcholine receptors function at the *C. elegans* neuromuscular junction. *Nature Neurosci.* 2, 791–797. [Abstract Article](#)
- Reiner, D.J., Weinschenker, D., and Thomas, J.H. (1995). Analysis of dominant mutations affecting muscle excitation in *Caenorhabditis elegans*. *Genetics* 141, 961–976. [Abstract](#)
- Reiner, D.J., Newton, E.M., Tian, H., and Thomas, J.H. (1999). Diverse behavioral defects caused by mutations in *Caenorhabditis elegans unc-43* CaM kinase II. *Nature* 402, 199–203. [Abstract Article](#)
- Rubin, G.M., Yandell, M.D., Wortman, J.R., Gabor Miklos, G.L., Nelson, C.R., Hariharan, I.K., Fortini, M.E., Li, P.W., Apweiler, R., Fleischmann, W., Cherry, J.M., Henikoff, S., Skupski, M.P., Misra, S., Ashburner, M., Birney, E., Boguski, M.S., Brody, T., Brokstein, P., Celniker, S.E., Chervitz, S.A., Coates, D., Cravchik, A., Gabrielian, A., Galle, R.F., Gelbart, W.M., George, R.A., Goldstein, L.S., Gong, F., Guan, P., Harris, N.L., Hay, B.A., Hoskins, R.A., Li, J., Li, Z., Hynes, R.O., Jones, S.J., Kuehl, P.M., Lemaitre, B., Littleton, J.T., Morrison, D.K., Mungall, C., O'Farrell, P.H., Pickeral, O.K., Shue, C., Vosshall, L.B., Zhang, J., Zhao, Q., Zheng, X.H., and Lewis, S. (2000). Comparative genomics of the eukaryotes. *Science* 287, 2204–2215. [Abstract Article](#)
- Saimi, Y., Loukin, S.H., Zhou, X.L., Martinac, B., and Kung, C. (1999). Ion channels in microbes. *Methods Enzymol.* 294, 507–524.
- Salkoff, L., Kunkel, M., Wang, Z., Butler, A., Nonet, M., and Wei, A. (1998). The impact of the *C. elegans* genome sequencing project on K<sup>+</sup> channel biology. In: *Function Potassium Channels: Molecular Structure, and Diseases. Current Topics in Membranes*, Academic Press, New York, Chapter 46, pp. 9–27.
- Salkoff, L., Butler, A., Fawcett, G., Kunkel, M., McArdle, C., Paz-y-Mino, G., Walton, N., Yuan, A., Wang, Z.W., and Wei, A. (2001). Evolution tunes the excitability of individual neurons. *Neuroscience* 103(4), 853–859. [Abstract Article](#)
- Sanguinetti, M.C., Jiang, C., Curran, M.E., and Keating, M.T. (1995). A mechanistic link between an inherited and an acquired cardiac arrhythmia: HERG encodes the IKr potassium channel. *Cell* 81, 299–307. [Abstract Article](#)
- Santi, C.M., Yuan, A., Fawcett, G., Wang, Z.W., Butler, A., Nonet, M.L., Wei, A., Rojas, P., and Salkoff, L. (2003). Dissection of K<sup>+</sup> currents in *Caenorhabditis elegans* muscle cells by genetics and RNA interference. *Proc. Natl. Acad. Sci. USA* 100, 14391–14396. [Abstract Article](#)
- Sentenac, H., Bonneaud, N., Minet, M., Lacroux, F., Salmon, J.M., Gaymard, F., and Grignon, C. (1992). Cloning and expression in yeast of a plant potassium ion transport system. *Science* 256, 663–665. [Abstract](#)
- Shtonda, B., and Avery, L. (2005). CCA-1, EGL-19 and EXP-2 currents shape action potentials in the *Caenorhabditis elegans* pharynx. *J. Exp. Biol.* 208, 2177–2190. [Abstract Article](#)
- Shyng, S., and Nichols, C.G. (1997). Octameric stoichiometry of the KATP channel complex. *J. Gen. Physiol.* 110, 655–664. [Abstract Article](#)
- Silverman, S.K., Kofuji, P., Dougherty, D.A., Davidson, N., and Lester, H.A. (1996). A regenerative link in the ionic fluxes through the weaver potassium channel underlies the pathophysiology of the mutation. *Proc. Natl. Acad. Sci. USA* 93, 15429–15434. [Abstract Article](#)
- Singh, N.A., Charlier, C., Stauffer, D., DuPont, B.R., Leach, R.J., Melis, R., Ronen, G.M., Bjerre, I., Quattlebaum, T., Murphy, J.V., et al. (1998). A novel potassium channel gene, KCNQ2, is mutated in an inherited epilepsy of newborns. *Nat. Genet.* 18, 25–29. [Abstract Article](#)
- Singh, N.A., Westenskow, P., Charlier, C., Pappas, C., Leslie, J., Dillon, J., Anderson, V.E., Sanguinetti, M.C., and Leppert, M.F. (2003). KCNQ2 and KCNQ3 potassium channel genes in benign familial neonatal convulsions: expansion of the functional and mutation spectrum. *Brain* 126, 2726–2737. [Abstract Article](#)
- Smith, P.L., Baukrowitz, T., and Yellen, G. (1996). The inward rectification mechanism of the HERG cardiac potassium channel. *Nature* 379, 833–836. [Abstract Article](#)



- Thomas, J.H. (1990). Genetic analysis of defecation in *Caenorhabditis elegans*. *Genetics* 124, 855–872. [Abstract](#)
- Trent, C., Tsuing, N., and Horvits, H.R. (1983). Egg-laying defective mutants of the nematode *Caenorhabditis elegans*. *Genetics* 104, 619–647. [Abstract](#)
- Trudeau, M.C., Warmke, J.W., Ganetzky, B., and Robertson, G.A. (1995). HERG, a human inward rectifier in the voltage-gated potassium channel family. *Science* 269, 92–95. [Abstract](#)
- Wang, Z.W., Saifee, O., Nonet, M.L., and Salkoff, L. (2001). Slo-1 Potassium channels control quantal content of neurotransmitter release at the *C. elegans* neuromuscular junction. *Neuron* 32, 867–881. [Abstract Article](#)
- Warmke, J., Drysdale, R., and Ganetzky, B. (1991). A distinct potassium channel polypeptide encoded by the *Drosophila eag* locus. *Science* 252, 1560–1562. [Abstract](#)
- Warmke, J., and Ganetzky, B. (1994). A family of potassium channel genes related to *eag* in *Drosophila* and mammals. *PNAS* 91, 3438–3442. [Abstract](#)
- Wei, A., Butler, A., and Salkoff, L. (2005). KCNQ-like potassium channels in *Caenorhabditis elegans*: conserved properties and modulation. *J. Biol. Chem.* 280, 21337–21345. [Abstract Article](#)
- Wei, A., Jegla, T., and Salkoff, L. (1996). Eight potassium channel families revealed by the *C. elegans* genome project. *Neuropharmacology* 35, 805–829. [Abstract Article](#)
- Wei, A., Yuan, A., Fawcett, G., Butler, A., Davis, T., Xu, S.Y., and Salkoff, L. (2002). Efficient isolation of targeted *Caenorhabditis elegans* deletion strains using highly thermostable restriction endonucleases and PCR. *Nucl. Acids. Res.* 30, e110–e117. [Abstract Article](#)
- Weinshenker, D., Wei, A., Salkoff, L., and Thomas, J. (1999). Block of an ether-a-go-go-like K<sup>+</sup> channel by imipramine rescues *egl-2* excitation defects in *Caenorhabditis elegans*. *J. Neurosci.* 19, 9831–9840. [Abstract](#)
- Yuan, A., Dourado, M., Butler, A., Blanton, N., Wei, A., and Salkoff, L. (2000). Slo-2, A K<sup>+</sup> channel with a unique C1- dependence. *Nature Neurosci.* 3, 771–779. [Abstract Article](#)
- Yuan, A., Santi, C.M., Wei, A., Wang, Z.W., Pollak, K., Nonet, M., Kaczmarek, L., Crowder, C.M., and Salkoff, L. (2003). The sodium-activated potassium channel is encoded by a member of the *Slo* gene family. *Neuron* 37, 765–763. [Abstract Article](#)
- Zhu, X.L., Vaillant, B., Loukin, S.H., Kung, C., and Saimi, Y. (1995). YKC1 encodes the depolarization-activated K<sup>+</sup> channel in the plasma membrane of yeast. *FEBS Lett.* 373, 170–176. [Abstract Article](#)

