

2014

Chronic lung allograft dysfunction following lung transplantation: challenges and solutions

Bradford C. Bemiss

Washington University School of Medicine in St. Louis

Chad A. Witt

Washington University School of Medicine in St. Louis

Follow this and additional works at: http://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Bemiss, Bradford C. and Witt, Chad A., "Chronic lung allograft dysfunction following lung transplantation: challenges and solutions." *Transplant Research and Risk Management*. 2014, 6. 87-97. (2014).
http://digitalcommons.wustl.edu/open_access_pubs/3636

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Chronic lung allograft dysfunction following lung transplantation: challenges and solutions

Bradford C Bemiss
Chad A Witt

Department of Internal Medicine,
Division of Pulmonary and Critical
Care Medicine, Washington University
School of Medicine, St Louis, MO, USA

Abstract: Chronic rejection is a major cause of death after the first year following lung transplantation. Bronchiolitis obliterans (BO) is the most common pathologic finding on biopsy, characterized by fibrous granulation tissue, which obliterates the lumen of the bronchiole. Clinically, in the absence of tissue for pathology, BO syndrome refers to a progressive irreversible drop in the forced expiratory volume in 1 second. Recently, a broader definition of chronic rejection, termed “chronic lung allograft dysfunction”, has been used to encompass a more inclusive definition of posttransplant dysfunction. Recently, the lung transplant community has come to realize that chronic rejection may be the final common result after repetitive epithelial insults. Acute rejection, infection, and alloreactivity to mismatched HLA antigens are a few of these insults that damage the surface of the bronchioles. Recent evidence of autoimmunity to the normally hidden structural proteins collagen V and K- α 1 tubulin have been correlated with a BO phenotype as well, perhaps correlating the epithelial damage with a mechanism for developing BO lesions. Many immunomodulatory medications and treatments have been studied for effectiveness for the treatment of chronic lung allograft dysfunction. New drugs, which more precisely target the immune system, are being developed and tested. Further study is required, but recent advances have improved our understanding of the pathogenesis and potential intervention for this common and deadly complication of lung transplantation.

Keywords: lung transplant, bronchiolitis obliterans syndrome, rejection

Introduction

Dr Bruce Reitz performed the first successful heart–lung transplants at Stanford University (Stanford, CA, USA), in 1981, for idiopathic pulmonary hypertension and congenital heart disease.¹ Following this achievement, the first single-lung and double-lung transplant were performed by Dr Joel Cooper in Toronto, ON, Canada, in 1983 and 1986, respectively.² Beginning in the mid-1990s, the number of lung transplants has increased yearly to a total of 3,640 patients in 2011. From 1994 through 2011, the median survival of lung transplant recipients was 5.6 years. Improved survival early after lung transplant has resulted in better survival in the most recent era of lung transplantation. However, the slope of overall mortality after the first year during the most recent era mirrors that of previous eras. Following the first year, the most common causes of death are chronic rejection and non-Cytomegalovirus (CMV) infection.³

Chronic lung transplant rejection has classically been described as obliterative bronchiolitis (also called bronchiolitis obliterans [BO]). This was first described at Stanford in 1984. Of the 19 heart–lung transplants performed for end-stage pulmonary vascular disease, 14 patients were long-term survivors. Five of these patients demonstrated a

Correspondence: Chad A Witt
Department of Internal Medicine,
Division of Pulmonary and Critical Care
Medicine, Washington University School
of Medicine, 660 S Euclid Ave Campus
Box 8052, St Louis, MO 63110, USA
Tel +1 314 454 8762
Fax +1 314 454 7524
Email cwitt@dom.wustl.edu

typical initial restrictive ventilatory defect on spirometry, which improved over the first few months. However, this was followed by the development of a progressive obstructive ventilatory defect with a decline in the forced expiratory volume in 1 second (FEV_1). On biopsy, these patients had findings consistent with BO.⁴

BO develops in a nonuniform pattern and is difficult to sample bronchoscopically by transbronchial biopsy. Therefore, the clinical diagnosis of BO syndrome (BOS) is used to describe a progressive obstructive ventilatory defect without evidence of another explanation (acute rejection, infection, airway complication, or other cause). Recently, a paradigm shift has occurred in the thinking about chronic lung transplant rejection. Namely, BOS is now thought of as a large, but specific, form of chronic rejection. The term “chronic lung allograft dysfunction” (CLAD) has been introduced to include specific forms of allograft dysfunction (anastomotic stricture, azithromycin-responsive allograft dysfunction, disease recurrence, etc) in addition to a clinical description of both obstructive CLAD (BOS) and restrictive allograft syndrome (RAS) (Figure 1).⁵

Throughout this article, we will review the challenges inherent to the management of CLAD from diagnosis to

understanding the pathogenesis and reducing risk factors. In addition, we will examine the available treatments for CLAD and their efficacy. This review will form a basic framework for understanding CLAD and discuss potential future solutions for reducing chronic rejection through new, more successful treatments.

BOS

BOS is the most common form of CLAD, accounting for approximately 70% of cases.⁶ Pathologically, early BO is characterized by lymphocytic inflammation in the submucosa of respiratory bronchioles. This leads to late proliferation of dense fibromyxoid granulation tissue, which organizes to partially or completely fill the lumen of the airway (Figure 2).^{7,8} On biopsy interpretation, these findings are denoted as “C1”. These fibrotic lesions are thought to be the histologic result of repetitive epithelial injury by multiple different insults. However, obtaining sufficient histological specimens by transbronchial biopsy documenting these findings is unreliable.⁹

In the absence of histopathology from a biopsy specimen, BOS is more typically diagnosed on routine spirometry. In 1993, the International Society for Heart and Lung Transplantation (ISHLT) proposed criteria for the clinical diagnosis of BOS based on spirometry (Table 1). Measurements are made of post-transplant spirometry. When patients reach a baseline maximal FEV_1 (average of the two best measurements taken at least 3 weeks apart without administration of bronchodilators), this is termed “BOS Stage 0”. Progressive decline in FEV_1 is staged as BOS 1 through 3, with a higher stage indicating a worsened obstruction. This schema was updated in 2001 to reflect new findings that a drop in FEV_1 and forced expiratory flow (FEF) 25%–75% is a more sensitive marker of early obstruction.^{9,10} The new BOS stage 0p is defined as a decrease in the FEV_1 to <90% and a decrease in FEF 25%–75% to $\leq 75\%$ of the posttransplant baseline. It should be noted that, in children, by convention, the establishment of a reference value of lung function and subsequent decline are expressed in terms of percent predicted FEV_1 .

Patients may present with nonproductive cough or dyspnea on exertion early in the course of BOS. However, patients frequently have few complaints, but an asymptomatic decline in their FEV_1 is discovered. In the more advanced stages of BOS, symptoms may progress to dyspnea at rest and productive cough with evidence of severe obstruction on spirometry. In an evaluation of health-related quality of life, Gerbase et al prospectively evaluated 58 patients and reported a decrease in the St George’s Respiratory Questionnaire




Figure 1 CLAD Phenotypes.

Notes: Chronic lung allograft dysfunction (CLAD) is a descriptive term for chronic lung transplant rejection which encompasses multiple different phenotypes, including bronchiolitis obliterans syndrome (BOS), restrictive allograft syndrome (RAS), recurrence of primary disease, anastomotic stricture, and azithromycin-responsive allograft dysfunction (ARAD), as well as other specific causes of decline in lung function.




Figure 2 Lung biopsy showing bronchiolitis obliterans in the setting of chronic lung transplant rejection.

Notes: Hematoxylin and eosin stain was used.

(a standardized questionnaire used to measure health and perception of well-being¹¹) among patients who developed BOS.¹² BOS is a major cause of both morbidity and mortality among lung transplant recipients. The most recent ISHLT data show that, 5 years after transplant, 49% of lung transplant recipients develop BOS, increasing to 76% at 10 years (Figure 3).³ One retrospective analysis of actuarial survival placed the 3-year mortality after the onset of BOS at 51%.¹³ Another study, from Denmark, showed a threefold increased risk of death after progression from BOS Stage 1 to Stage 2 and from BOS Stage 2 to Stage 3.¹⁴

When a lung transplant recipient is found to have a drop in their FEV₁, this should be investigated to determine if another cause is responsible. Typically, this includes chest radiography and bronchoscopy with bronchoalveolar lavage

Table 1 BOS classification

Stage	1994 definition	2001 definition
BOS 0	FEV ₁ ≥80% of baseline	FEV ₁ >90% of baseline FEF 25%–75% >75% of baseline
BOS 0p		FEV ₁ 81%–90% of baseline FEF 25%–75% ≤75% of baseline
BOS 1	FEV ₁ 66%–80% of baseline	FEV ₁ 66%–80% of baseline
BOS 2	FEV ₁ 51%–65% of baseline	FEV ₁ 51%–65% of baseline
BOS 3	FEV ₁ ≤50% of baseline	FEV ₁ ≤50% of baseline

Abbreviations: BOS, bronchiolitis obliterans syndrome; FEF, forced expiratory flow; FEV₁, forced expiratory volume in 1 second.

(BAL) and transbronchial biopsies. The chest X-ray is not a useful tool for screening for early BOS, but is useful in establishing a differential diagnosis for the change in spirometry. Expiratory chest computed tomography reliably demonstrates air trapping and mosaic attenuation in patients with BOS.^{15,16} However, when using the revised criteria to include BOS Stage 0p, which is more effective at early detection of BOS, thin-slice computed tomography was not helpful in making the diagnosis.¹⁷ Bronchoscopy should be performed to rule out airway abnormalities and infectious causes of declining FEV₁. Aside from assessing




Figure 3 Adult lung transplant freedom from bronchiolitis obliterans syndrome conditional on survival to 14 days (April 1994–June 2012).

Notes: Reprinted from *J Heart Lung Transplant*, 32(10), Yusen RD, Christie JD, Edwards LB, et al; International Society for Heart and Lung Transplantation. The Registry of the International Society for Heart and Lung Transplantation: Thirtieth Adult Lung And Heart-Lung Transplant Report – 2013; focus theme: age, 965–978, Copyright 2013, with permission from Elsevier.³

infection, the BAL fluid has been studied to determine its utility in predicting onset of BOS. In a retrospective study, BAL neutrophilia $\geq 20\%$ was a significant predictor for BOS Stage 1 or greater in a multivariate regression model. Neurohr et al also demonstrated increased IL-8 levels and reduced secretory leukocyte protease inhibitor (SLPI) within the BAL fluid. These markers were present independent of airway bacterial colonization.¹⁸ Other studies have demonstrated elevated levels of other cytokines and inflammatory proteins, such as MPO, IL-12, and IL-17; however, it is unclear what predictive value these or the presence of airway neutrophilia have in the diagnosis of BOS due to multiple possible confounding factors.^{19–21}

RAS

While BOS is the most common form of CLAD, increasing recognition of a restrictive form of chronic rejection has been reported by several centers. This restrictive ventilatory defect has been termed “RAS”. The initial report by Sato et al, a retrospective review, reports their experience over a 13-year period. Patients were screened for CLAD by routine spirometry and diagnosed when their FEV₁ declined to $< 80\%$ of previous baseline. RAS was diagnosed when a follow-up total lung capacity (TLC) measurement declined to $< 90\%$ of baseline. Of the 156 episodes of CLAD diagnosed, 47% were characterized as RAS. Patients with RAS had a higher incidence of CMV mismatch but were otherwise similar regarding donor age, recipient age, primary diagnosis, and sex. Most notably, Sato et al discovered that patients with RAS had a median survival that was less than half that of BOS (541 days versus 1,421 days).⁶ This definition of RAS was limited to bilateral lung transplant recipients only.

These investigators evaluated the progression of RAS in a separate study. During this retrospective review, they found that RAS, unlike BOS, was characterized by sporadic acute exacerbations and a stepwise decline in FEV₁ over time. Patients primarily presented with symptoms of shortness of breath, fever, and cough during an acute exacerbation. The patients they studied had between one and four exacerbations, after which they did not recover to their baseline FEV₁. Their outcomes included 32% of patients that had a partial recovery, 27% that had stable disease, and 41% that had progressive disease. Radiologic findings during acute exacerbation were principally characterized by asymmetric bilateral ground-glass opacities. However, during the period between acute exacerbations, interstitial reticular shadows, consolidation, and traction bronchiectasis were more often present.²²

While definitive pathologic findings have not been thoroughly described, initial work from Toronto has demonstrated consistent findings within the cohort of RAS patients described previously. Of the 47 patients diagnosed with RAS, 16 had histopathologic specimens available for examination. Fifteen of these sixteen had findings of parenchymal fibroelastosis in a subpleural distribution. There were often coincident findings of BO and diffuse alveolar damage.²³ Diffuse alveolar damage was the predominant histopathologic finding within the acute exacerbation group described previously.²² As a newly described phenomena, the characterization of RAS continues to evolve, but it undoubtedly represents a significant and potentially severe posttransplant complication.

Risk factors for the development of CLAD

Initially, after BOS was recognized to be a major contributor to posttransplant mortality, work began to determine what factors increased the risk of developing this progressive and, in many cases, untreatable complication. In 1998, a report from Heng et al did not show any relation to sex, age, primary disease, or graft ischemic time. Early acute rejection, CMV status, pulmonary infection, organizing pneumonia, and human leukocyte antigen (HLA) mismatch seemed to be significantly associated with onset of BOS.¹³ A common thread among these risk factors may be the immune system's exposure to injured pulmonary epithelium. It has been hypothesized that this injury response via alloimmune and autoimmune mechanisms could play a major role in the development of BOS. Further delineation of the major risk factors for BOS, proposed underlying mechanisms of injury, and the immune system response will be reviewed here.

Acute cellular rejection

An early retrospective investigation by Bando et al identified, by multivariate analysis, that three or more episodes of acute cellular rejection graded mild to moderate (grade II to III, or A2 to A3) or greater significantly increased the risk of developing BOS.²⁴ Kroshus et al also showed that the cumulative number of episodes of acute rejection after 90 days increased the relative risk of BOS.²⁵ While these analyses showed that moderate-to-severe rejection and multiple episodes of rejection were risk factors for BOS, Hachem et al²⁶ found that even a single episode of early or late minimal (A1) acute cellular rejection was associated with a hazard ratio (HR) of 2.64 for the development of BOS 1 stage. Khalifah et al showed that

a single episode of A2 was associated with more frequent progressive BOS.²⁶ Recent work from San Segundo et al has demonstrated that an increased number of peripherally circulating CD8+ effector T-cells, both pre-transplant and during the first year after lung transplant, is associated with a higher risk of developing acute rejection.²⁸ Additionally, T-cell activity, in the form of granzyme B production and other cytokine profiles, has been shown to be elevated in lung recipients with acute rejection and prior to or at onset of BOS, leading to the concept that physiologic measurements and assessments of T-cell function may be a better way of monitoring the level of immunosuppression in lung recipients than the current method of adjusting medications based on serum drug levels.^{29,30}

Lymphocytic bronchiolitis

In 1992, Yousem described submucosal lymphocytic and plasma cell infiltration around the airways and deep to the smooth muscle layer, which was called lymphocytic bronchitis and bronchiolitis (LBB, or B-grade) (Figure 4). This lymphocytic infiltrate was present more commonly in patients who later went on to develop BO.³¹ Husain et al showed that LBB was, similarly to acute perivascular rejection, associated with onset of BOS. At 1 year, patients who went on to develop BOS had over twice as many episodes of LBB as did those without BOS.³²

CMV infection

CMV infection is a common infectious complication in the posttransplant period in patients with both reactivated or donor-derived CMV infection. An initial study by Keenan et al, from Pittsburgh, PA, USA, associated BOS with pre-transplant seropositive status, seropositive donor status, and posttransplant CMV infection.³³ Further work by Smith et al found that CMV mismatch (donor positive/recipient negative) may be a risk factor for developing BOS within 3 years of transplantation.³⁴ However, other groups have found diverging data regarding the risk of BOS based on CMV serostatus and infection.^{35–37} Tamm et al reviewed their center's experience with CMV pneumonia and CMV serostatus, finding no significant differences in development of BOS, and suggested that treatment of CMV pneumonia with intravenous ganciclovir is protective.³⁸ When ganciclovir is ineffective, such as in cases of ganciclovir-resistant CMV, an earlier onset of BOS has been reported.³⁹

Non-CMV infections

While CMV pneumonitis is perhaps the most consistently implicated infection in the development of BOS, multiple studies have evaluated whether other infections are significant risk factors. An early study by Hohlfield et al found that 47% of the onset of BOS seemed to cluster in the first quarter of the year, January to March, suggesting a possible relationship




Figure 4 Lung biopsy showing lymphocytic bronchiolitis in the setting of acute lung transplant rejection.

Notes: Hematoxylin and eosin stain was used.

with viral infections.⁴⁰ Community-acquired respiratory viruses ([CARVs] including rhinovirus, coronavirus, respiratory syncytial virus, parainfluenza, human metapneumovirus, influenza A and B, and adenovirus) have been implicated in the development of BOS.^{41,42} Kumar et al⁴² reported that of nine of 50 patients with a decline in FEV₁ after CARV infection, six had persistent disease classified as BOS. In a retrospective review by Khalifah et al, 259 consecutive lung transplant patients were found to have 21 episodes of CARV infection, which were associated with an HR of 2.05 for development of BOS Stage 1, and lower respiratory tract involvement increased the HR to 3.03.⁴³ This was verified prospectively in a mouse model using orthotopic tracheal transplant.⁴⁴ In one series, 0 of 18 patients developed BOS after treatment with intravenous ribavirin, suggesting that treatment of CARV could prevent the onset of chronic rejection.⁴⁵ Other viruses, such as Epstein–Barr virus and human herpes virus-6, have also been implicated in the development of BOS, though with less supporting evidence.^{46,47}

In addition to viral infection, bacterial and fungal infections predispose patients to increased rates of BOS as well. Valentine et al published their experience, finding that Gram-negative, Gram-positive, and fungal pneumonias all significantly increased the risk for BOS. In fact, compared with uninfected patients, those with bacterial pneumonia and fungal pneumonia developed BOS 6 months and 2 years earlier, respectively.⁴⁸ Further, not just infection, but colonization of the airways seems to be a significant risk factor for the development of BOS. Botha et al found that *Pseudomonas aeruginosa* colonization of the airways in lung transplant recipients increased the risk of BOS at 2 years from 7.7% to 23.4%.⁴⁹ Similarly, Weigt et al examined the effect of *Aspergillus* spp. colonization on the incidence of BOS. Using a multivariate analysis, *Aspergillus* colonization was independently associated with onset and mortality due to BOS.⁵⁰

Primary graft dysfunction (PGD)

PGD is a common early complication after lung transplantation manifested by infiltrates that can be seen on chest radiograph and impairment of oxygenation. Whitson et al showed that PGD grade 3 (bilateral infiltrates and PaO₂/FiO₂ <200 mmHg) was associated with BOS and a consistently lower FEV₁ at all time points in bilateral lung transplant recipients.⁵¹ In a retrospective review, Daud et al found that PGD was significantly associated with onset of BOS independent of acute rejection, lymphocytic bronchiolitis, and CARV. With increasing grade of PGD, there was a progressive

increase in relative risk for development of BOS (PGD grade 1=1.68, grade 2=2.04, grade 3=2.61).⁵²

Gastroesophageal reflux (GER)

D'Ovidio et al investigated the hypothesis that GER could be a contributing factor for development of BOS. Using esophageal pH <4, they showed that patients with abnormal pH had an increased rate of BOS. In addition, they correlated presence of bile acid and subsequent decreased levels of surfactant protein A in the BAL fluid.⁵³ Blondeau et al⁵⁴ found that both pepsin and bile acid were routinely found in the BAL of lung transplant recipients. However, they did not find a correlation with GER, but instead an association with both acid and non-acid reflux, suggesting this could be the true risk factor for BOS. In a single center study, Cantu et al described a strategy for surgical treatment, via fundoplication, for lung transplant patients. At their center, all transplant candidates are screened with esophageal pH monitoring and surgery is performed posttransplant for any patient with evidence of reflux and without a contraindication. Their retrospective review evaluated the results of these efforts over time. Of 202 patients screened for reflux, 85 had either early (average time after transplant =43 days) or late (average time after transplant =684 days) surgical treatment aimed at control of reflux. Freedom from BOS at 1 year was observed in 100% of those with early surgery (n=14), compared with 91% of those without reflux (n=180) and 92% of those in the late surgery group (n=62). At 3 years, that difference was greater: freedom from BOS was 100% in the early surgery group (n=5), compared with 62% in those without reflux (n=93) and 47% in the late surgery group (n=30).⁵⁵ The benefit of surgical correction of GER in asymptomatic lung transplant recipients is an ongoing area of investigation; there remains no consensus between transplant centers on the use of fundoplication for GER in lung recipients. The preliminary data from the multicenter Reflux Surgery in Lung Transplantation (RESULT) study was presented at the ISHLT meeting in April 2014 and, at that time, did not reveal any significant benefit to fundoplication. The final report of this study is pending at this time.⁵⁶

HLA mismatching

Given the hypothesis of alloimmune activation as a primary cause for CLAD, HLA mismatch between donor and recipient has been extensively studied. Previous studies in heart and kidney transplantation revealed that HLA mismatch was a significant risk factor for graft loss.^{57,58} Single-center studies revealed an association between both HLA class I (HLA-A)

and class II (HLA-DR) mismatches and development of BOS.^{59,60} A large, United Network for Organ Sharing (UNOS)/ISHLT database study revealed a significant increase in the risk of acute rejection and graft failure associated with HLA-A and HLA-DR, but, due to limited follow-up, did not find a significant increase in the rate of BOS.⁵⁸ Many other contradictory studies, which make this risk less clear, have also been published.^{13,35,60}

The evidence is more consistent in the study of anti-HLA antibodies that develop posttransplant. Sundaresan et al showed this was a much stronger signal for development of BOS than HLA-A mismatch at the time of transplantation.⁵⁹ These results have been duplicated as well.⁶¹ Improvement in the sensitivity of diagnostic techniques has allowed for better detection of anti-HLA antibodies and increased ability to correlate those results with BOS.⁶² Previous work by Jaramillo et al has demonstrated that anti-HLA class I antibodies trigger the activation and proliferation of airway epithelial cells as well as production of growth factors, fibroblast proliferation, and apoptosis.⁶³

Further work by Hachem et al examined the development of donor-specific anti-HLA antibodies (DSA) in a single-center prospective trial. Of the 116 patients examined, 56% developed DSA (15% class I, 63% class II, 22% both classes). This study was designed to examine the effect of treatment for DSA with either intravenous immunoglobulin (IVIG) or IVIG plus rituximab on the development of antibody-mediated acute and chronic rejection. They found that, irrespective of the treatment received, patients that cleared their DSA after treatment had a lower incidence of BOS and prolonged survival compared to those with persistent DSA.⁶⁴ Newer modalities of therapy for antibody-mediated rejection and depletion of anti-HLA antibodies have been studied more extensively in kidney transplantation, including bortezomib, a 26S proteasome inhibitor that can block NF- κ B signaling within mature plasma cells.⁶⁵

Non-HLA autoimmune response

In addition to the alloimmune response to donor HLA, increased risk of BOS due to a similar autoimmune response has been observed in non-HLA antibodies such as collagen V and K- α 1 tubulin.^{66,67} Specifically, these anti-airway epithelial cell antibodies have been found in up to 31% of lung transplant recipients with BOS. Downstream signaling via increased calcium influx leads to cell proliferation and increased transcription of fibrogenic growth factors, similar to the anti-HLA response.⁶⁸ This autoimmune activation seems to be mediated by a Th17 response.⁶⁹ Other labs have

also shown that a failure of maturation of regulatory T cells may contribute to the autoimmune reaction seen in chronic lung transplant rejection.⁷⁰

Treatment

Typical immunosuppression for lung transplant recipients includes a calcineurin inhibitor, a purine synthesis inhibitor, and a corticosteroid. Based on the major risk factors for CLAD identified to date, it is clear that immunosuppression plays a major role in its prevention. Aside from ensuring adequate dosing and blood levels of immunosuppressive drugs, a switch between these drugs has been shown to potentially treat BOS. In one of the first trials to assess the change in pulmonary function when switching from cyclosporine to tacrolimus, Cairn et al showed that, in patients with BOS, the rate of decline in FEV₁ and FEF 25%–75% decreased significantly in the 70% of patients that responded.⁷¹ Hachem et al conducted a randomized controlled trial to evaluate the outcomes after treating patients with one of the calcineurin inhibitors, either cyclophosphamide or tacrolimus. There was a nonstatistically significant decline in the occurrence of BOS in the tacrolimus group compared to the cyclosporine group.⁷² Similar stabilization of spirometry has also been shown when switching from azathioprine to mycophenolate mofetil.⁷³ Other immunosuppressive agents, such as methotrexate and cyclophosphamide, have also been studied in small case series with successful stabilization of FEV₁.^{74,75} However, given their potentially severe side effects, they have not gained wide acceptance.

Aerosolized cyclosporine and tacrolimus have also been studied to try to maximize drug delivery to the lungs while minimizing toxicity.^{76–78} Initially, cyclosporine was studied as a means of reducing acute rejection; however, Iacono et al noted increased chronic rejection-free survival instead.⁷⁸ The follow-up study by Groves et al, which randomized patients to inhaled cyclosporine versus inhaled placebo, determined that, in addition to standard immunosuppression, inhaled cyclosporine yielded improved FEV₁ and FEF 25%–75%.⁷⁶

The macrolide antibiotic azithromycin has been extensively studied for the treatment and prevention of CLAD. An initial pilot study of six patients demonstrated that, after initiation of azithromycin, five of six had improvement in FEV₁ of approximately 17% of predicted values.⁷⁹ Yates et al used azithromycin to treat BOS in 20 patients, resulting in a mean improvement in FEV₁ of 110 mL at 3 months. This improvement was maintained in 12 of the 17 patients that showed initial improvement.⁸⁰ Further evidence suggests that a survival advantage is obtained when starting azithromycin

during BOS Stage 1, rather than BOS at stages higher than Stage 1.⁸¹ Attempting to determine a predictive model in which patients will benefit from azithromycin therapy, Verleden et al found that responders had a significantly higher pretreatment BAL neutrophilia and elevated IL-8 mRNA levels.⁸² Azithromycin has also been investigated for prevention of BOS in a randomized controlled trial at the Leuven University Hospital (Leuven, Belgium). Over 2 years, BOS occurred less frequently in those on azithromycin (12.5%) versus those on placebo (44.2%).⁸³

Newer antiproliferative agents such as sirolimus and everolimus have been investigated for their unique mechanism of action. Typically, they are substituted for azathioprine, as in the study by Hachem et al, who made the substitution after patients developed a composite endpoint of cumulative acute rejection, lymphocytic bronchiolitis, or BOS.⁷² While these patients had a decrease in their cumulative acute rejection score, they did not have less incidence of lymphocytic bronchiolitis or BOS. In fact, approximately 60% of patients had to discontinue therapy due to adverse reaction. In another study, Roman et al found improved renal insufficiency but no change in FEV₁ in patients who were converted to everolimus from either a calcineurin inhibitor or a purine synthesis inhibitor.⁸⁴

Immunodepleting and cytotoxic therapies have also been investigated. Anti-thymocyte globulin, which depletes circulating T-cells, was found to slow the decline of FEV₁ over the first 3 months after treatment. However, in all but two cases, BOS ultimately progressed.⁸⁵ Alemtuzumab, an anti-CD52 antibody, was studied in the setting of refractory acute rejection by Reams et al. CD52 is a surface protein expressed on B-cells, T-cells, monocytes, macrophages, and platelets. Their study observed ten patients with BOS who had previously failed increased prednisone and anti-thymocyte globulin. Four of the ten patients had improved FEV₁. However, during the study, 73% of patients experienced an infectious complication.⁸⁶

While medications to modify the immune system represent the majority of the treatments that have been used for BOS, certain procedures to modify the immune system have also been studied. One such therapy is extracorporeal photopheresis (ECP). This procedure involves administering 8-methoxypsoralen, performing leukapheresis through an extracorporeal circuit, exposure of these leukocytes to ultraviolet A light, and infusing the leukocytes back into the patient. DNA in the leukocytes is cross-linked by the ultraviolet A light, though this only affects 2%–5% of the peripheral leukocytes per treatment. Other proposed mechanisms

by which ECP works include a release of inflammatory mediators, apoptosis of peripheral T-cell populations, and, most importantly, the production of a suppressor T-cell response and induction of immune tolerance.⁸⁷ Preliminary reports showed that ECP can stabilize the decline in FEV₁ in some patients who were refractory to all other available treatments.^{87,88} Based on reports from Meloni et al, responders to ECP tend to have increased circulating T-regulatory cells in response to treatment.⁸⁹ In the largest study to date, Morrell et al¹ showed that, after initiation of ECP, rate of decline in FEV₁ slowed and stabilized with minimal related complications. Other nonpharmacological therapies, such as total lymphoid irradiation, were initially successful in small series of patients, but there was significant drop out due to progressive BOS or bone marrow suppression.^{90,91}

Treatments for RAS mirror the treatments for BOS described as above. While understanding of the risk factors and underlying immune response for BOS has slowly progressed, the questions regarding the etiology of RAS are even more numerous. In initial studies, neither augmented corticosteroids nor antibiotics showed improved survival for patients with RAS.²² Other investigators have attempted to use drugs being investigated for pulmonary fibrosis, such as pirfenidone, for treatment, with little success.⁹²

The last line of treatment for refractory CLAD is repeat lung transplantation. However, ethical questions regarding the appropriateness of retransplantation in the setting of large waiting lists for first-time transplantation have been raised. In addition, early data from a pulmonary retransplant registry showed worse survival and earlier onset of BOS compared with first-time lung transplants.^{1,93} More recently, Hayes reviewed analyses of retransplantation and found a comparable 1-year and 5-year survival rates.⁹⁴ The role of retransplantation as a treatment for chronic lung allograft dysfunction remains controversial and policies vary by center on the frequency of its use.

Conclusion

CLAD is the most common cause of death among lung transplant recipients after the first year. Through 3 decades of experience, the rate of mortality within the first year after transplantation has decreased. While there are still many unanswered questions regarding the development of chronic lung transplant rejection, recognition of subgroups within CLAD allows more specific pathophysiologic investigations and the potential to develop effective treatment modalities for patients as we begin to practice precision medicine. With the knowledge accumulated over time, the major risk factors for the development of chronic rejection have been clarified,

a greater understanding of the immunobiology related to the progression of CLAD has been achieved, and new immunomodulatory treatments have been developed to take advantage of knowledge gained. We can imagine that, in the decades to come, CLAD will become better understood and that newer, more precise therapies will be implemented, allowing lung transplant recipients to enjoy longer survival than those lung recipients who have gone before them.

Disclosure

The authors report no conflicts of interest in this work.

References

- Morrell MR, Despotis GJ, Lublin DM, Patterson GA, Trulock EP, Hachem RR. The efficacy of photopheresis for bronchiolitis obliterans syndrome after lung transplantation. *J Heart Lung Transplant.* 2010;29(4):424–431.
- Linden PK. History of solid organ transplantation and organ donation. *Crit Care Clin.* 2009;25(1):165–184, ix.
- Yusen RD, Christie JD, Edwards LB, et al; International Society for Heart and Lung Transplantation. The Registry of the International Society for Heart and Lung Transplantation: Thirtieth Adult Lung And Heart-Lung Transplant Report – 2013; focus theme: age. *J Heart Lung Transplant.* 2013;32(10):965–978.
- Burke CM, Theodore J, Dawkins KD, et al. Post-transplant obliterative bronchiolitis and other late lung sequelae in human heart-lung transplantation. *Chest.* 1984;86(6):824–829.
- Verleden GM, Raghu G, Meyer KC, Glanville AR, Corris P. A new classification system for chronic lung allograft dysfunction. *J Heart Lung Transplant.* 2014;33(2):127–133.
- Sato M, Waddell TK, Wagnetz U, et al. Restrictive allograft syndrome (RAS): a novel form of chronic lung allograft dysfunction. *J Heart Lung Transplant.* 2011;30(7):735–742.
- Stewart S, Fishbein MC, Snell GL, et al. Revision of the 1996 working formulation for the standardization of nomenclature in the diagnosis of lung rejection. *J Heart Lung Transplant.* 2007;26(12):1229–1242.
- Yousem SA, Duncan SR, Griffith BP. Interstitial and airspace granulomatous tissue reactions in lung transplant recipients. *Am J Surg Pathol.* 1992;16(9):877–884.
- Estenne M, Maurer JR, Boehler A, et al. Bronchiolitis obliterans syndrome 2001: an update of the diagnostic criteria. *J Heart Lung Transplant.* 2002;21(3):297–310.
- Estenne M, Van Muylem A, Knoop C, Antoine M. Detection of obliterative bronchiolitis after lung transplantation by indexes of ventilation distribution. *Am J Respir Crit Care Med.* 2000;162(3 Pt 1):1047–1051.
- Jones PW, Quirk FH, Baveystock CM. The St George's Respiratory Questionnaire. *Respir Med.* 1991;85 Suppl B:25–31.
- Gerbase MW, Soccia PM, Spiliopoulos A, Nicod LP, Rochat T. Long-term health-related quality of life and walking capacity of lung recipients with and without bronchiolitis obliterans syndrome. *J Heart Lung Transplant.* 2008;27(8):898–904.
- Heng D, Sharples LD, McNeil K, Stewart S, Wreghitt T, Wallwork J. Bronchiolitis obliterans syndrome: incidence, natural history, prognosis, and risk factors. *J Heart Lung Transplant.* 1998;17(12):1255–1263.
- Burton CM, Carlsen J, Mortensen J, Andersen CB, Milman N, Iversen M. Long-term survival after lung transplantation depends on development and severity of bronchiolitis obliterans syndrome. *J Heart Lung Transplant.* 2007;26(7):681–686.
- Siegel MJ, Bhalla S, Gutierrez FR, Hildebolt C, Sweet S. Post-lung transplantation bronchiolitis obliterans syndrome: usefulness of expiratory thin-section CT for diagnosis. *Radiology.* 2001;220(2):455–462.
- Bankier AA, Van Muylem A, Knoop C, Estenne M, Gevenois PA. Bronchiolitis obliterans syndrome in heart-lung transplant recipients: diagnosis with expiratory CT. *Radiology.* 2001;218(2):533–539.
- Konen E, Gutierrez C, Chaparro C, et al. Bronchiolitis obliterans syndrome in lung transplant recipients: can thin-section CT findings predict disease before its clinical appearance? *Radiology.* 2004;231(2):467–473.
- Neurohr C, Huppmann P, Samweber B, et al; Munich Lung Transplant Group. Prognostic value of bronchoalveolar lavage neutrophilia in stable lung transplant recipients. *J Heart Lung Transplant.* 2009;28(5):468–474.
- Meloni F, Vitulo P, Cascina A, et al. Bronchoalveolar lavage cytokine profile in a cohort of lung transplant recipients: a predictive role of interleukin-12 with respect to onset of bronchiolitis obliterans syndrome. *J Heart Lung Transplant.* 2004;23(9):1053–1060.
- Riise GC, Andersson BA, Kjellström C, et al. Persistent high BAL fluid granulocyte activation marker levels as early indicators of bronchiolitis obliterans after lung transplant. *Eur Respir J.* 1999;14(5):1123–1130.
- Vanaudenaerde BM, De Vleeschauwer SI, Vos R, et al. The role of the IL23/IL17 axis in bronchiolitis obliterans syndrome after lung transplantation. *Am J Transplant.* 2008;8(9):1911–1920.
- Sato M, Hwang DM, Waddell TK, Singer LG, Keshavjee S. Progression pattern of restrictive allograft syndrome after lung transplantation. *J Heart Lung Transplant.* 2013;32(1):23–30.
- Ofek E, Sato M, Saito T, et al. Restrictive allograft syndrome post lung transplantation is characterized by pleuroparenchymal fibroelastosis. *Mod Pathol.* 2013;26(3):350–356.
- Bando K, Paradis IL, Similo SL, et al. Obliterative bronchiolitis after lung and heart-lung transplantation. An analysis of risk factors and management. *J Thorac Cardiovasc Surg.* 1995;110:4–14.
- Kroshus TJ, Kshetry VR, Savik K, John R, Hertz MI, Bolman RM 3rd. Risk factors for the development of bronchiolitis obliterans syndrome after lung transplantation. *J Thorac Cardiovasc Surg.* 1997;114:195–202.
- Hachem RR, Khalifah AP, Chakinala MM, et al. The significance of a single episode of minimal acute rejection after lung transplantation. *Transplantation.* 2005;80(10):1406–1413.
- Khalifah AP, Hachem RR, Chakinala MM, et al. Minimal acute rejection after lung transplantation: a risk for bronchiolitis obliterans syndrome. *Am J Transplant.* 2005;5(8):2022–2030.
- San Segundo D, Ballesteros MÁ, Naranjo S, Zurbano F, Miñambres E, López-Hoyos M. Increased numbers of circulating CD8 effector memory T cells before transplantation enhance the risk of acute rejection in lung transplant recipients. *PLoS One.* 2013;8(11):e80601.
- Hodge S, Hodge G, Ahern J, et al. Increased levels of T cell granzyme b in bronchiolitis obliterans syndrome are not suppressed adequately by current immunosuppressive regimens. *Clin Exp Immunol.* 2009;158(2):230–236.
- Hodge G, Hodge S, Ahern J, Holmes-Liew C-L, Reynolds PN, Holmes M. Targeting steroid resistant peripheral blood pro-inflammatory CD28null T cells by inhibiting CD137 expression: relevance to BOS. *J Heart Lung Transplant.* 2013;32(4):S246.
- Yousem SA. Lymphocytic bronchitis/bronchiolitis in lung allograft recipients. *Am J Surg Pathol.* 1993;17(5):491–496.
- Husain AN, Siddiqui MT, Holmes EW, et al. Analysis of risk factors for the development of bronchiolitis obliterans syndrome. *Am J Respir Crit Care Med.* 1999;159(3):829–833.
- Keenan RJ, Lega ME, Dummer JS, et al. Cytomegalovirus serologic status and postoperative infection correlated with risk of developing chronic rejection after pulmonary transplantation. *Transplantation.* 1991;51(2):433–438.
- Smith MA, Sundaresan S, Mohanakumar T, et al. Effect of development of antibodies to HLA and cytomegalovirus mismatch on lung transplantation survival and development of bronchiolitis obliterans syndrome. *J Thorac Cardiovasc Surg.* 1998;116(5):812–820.

35. Ettinger NA, Bailey TC, Trulock EP, et al. Cytomegalovirus infection and pneumonitis. Impact after isolated lung transplantation. Washington University Lung Transplant Group. *Am Rev Respir Dis*. 1993;147:1017–1023.
36. Snyder LD, Finlen-Copeland CA, Turbyfill WJ, Howell D, Willner DA, Palmer SM. Cytomegalovirus pneumonitis is a risk for bronchiolitis obliterans syndrome in lung transplantation. *Am J Respir Crit Care Med*. 2010;181(12):1391–1396.
37. Duncan SR, Paradis IL, Yousem SA, et al. Sequelae of cytomegalovirus pulmonary infections in lung allograft recipients. *Am Rev Respir Dis*. 1992;146:1419–1425.
38. Tamm M, Aboynou CL, Chhajer PN, Rainer S, Malouf MA, Glanville AR. Treated cytomegalovirus pneumonia is not associated with bronchiolitis obliterans syndrome. *Am J Respir Crit Care Med*. 2004;170(10):1120–1123.
39. Kruger RM, Shannon WD, Arens MQ, Lynch JP, Storch GA, Trulock EP. The impact of ganciclovir-resistant cytomegalovirus infection after lung transplantation. *Transplantation*. 1999;68(9):1272–1279.
40. Hohlfeld J, Niedermeyer J, Hamm H, Schafers H, Wagner T, Fabel H. Seasonal onset of bronchiolitis obliterans syndrome in lung transplant recipients. *J Heart Lung Transplant*. 1996;15(9):888–894.
41. Vilchez RA, Dauber J, Kusne S. Infectious etiology of bronchiolitis obliterans: the respiratory viruses connection – myth or reality? *Am J Transplant*. 2003;3(3):245–249.
42. Kumar D, Erdman D, Keshavjee S, et al. Clinical impact of community-acquired respiratory viruses on bronchiolitis obliterans after lung transplant. *Am J Transplant*. 2005;5(8):2031–2036.
43. Khalifah AP, Hachem RR, Chakinala MM, et al. Respiratory viral infections are a distinct risk for bronchiolitis obliterans syndrome and death. *Am J Respir Crit Care Med*. 2004;170(2):181–187.
44. Kuo E, Bharat A, Goers T, et al. Respiratory viral infection in obliterative airway disease after orthotopic tracheal transplantation. *Ann Thorac Surg*. 2006;82(3):1043–1050.
45. Glanville AR, Scott AI, Morton JM, et al. Intravenous ribavirin is a safe and cost-effective treatment for respiratory syncytial virus infection after lung transplantation. *J Heart Lung Transplant*. 2005;24(12):2114–2119.
46. Engelmann I, Hesse N, Fegbeutel C, et al. Incidence and impact of herpes simplex and cytomegalovirus detection in the respiratory tract after lung transplantation. *Transpl Infect Dis*. 2011;13(3):259–265.
47. Neurohr C, Huppmann P, Leuchte H, et al. Human herpesvirus 6 in bronchoalveolar lavage fluid after lung transplantation: a risk factor for bronchiolitis obliterans syndrome? *Am J Transplant*. 2005;5(12):2982–2991.
48. Valentine VG, Gupta MR, Walker JE Jr, et al. Effect of etiology and timing of respiratory tract infections on development of bronchiolitis obliterans syndrome. *J Heart Lung Transplant*. 2009;28(2):163–169.
49. Botha P, Archer L, Anderson RL, et al. Pseudomonas aeruginosa colonization of the allograft after lung transplantation and the risk of bronchiolitis obliterans syndrome. *Transplantation*. 2008;85(5):771–774.
50. Weigt SS, Elashoff RM, Huang C, et al. Aspergillus colonization of the lung allograft is a risk factor for bronchiolitis obliterans syndrome. *Am J Transplant*. 2009;9(8):1903–1911.
51. Whitson BA, Prekker ME, Herrington CS, et al. Primary graft dysfunction and long-term pulmonary function after lung transplantation. *J Heart Lung Transplant*. 2007;26(10):1004–1011.
52. Daud SA, Yusef RD, Meyers BF, et al. Impact of immediate primary lung allograft dysfunction on bronchiolitis obliterans syndrome. *Am J Respir Crit Care Med*. 2007;175(5):507–513.
53. D'Ovidio F, Mura M, Ridsdale R, et al. The effect of reflux and bile acid aspiration on the lung allograft and its surfactant and innate immunity molecules SP-A and SP-D. *Am J Transplant*. 2006;6(8):1930–1938.
54. Blondeau K, Mertens V, Vanaudenaerde BA, et al. Gastro-oesophageal reflux and gastric aspiration in lung transplant patients with or without chronic rejection. *Eur Respir J*. 2008;31(4):707–713.
55. Cantu E 3rd, Appel JZ 3rd, Hartwig MG, et al. J Maxwell Chamberlain Memorial Paper. Early fundoplication prevents chronic allograft dysfunction in patients with gastroesophageal reflux disease. *Ann Thorac Surg*. 2004;78(4):1142–1151; discussion 1142–1151.
56. Davis D, Green C, Keshavjee S, et al. RESULT (Reflux Surgery in Lung Transplantation) to determine if prevention of GERD related aspiration by surgical fundoplication improves lung allograft function. Sponsor: National Heart, Lung, and Blood Institute (NHLBI). *J Heart Lung Transplant*. 2014;33(4):S41.
57. Opelz G, Wujciak T. The influence of HLA compatibility on graft survival after heart transplantation. *N Engl J Med*. 1994;330:816–819.
58. Quantz MA, Bennett LE, Meyer DM, Novick RJ. Does human leukocyte antigen matching influence the outcome of lung transplantation? An analysis of 2,549 lung transplantations. *J Heart Lung Transplant*. 2000;19:473–479.
59. Sundaresan S, Mohanakumar T, Smith MA, et al. HLA-A locus mismatches and development of antibodies to HLA after lung transplantation correlate with the development of bronchiolitis obliterans syndrome. *Transplantation*. 1998;65(5):648–653.
60. van den Berg JW, Hepkema BG, Geertsma A, et al. Long-term outcome of lung transplantation is predicted by the number of HLA-DR mismatches. *Transplantation*. 2001;71(3):368–373.
61. Girnita AL, Duquesnoy R, Yousem SA, et al. HLA-specific antibodies are risk factors for lymphocytic bronchiolitis and chronic lung allograft dysfunction. *Am J Transplant*. 2005;5:131–138.
62. Jaramillo A, Smith MA, Phelan D, et al. Development of ELISA-detected anti-HLA antibodies precedes the development of bronchiolitis obliterans syndrome and correlates with progressive decline in pulmonary function after lung transplantation. *Transplantation*. 1999;67(8):1155–1161.
63. Jaramillo A, Smith CR, Maruyama T, Zhang L, Patterson GA, Mohanakumar T. Anti-HLA class I antibody binding to airway epithelial cells induces production of fibrogenic growth factors and apoptotic cell death: a possible mechanism for bronchiolitis obliterans syndrome. *Hum Immunol*. 2003;64(5):521–529.
64. Hachem RR, Yusef RD, Meyers BF, et al. Anti-human leukocyte antigen antibodies and preemptive antibody-directed therapy after lung transplantation. *J Heart Lung Transplant*. 2010;29(9):973–980.
65. Lemy A, Toungouz M, Abramowicz D. Bortezomib: a new player in pre- and post-transplant desensitization? *Nephrol Dial Transplant*. 2010;25(11):3480–3489.
66. Fukami N, Ramachandran S, Saini D, et al. Antibodies to MHC class I induce autoimmunity: role in the pathogenesis of chronic rejection. *J Immunol*. 2008;182(1):309–318.
67. Goers TA, Ramachandran S, Aloush A, Trulock E, Patterson GA, Mohanakumar T. De novo production of K-alpha1 tubulin-specific antibodies: role in chronic lung allograft rejection. *J Immunol*. 2008;180(7):4487–4494.
68. Jaramillo A, Naziruddin B, Zhang L, et al. Activation of human airway epithelial cells by non-HLA antibodies developed after lung transplantation: a potential etiological factor for bronchiolitis obliterans syndrome. *Transplantation*. 2001;71(7):966–976.
69. Burlingham WJ, Love RB, Jankowska-Gan E, et al. IL-17-dependent cellular immunity to collagen type V predisposes to obliterative bronchiolitis in human lung transplants. *J Clin Invest*. 2007;117(11):3498–3506.
70. Sumpter TL, Wilkes DS. Role of autoimmunity in organ allograft rejection: a focus on immunity to type V collagen in the pathogenesis of lung transplant rejection. *Am J Physiol Lung Cell Mol Physiol*. 2004;286(6):L1129–L1139.
71. Cairn J, Yek T, Banner NR, Khaghani A, Hodson ME, Yacoub M. Time-related changes in pulmonary function after conversion to tacrolimus in bronchiolitis obliterans syndrome. *J Heart Lung Transplant*. 2003;22(1):50–57.
72. Hachem RR, Yusef RD, Chakinala MM, et al. A randomized controlled trial of tacrolimus versus cyclosporine after lung transplantation. *J Heart Lung Transplant*. 2007;26(10):1012–1018.

73. Whyte RI, Rossi SJ, Mulligan MS, et al. Mycophenolate mofetil for obliterative bronchiolitis syndrome after lung transplantation. *Ann Thorac Surg*. 1997;64(4):945–948.
74. Verleden GM, Buyse B, Delcroix M, et al. Cyclophosphamide rescue therapy for chronic rejection after lung transplantation. *J Heart Lung Transplant*. 1999;18(11):1139–1142.
75. Dusmet M, Maurer J, Winston T, Kesten S. Methotrexate can halt the progression of bronchiolitis obliterans syndrome in lung transplant recipients. *J Heart Lung Transplant*. 1996;15(9):948–954.
76. Groves S, Galazka M, Johnson B, et al. Inhaled cyclosporine and pulmonary function in lung transplant recipients. *J Aerosol Med Pulm Drug Deliv*. 2010;23(1):31–39.
77. Hayes D Jr, Zwischenberger JB, Mansour HM. Aerosolized tacrolimus: a case report in a lung transplant recipient. *Transplant Proc*. 2010;42(9):3876–3879.
78. Iacono AT, Johnson BA, Grgurich WF, et al. A randomized trial of inhaled cyclosporine in lung-transplant recipients. *N Engl J Med*. 2006;354(2):141–150.
79. Gerhardt SG, McDyer JF, Grgis RE, Conte JV, Yang SC, Orens JB. Maintenance azithromycin therapy for bronchiolitis obliterans syndrome: results of a pilot study. *Am J Respir Crit Care Med*. 2003;168(1):121–125.
80. Yates B, Murphy DM, Forrest IA, et al. Azithromycin reverses airflow obstruction in established bronchiolitis obliterans syndrome. *Am J Respir Crit Care Med*. 2005;172(6):772–775.
81. Jain R, Hachem RR, Morrell MR, et al. Azithromycin is associated with increased survival in lung transplant recipients with bronchiolitis obliterans syndrome. *J Heart Lung Transplant*. 2010;29(5):531–537.
82. Verleden GM, Vanaudenaerde BM, Dupont LJ, Van Raemdonck DE. Azithromycin reduces airway neutrophilia and interleukin-8 in patients with bronchiolitis obliterans syndrome. *Am J Respir Crit Care Med*. 2006;174(5):566–570.
83. Vos R, Vanaudenaerde BM, Verleden SE, et al. A randomised controlled trial of azithromycin to prevent chronic rejection after lung transplantation. *Eur Respir J*. 2011;37(1):164–172.
84. Roman A, Ussetti P, Zurbano F, et al. A retrospective 12-month study of conversion to everolimus in lung transplant recipients. *Transplant Proc*. 2011;43(7):2693–2698.
85. Date H, Lynch JP, Sundaresan S, Patterson GA, Trulock EP. The impact of cytolytic therapy on bronchiolitis obliterans syndrome. *J Heart Lung Transplant*. 1998;17(9):869–875.
86. Reams BD, Musselwhite LW, Zaas DW, et al. Alemtuzumab in the treatment of refractory acute rejection and bronchiolitis obliterans syndrome after human lung transplantation. *Am J Transplant*. 2007;7(12):2802–2808.
87. Salerno CT, Park SJ, Kreykes NS, et al. Adjuvant treatment of refractory lung transplant rejection with extracorporeal photopheresis. *J Thorac Cardiovasc Surg*. 1999;117(6):1063–1069.
88. O'Hagan AR, Stillwell PC, Arroliga A, Koo A. Photopheresis in the treatment of refractory bronchiolitis obliterans complicating lung transplantation. *Chest*. 1999;115(5):1459–1462.
89. Meloni F, Cascina A, Miserere S, Perotti C, Vitulo P, Fietta AM. Peripheral CD4(+)/CD25(+) TREG cell counts and the response to extracorporeal photopheresis in lung transplant recipients. *Transplant Proc*. 2007;39(1):213–217.
90. Diamond DA, Michalski JM, Lynch JP, Trulock EP 3rd. Efficacy of total lymphoid irradiation for chronic allograft rejection following bilateral lung transplantation. *Int J Radiat Oncol Biol Phys*. 1998;41(4):795–800.
91. Fisher AJ, Rutherford RM, Bozzino J, Parry G, Dark JH, Corris PA. The safety and efficacy of total lymphoid irradiation in progressive bronchiolitis obliterans syndrome after lung transplantation. *Am J Transplant*. 2005;5(3):537–543.
92. Vos R, Verleden SE, Ruttens D, et al. Pirfenidone: a potential new therapy for restrictive allograft syndrome? *Am J Transplant*. 2013;13(11):3035–3040.
93. Novick RJ, Stitt LW, Al-kattan K, et al. Pulmonary retransplantation: predictors of graft function and survival in 230 patients. *Ann Thorac Surg*. 1998;65(1):227–234.
94. Hayes D Jr. A review of bronchiolitis obliterans syndrome and therapeutic strategies. *J Cardiothorac Surg*. 2011;6:92.

Transplant Research and Risk Management

Publish your work in this journal

Transplant Research and Risk Management is an international, peer-reviewed open access journal focusing on all aspects of transplantation and risk management to achieve optimal outcomes in the recipient improving survival and quality of life. The journal welcomes submitted papers covering original research, basic science, clinical studies,

Submit your manuscript here: <http://www.dovepress.com/transplant-research-and-risk-management-journal>

reviews & evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.