

Washington University School of Medicine Digital Commons@Becker

Posters

2005: Alcoholism and Comorbidity

2005

Evidence for specificity of transmission of alcohol and nicotine dependence in an offspring of twins sample

Heather E. Volk

Jeffrey F. Scherrer

Kathleen K. Bucholz

Andrew C. Heath

Theodore Jacob

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.wustl.edu/guzeposter2005>

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Volk, Heather E.; Scherrer, Jeffrey F.; Bucholz, Kathleen K.; Heath, Andrew C.; Jacob, Theodore; and True, William R., "Evidence for specificity of transmission of alcohol and nicotine dependence in an offspring of twins sample" (2005). *Posters*. Paper 15 Samuel B. Guze Symposium on Alcoholism.
<http://digitalcommons.wustl.edu/guzeposter2005/15>

This Poster is brought to you for free and open access by the 2005: Alcoholism and Comorbidity at Digital Commons@Becker. It has been accepted for inclusion in Posters by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Authors

Heather E. Volk, Jeffrey F. Scherrer, Kathleen K. Bucholz, Andrew C. Heath, Theodore Jacob, and William R. True

Evidence for Specificity of Transmission of Alcohol and Nicotine Dependence in an Offspring of Twins Sample

Heather E Volk MPH, Jeffrey F Scherrer PhD,
Kathleen K Buchloz PhD, Andrew C Heath DPhil,
Theodore Jacob PhD, William R True, PhD MPH



Background

- Alcohol dependence (AD) and nicotine dependence (ND) frequently co-occur
 - 22.8% of ND have alcohol use disorder¹
 - 34.5% with alcohol use disorder are ND¹
- AD and ND are both influenced by genes
 - 50-60% of risk for AD²⁻⁴
 - 30-60% of risk for smoking initiation^{2,5}
 - 58-74% of risk for smoking persistence^{2,6}

Background

- AD and ND share genetic vulnerability
 - Genetic Correlation (r^A)
= 0.68 (95%CI 0.61-0.74)
 - Unique Environmental Correlation (r^E)
= 0.23 (95%CI 0.14-0.32)²
- However, this overlap is incomplete. Risk may still be transmitted for AD only or ND only as well as for both substances together.
- Factors such as gender, age, and externalizing or internalizing disorders may moderate risk

Objective

- To test for specificity of transmission of AD and ND extending existing results on overlap of AD and ND using an offspring of twins design.

Methods (sample)

- Data from 2000-2002 study of adolescent and adult offspring of twin fathers sampled from the Vietnam Era Twin Registry
 - 730 twin fathers, 904 biologic and/or rearing mothers, 1,356 offspring
 - Lifetime diagnoses derived from structured diagnostic interview

Methods (measures)

- Twin father's lifetime AD and ND diagnoses
- Offspring AD, ND, conduct disorder, panic attack, major depression, generalized anxiety disorder
- Maternal report of offspring attention deficit hyperactivity disorder (ADHD) and oppositional defiant disorder (ODD)

Methods (measures)

- Offspring disorders comorbid with AD or ND collapsed
 - Externalizing disorders: conduct disorder, ADHD, ODD
 - Internalizing disorders: panic attack, major depression, generalized anxiety disorder

Analysis

- Multinomial logistic regression to predict risk for comorbid AD and ND, AD only, or ND only
- 4 group offspring of twin design created for paternal AD and ND separately
 - Group 1: MZ/DZ affected
 - Group 2: MZ unaffected, cotwin affected
 - Group 3: DZ unaffected, cotwin affected
 - Group 4: MZ/DZ unaffected

Analysis

- Repeated multinomial logistic regression to examine effect of offspring gender, age, comorbid externalizing and internalizing psychopathology

Results

Table 1: Multinomial Logistic Regression to Examine the Co-transmission of AD and ND due to Genetic Factors

	Comorbid AD and ND N=116	AD N=100	ND N=196
MZ/DZ AD (Group 1)	1.79 (1.10-2.94)	2.09 (1.20-3.64)	1.06(0.70-1.62)
MZ unaffected, cotwin AD (Group 2)	1.01 (0.40-2.57)	2.22 (1.08-4.53)	0.94 (0.50-1.78)
DZ unaffected, cotwin AD (Group 3)	1.25 (0.61-2.56)	1.64 (0.74-3.60)	0.69 (0.36-1.34)
MZ/DZ ND (Group 1)	1.93 (1.10-3.37)	0.79 (0.42-1.46)	2.56 (1.62-4.06)
MZ unaffected, cotwin ND (Group 2)	1.68 (0.72-3.93)	0.99 (0.50-1.98)	2.21 (1.25-3.92)
DZ unaffected, cotwin ND (Group 3)	1.65 (0.91-2.99)	0.83 (0.44-1.58)	1.33 (0.78-2.27)

Table 2: Multinomial Logistic Regression to Examine the Co-transmission of AD and ND due to Genetic Factors Adjusting for Covariates

	Comorbid AD and ND N=116	AD N=100	ND N=196
MZ/DZ AD (Group 1)	1.54 (0.93-2.56)	2.07 (1.17-3.68)	0.96 (0.63-1.47)
MZ unaffected, cotwin AD (Group 2)	0.82 (0.32-2.12)	2.24 (1.09-4.63)	0.82 (0.42-1.63)
DZ unaffected, cotwin AD (Group 3)	1.28 (0.62-2.65)	1.73 (0.78-3.87)	0.72 (0.37-1.38)
MZ/DZ ND (Group 1)	1.81 (1.01-3.25)	0.77 (0.42-1.44)	2.47 (1.57-3.90)
MZ unaffected, cotwin ND (Group 2)	1.49 (0.63-3.52)	0.97 (0.48-1.98)	2.04 (1.13-3.70)
DZ unaffected, cotwin ND (Group 3)	1.32 (0.71-2.44)	0.79 (0.41-1.53)	1.14 (0.79-1.64)
Male Gender	1.10 (0.70-1.75)	2.45 (1.53-3.92)	1.14 (0.79-1.64)
Household Income >\$50K/year	0.74 (0.45-1.23)	1.03 (0.62-1.72)	0.88 (0.59-1.32)
Age \geq 18 years	3.14 (1.81-5.46)	10.22 (4.82-21.64)	3.17 (2.04-4.92)
Externalizing Disorders	6.91 (4.33-11.02)	1.95 (1.17-3.24)	3.34 (2.33-4.78)
Internalizing Disorders	1.91 (1.21-3.02)	0.98 (0.52-1.81)	2.04 (1.39-2.99)

Discussion

- Paternal AD and ND are associated with offspring AD and ND, respectively
- Paternal AD and ND predict comorbid AD and ND in offspring
- Specific genetic effects exist for transmission of AD and ND despite genetic correlation between the disorders

Discussion

- After controlling for genetic factors:
 - Age \geq 18 years and externalizing psychopathology increase risk for all outcomes
 - Internalizing disorders are associated with increased risk for comorbid AD and ND and ND alone
 - Male gender is associated with increased risk for AD alone

References

- 1) Grant BF, Hasin DH, Chou SP, Stinson FS, Dawson DA (2004) Nicotine dependence and psychiatric disorders in the United States. *Arch Gen Psychiatry* 61: 1107-1115.
- 2) True W, Xian H, Scherrer J, Madden P, Bucholz KK, Heath AC, Eisen S, Lyons M, Goldberg J, Tsuang M (1999) Common genetic vulnerability for nicotine and alcohol dependence in men. *Arch Gen Psychiatry* 56: 655-661.
- 3) Heath AC, Bucholz KK, Madden P, Dinwiddie SH, Slutske WS, Statham DJ, Dunne MP, Whitfield J, Martin N (1997) Genetic and environmental contributions to alcohol dependence risk in a national sample: Consistency of findings in men and women. *Psychol Med* 27: 1381-1896.
- 4) Kendler KS, Prescott CA, Neale MC, Pedersen NL (1997) Temperance board registration for alcohol abuse in a national sample of Swedish male twins, born 1902 to 1949. *Arch Gen Psychiatry* 54: 178-184.
- 5) Heath AC and Martin N (1993) Genetic models for natural history of smoking: evidence for a genetic influence on smoking persistence. *Addict Behav* 18: 19-34
- 6) Heath AC and Madden P (1995) Genetic Influences on smoking behavior. In: Turner JR, Cardon LR, Hweitt JK, eds. *Behavior Genetic Approaches in Behavioral Medicine*. New York, NY: Plenum Publishing Corp; 45-66.