

Washington University School of Medicine Digital Commons@Becker

Open Access Publications

2006

The fidelity of DNA synthesis by yeast DNA polymerase zeta alone and with accessory proteins

Xuejun Zhong

National Institutes of Health

Parie Garg

Washington University School of Medicine in St. Louis

Stephanie A. Nick McElhinny

Washington University School of Medicine in St. Louis

Carrie M. Stith

Washington University School of Medicine in St. Louis

Grace E. Kissling

National Institutes of Health

See next page for additional authors

Follow this and additional works at: http://digitalcommons.wustl.edu/open_access_pubs

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Zhong, Xuejun; Garg, Parie; Nick McElhinny, Stephanie A.; Stith, Carrie M.; Kissling, Grace E.; Burgers, Peter M.J.; and Kunkel, Thomas A., "The fidelity of DNA synthesis by yeast DNA polymerase zeta alone and with accessory proteins." *Nucleic Acids Research*.34,17. 4731-4742. (2006).

http://digitalcommons.wustl.edu/open_access_pubs/111

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Authors

Xuejun Zhong, Parie Garg, Stephanie A. Nick McElhinny, Carrie M. Stith, Grace E. Kissling, Peter M.J. Burgers, and Thomas A. Kunkel

The fidelity of DNA synthesis by yeast DNA polymerase zeta alone and with accessory proteins

Xuejun Zhong, Parie Garg², Carrie M. Stith², Stephanie A. Nick McElhinny, Grace E. Kissling¹, Peter M. J. Burgers² and Thomas A. Kunkel*

Laboratory of Molecular Genetics and Laboratory of Structural Biology and ¹Biostatistics Branch, National Institute of Environmental Health Sciences, NIH, DHHS, Research Triangle Park, NC 27709, USA and ²Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine, St Louis, MO 63110, USA

Received May 2, 2006; Revised and Accepted June 15, 2006

ABSTRACT

DNA polymerase zeta (pol ζ) participates in several DNA transactions in eukaryotic cells that increase spontaneous and damage-induced mutagenesis. To better understand this central role in mutagenesis *in vivo*, here we report the fidelity of DNA synthesis *in vitro* by yeast pol ζ alone and with RFC, PCNA and RPA. Overall, the accessory proteins have little effect on the fidelity of pol ζ . Pol ζ is relatively accurate for single base insertion/deletion errors. However, the average base substitution fidelity of pol ζ is substantially lower than that of homologous B family pols α , δ and ϵ . Pol ζ is particularly error prone for substitutions in specific sequence contexts and generates multiple single base errors clustered in short patches at a rate that is unprecedented in comparison with other polymerases. The unique error specificity of pol ζ *in vitro* is consistent with Pol ζ -dependent mutagenic specificity reported *in vivo*. This fact, combined with the high rate of single base substitution errors and complex mutations observed here, indicates that pol ζ contributes to mutagenesis *in vivo* not only by extending mismatches made by other polymerases, but also by directly generating its own mismatches and then extending them.

INTRODUCTION

In order to copy the many different types of DNA substrates encountered during replication, repair and recombination, eukaryotes encode multiple DNA polymerases. Among these, DNA polymerase ζ (pol ζ) has a particularly important role in cellular processes that result in mutagenesis (1). Pol ζ

is a heterodimer composed of proteins encoded by the *REV3* and *REV7* genes (2). *REV3* encodes the catalytic subunit, which contains a B family DNA polymerase domain at its C-terminus. Genetic studies reveal that cells lacking Rev3p or Rev7p have reduced levels of mutagenesis induced by a variety of physical and chemical agents that damage DNA (1). This strongly implicates pol ζ activity in translesion DNA synthesis. Pol ζ function is also required for the majority of spontaneous mutagenesis in wild-type yeast cells (1,3–7), as well as for mutagenesis associated with transcription (8), with double-strand break repair (9–11), and with defective DNA repair (12–15). Vertebrate pol ζ also modulates base substitution mutagenesis during somatic hypermutation of immunoglobulin genes (16,17).

Unlike B family enzymes such as pol δ and pol ϵ that have intrinsic 3' exonuclease activities and are highly accurate (18–20), pol ζ lacks intrinsic 3' exonuclease activity (2), and cannot proofread any misinsertions it may make. Steady-state kinetic analyses of insertion of individual dNTPs at specific primer-templates indicate that, depending on the mismatch, yeast pol ζ discriminates against dNTP misinsertion by factors of $0.19\text{--}41 \times 10^{-4}$, leading to the suggestion that pol ζ synthesizes DNA with accuracy approaching that of exonuclease proficient pol δ (21). Also, with the exception of a C–C mismatch, pol ζ (21) and pol α (22) exhibit a similar range of dNTP misinsertion efficiencies, suggesting that these two naturally exonuclease-deficient polymerases could have similar base substitution fidelity.

Of importance to understanding the basis of mutagenesis were steady-state kinetic analyses indicating that pol ζ discriminates poorly against the insertion of correct dNTPs onto mismatched primer termini (21,23,24). The promiscuity of pol ζ in mismatch extension greatly exceeds that of pol α , the other non-proofreading B-class enzyme (22,25). Along with reports that pol ζ has a limited ability to insert nucleotides opposite damaged template bases (26), this mismatch extension promiscuity has led to the idea that the role of pol ζ in translesion DNA synthesis is primarily to

*To whom correspondence should be addressed at: Bldg 101, Room E342B, 111 T.W. Alexander Drive, Research Triangle Park, NC 27709-2233, USA. Tel: +1 919 541 2644; Fax: +1 919 541 7613; Email: kunkel@niehs.nih.gov

The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors

extend mismatches made by other DNA polymerases (1,27). Those other polymerases could include pols α , δ and ϵ , which have primary responsibility for replicating the undamaged nuclear genome. In fact, among those polymerases examined to date, the promiscuity of pol ζ in mismatch extension is shared only by Y-family enzymes such as pol η (28) and pol κ (27) and by X family enzymes such as pol λ (29) and, for some mismatches, pol β (30,31). The X- and Y-family polymerases are considerably less accurate than the three major replicative polymerases in family B (see further discussions below).

The studies mentioned above, and genetic evidence indicating that pol ζ also participates in forming spontaneous single base insertions and deletions and more complex mutations (5,6,12,32), prompted the present study to define the fidelity of DNA synthesis by *Saccharomyces cerevisiae* pol ζ when copying undamaged DNA templates. Here we do so in reactions containing all four dNTPs in direct competition with each other, using an assay that defines rates for single base errors and detects more complex errors in a variety of sequence contexts, similar to those encountered by pol ζ *in vivo*. This assay has been used previously to describe the fidelity of many other DNA polymerases, allowing direct comparisons to be made for copying the same template bases. Our fidelity studies have been carried out with the *S.cerevisiae* pol ζ heterodimer composed of Rev3p (173 kDa) and Rev7p (29 kDa). Motivated by a recent study (33) demonstrating that the efficiency of translesion synthesis by pol ζ is strongly enhanced in the presence of the PCNA processivity clamp and the single-strand DNA-binding protein complex RPA, we also report the effects of these accessory proteins on the fidelity of pol ζ .

MATERIALS AND METHODS

Enzymes and reagents

S.cerevisiae Pol ζ was purified as described previously (33). RPA, PCNA and RFC were purified from *Escherichia coli* overproduction strains as described previously (34,35). Materials for the fidelity assay were from previously described sources (19,36).

Preparation of complementary DNA substrates

In order to measure the fidelity of DNA synthesis when copying complementary strands of the same DNA sequence, two new M13lac DNA substrates were constructed for the M13-based forward mutation assay as follows. Double-stranded M13mp2 DNA was digested with restriction endonuclease HhaI and the fragment spanning nt -83 through $+196$ (where $+1$ is the first transcribed nucleotide of the *LacZ* gene) was isolated by electroelution from a 2.0% agarose gel. This fragment was treated with T4 DNA polymerase in a reaction containing dGTP to generate a 275 bp blunt-ended fragment whose ends were bp -81 and $+194$. Double-stranded M13 DNA was digested with restriction endonuclease Aval, which cuts once at position 5825 in the

M13 genome. This DNA was similarly treated with T4 DNA polymerase in a reaction containing dGTP to generate a linear, blunt-ended M13 vector. The two blunt-ended fragments were ligated and introduced into competent *E.coli* cells to score plaques having a blue-plaque phenotype. DNA from several blue plaques was analyzed by DNA sequence analysis to determine the sequence and orientation of the *LacZ* insert. Two new vectors, one having an insert in the orientation identical to that found in the original series of M13 lac vectors and the other having the same insert in the opposite orientation, were chosen for preparation of two different gapped DNA substrates for fidelity assays. The restriction enzymes used for gap construction were BanII and Aval. These cut both new constructs to yield a large 6218 bp fragment. This fragment was purified using three cycles of PEG precipitation at 37°C , in order to completely remove the smaller DNA fragments with sticky ends. The 6218 bp primer fragment was then hybridized to complementary single-stranded circular M13lac DNAs, as described previously (36). This generates duplex circular M13lac substrates containing a 461 nt single-stranded DNA gap. These gapped substrates were gel-purified and used in polymerization reactions. One gap contains the (+) strand of the 275 base *lacZ* α -complementation sequence, such that the template being copied (Figure 1) is identical to that used in previous studies of polymerase fidelity that involved the M13mp2 forward mutation assay. The other gap contains the (–) strand as a template, thereby allowing parallel measurements of error rates for synthesis instructed by each of the two complementary DNA strands.


Gap-filling DNA synthesis reactions and product analysis

Reactions (25 μl) contained 20 mM Tris-HCl, pH 7.7, 8 mM MgAc_2 , 60 mM NaCl, 0.5 mM ATP, 100 μM each dNTPs, 1 mM DTT, 100 $\mu\text{g/ml}$ BSA, 25 fmol (1 nM) gapped DNA, ~ 1 pmol pol ζ and 0.1% Triton X-100. When included, the amounts of accessory proteins used were 500 fmol PCNA, 200 fmol RFC and 5 pmol RPA. Reaction mixtures without accessory factors were incubated for 45 min at 30°C , and those with accessory factors for 30 min, and DNA products were then analyzed by agarose gel electrophoresis as described previously (36). All reactions filled the gap [data not shown, but for typical result see Figure 3 in Ref. (36)]. DNA products of gap-filling reactions were introduced into *E.coli* cells and plated as described previously (1) to score blue M13 plaques resulting from correct synthesis and light blue and colorless plaques containing polymerization errors. The types of errors were determined by sequencing the *lacZ* α -complementation gene in single-stranded DNA isolated from independent mutant M13 plaques. Error rates were calculated according to following equation: $\text{ER} = (N_i/N) \times \text{MF} / (D \times 0.6)$, where N_i is the number of mutations of a particular type, N is the total number of mutants analyzed, MF is the frequency of *lacZ* mutants, D is the number of detectable sites for the particular type of mutation and 0.6 is the probability of expressing a mutant allele in *E.coli* (36). Error rates for phenotypically


silent nucleotide changes were calculated as the number of changes observed divided by the total number of bases analyzed, i.e. number of mutants sequenced times the number of bases in the analyzed region (from position 198, the

first nucleotide incorporated, to position -81, see spectra in Figure 1). The statistical significance of differences in pairs of error rates was calculated using the Fisher's exact test (37).

(A) Pol ζ, (+) FWD



(B) Pol ζ, (-) REV




RESULTS AND DISCUSSION

Fidelity measurements and calculation of error rates


The fidelity of pol ζ with and without accessory proteins was determined for synthesis to fill a single-stranded gap in a circular duplex M13*lac* DNA substrate. The gap contains the *lacZ* α-complementation template sequence (Figure 1)

that when copied correctly leads to a blue M13 plaque phenotype. Polymerization errors are detected as light blue and colorless plaques. Gap-filling reactions were conducted with the pol ζ heterodimer in four conditions: alone, with RPA, with RFC and PCNA, or with RPA, RFC and PCNA. All four reactions filled the gap (data not shown). When the DNA products were analyzed for *lacZ* mutant frequencies,


(C) Pol ζ +PCNA/RFC, (+) FWD



(D) Pol ζ +RPA, (+) FWD



(E) Pol ζ +PCNA/RFC+RPA, (+) FWD



(F) Pol ζ +PCNA/RFC+RPA, (-) REV





Figure 1. Error spectra of single base change by pol ζ alone and with accessory proteins. (A) Pol ζ alone with lacZα (+) strand as template in the forward direction: DNA synthesis fidelity assay was performed with gapped M13mp2 substrate that contained a portion of the lacZ gene in the 461 nt gap region. Error changes generated by pol ζ near the M13mp2 lacZ target sequence are shown above the bottom template sequence. The base substitution mutations are shown as the base changed to, single nucleotide deletions are shown as triangles and additions are shown as the plus sign and the base added. The lacZ(+) strand in forward direction is shown, with the transcriptional start site designated as position 1 and the first 53 codons displayed as triplets except the bases introduced during cloning (positions 194–198). The changes colored in gray are phenotypically silent and were found as hitchhikers in the mutants with detectable mutations. The numbers next to the A175G and C196G changes are base substitution error rates at these positions. (B) Pol ζ alone with lacZα(-) strand as template in the reverse direction. (C) Pol ζ and PCNA/RFC with lacZα (+) strand as template in the forward direction. (D) Pol ζ and RPA with lacZα(+) strand as template in the forward direction. (E) Pol ζ, PCNA/RFC and RPA with lacZα(+) strand as template in the forward direction. (F) Pol ζ, PCNA/RFC and RPA with lacZα(-) strand as template in the reverse direction.

Table 1. Error rates of DNA polymerase ζ with and without accessory proteins

Enzyme	Pol ζ		Pol ζ +PCNA/RFC	Pol ζ +RPA	Pol ζ +PCNA/RFC+RPA		
DNA template strand and orientation	(+) FWD	(-) REV	(+) FWD	(+) FWD	(+) FWD	(-) REV	
Mutant frequency	12%	10%	7.9%	7.1%	6.0%	5.2%	
No. of mutants analyzed	133	132	116	124	159	133	
Error rates ($\times 10^{-5}$)	Frameshift	-1	4.4 (6)	5.7 (10)	4.3 (9)	2.8 (9)	1.0 (3)
		+1	0.7 (1)	5.1 (8)	2.3 (4)	1.4 (3)	2.0 (6)
Base substitution	Detectable sites	130 (131)	98 (118)	93 (120)	84 (129)	64 (115)	56 (131)
	Non-detectable sites	140 (51)	150 (55)	80 (26)	120 (43)	74 (33)	49 (18)
Mutant frequency of complex mutations		1.2% (14)	3.0% (39)	0.75% (11)	0.63% (11)	0.26% (7)	0.35% (9)

Notes: The numbers in the parentheses are the corresponding number of mutation events detected. The complex mutations are defined as multiple mutations occurred within 6 bases (Table 3). The single base substitutions/frameshift changes that observed in complex mutations and separated by one or more bases were counted in the error calculation of both complex mutation and base substitution/frameshift events. The tandem mutations were counted in complex mutation only. The base substitutions at non-detectable sites were recovered as 'hitchhikers' in *lacZ* mutants (shown as gray in Figure 1). For details of the template strand and orientation of the DNA gap substrates see Materials and Methods and Figure 1A–F.

**Figure 2.** Comparison of pol ζ base substitution error rates with and without accessory proteins for all 12 possible mismatches. Some error rates are reported as less than or equal to values (indicated by an asterisk above the bar).

the four reactions yielded mutant frequencies of 12, 7.1, 7.9 and 6.0%, respectively. A total of 133, 116, 124 and 159 mutant plaques, respectively, were sequenced to identify the nucleotide changes responsible for the phenotypes. Sequence analysis also revealed phenotypically silent nucleotide changes (light gray in Figure 1) present as hitchhikers in *lacZ* mutants that contain one or more detectable changes. Figure 1A and C–E shows the distribution of single base changes made by Pol ζ when copying the (+) strand template. The mutant frequency data and sequencing results were used to calculate rates for the different types of errors. These rates are summarized in Table 1 and Figure 2, and compared to average single base error rates for other DNA polymerases in Table 2. Also shown in Table 1 are mutant frequencies for more complex sequence changes, as listed in Table 3.

Parallel reactions were performed with the gapped substrate containing the complementary (–) strand of the *lacZ*

Table 2. Single base error rates of pol ζ compared to other DNA polymerases

DNA polymerase	Family	Error rate ($\times 10^{-5}$)		Reference
		Substitution	Deletion	
hpol η	pol Y	3500	240	(47)
hpol κ	pol Y	580	180	(48)
ypol ζ	pol B	130	4.4	This study
ypol α	pol B	9.6	3.1	(50,19) ^a
ypol δ (exo-)	pol B	13	5.7	(20)
ypol ϵ (exo-)	pol B	24	5.6	(19)
ypol δ	pol B	≤ 1.3	1.3	(20)
ypol ϵ	pol B	≤ 2	≤ 0.05	(19)

Notes: The abbreviations used in Table 2 are h, human; y, yeast.

^aData are from Ref. (50) and recalculated in Ref. (19) based on new detectable sites found.

α -complementation sequence. When copied either by pol ζ alone or by pol ζ plus all accessory proteins, the mutant frequencies were 10 and 5.2%, respectively, similar to values obtained with the (+) strand template (Table 1). DNA sequence analysis of 132 and 133 *lacZ* mutants respectively (Figure 1B and F), from those reactions resulted in the average pol ζ error rates shown in Table 1. Informative comparisons of error rates with the (+) and (–) strand templates are considered in the following discussion of the three major categories of errors made by pol ζ .

Pol ζ has relatively high single base insertion/deletion fidelity

When copying the (+) strand template, yeast pol ζ generated single base insertion and single base deletion errors at average rates of 0.7×10^{-5} and 4.4×10^{-5} , respectively (Table 1). Values when copying the (–) strand template were similar but slightly higher (one-tailed $P = 0.033$ for insertions; $P = 0.056$ for deletions), perhaps reflecting small sequence context effects on insertion/deletion fidelity, since the two templates are complementary rather than identical. Alternatively, small differences in error rates for specific subsets of errors mentioned here and below may partly reflect experimental fluctuation, given that the number of events detected by sequence analysis is sometimes small for certain errors.

Table 3. Complex mutations generated by DNA polymerase ζ

Location	Change
Pol ζ , (+) FWD	
-63 to -57	<u>TTAGCTC</u> to <u>GTAGCTG</u>
-58 to -53	<u>TCACTC</u> to <u>CCCCCCT</u>
-15 to 26	<u>T-15A; Δ-11-25; A26C</u>
-13 to 25	Δ -13-25
-9 to 26	Δ -9-26
25 to 26	CA to AT
76 to 80	<u>A76G; Δ78-80</u>
139 to 140	TC to G
161 to 163	<u>AGA</u> to G
161 to 163	<u>AGA</u> to <u>GGG</u>
161 to 163	<u>AGA</u> to <u>GGG</u>
166 to 170	<u>CCCGC</u> to <u>GCCGG</u>
178 to 181	<u>GCCCC</u> to <u>TCCT</u>
193 to 194	<u>TC</u> to <u>AG</u>
Pol ζ , (-) REV	
-65 to -61	TCAAT to G
-65 to -61	<u>TCAAT</u> to <u>G</u>
-62 to -56	<u>ATCGAGT</u> to <u>GTCGAGG</u>
-60 to -58	<u>CGA</u> to <u>GGG</u>
-55 to -53	<u>GAG</u> to <u>TT</u>
-54 to -48	<u>AGTAAATC</u> to <u>GG</u>
-47 to -45	<u>CGT</u> to <u>GTG</u>
-42 to -41	<u>GG</u> to C
-41 to -39	<u>GTC</u> to <u>CT</u>
-40	T to <u>GG</u>
-38 to -36	<u>CGA</u> to T
-37 to -34	<u>GAAA</u> to <u>TTG</u>
-34 to -33	Δ AT
-34 to -33	<u>AT</u> to <u>GG</u>
-34 to -33	<u>AT</u> to <u>GG</u>
-24 to -21	<u>CGAA</u> to G
-22 to -21	AA to <u>GAG</u>
-17 to -11	<u>CGAGCAT</u> to <u>GGCGCA</u>
-12 to -10	<u>ATA</u> to <u>GTC</u>
22 to 24	<u>AGT</u> to <u>GG</u>
27 to 28	Δ GT
40 to 42	<u>ACT</u> to <u>CC</u>
41 to 52	<u>TCCGAAATGTGA</u> to <u>GCCGAAAGTGTGG</u>
52 to 54	<u>GCT</u> to <u>CCC</u>
52 to 87	G52T; Δ 53-87
54 to 55	<u>TT</u> to <u>AA</u>
55 to 56	<u>TA</u> to <u>CC</u>
65 to 67	<u>GCA</u> to <u>CC</u>
67 to 70	<u>AGCT</u> to <u>GGCGC</u>
73 to 77	<u>ATGTT</u> to <u>GGGTG</u>
83 to 85	<u>ACT</u> to <u>CCG</u>
107 to 109	<u>GGT</u> to <u>TA</u>
108 to 111	<u>GTTG</u> to <u>ATTT</u>
109 to 110	<u>TT</u> to <u>GTG</u>
109 to 110	<u>TT</u> to <u>GTG</u>
117 to 122	<u>GCCGAA</u> to <u>CCCGGG</u>
129 to 132	<u>GTAG</u> to <u>AT</u>
136 to 137	+GG
153 to 155	<u>TTA</u> to <u>GTC</u>
Pol ζ + PCNA/RFC, (+) FWD	
-69 to -68	Δ TG
-13 to 25	Δ -13-25
33 to 39	<u>ACAGCTA</u> to <u>GCAGCTG</u>
105	A to <u>TT</u>
111 to 115	<u>CITAA</u> to <u>TTTAG</u>
126 to 136	<u>GCACATCCCC</u> to <u>ACCCTTCCCTT</u>
139 to 141	<u>TCG</u> to <u>CC</u>
149 to 151	<u>GCCG</u> to <u>CC</u>
150 to 156	<u>CGTAAATA</u> to <u>AGTAATG</u>
165 to 171	<u>GCCCGCA</u> to <u>CCCCCCG</u>
195 to 196	<u>TC</u> to <u>CG</u>
Pol ζ +RPA, (+) FWD	
61 to 64	<u>TGGC</u> to <u>GGGG</u>
71 to 83	Δ 71-83

Table 3. Continued

Location	Change
79 to 82	<u>GTCG</u> to <u>TTCT</u>
112 to 115	<u>TTAA</u> to <u>CTAG</u>
131 to 136	<u>TCCCC</u> to <u>ACCCCCT</u>
131 to 137	<u>TCCCCCT</u> to <u>ACCCCCC</u>
156 to 161	<u>AGCGAA</u> to <u>GGCGAG</u>
160 to 161	AA to <u>GG</u>
194 to 196	<u>CTC</u> to <u>TTG</u>
195 to 196	<u>TC</u> to <u>CG</u>
195 to 196	<u>TC</u> to <u>CG</u>
Pol ζ + PCNA/RFC+RPA, (+) FWD	
-56 to -55	Δ AC
61 to 65	<u>TGGCC</u> to <u>GGCGC</u>
111 to 115	<u>CTTAA</u> to <u>TTTAG</u>
152 to 156	<u>TAATA</u> to <u>GAAATG</u>
161 to 163	AGA to <u>GGG</u>
178 to 184	<u>GCCCTT</u> to <u>CCCCTTC</u>
195 to 196	<u>TC</u> to <u>CTG</u>
Pol ζ + PCNA/RFC+RPA, (-) REV	
-74 to -69	<u>TAATTA</u> to <u>GAATTT</u>
-73 to -69	<u>AATTA</u> to <u>TATTT</u>
-73 to -69	<u>AATTA</u> to <u>TATTT</u>
-73 to -69	<u>AATTA</u> to <u>TATTT</u>
-73 to -69	<u>AATTA</u> to <u>TAATTT</u>
-69 to -65	<u>ACACT</u> to <u>TCACA</u>
-25 to -20	<u>ACGAA</u> to <u>GCGAAG</u>
42 to 44	<u>TGG</u> to <u>GGT</u>
43 to 44	<u>GG</u> to <u>TT</u>

Notes: Base changes are underlined. Δ , deletions; +, insertions; (+) FWD and (-) REV, substrate strand and orientation. Multiple mutations within 6 bases were counted.

The ability of pol ζ to generate single base insertions and deletions is consistent with genetic studies implicating pol ζ in frameshift mutagenesis *in vivo* (12,38-41). Nonetheless, yeast pol ζ is substantially more accurate than the Y-family DNA polymerases pol η and pol κ (Table 2) or the X family polymerases involved in DNA repair (42), including yeast pol IV (43). In fact, pol ζ insertion/deletion error rates are similar to those for yeast pol α and exonuclease-deficient derivatives of yeast pol δ and pol ϵ (Table 2). Thus all four eukaryotic family B polymerases interact with undamaged primer-templates in a manner that modulates to similar extents misalignments that contain mismatches with a single unpaired base in the template strand (for deletions) or primer strand (for insertions).

The pol ζ error rate for deletions is marginally higher than for insertions [Table 1, one-tailed $P = 0.063$ for (+) strand template; $P = 0.076$ for (-) strand template], which is similar to observations with other polymerases [reviewed in (44)]. Both types of errors were more frequently observed in repetitive sequences (e.g. 22/31 deletions, 12/15 insertions, Figure 1) as compared with non-iterated sequences. This was also the case when copying the complementary strand substrate (data not shown), and is consistent with the strand slippage hypothesis proposed 40 years ago (45) and reviewed recently in (42). With one exception (higher deletion rate (10×10^{-5}) in the (-) strand template for pol ζ alone than for pol ζ with accessory proteins (1.0×10^{-5}), one-tailed $P < 0.0001$), pol ζ error rates for single base insertions and deletions were similar in the presence of the accessory proteins (Table 1). The limited number of mutants sequenced, and the infrequency of their occurrence, does not exclude that

accessory proteins might differentially influence error rates for specific insertion or deletion errors depending on the type or composition of the misalignment and/or its location. Nonetheless, any effects of RPA, RFC and PCNA on insertion and deletion error rates are small in comparison with the >10 000-fold selectivity against misalignments (i.e. error rates $<10^{-4}$) conferred by pol ζ alone. Thus pol ζ itself is by far the primary determinant of selectivity against single base errors resulting from substrate misalignments occurring when it copies undamaged DNA. Among many different exonuclease-deficient eukaryotic DNA polymerases (46), pol ζ has relatively high insertion/deletion fidelity (Table 2).

Pol ζ generates lacZ mutants with complex multiple sequence changes

Pol ζ generated many *lacZ* mutants that contained more than a single base change. This includes six tandem double base substitutions and one tandem triple base substitution. These were recovered among 265 sequenced *lacZ* mutants [Table 1 and Table 3, Pol ζ (+) FWD and Pol ζ , (-) REV], yielding an error rate for tandem double substitutions of $\sim 3 \times 10^{-5}$ (estimated by treating them as single detectable base substitutions). This is an extraordinary rate given that forming a tandem double base substitution requires three consecutive rare incorporation events: an initial misinsertion, a second misinsertion from the resulting mismatch and extension of doubly mismatched primer terminus. Pol ζ infidelity for tandem double base substitution errors is only exceeded by pol η (47) and pol κ (48), both of which also have a lower fidelity than pol ζ for single base substitutions (Table 2). Interestingly, tandem double base substitutions, including those unlikely to result from cytosine deamination in photodimers, are reported in the spectra of mutations induced by UV light (49).

Many of the *lacZ* mutants generated by pol ζ during DNA synthesis *in vitro* contained two or more non-tandem but closely spaced sequence changes (Tables 1 and 3). Many of the sequence changes present in these complex *lacZ* mutants are base substitutions, while others are deletion or insertion of one or more nucleotides, and these events are found in various combinations (Table 3). Importantly, the proportion of such mutants among the total number of *lacZ* mutants analyzed, and the overall frequency with which these complex events are generated (Table 1), far exceeds that observed in previous studies of other family B polymerases, including yeast pol α ($\leq 0.06\%$) (50), pol δ ($\leq 0.05\%$) (20) and pol ϵ ($\leq 0.04\%$) (19). These data lead to two important conclusions. First, they clearly demonstrate that yeast pol ζ can both create and then promiscuously extend primer-templates containing multiple, closely spaced mismatches. Second, they strongly imply that it is indeed pol ζ that is directly responsible for generating complex mutations in yeast that include a single base frameshift combined with nearby base substitutions (5,32,41).

Also observed in the collection of non-single base errors generated by pol ζ alone were deletions of larger numbers of nucleotides (Table 3). Previous studies have shown that two other B family enzymes, pol α (50) and pol δ (20), also delete large numbers of nucleotides. Those deletions involved loss of one of two direct repeats of 2–9 bases plus

the intervening nucleotides. They can be explained by a model (51) in which the first repeat sequence encountered by a polymerase during gap filling is copied, then the primer containing the repeat frays and relocates to the second repeat sequence. This provides a primer terminus adequate for continued synthesis from the misaligned intermediate that contains a loop of unpaired, template strand bases that are eventually deleted. Interestingly, the endpoints of the deletions generated here by pol ζ do not involve obvious direct repeat sequences. Thus, they may be generated by a mechanism that is distinct from large deletions generated by pol α and pol δ . However, the deletions generated by pol ζ are sometimes associated with base substitutions adjacent to the deletion endpoints. Thus, the general misalignment process may be similar, but by virtue of its promiscuous mismatch extension capacity, pol ζ may be able to extend misaligned large deletion intermediates containing less homology at the primer terminus than is required for extension by pol α and pol δ . The capacity to extend such substrates may be relevant to participation of pol ζ in the mutagenic repair of double-strand breaks in DNA (9,10).

Pol ζ exhibits low base substitution fidelity

The single base substitution error rates of pol ζ alone when copying the (+) and (-) strand templates were 130×10^{-5} and 97×10^{-5} (Table 1). These rates represent the number of substitutions generated per phenotypically detectable nucleotide polymerized, i.e. those errors known to result in a M13 plaque color phenotype. The values in Table 1 are average error rates for the 12 possible single base–base mismatches scored in many different sequence contexts. Similar rates (Table 1) were calculated (Materials and Methods) for errors at phenotypically silent locations, recovered as ‘hitchhiker’ substitutions in *lacZ* mutants.

A comparison of base substitution errors made by yeast pol ζ with those made by other DNA polymerases in the same target shows that pol ζ is much less accurate than the homologous B family yeast polymerases that conduct the bulk of chain elongation during replication of the yeast nuclear genome. Pol δ and pol ϵ are at least 100-fold more accurate than pol ζ (Table 2). Although this is partly due to their intrinsic proofreading activities (19,20), even exonuclease-deficient derivatives of pol δ and pol ϵ , as well as naturally exonuclease-deficient pol α , have substantially higher base substitution fidelity than pol ζ . The lower base substitution fidelity of pol ζ must necessarily result from lower average discrimination against dNTP misinsertion, because pol ζ is the only enzyme in the *in vitro* assay. However, the average base substitution fidelity of yeast pol ζ is higher than those for human pol κ and human pol η , both are Y-family DNA polymerases (Table 2). This can be rationalized by structural studies suggesting that Y-family polymerases have larger and more solvent accessible nascent base pair binding pockets than do B family polymerases (52).

Pol ζ has highly unusual base substitution error specificity

Error rates for the 12 different single base–base mismatches made when copying the (+) strand templates are shown in

Figure 2. Similar rates were observed for copying the (–) strand template (data not shown). Many of these error rates differ from steady-state kinetic values for individual dNTP misinsertions, as well as for correct dNTP incorporation onto mismatched primer termini (21). Logically, mismatches resulting from higher misinsertion rates that are also extended with higher efficiencies might be expected to give rise to the highest base substitution error rates. For example, the kinetic results (21) imply that the highest error rate among the 12 mismatches should be for the T-dGMP mismatch. However, the highest base substitution error rate detected in the forward mutation assay is for stable misincorporation of dCMP opposite template A. Notably, this high rate is not due to a single mutational hotspot in the lacZ target because misincorporation of dCMP was detected multiple times at nine distinct template A positions (Figure 1A). The disparity with the steady-state kinetics are likely due to the fact that the kinetic experiments were performed within a single sequence context for each mismatch, while the base substitution error rates determined here using the forward mutation assay include a wide variety of sequence contexts (Figure 1), with error rates highly dependent on sequence context (discussed below).

The present data reveal that average pol ζ error rates vary over 30-fold, depending on the mismatch. Unexpectedly, two aspects of pol ζ error specificity differ from all other polymerases characterized to date, including pol ζ 's B family homologs pol α (50), pol δ (20) and pol ϵ (19). Pol ζ is the only polymerase whose error rate for the A-dCMP mismatch exceeds that of the 11 other mismatches (Figure 2), and it is the only enzyme whose error rate for the C-dCMP mismatch is second highest among all mismatches. The error rates for these and other mismatches are only reduced by ≤ 3 -fold in the presence of the accessory proteins, implying that these error signatures should be useful for discerning the contribution of pol ζ to base substitution mutagenesis *in vivo*. For example, the unusually high relative rate for the C–C mismatch, and the fact that the third highest rate is for the G-dGMP mismatch, both suggest that pol ζ generates C–G to G–C transversions *in vivo*. Strong support for this prediction comes from the study of a yeast strain harboring a point mutation in the active site of pol δ that results in replicative stress (53). This strain has a spontaneous mutator phenotype that includes a high mutation rate for C–G to G–C transversions. Importantly, this rate is reduced by ≥ 12 -fold upon inactivation of Rev3, just as predicted by the error specificity of pol ζ in Figure 2.

Strong sequence context effects on pol ζ base substitution error rates

Pol ζ base substitutions are distributed non-randomly within the lacZ target sequence (Figure 1). For example, at template position 175 (Figure 1A), the error rate for incorporation of dCMP opposite A is 3.5% (1/29) (Figure 1A). An even higher rate of 4.5% (1/22) is observed for incorporation of dCMP opposite C at C196, a phenotypically silent location. Thus, in some sequence contexts, pol ζ is among the least accurate of DNA polymerases, rivaling the remarkable infidelity of the Y-family polymerases. The types and locations of sites of

most frequent substitutions shown in Figure 1 are distinct for those of pol η (47) and other DNA polymerases like pol β (54) when copying the same template. They are not limited to one specific sequence or to one mismatch. Initial examination of the local sequence environments of the most versus least frequent sites did not reveal an obvious pattern that could explain the distribution.

Pol ζ error rates are highly asymmetric

Inspection of the error specificity of pol ζ reveals two types of asymmetric misincorporation. First, error rates for reciprocal mismatches of the same base composition are not the same. For example, the error rate for misincorporation of dCMP opposite template A is 11-fold higher than for misincorporation of dAMP opposite template C (Figure 2, one-tailed $P < 0.0001$). Other asymmetries of this type are seen in Figure 2 and were also observed for copying the (–) strand template (data not shown). Thus the ability of pol ζ to discriminate against misincorporation depends not only on the base composition of the mismatch, but also mismatch symmetry with respect to which base is in the template or incoming as a dNTP. This is consistent with structural studies showing different polymerase side chain interactions with the template base and the incoming dNTP.

A second type of asymmetry involves differences in error rates for the two mismatches that could theoretically explain a base substitution arising during replication of double-stranded DNA *in vivo*. For example, the error rate for incorporating dCMP opposite A175 in the (+) template is 3.5%, but the error rate for incorporating dGMP opposite T175 in the complementary (–) template is $\leq 0.13\%$, a difference of at least 27-fold. Additional examples of this type of asymmetry are seen in different sequence contexts, resulting in an average error rate for misincorporation of dCMP opposite template A that is 8-fold higher than for misincorporation of dGMP opposite template T (Figure 2, one-tailed $P < 0.0001$). These asymmetries and the variations in error rates by mismatch and sequence context may be useful for assigning a role for pol ζ in specific mutagenic transactions *in vivo*.

Accessory proteins have modest effects on single base substitution fidelity

The overall average base substitution fidelity of pol ζ was increased by ~ 2 fold in the presence of the accessory proteins (Table 1). When individual single base mismatches (Figure 2) or individual template locations (Table 1 and Figure 1) are compared, the accessory proteins either have no effect or they increased (e.g. C-dCMP mismatch) or decreased (G-dATP) base substitution fidelity by a few-fold. The one exception is at template nucleotide –69 in the (–) strand template, where the accessory proteins increased the rate of A-dATP errors (Figure 1B and F). The results suggest that the accessory proteins may modulate the fidelity of Pol ζ either up or down, in a mismatch and/or sequence-specific manner. Similar effects of accessory proteins have also been observed previously with two highly accurate B family enzymes, RB69 DNA polymerase (55,56) and yeast pol δ (57). The present study shows that the effects

of accessory proteins are small compared to the level of discrimination imparted by the polymerase. In other words, the pol ζ heterodimer itself is by far the prime determinant of fidelity for both single base substitutions and single base-insertion/deletions errors. In the future, it will be interesting to test whether other accessory proteins influence the fidelity of pol ζ . Candidate proteins include Rev1p, which is implicated by genetic evidence to participate in pol ζ -dependent mutagenesis *in vivo* (1), and the 9-1-1-checkpoint clamp, which has been shown recently to physically interact with Pol ζ and is partially required for its contribution to spontaneous mutagenesis in yeast (7).

Dual roles for pol ζ in mutagenic synthesis

Based on steady-state kinetic analysis, it has been suggested previously that pol ζ is a highly accurate DNA polymerase whose primary role in mutagenesis is mismatch extension (21). The present study demonstrates that pol ζ actually has low average fidelity for single base substitutions, and very low fidelity in specific sequence contexts. Such low fidelity, and the similarities in *in vitro* error signatures and *in vivo* mutagenesis described above, strongly implicates pol ζ in both nucleotide misinsertion and mismatch extension during mutagenic synthesis *in vivo*. It is important to note that this idea does not exclude the current 'two polymerase model' for TLS (58,59). A large number of structurally diverse intermediates can be envisioned for both spontaneous and damage-induced mutagenesis. It may well be that the relative contribution of misinsertion and mismatch extension by pol ζ versus other low fidelity polymerases, several of which are also promiscuous for mismatch extension (27,28,48) may depend on the nature of the mutagenic intermediate that must be created and then extended. Consistent with a role for pol ζ in both misinsertion and mismatch extension during TLS is the demonstration that accessory proteins stimulate pol ζ to perform TLS at rates comparable to copying undamaged DNA (33). That study clearly indicates that pol ζ can insert nucleotides opposite lesions and then extend those insertions to complete bypass. Moreover, *in vivo* results on the bypass of abasic sites, T-T (6-4) photoadducts and T-T *cis-syn* cyclobutane dimers, have led to the suggestion that pol ζ , not pol δ , is responsible for insertion during TLS when pol η does not perform this function (41,60).

It is currently unknown whether pol ζ error rates during TLS are similar to, lower, or higher than those observed here for copying undamaged DNA. Nonetheless, it is particularly interesting that pol ζ has higher base substitution fidelity when copying undamaged DNA than does pol η . During complete bypass of a *cis-syn* thymine-thymine dimer, pol η generates base substitution errors at the 3' T at rates that are high, and similar to those observed for copying the equivalent undamaged T (61). However, yeast pol ζ can also clearly conduct complete TLS reactions, inefficiently without accessory proteins (2,26) and efficiently when the accessory proteins are present (33). Is pol ζ less accurate during lesion bypass than seen here with undamaged templates? If not, how is damage-induced mutagenesis *in vivo* suppressed by pol η , an enzyme with very low intrinsic fidelity, whereas damage-induced mutagenesis is promoted by

pol ζ , a polymerase with at least 10-fold higher base substitution fidelity (Table 2). Among several possibilities, it may be that TLS errors made by pol η are subjected to error correction mechanisms to a greater extent than are errors made by pol ζ . For example, if pol η conducts bypass at the replication fork (27,62), those mismatches may be proofread by pol δ or pol ϵ or corrected by mismatch repair (63), a process that may be physically coupled to the replication fork (64,65). However, if pol ζ were to participate in TLS during gap-filling DNA synthesis after the fork has moved on (66-68), those mismatches might not be as efficiently proofread or repaired by mismatch repair, resulting in mutagenesis. The potential of pol ζ to replace pol δ as a TLS gap-filling replicase in order to avoid replication fork stalling could explain why pol ζ is responsible for the majority of both spontaneous and damage-induced mutagenesis observed *in vivo*.

ACKNOWLEDGEMENTS

We thank Yuri Pavlov, Marilyn Diaz, John Drake and Zachary Pursell for thoughtful suggestions on this manuscript, and Dinh Nguyen for assistance in DNA substrate preparation. This research was supported in part by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences to T.A.K., and partly by NIH grant number GM3243 to P.M.J.B. Funding to pay the Open Access publication charges for this article was provided by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences.

Conflict of interest statement. None declared.

REFERENCES

- Lawrence,C.W. (2004) Cellular functions of DNA polymerase zeta and Rev1 protein. *Adv. Protein Chem.*, **69**, 167-203.
- Nelson,J.R., Lawrence,C.W. and Hinkle,D.C. (1996) Thymine-thymine dimer bypass by yeast DNA polymerase zeta. *Science*, **272**, 1646-1649.
- Quah,S.K., von Borstel,R.C. and Hastings,P.J. (1980) The origin of spontaneous mutation in *Saccharomyces cerevisiae*. *Genetics*, **96**, 819-839.
- Cassier,C., Chanet,R., Henriques,J.A. and Moustacchi,E. (1980) The effects of three PSO genes on induced mutagenesis: a novel class of mutationally defective yeast. *Genetics*, **96**, 841-857.
- Minesinger,B.K., Abdulovic,A.L., Ou,T.M. and Jinks-Robertson,S. (2006) The effect of oxidative metabolism on spontaneous Pol zeta-dependent translesion synthesis in *Saccharomyces cerevisiae*. *DNA Repair (Amst.)*, **5**, 226-234.
- Minesinger,B.K. and Jinks-Robertson,S. (2005) Roles of RAD6 epistasis group members in spontaneous polzeta-dependent translesion synthesis in *Saccharomyces cerevisiae*. *Genetics*, **169**, 1939-1955.
- Sabbioneda,S., Minesinger,B.K., Giannattasio,M., Plevani,P., Muzi-Falconi,M. and Jinks-Robertson,S. (2005) The 9-1-1 checkpoint clamp physically interacts with polzeta and is partially required for spontaneous polzeta-dependent mutagenesis in *Saccharomyces cerevisiae*. *J. Biol. Chem.*, **280**, 38657-38665.
- Datta,A. and Jinks-Robertson,S. (1995) Association of increased spontaneous mutation rates with high levels of transcription in yeast. *Science*, **268**, 1616-1619.
- Holbeck,S.L. and Strathern,J.N. (1997) A role for REV3 in mutagenesis during double-strand break repair in *Saccharomyces cerevisiae*. *Genetics*, **147**, 1017-1024.
- Ratray,A.J., Shafer,B.K., McGill,C.B. and Strathern,J.N. (2002) The roles of REV3 and RAD57 in double-strand-break-repair-induced mutagenesis of *Saccharomyces cerevisiae*. *Genetics*, **162**, 1063-1077.

11. Hirano, Y. and Sugimoto, K. (2006) ATR homolog Mec1 controls association of DNA polymerase zeta-Rev1 complex with regions near a double-strand break. *Curr. Biol.*, **16**, 586–590.
12. Heidenreich, E., Holzmann, V. and Eisler, H. (2004) Polymerase zeta dependency of increased adaptive mutation frequencies in nucleotide excision repair-deficient yeast strains. *DNA Repair (Amst.)*, **3**, 395–402.
13. Roche, H., Gietz, R.D. and Kunz, B.A. (1994) Specificity of the yeast rev3 delta antimutator and REV3 dependency of the mutator resulting from a defect (rad1 delta) in nucleotide excision repair. *Genetics*, **137**, 637–646.
14. Roche, H., Gietz, R.D. and Kunz, B.A. (1995) Specificities of the *Saccharomyces cerevisiae* rad6, rad18, and rad52 mutators exhibit different degrees of dependence on the REV3 gene product, a putative nonessential DNA polymerase. *Genetics*, **140**, 443–456.
15. Sarkar, S., Davies, A.A., Ulrich, H.D. and McHugh, P.J. (2006) DNA interstrand crosslink repair during G1 involves nucleotide excision repair and DNA polymerase zeta. *EMBO J.*, **25**, 1285–1294.
16. Diaz, M. and Lawrence, C. (2005) An update on the role of translesion synthesis DNA polymerases in Ig hypermutation. *Trends Immunol.*, **26**, 215–220.
17. Diaz, M., Watson, N.B., Turkington, G., Verkoczy, L.K., Klinman, N.R. and McGregor, W.G. (2003) Decreased frequency and highly aberrant spectrum of ultraviolet-induced mutations in the hprt gene of mouse fibroblasts expressing antisense RNA to DNA polymerase zeta. *Mol. Cancer Res.*, **1**, 836–847.
18. Thomas, D.C., Roberts, J.D., Sabatino, R.D., Myers, T.W., Tan, C.K., Downey, K.M., So, A.G., Bambara, R.A. and Kunkel, T.A. (1991) Fidelity of mammalian DNA replication and replicative DNA polymerases. *Biochemistry*, **30**, 11751–11759.
19. Shcherbakova, P.V., Pavlov, Y.I., Chilkova, O., Rogozin, I.B., Johansson, E. and Kunkel, T.A. (2003) Unique error signature of the four-subunit yeast DNA polymerase epsilon. *J. Biol. Chem.*, **278**, 43770–43780.
20. Fortune, J.M., Pavlov, Y.I., Welch, C.M., Johansson, E., Burgers, P.M. and Kunkel, T.A. (2005) *Saccharomyces cerevisiae* DNA polymerase delta: high fidelity for base substitutions but lower fidelity for single- and multi-base deletions. *J. Biol. Chem.*, **280**, 29980–29987.
21. Johnson, R.E., Washington, M.T., Haracska, L., Prakash, S. and Prakash, L. (2000) Eukaryotic polymerases iota and zeta act sequentially to bypass DNA lesions. *Nature*, **406**, 1015–1019.
22. Mendelman, L.V., Petruska, J. and Goodman, M.F. (1990) Base mispair extension kinetics. Comparison of DNA polymerase alpha and reverse transcriptase. *J. Biol. Chem.*, **265**, 2338–2346.
23. Lawrence, C.W., Gibbs, P.E., Murante, R.S., Wang, X.D., Li, Z., McManus, T.P., McGregor, W.G., Nelson, J.R., Hinkle, D.C. and Maher, V.M. (2000) Roles of DNA polymerase zeta and Rev1 protein in eukaryotic mutagenesis and translesion replication. *Cold Spring Harb. Symp. Quant. Biol.*, **65**, 61–69.
24. Lawrence, C.W. and Hinkle, D.C. (1996) DNA polymerase zeta and the control of DNA damage induced mutagenesis in eukaryotes. *Cancer Surv.*, **28**, 21–31.
25. Perrino, F.W. and Loeb, L.A. (1990) Hydrolysis of 3'-terminal mispairs *in vitro* by the 3'-5' exonuclease of DNA polymerase delta permits subsequent extension by DNA polymerase alpha. *Biochemistry*, **29**, 5226–5231.
26. Guo, D., Wu, X., Rajpal, D.K., Taylor, J.S. and Wang, Z. (2001) Translesion synthesis by yeast DNA polymerase zeta from templates containing lesions of ultraviolet radiation and acetylaminofluorene. *Nucleic Acids Res.*, **29**, 2875–2883.
27. Prakash, S., Johnson, R.E. and Prakash, L. (2005) Eukaryotic translesion synthesis DNA polymerases: specificity of structure and function. *Annu. Rev. Biochem.*, **74**, 317–353.
28. Bebenek, K., Matsuda, T., Masutani, C., Hanaoka, F. and Kunkel, T.A. (2001) Proofreading of DNA polymerase eta-dependent replication errors. *J. Biol. Chem.*, **276**, 2317–2320.
29. Picher, A.J., Garcia-Diaz, M., Bebenek, K., Pedersen, L.C., Kunkel, T.A. and Blanco, L. (2006) Promiscuous mismatch extension by human DNA polymerase lambda. *Nucleic Acids Res.*, **34**, 3259–3266.
30. Dalal, S., Kosa, J.L. and Sweasy, J.B. (2004) The D246V mutant of DNA polymerase beta misincorporates nucleotides: evidence for a role for the flexible loop in DNA positioning within the active site. *J. Biol. Chem.*, **279**, 577–584.
31. Beard, W.A., Shock, D.D. and Wilson, S.H. (2004) Influence of DNA structure on DNA polymerase beta active site function: extension of mutagenic DNA intermediates. *J. Biol. Chem.*, **279**, 31921–31929.
32. Harfe, B.D. and Jinks-Robertson, S. (2000) DNA polymerase zeta introduces multiple mutations when bypassing spontaneous DNA damage in *Saccharomyces cerevisiae*. *Mol. Cell*, **6**, 1491–1499.
33. Garg, P., Stith, C.M., Majka, J. and Burgers, P.M. (2005) Proliferating cell nuclear antigen promotes translesion synthesis by DNA polymerase zeta. *J. Biol. Chem.*, **280**, 23446–23450.
34. Henriksen, L.A., Umbricht, C.B. and Wold, M.S. (1994) Recombinant replication protein A: expression, complex formation, and functional characterization. *J. Biol. Chem.*, **269**, 11121–11132.
35. Ayyagari, R., Gomes, X.V., Gordenin, D.A. and Burgers, P.M. (2003) Okazaki fragment maturation in yeast. I. Distribution of functions between FEN1 AND DNA2. *J. Biol. Chem.*, **278**, 1618–1625.
36. Bebenek, K. and Kunkel, T.A. (1995) Analyzing fidelity of DNA polymerases. *Methods Enzymol.*, **262**, 217–232.
37. Fleiss, J.L. (1973) *Statistical Methods for Rates and Proportions*. John Wiley and Sons, NY.
38. Lawrence, C.W. (2002) Cellular roles of DNA polymerase zeta and Rev1 protein. *DNA Repair (Amst.)*, **1**, 425–435.
39. Lawrence, C.W., O'Brien, T. and Bond, J. (1984) UV-induced reversion of his4 frameshift mutations in rad6, rev1, and rev3 mutants of yeast. *Mol. Gen. Genet.*, **195**, 487–490.
40. Heidenreich, E., Eisler, H. and Steinboeck, F. (2006) Epistatic participation of REV1 and REV3 in the formation of UV-induced frameshift mutations in cell cycle-arrested yeast cells. *Mutat Res.*, **593**, 187–195.
41. Abdulovic, A.L. and Jinks-Robertson, S. (2006) The *in vivo* characterization of translesion synthesis across UV-induced lesions in *Saccharomyces cerevisiae*: novel insights into Pol{zeta} and Pol{eta} dependent frameshift mutagenesis. *Genetics*, **172**, 1487–1498.
42. Garcia-Diaz, M., Bebenek, K., Gao, G., Pedersen, L.C., London, R.E. and Kunkel, T.A. (2005) Structure–function studies of DNA polymerase lambda. *DNA Repair (Amst.)*, **4**, 1358–1367.
43. Bebenek, K., Garcia-Diaz, M., Patishall, S.R. and Kunkel, T.A. (2005) Biochemical properties of *Saccharomyces cerevisiae* DNA polymerase IV. *J. Biol. Chem.*, **280**, 20051–20058.
44. Bebenek, K. and Kunkel, T.A. (2000) Streisinger revisited: DNA synthesis errors mediated by substrate misalignments. *Cold Spring Harb. Symp. Quant. Biol.*, **65**, 81–91.
45. Streisinger, G., Okada, Y., Emrich, J., Newton, J., Tsugita, A., Terzaghi, E. and Inouye, M. (1966) Frameshift mutations and the genetic code. This paper is dedicated to Professor Theodosius Dobzhansky on the occasion of his 66th birthday. *Cold Spring Harb. Symp. Quant. Biol.*, **31**, 77–84.
46. Bebenek, K. and Kunkel, T.A. (2004) Functions of DNA polymerases. *Adv. Protein Chem.*, **69**, 137–165.
47. Matsuda, T., Bebenek, K., Masutani, C., Rogozin, I.B., Hanaoka, F. and Kunkel, T.A. (2001) Error rate and specificity of human and murine DNA polymerase eta. *J. Mol. Biol.*, **312**, 335–346.
48. Ohashi, E., Bebenek, K., Matsuda, T., Feaver, W.J., Gerlach, V.L., Friedberg, E.C., Ohmori, H. and Kunkel, T.A. (2000) Fidelity and processivity of DNA synthesis by DNA polymerase kappa, the product of the human DINB1 gene. *J. Biol. Chem.*, **275**, 39678–39684.
49. Kozmin, S.G., Pavlov, Y.I., Kunkel, T.A. and Sage, E. (2003) Roles of *Saccharomyces cerevisiae* DNA polymerases Poleta and Polzeta in response to irradiation by simulated sunlight. *Nucleic Acids Res.*, **31**, 4541–4552.
50. Kunkel, T.A., Hamatake, R.K., Motto-Fox, J., Fitzgerald, M.P. and Sugino, A. (1989) Fidelity of DNA polymerase I and the DNA polymerase I-DNA primase complex from *Saccharomyces cerevisiae*. *Mol. Cell Biol.*, **9**, 4447–4458.
51. Kunkel, T.A. and Soni, A. (1988) Mutagenesis by transient misalignment. *J. Biol. Chem.*, **263**, 14784–14789.
52. Yang, W. (2005) Portraits of a Y-family DNA polymerase. *FEBS Lett.*, **579**, 868–872.
53. Pavlov, Y.I., Shcherbakova, P.V. and Kunkel, T.A. (2001) *In vivo* consequences of putative active site mutations in yeast DNA polymerases alpha, epsilon, delta, and zeta. *Genetics*, **159**, 47–64.

54. Osheroff, W.P., Jung, H.K., Beard, W.A., Wilson, S.H. and Kunkel, T.A. (1999) The fidelity of DNA polymerase beta during distributive and processive DNA synthesis. *J. Biol. Chem.*, **274**, 3642–3650.
55. Bebenek, A., Carver, G.T., Kadyrov, F.A., Kissling, G.E. and Drake, J.W. (2005) Processivity clamp gp45 and ssDNA-binding-protein gp32 modulate the fidelity of bacteriophage RB69 DNA polymerase in a sequence-specific manner, sometimes enhancing and sometimes compromising accuracy. *Genetics*, **169**, 1815–1824.
56. Bebenek, A., Carver, G.T., Dressman, H.K., Kadyrov, F.A., Haseman, J.K., Petrov, V., Konigsberg, W.H., Karam, J.D. and Drake, J.W. (2002) Dissecting the fidelity of bacteriophage RB69 DNA polymerase: site-specific modulation of fidelity by polymerase accessory proteins. *Genetics*, **162**, 1003–1018.
57. Fortune, J.M., Stith, C.M., Kissling, G.E., Burgers, P.M.J. and Kunkel, T.A. (2006) RPA and PCNA suppress formation of large deletion errors by yeast DNA polymerase delta. *Nucleic Acids Res.*, in press.
58. Woodgate, R. (2001) Evolution of the two-step model for UV-mutagenesis. *Mutat. Res.*, **485**, 83–92.
59. Prakash, S. and Prakash, L. (2002) Translesion DNA synthesis in eukaryotes: a one- or two-polymerase affair. *Genes Dev.*, **16**, 1872–1883.
60. Gibbs, P.E., McDonald, J., Woodgate, R. and Lawrence, C.W. (2005) The relative roles *in vivo* of *Saccharomyces cerevisiae* Pol eta, Pol zeta, Rev1 protein and Pol32 in the bypass and mutation induction of an abasic site, T–T (6-4) photoadduct and T–T *cis-syn* cyclobutane dimer. *Genetics*, **169**, 575–582.
61. McCulloch, S.D., Kokoska, R.J., Masutani, C., Iwai, S., Hanaoka, F. and Kunkel, T.A. (2004) Preferential *cis-syn* thymine dimer bypass by DNA polymerase eta occurs with biased fidelity. *Nature*, **428**, 97–100.
62. Vaisman, A., Lehmann, A.R. and Woodgate, R. (2004) DNA polymerases eta and iota. *Adv. Protein Chem.*, **69**, 205–228.
63. McCulloch, S.D., Kokoska, R.J. and Kunkel, T.A. (2004) Efficiency, fidelity and enzymatic switching during translesion DNA synthesis. *Cell Cycle*, **3**, 580–583.
64. Umar, A. and Kunkel, T.A. (1996) DNA-replication fidelity, mismatch repair and genome instability in cancer cells. *Eur. J. Biochem.*, **238**, 297–307.
65. Kunkel, T.A. and Erie, D.A. (2005) DNA mismatch repair. *Annu. Rev. Biochem.*, **74**, 681–710.
66. Rupp, W.D. and Howard-Flanders, P. (1968) Discontinuities in the DNA synthesized in an excision-defective strain of *Escherichia coli* following ultraviolet irradiation. *J. Mol. Biol.*, **31**, 291–304.
67. Lopes, M., Foiani, M. and Sogo, J.M. (2006) Multiple mechanisms control chromosome integrity after replication fork uncoupling and restart at irreparable UV lesions. *Mol. Cell*, **21**, 15–27.
68. Heller, R.C. and Marians, K.J. (2006) Replication fork reactivation downstream of a blocked nascent leading strand. *Nature*, **439**, 557–562.