

Washington University School of Medicine Digital Commons@Becker

Open Access Publications

2015

Mobility and safety issues in drivers with dementia

David B. Carr

Washington University School of Medicine in St. Louis

Desmond O'Neill

Tallaght Hospital and Trinity College Dublin

Follow this and additional works at: http://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Carr, David B. and O'Neill, Desmond, "Mobility and safety issues in drivers with dementia." *International Psychogeriatrics*.27,10. 1613-22. (2015).

http://digitalcommons.wustl.edu/open_access_pubs/4895

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

REVIEW

Mobility and safety issues in drivers with dementia

David B. Carr¹ and Desmond O'Neill²

¹Division of Geriatrics and Nutritional Science, Department of Medicine and Neurology, Washington University in St. Louis, 660 Euclid Dr. St. Louis, MO 63110, USA

²Tallaght Hospital and Trinity College Dublin, Trinity Centre for Health Sciences, Tallaght Hospital, Dublin 24, Ireland

ABSTRACT

Although automobiles remain the mobility method of choice for older adults, late-life cognitive impairment and progressive dementia will eventually impair the ability to meet transport needs of many. There is, however, no commonly utilized method of assessing dementia severity in relation to driving, no consensus on the specific types of assessments that should be applied to older drivers with cognitive impairment, and no gold standard for determining driving fitness or approaching loss of mobility and subsequent counseling. Yet, clinicians are often called upon by patients, their families, health professionals, and driver licensing authorities to assess their patients' fitness-to-drive and to make recommendations about driving privileges. We summarize the literature on dementia and driving, discuss evidence-based assessments of fitness-to-drive, and outline the important ethical and legal concerns. We address the role of physician assessment, referral to neuropsychology, functional screens, dementia severity tools, driving evaluation clinics, and driver licensing authority referrals that may assist clinicians with an evaluation. Finally, we discuss mobility counseling (e.g. exploration of transportation alternatives) since health professionals need to address this important issue for older adults who lose the ability to drive. The application of a comprehensive, interdisciplinary approach to the older driver with cognitive impairment will have the best opportunity to enhance our patients' social connectedness and quality of life, while meeting their psychological and medical needs and maintaining personal and public safety.

Key words: older adult, dementia, Alzheimer's disease, cognitive impairment, driving, automobile, transportation, assessment

Introduction

Driving is the most important method of transportation for adults across the lifespan. In 2009, there were 33 million licensed drivers over the age of 65 years in the US (USD Department of Transportation, 2010) and 40,000,000 drivers over age of 70 years in the UK (BBC News, September 2013). Progressive disease that impairs driving in older adults has at least two serious adverse outcomes: injury or death from a motor vehicle accident (MVA) or driving cessation. However, older drivers are the safest demographic group in terms of crashes per year and older drivers with medical conditions relevant to driving are also safer than younger groups with medical conditions relevant to driving (Redelmeier et al., 2012; Papa et al., 2014). This occurs despite higher levels of multimorbidity, and points to factors in addition to reduced exposure as yet inadequately described

such as prudence and wisdom, which may mediate the impact of multiple illnesses on driver safety. However, older adults aged 70–74 years will be dependent on alternative sources of transportation for about seven years and for women about 10 years after stopping driving (Foley et al., 2002). Sharp declines in health have been associated with loss of mobility in older adults (Edwards et al., 2009). The US due to its size and rural geographic areas makes transportation options for older adults especially challenging when compared to European countries that may have more developed and cost effective public transportation systems. Thus, developed countries will be increasingly faced with an increased transportation burden and need from their elders.

Dementia and specifically common neurodegenerative diseases such as Alzheimer's disease (AD) likely contribute to both issues of traffic safety and loss of mobility. Perhaps, 4% of current drivers over age 75 years have a dementia (Foley et al., 2000) and many of older adults continue to drive well into the disease process (Odenheimer, 1993). In a study where older adults were administered a well-validated brief cognitive screen to detect dementia (e.g. the Short Blessed Test) (Katzman et al., 1983),

Correspondence should be addressed to: David B. Carr, Division of Geriatrics and Nutritional Science, Department of Medicine and Neurology, Washington University in St. Louis, 660 Euclid Dr. St. Louis, MO 63110, USA. Phone: 314-286-2706; Fax: 314-286-2701. Email: dcarr@wustledu. Received 14 Sep 2014; revision requested 7 Jan 2015; revised version received 20 Apr 2015; accepted 10 May 2015. First published online 26 June 2015.

nearly 20% of those over age 80 years failed (Stutts et al., 1998). The province of Ontario in Canada estimates that they will have over 100,000 drivers with dementia on the road by 2,028 (Hopkins et al., 2004). These studies probably underestimate the actual number of drivers with dementia on the road, since some older adults with memory loss continue to drive even after they are reported to have stopped or even if they failed to renew their licenses.

In this review, we present an evidenced-based approach to the evaluation, assessment, and counseling of older drivers with cognitive impairment. We review studies that have examined functional abilities and traffic skills in drivers with dementia, identify comorbidities that can further reduce driving competence, examine options for driving evaluation, and finally, discuss key aspects of mobility counseling to inform patients of transportation alternatives.

Definitions

Mild cognitive impairment (MCI) (termed "mild neurocognitive disorder" under the DSM-5 classification) is a syndrome defined by one or more abnormalities in a specific cognitive domain (e.g. memory, executive function), a deviation from the norm on a standardized psychometric test related to the same, and usually the absence of significant impairment in daily activities (Albert et al., 2011). Two preliminary studies indicate there may be possible impairment in driving skills MCI (Frittelli et al., 2009; Wadley et al., 2009). However, one study combined dementia and MCI together which makes interpretation of the analysis difficult and the other showed less than optimal performance rather than severe impairment. More research is needed on MCI. The diagnosis remains somewhat controversial since definitions may vary as to what level of intermittent or very mild functional impairment is allowed, if at all. In addition, many patients obviously fit a phenotype of early AD, even if they do not yet have definite functional impairment.

In contrast, dementia (termed "major neurocognitive disorder" under the DSM-5 classification) is manifested by the onset of impairment in a specific domain (e.g. often memory in the most common form of this syndrome), requires the presence of impairment in an additional cognitive domain (e.g. attention, executive function), and the deficits cause significant impairment in social and/or occupational functioning. Formal criteria to diagnose MCI, dementia, and specifically AD have recently been updated (McKhann et al., 2011) and have now been placed under the category of

Neurocognitive Disorders in the DSM-5 manual [16].

Dementia and driving outcomes

There are two major outcomes of importance in studying the impact of dementia on driving: mobility and safety. It is clear that one of the most devastating consequences of dementia is eventual loss of mobility (driving cessation), and loss of mobility in dementia is associated with mismatch between transportation needs and available resources. This important issue is discussed later in this paper. One key outcome relevant to driving safety is a crash causing injury or death, but a challenge of this outcome is the relative infrequency of crashes. In general, most studies have documented a 2-5-fold increase in crash rate compared to age-matched controls (>70+ years), although two studies have documented no differences (Carr et al., 2010).

Performance-based road tests are often used as a measure of driving competency. The majority of studies report on qualitative outcomes (e.g. "pass/fail" rates) in comparison to controls, but some studies have tracked specific types of errors (Odenheimer et al., 1994; Hunt et al., 1997). Drivers with dementia have been documented to have particular difficulties with lane checking and changing, merging, left turns, signaling to park, and route following (Akinwuntan et al., 2005). The Clinical Dementia Rating (CDR), a global measure of dementia severity, uses a semi-structured interview and exam to rate the severity of the dementia (Morris, 1993). Pooled data from two longitudinal studies involving a total of 134 individual drivers with dementia (Duchek et al., 2003; Ott et al., 2008b) reveal that 88% of drivers with very mild dementia (CDR = 0.5) and 69% of drivers with mild dementia (CDR = 1.0) were still able to pass a formal road test.

In driving simulation studies, drivers with AD in general perform more poorly than do controls without dementia (Rizzo et al., 2001; Freund et al., 2002) make slower left hand turns, are more likely to drive off the road, and drive more slowly than the speed limit (Cox et al., 1998). Studies from the National Advanced Driving Simulator at the University of Iowa noted that slow or inappropriate responses were major factors leading to simulator accidents (Rizzo et al., 2001).

The majority of studies on dementia and driving have focused on AD; however, other degenerative dementias are not uncommon and may impact driving fitness. Indeed, pure dementia syndromes are less common in advanced age justifying the

need for more research in this age group (Brayne and Davis, 2012). In a road test study of controls and vascular dementia (VD) and/or AD dementia, driving performance errors were comparable between AD and VD patients (Fitten et al., 1995). Similar to MCI, vascular dementia definitions that is not related to discrete subsequent strokes remain controversial since the minimum amount and location of white matter disease to establish this diagnosis is still unknown. Disinhibited behaviors in patients with frontotemporal dementia (FTD) have been associated with impaired driving (Caselli and Yaari, 2007; de Simone et al., 2007). Lewy Body dementia (LBD) patients may have prominent visuospatial and attention deficits, underlying psychosis and fluctuating levels of alertness, any of which may impair driving abilities early in the course of the disease. However, we are currently unaware of any specific studies on LBD and driving.

Evaluating mobility and driving safety

A preliminary enquiry to all patients who are being assessed for memory disorders and as to whether or not they drive is an important first step in clarifying the impact of dementia on both mobility and safety. The clinical opinion of a primary care physician or subspecialist, evidence of a recent crash, new onset of impaired driving behaviors noted by caregivers, decline in key cognitive domains (e.g. attention, visuospatial skills), impairment in higher order level (executive function), activities of daily living, impairment in performance-based evaluations such as road tests, and difficulty with simulator scenarios have all been used in various settings to risk stratify or assess fitness-to-drive in individuals with a dementing illness. There is no accepted gold standard for an approach to assessing driving safety, although two recent evidence-based reviews provide clinicians some guidelines based on dementia and driving studies in the literature (Carr and Ott, 2010; Iverson et al., 2010).

A useful approach for clinicians that are providing fitness-to-drive recommendation in older adults with a dementing illness should consider the following three steps: (a) confirming a diagnosis and treating reversible causes for cognitive decline (e.g. sleep apnea, discontinue sedating medications); (b) rating the severity of the dementia; and (c) identify additional comorbidities that have the potential to further decrease the ability to operate an automobile. Additional queries could focus on identifying any decrements in driving behavior that have occurred during the course of the dementing illness, inquire about the new onset of impairment in other higher order instrumental activities of

daily living which could be a proxy for impaired driving in patient, and to document the presence of specific cognitive domains (e.g. attention, visuospatial skill) that have been associated with impaired driving outcomes via psychometric testing.

Although some individuals with dementia have consistent difficulty in driving situations, many patients early in the course of dementia are still able to pass a driving performance test and are likely still relatively safe to drive. One caveat that should be considered is that simply having a diagnosis of dementia should not be the sole justification for the revocation of a driver's license (Iverson et al., 2010). However, with any new diagnosis of a progressive neurodegenerative dementia, clinicians should immediately begin a conversation about the inevitability of future driving cessation. This discussion should include mobility counseling (a discussion of transportation alternatives and/or barriers to cessation). These discussions should be repeated with both the patient and caregiver to reduce the possibility of resistance or non-compliance with future recommendations.

Clinician evaluations

Family members have expressed their desire or wish that physicians provide guidance in this area (Perkinson et al., 2005). Thus, the primary care physician or subspecialist may be the only opinion available or acceptable to the patient, caregiver, or community in regards to fitness-to-drive in an older adult with dementia. In one study, most accurate were clinicians specially trained in dementia assessment (Ott et al., 2005).

Professional guidelines and consensus statements

A consensus among national medical, transportation, and elder advocacy societies is that drivers with moderately severe dementia should not drive. Unfortunately, clinicians are rarely taught how to assess dementia severity and would be hard pressed to quantify or rate patients with a moderate dementia. The CDR, used more in research than in clinical practice, takes specialized training and often takes 45 minutes or more to complete, rendering it often impractical for many office settings. Although there are limitations to the use of psychometric testing in assessing dementia severity, specific scores on global psychometric cognitive screens or specific tests may give useful ranges that could place drivers in "at-risk" categories (see Table 1).

Comorbidities and medications

The influence of multiple medical illnesses or comorbidities on further impairing driving ability in

Table 1. Association of global cognitive screens and psychometric tests with dementia severity levels

CLINICAL MEASURE OF DEMENTIA SEVERITY	QUESTIONABLE OR VERY MILD DEMENTIA			
	NO DEMENTIA (CDR = 0)	VERY MILD DEMENTIA (CDR = 0.5)	MILD DEMENTIA (CDR = 1.0)	MODERATE TO SEVERE DEMENTIA (CDR = 2.0)
For the Dementia Specialist: Clinical dementia rating	No memory loss or inconsistent memory loss Fully oriented Judgment intact Function intact Personal care intact	Consistent slight forgetfulness Slight difficulty with orientation or judgment Slight impairment in community activities or home activities Personal care intact	Memory loss interferes with everyday activities Geographic disorientation Moderate impairment in judgment Mild but definite impairment of community or home activities Needs prompting for personal care	Severe memory loss Severe difficulty with time relationships and judgment No longer independent in activities Only simple chores preserved Needs assistance in personal effects
	N (SD)	N (SD)	N (SD)	N (SD)
For the Clinician: Short Blessed test	1.2 (1.9)	4.8 (5.9)	15.4 (5.2)	18.5 (5.5)
Mini-Mental State Examination	28.9 (1.3)	23.1 (2.5)	20 (3.9)	16.1 (4.7)
For the Psychologist:				
Logical memory	8.8 (2.9)	4.3 (2.7)	1.9 (1.7)	1.5 (2.3)
Block design	30.1 (8.6)	22.2 (9.8)	12.0 (9.6)	3.2 (6.6)
Digit symbol	45.6 (11.5)	31.7 (13.6)	17.0 (13.3)	8.3 (8.7)
Trail making test A	40.9 (20.0)	70.2 (39.2)	108.3 (50.5)	Not available
Trail making test B	81.5 (36.1)	136.0 (78.0)	190.8 (81.6)	Not available
Benton copy	9.6 (.88)	9.1 (1.6)	7.3 (2.7)	Not available

(Based on samples that average 75 years of age and 14 years education)
 From Johnson et al. (2009), Hill et al. (1992), Kanne et al. (1998), Kemper et al. (1993), Morris et al. (1989), Nourhashemi et al. (2008).

patients with dementia has not been well studied, but should be considered when evaluating driving competency. A comprehensive publication in the past five years summarizes the extensive literature on medical conditions and crash risk (Charlton et al., 2010). However, many medical conditions have not been studied.

Medical conditions that are age-related and associated with impaired driving ability – but which also hold the potential for remediation – include: diseases affecting vision (e.g. macular degeneration, glaucoma, cataracts); cardiovascular diseases (e.g. implantable defibrillators, arrhythmias); respiratory diseases (e.g. COPD, sleep apnea); neurologic diseases (e.g. AD, multiple sclerosis, stroke, Parkinson disease); psychiatric diseases (e.g. psychosis, anxiety, depression); metabolic diseases (e.g. hyper or hypoglycemia); and musculoskeletal diseases (e.g. cervical arthritis with restrictive range of motion). Medications should be reviewed and sedating drugs discontinued if safer alternatives exist. It has been difficult to consistently show

associations between some drug classes and driving impairment. The gap between theoretical and real risk has recently been illustrated in a major European study, which showed reduced crash risk with antihistamines and another that showed only significant risk with the use of benzodiazepines (Orriols et al., 2009). Yet, a recent evidenced-based review did find many associations of driving impairment with certain medications and these should be avoided or minimized when operating a motor vehicle especially in older adults with cognitive impairment: sedating antihistamines, anti-psychotics, tricyclic antidepressants, bowel/bladder antispasmodics, benzodiazepines, muscle relaxants, and narcotics (Hietland and Carr, 2014).

Psychometric tests

Studies regarding the utility of global cognitive measures like Mini-Mental State Examination (MMSE) for estimating driving impairment have been mixed (Fitten et al., 1995; Lesikar et al.,

2002). Although the MMSE may correlate with degree of driving impairment on road tests and history of crashes, it does not appear to predict future involvement in crashes and valid cut-off scores have not been defined (Molnar et al., 2006). Given its limitation in predicting performance, we do not recommend the MMSE for determining driving privileges. However, the scores can provide a rough estimate of dementia severity and possibly at-risk driving (e.g. need for further assessment) given known limitations for race, education, sensory deprivation, etc.

In 2004, a meta-analysis of neuropsychological tests of driving performance in patients with dementia concluded that tests of visuospatial skills are the most relevant predictor of driving impairment (Reger et al., 2004). More recently, visuomotor and executive function tests such as Trailmaking Tests and maze completion (Whelihan et al., 2005; Ott et al., 2008a) have been associated with driving impairment in older adults with dementia. One study that evaluated a dementia education program modeled after the American Medical Association older driver curriculum suggested that physicians may be willing to adopt such tests (Muser et al., 2006). Psychometric tests may serve to identify drivers at-risk (e.g. those that may warrant further evaluation by a performance-based road test), but should not be the sole determinants in deciding to continue or revoke driving privileges (Freund, 2006; Molnar et al., 2006). This was recently confirmed in an article showing the lack of predictive value for prospective crashes in older adults that completed the MMSE (Joseph et al., 2014).

However, some recognize that standard clinical psychometric tests have little congruence with modern models of driving behavior (Fuller, 2005) and a number of innovative approaches have been developed which seek to draw on these insights, including the Adelaide Driving Self-Efficacy Scale (George et al., 2007) and a scale of strategic and tactical compensation outlined by de Raedt in 2000 (de Raedt and Ponjaert-Kristofferson, 2000). However, neither of these approaches has yet been tested for utility with drivers with dementia.

Referral

In the absence of a gold standard or consensus for determining driving competency, clinicians may request assistance from a driving clinic or refer to other subspecialists in the community such as a geriatrician, psychiatrist, neurologist, or neuropsychologist. A Driver Rehabilitation Specialist (DRS) evaluates, develops, and implements driving

services for individuals with disabilities. DRSs in the US are often occupational therapists with additional training in driver evaluation, vehicle modification, and rehabilitation. Occupational therapy practice guidelines for these evaluations have been published, but a recent review of practices across the US and Canada indicates assessments vary significantly across programs and few have adopted standardized tools (Komer-Bitensky et al., 2006). A European project found similar results (Middleton et al., 2005). In addition, there are relatively few DRSs trained and available in smaller communities.

A driving evaluation in the US costs approximately \$350–\$800 and is generally an out-of-pocket expense, but the situation varies in other countries. Clinicians who are interested in this service can contact the occupational therapy departments in local hospitals or rehabilitation centers or the ADED directory (see, online Web resources). Many local chapters of the Alzheimer's advocacy organizations (such as the Alzheimer's Association in the USA) may provide referral sources for area driving evaluation programs.

A performance-based road test for drivers with dementia could be considered to assist with risk stratification when; (a) there is observation of new impairments in traffic skills (e.g. near misses, failure to scan, etc.), (b) prominent impairments in key cognitive domains (e.g. attention, executive function, visuospatial skills), and/or (c) the presence of a mild dementia severity rating ($CDR = 1$). Private or university-based driving clinics may not be available to everyone, but many driver licensing authorities can provide or facilitate on-road tests: in the USA, every state Department of Motor Vehicles (DMV) conducts such tests. In Europe, there is a much broader range of options through state, not-for-profit, and private organizations (Sanders et al., 2006).

A review of this topic concluded there was no evidence to demonstrate the benefit of driving evaluations with respect to the preservation of mobility or a reduction in crashes (Martin et al., 2009). Yet, a recent longitudinal study noted crash rates for drivers with dementia declined to the levels of healthy control drivers during a period of 3 years when evaluated with road tests every 6 months (Ott et al., 2008b). This finding was probably due to the removal of AD drivers who had failed road tests and were no longer driving and/or to changes in behavior of AD drivers who continued to drive.

Mobility counseling

Patients may stop driving based on physician advice (Persson, 1993). There is little data to suggest that

clinicians should recommend reducing exposure in patients with dementia (e.g. limiting trips) or mandating a co-pilot to significantly reduce driving risk. In fact, data would suggest the older drivers at highest risk for a crash are the infrequent drivers that are on the road less than 3,000 km (2,000 miles) a year (Langford et al., 2006). Two recent education interventions for health professionals were positively associated with improving communication when discussing driving with patients with dementia and/or using tools that might be of help in the assessment process (Byszewski et al., 2003; Mueser et al., 2006).

Driving cessation has been associated with a decrease in social integration (Mezuk and Rebok, 2008), decreased out-of-home activities (Marottoli et al., 2000), an increase in depressive and anxiety symptoms in the elderly (Fonda et al., 2001), and an increased risk of nursing home placement (Freeman et al., 2006). Thus, a recommendation to stop driving should not be taken lightly by the clinician. Often, the situation is not urgency and there is time to work through the process of mobility loss. The brochures "We Need to Talk" and "At the Crossroads" from The Hartford Foundation (see Web resources) could enhance communication with the patient, their families, and their clinicians and assist the patient and the family member to reach important driving decisions and maintain linkage with key destinations. Educational interventions with these materials and group meetings may be effective ways to prepare caregivers for dealing with this difficult issue (Stem et al., 2008). Referral to social workers or gerontological care managers may provide important local and regional transportation options. Caregiver support groups for families also have proven efficacy with driving cessation (Dobbs et al., 2009). Finally, the Independent Transportation Network (ITN) America is a model program that assists seniors with transportation to needed destinations and now has many sites across the US (see Web resources).

The physician's legal and ethical obligations

Many physicians are uncertain of their legal responsibility to report unsafe drivers to the state or local licensing authorities (Miller and Morley, 1993; Kelly et al., 1999). In Europe, all jurisdictions rely on reporting by the driver or family, with third-party reporting the exception from a physician. The AMA's policy states, "in situations where clear evidence of substantial driving impairment implies a strong threat to patient and public safety, and where the physician's advice to discontinue driving privileges is ignored, it is desirable and ethical to

notify the DMV" (American Medical Association, 1999). Obviously, it is preferred that referrals to the license authority be done with the patient's knowledge, and that the report be documented in the medical record. However, many primary care physicians, fearing the deterioration of a long-standing relationship with their patient, may be reluctant to be this forthcoming. If a physician decides to report an unsafe driver, most jurisdictions in the US will accept a formal letter.

Physicians may be in a "double bind," concerned that they will be liable for breach of confidentiality, but also fearing legal action if they fail to report an unsafe cognitively impaired driver who is involved in an injurious crash. Most legal experts land on the side of reporting where the physician believes that the patient and community are at high risk for a crash. Since laws and regulations on driving often change, clinicians should review their laws, statutes, and regulations in their own jurisdiction to determine current requirements. Development of specific policies regarding reporting should be vetted by legal advice. In jurisdictions with voluntary reporting laws, we recommend formal referral to the driver licensing authority for refractory cases or for those patients deemed to be at a very high risk for a crash and/or injury.

Studies are needed to compare the benefits and costs of mandatory reporting to driver licensing authorities with voluntary reporting, although studies in other illnesses are not supportive of mandatory reporting (McLachlan et al., 2007). Decision analysis studies have not shown the benefits of systematically screening and evaluating drivers with dementia (Retchin and Hilner, 1994; Leproust et al., 2008). Some studies have noted positive impact on driving of cognitive stimulation (Edwards et al., 2005) and exercise interventions on older adults (Mameleira et al., 2009), although these studies were not performed in patients with dementia. Intervention studies are needed in the earliest stages of the disease to determine whether driving could be maintained safely. As the baby boom generation comes of age, there will be a pressing need to develop comprehensive interventions to maintain driving life expectancy and to develop alternative transportation systems for our cognitively impaired older drivers. The new era of smart cars and smart roads may be of assistance to those older adults with cognitive and/or physical frailty as they try to navigate down the road in the future.

Conflict of interest

Dr Carr has support from the NIA, Missouri Department of Transportation Division of Traffic

and Highway Safety. He has been a paid consultant for the American Medical Association Older Driver Project, AD EPT, TIRF, and Medscape in the past two years. Prof O'Neill is Director of the National Programme Office for Traffic Medicine, Ireland.

Description of authors' roles

Each author wrote paragraphs, reviewed the literature, and modified each other's contributions.

Web Resources on Dementia and Driving (All accessed 04/12/15)

1. General guidelines for the clinician that include dementia and additional medical conditions
 - a. CMA Fitness to Drive
<https://www.cma.ca/En/Pages/drivers-guide.aspx>
 - b. Austroads Fitness to Drive
<https://www.onlinepublications.austroads.com.au/items/AP-G56-13>
 - c. Irish Medical Guidelines
http://www.rsa.ie/Documents/Licensed%20Drivers/Medical_Issues/Slaite_agus_Tiomaint_Medical_Fitness_to_Drive_Guidelines.pdf
 - d. AMA Older Driver Curriculum on AGS website
<http://geriatricsonline.org/ProductAbstract/physicians-guide-to-assessing-and-counseling-older-drivers/B013>
2. For difficult cases, consider referring to a driving rehabilitation specialist
 - a. AOTA
http://myaota.aota.org/Driver_search/index.aspx
 - b. Association of Driver Rehabilitation Specialists: ADED
<http://aded.site-ym.com/search/custom.asp?id=1984>
3. For refractory cases, consider referral to Department of Revenue or your Countries License Authority
 - a. Summary of United State Guidelines
<http://www.iihs.org/iihs/topics/laws/olderdrivers?topicName=older-drivers>
4. Consider Web Resources and/or Office Handouts
 - a. We Need to Talk and At the Crossroads
<http://www.thehartford.com/feature-market-excellence/publications-on-aging>
 - b. Alzheimer's Association: Dementia and Driving Resource Center
<http://www.alz.org/care/alzheimers-dementia-and-driving.as>
5. Transportation Alternatives/Social Workers, Case Managers or Local Area on Aging.

References

Akinwuntan, A.E., De Weerd, W., Feys, H., Baten, G., and Kiekens, C. (2005). The validity of a road test after stroke. *Archives of Physical Medicine and Rehabilitation*, 86, 421-426. doi:10.1016/j.apmr.2004.04.047.

- Albert, M.S. et al. (2011). The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the national institute on aging-Alzheimer's association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's and Dementia*, 7, 270-279. doi:10.1016/j.jalz.2011.03.008.
- American Medical Association (1999). Impaired Drivers and Their Physicians [Online]. AMA. Available at: <http://www.ama-assn.org/ama/pub/physician-resources/medical-ethics/code-medical-ethics/opinion224.page>; last accessed June 12, 2015.
- BBC News (September 2013). Who, what, why: How Dangerous are Elderly Drivers? [Online]. Available at: <http://www.bbc.com/news/magazine-24204489>; last accessed June 12, 2015.
- Rayne, C. and Davis, D. (2012). Making Alzheimer's and dementia research fit for populations. *Lancet*, 380, 1441-1443. doi:10.1016/S0140-6736(12)61803-0.
- Byszewski, A.M. et al. (2003). A continuing medical education initiative for Canadian primary care physicians: the driving and dementia toolkit: a pre- and post-evaluation of knowledge, confidence gained, and satisfaction. *Journal of the American Geriatrics Society*, 51, 1484-1489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14511173>
- Carr, D.B. and Ott, B.R. (2010). The older adult driver with cognitive impairment: "It's a very frustrating life." *JAMA*, 303, 1632-1641. doi:10.1001/jama.2010.481.
- Caselli, R.J. and Yaari, R. (2007). Medical management of frontotemporal dementia. *American Journal of Alzheimer's Disease and Other Dementias*, 22, 489-498. doi:10.1177/1533317507306654.
- Charlton, J. et al. (2010). Influence of chronic illness on crash involvement of motor vehicle drivers: 2nd. Report No. 300. Clayton, Australia: Monash University Accident Research Centre. Accessed from <http://www.monash.edu.au/miri/research/reports/muar300.html>; last accessed June 12, 2015.
- Cox, D.J., Quillian, W.C., Thorndike, F.P., Kovatchev, B.P., and Hanna, G. (1998). Evaluating driving performance of outpatients with Alzheimer disease. *The Journal of the American Board of Family Practice/American Board of Family Practice*, 11, 264-271. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9719348>.
- De Raedt, R. and Ponjaert-Kristoffersen, I. (2000). Can a strategic and tactical compensation reduce crash risk in older drivers? *Age and Ageing*, 29, 517-521. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11191244>.
- Desimone, V., Kaplan, L., Patronas, N., Wasserman, E.M., and Grafman, J. (2007). Driving abilities in frontotemporal dementia patients. *Dementia and Geriatric Cognitive Disorders*, 23, 1-7. doi:10.1159/000096317.
- Dobbs, B.M., Harper, L.A., and Wood, A. (2009). Transitioning from driving to driving cessation: the role of specialized support groups for individuals with dementia. *Topics in Geriatric Rehabilitation*, 25, 73-86.
- Duchek, J.M. et al. (2003). Longitudinal driving performance in early-stage dementia of the Alzheimer type. *Journal of the American Geriatrics Society*, 51, 1342-1347. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14511152>.

- Edwards, J. D., Wadley, V. G., Vance, D. E., Wood, K., Roenker, D. L. and Ball, K. K. (2005). The impact of speed of processing training on cognitive and everyday performance. *Aging and Mental Health*, 9, 262–271. doi:10.1080/13607860412331336788.
- Edwards, J. D., Lunsmann, M., Perkins, M., Rebok, G. W. and Roth, D. L. (2009). Driving cessation and health trajectories in older adults. *The Journals of Gerontology, Series A, Biological Sciences and Medical Sciences*, 64, 1290–1295. doi:10.1093/gerona/gfp114.
- Fitten, L. J. et al. (1995). Alzheimer and vascular dementias and driving: a prospective road and laboratory study. *JAMA*, 273, 1360–1365. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7715061>.
- Foley, D. J., Masaki, K. H., Ross, G. W. and White, L. R. (2000). Driving cessation in older men with incident dementia. *Journal of the American Geriatrics Society*, 48, 928–930. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10968296>.
- Foley, D. J., Heimovitz, H. K., Guralnik, J. M. and Rock, D. B. (2002). Driving life expectancy of persons aged 70 years and older in the United States. *American Journal of Public Health*, 92, 1284–1289. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12144985>.
- Fonda, S. J., Wallace, R. B. and Herzog, A. R. (2001). Changes in driving patterns and worsening depressive symptoms among older adults. *The Journals of Gerontology, Series B, Psychological Sciences and Social Sciences*, 56, S343–S351. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11682595>.
- Freeman, E. E., Gange, S. J., Muñoz, B. and West, S. K. (2006). Driving status and risk of entry into long-term care in older adults. *American Journal of Public Health*, 96, 1254–1259. doi:10.2105/AJPH.2005.069146.
- Freund, B. (2006). Office-based evaluation of the older driver. *Journal of the American Geriatrics Society*, 54, 1943–1944. doi:10.1111/j.1532-5415.2006.00966.x.
- Freund, B., Ravenstein, S., Ferris, R. and Shaheen, E. (2002). Evaluating driving performance of cognitively impaired and healthy older adults: a pilot study comparing on-road testing and driving simulation. *Journal of the American Geriatrics Society*, 50, 1309–1310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12133033>.
- Frittelli, C. et al. (2009). Effects of Alzheimer's disease and mild cognitive impairment on driving ability: a controlled clinical study by simulated driving test. *International Journal of Geriatric Psychiatry*, 24, 232–238. doi:10.1002/gps.2095.
- Fuller, R. (2005). Towards a general theory of driver behaviour. *Accident Analysis and Prevention*, 37, 461–472. doi:10.1016/j.aap.2004.11.003.
- George, S., Clark, M. and Crotty, M. (2007). Development of the Adelaide driving self-efficacy scale. *Clinical Rehabilitation*, 21, 56–61. doi:10.1177/0269215506071284.
- Hetland, A. and Carr, D. B. (2014). Medications and impaired driving. *The Annals of Pharmacotherapy*, 48, 494–506. doi:10.1177/1060028014520882.
- Hill, R. D., Storandt, M. and LaBarge, E. (1992). Psychometric discrimination of moderate senile dementia of the Alzheimer type. *Archives of Neurology*, 49, 377–380. doi:10.1001/archneur.1992.00530280065023.
- Hopkins, R. W., Kilik, L., Day, D. J., Rows, C. and Tseng, H. (2004). Driving and dementia in Ontario: a quantitative assessment of the problem. *Canadian Journal of Psychiatry. Revue Canadienne de Psychiatrie*, 49, 434–438. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15362247>.
- Hunt, L. A., Murphy, C. E., Carr, D., Duchek, J. M., Buckles, V. and Morris, J. C. (1997). Reliability of the Washington University road test: a performance-based assessment for drivers with dementia of the Alzheimer type. *Archives of Neurology*, 54, 707–712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9193205>.
- Iverson, D. J., Gronseth, G. S., Reger, M. A., Classen, S., Dubinsky, R. M., Quality Standards Subcommittee of the American Academy of Neurology. (2010). Practice parameter update: evaluation and management of driving risk in dementia: report of the quality standards subcommittee of the American Academy of Neurology. *Neurology*, 74, 1316–1324. doi:10.1212/WNL.0b013e3181da3b0f.
- Johnson, D. K., Storandt, M., Morris, J. C. and Galvin, J. E. (2009). Longitudinal study of the transition from healthy aging to Alzheimer disease. *Archives of Neurology*, 66, 1254–1259. doi:10.1001/archneurol.2009.158.
- Joseph, P. G. et al. (2014). The Mini-Mental State Examination, clinical factors, and motor vehicle crash risk. *Journal of the American Geriatrics Society*, 62, 1419–1426. doi:10.1111/jgs.12936.
- Kanne, S. M., Balota, D. A., Storandt, M., McKeel, D. W., Jr. and Morris, J. C. (1998). Relating anatomy to function in Alzheimer's disease: neuropsychological profiles predict regional neuropathology 5 years later. *Neurology*, 50, 979–985. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9566382>.
- Katzman, R., Brown, T., Fuld, P., Peck, A., Schechter, R. and Schinmehl, H. (1983). Validation of a short orientation-memory-concentration test of cognitive impairment. *The American Journal of Psychiatry*, 140, 734–739. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6846631>.
- Kelly, R., Warke, T. and Steele, I. (1999). Medical restrictions to driving: the awareness of patients and doctors. *Postgraduate Medical Journal*, 75, 537–539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10616686>.
- Kemper, S., LaBarge, E., Ferraro, F. R., Cheung, H., Cheung, H. and Storandt, M. (1993). On the preservation of syntax in Alzheimer's disease: evidence from written sentences. *Archives of Neurology*, 50, 81–86. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8418805>.
- Korner-Bitensky, N., Bitensky, J., Sofer, S., Man-Son-Hing, M. and Gelinas, I. (2006). Driving evaluation practices of clinicians working in the United States and Canada. *The American Journal of Occupational Therapy*, 60, 428–434. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16915873>.
- Langford, J., Metherst, R. and Hakamies-Bloomqvist, L. (2006). Older drivers do not have a high crash risk—a replication of low mileage bias. *Accident Analysis and Prevention*, 38, 574–578. doi:10.1016/j.aap.2005.12.002.

- Leproust, S., Lagarde, E. and Salmi, L.-R. (2008). Risks and advantages of detecting individuals unfit to drive: a markov decision analysis. *Journal of General Internal Medicine*, 23, 1796–803. doi:10.1007/s11606-008-0777-8.
- Lesikar, S.E., Gallo, J.J., Rebok, G.W. and Keyl, P.M. (2002). Prospective study of brief neuropsychological measures to assess crash risk in older primary care patients. *The Journal of the American Board of Family Practice/American Board of Family Practice*, 15, 11–19. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11841134>.
- Marmeleira, J.F., Godinho, M.B. and Fernandes, O.M. (2009). The effects of an exercise program on several abilities associated with driving performance in older adults. *Accident Analysis and Prevention*, 41, 90–97. doi:10.1016/j.aap.2008.09.008.
- Marottoli, R.A., de Leon, C.F.M., Glass, T.A., Williams, C.S., Cooney, L.M., Jr. and Berkman, L.F. (2000). Consequences of driving cessation: decreased out-of-home activity levels. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 55, S334–S340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11078110>.
- Martin, A.J., Marottoli, R. and O'Neill, D. (2009). Driving assessment form maintaining mobility and safety in drivers with dementia. *Cochrane Database Syst Rev*, CD006222. doi:10.1002/14651858.CD006222.pub2.
- Mckhann, G.M. et al. (2011). The diagnosis of dementia due to Alzheimer's disease: recommendations from the national institute on aging-Alzheimer's association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia*, 7, 263–269. doi:10.1016/j.jalz.2011.03.005.
- McLachlan, R.S., Starreveld, E. and Lee, M.A. (2007). Impact of mandatory physician reporting on accident risk in epilepsy. *Epilepsia*, 48, 1500–1505. doi:10.1111/j.1528-1167.2007.01051.x.
- Meuser, T.M., Carr, D.B., Berg-Weger, M., Newohner, P. and Morris, J.C. (2006). Driving and dementia in older adults: implementation and evaluation of a continuing education project. *The Gerontologist*, 46, 680–687. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17050760>.
- Mezuk, B. and Rebok, G.W. (2008). Social integration and social support among older adults following driving cessation. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 63, S298–S303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18818450>.
- Middleton, H., Westwood, D., Robson, J. and Kok, D. (2005). Assessment and decision criteria for driving competence in the elderly. In G.Underwood (ed.), *Traffic and Transport Psychology: Theory and Application*. London: Elsevier; 101–113.
- Miller, D.J. and Morley, J.E. (1993). Attitudes of physicians toward elderly drivers and driving policy. *Journal of the American Geriatrics Society*, 41, 722–724. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8315181>.
- Molnar, F.J., Patel, A., Marshall, S.C., Man-Son-Hing, M. and Wilson, K.G. (2006). Clinical utility of office-based cognitive predictors of fitness to drive in persons with dementia: a systematic review. *Journal of the American Geriatrics Society*, 54, 1809–1824. doi:10.1111/j.1532-5415.2006.00967.x.
- Morris, J.C. (1993). The clinical dementia rating (cdr): current version and scoring rules. *Neurology*, 43, 2412–2414. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8232972>.
- Morris, J.C. et al. (1989). The consortium to establish a registry for Alzheimer's disease (cerad). Part I. Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology*, 39, 1159–1165. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2771064>.
- Nourhashemi, F., Ousset, P.J., Gillette-Guyonnet, S., Cantet, C., Andrieu, S. and Vellas, B. (2008). A 2-year follow-up of 233 very mild (cdr 0.5) Alzheimer's disease patients (real Fr cohort). *International Journal of Geriatric Psychiatry*, 23, 460–465. doi:10.1002/gps.1904.
- Odenheimer, G.L. (1993). Dementia and the older driver. *Clinics in Geriatric Medicine*, 9, 349–364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8504384>.
- Odenheimer, G.L., Beaudet, M., Jette, A.M., Albert, M.S., Grande, L. and Minkner, K.L. (1994). Performance-based driving evaluation of the elderly driver: safety, reliability, and validity. *Journal of Gerontology*, 49, M153–M159. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8014389>.
- Orrison, L. et al. (2009). The impact of medicinal drugs on traffic safety: a systematic review of epidemiological studies. *Pharmacoepidemiology and Drug Safety*, 18, 647–658. doi:10.1002/pds.1763.
- Ott, B.R. et al. (2005). Clinician assessment of the driving competence of patients with dementia. *Journal of the American Geriatrics Society*, 53, 829–833. doi:10.1111/j.1532-5415.2005.53265.x.
- Ott, B.R., Festa, E.K., Amick, M.M., Grace, J., Davis, J.D. and Heindel, W.C. (2008a). Computerized maze navigation and on-road performance by drivers with dementia. *Journal of Geriatric Psychiatry and Neurology*, 21, 18–25. doi:10.1177/0891988707311031.
- Ott, B.R. et al. (2008b). A longitudinal study of drivers with Alzheimer disease. *Neurology*, 70, 1171–1178. doi:10.1212/01.wnl.0000294469.27156.30.
- Papa, M. et al. (2014). Comorbidities and crash involvement among younger and older drivers. *PLoS One*, 9, e94564. doi:10.1371/journal.pone.0094564.
- Perkinson, M.A. et al. (2005). Driving and dementia of the Alzheimer type: beliefs and cessation strategies among stakeholders. *The Gerontologist*, 45, 676–685. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16199403>.
- Persson, D. (1993). The elderly driver: deciding when to stop. *The Gerontologist*, 33, 88–91. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8440507>.
- Redelmeier, D.A., Yarnell, C.J., Thiruchelvam, D. and Tibshirani, R.J. (2012). Physicians' warnings for unfit drivers and the risk of trauma from road crashes. *The New England Journal of Medicine*, 367, 1228–1236. doi:10.1056/NEJMSa1114310.
- Reger, M.A., Welsh, R.K., Watson, G.S., Cholerton, B., Baker, L.D. and Craft, S. (2004). The relationship between neuropsychological functioning and driving ability in dementia: a meta-analysis. *Neuropsychology*, 18, 85–93. doi:10.1037/0894-4105.18.1.85.

- Retchin, S.M. and Hillner, B.E. (1994). The costs and benefits of a screening program to detect dementia in older drivers. *Medical Decision Making*, 14, 315-324. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7808207>.
- Rizzo, M., McGhee, D.V., Dawson, J.D. and Anderson, S.N. (2001). Simulated car crashes at intersections in drivers with Alzheimer disease. *Alzheimer Disease and Associated Disorders*, 15, 10-20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11236820>.
- Sanders, N. et al. (2006). *Eu medril project: medical testing for the driving licence: report No. Final Report*. Brussels: Cicca. Available at: <http://www.cicca.eu/download.asp?file=1FinalReportMEDRIL.pdf>; Accessed 14 April 2015
- Stem, R.A. et al. (2008). At the crossroads: development and evaluation of a dementia caregiver group intervention to assist in driving cessation. *Gerontology & Geriatrics Education*, 29, 363-382. doi:10.1080/02701960802497936.
- Stutts, J.C., Stewart, J.R. and Martell, C. (1998). Cognitive test performance and crash risk in an older driver population. *Accident Analysis and Prevention*, 30, 337-346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9663293>.
- US Department of Transportation (2010). *Highway statistics 2009* [Online]. Washington, DC: Federal Highway Administration. Available at: <http://www.fhwa.dot.gov/policyinformation/statistics/2009/022.cfm>; last accessed 25 February 2011.
- Wadley, V.G. et al. (2009). Mild cognitive impairment and everyday function: an investigation of driving performance. *Journal of Geriatric Psychiatry and Neurology*, 22, 87-94. doi:10.1177/0891988708328215.
- Welsh, W.M., Dickerson, M.A. and Paul, R.H. (2005). The relationship of neuropsychological functioning to driving competence in older persons with early cognitive decline. *Archives of Clinical Neuropsychology*, 20, 217-228. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15770793>.

