

**Washington University School of Medicine
Digital Commons@Becker**

Open Access Publications

2016

The myocardial and coronary histopathology and pathogenesis of hypoplastic left heart syndrome

Charles R. Cole

University of Cincinnati

Pirooz Eghtesady

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Cole, Charles R. and Eghtesady, Pirooz, "The myocardial and coronary histopathology and pathogenesis of hypoplastic left heart syndrome." *Cardiology in the Young*.26,1. 19-29. (2016).

https://digitalcommons.wustl.edu/open_access_pubs/4905

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Review Article

The myocardial and coronary histopathology and pathogenesis of hypoplastic left heart syndrome

Charles R. Cole,¹ Pirooz Eghtesady²

¹University of Cincinnati, Cincinnati, Ohio; ²Washington University in St. Louis, St. Louis, Missouri, United States of America

Abstract Hypoplastic left heart syndrome has the greatest mortality rate among all CHDs and without palliation is uniformly fatal. Despite noble efforts, the aetiology of this syndrome is unknown and a cure remains elusive. The genetic and anatomical heterogeneity of hypoplastic left heart syndrome supports a rethinking of old hypotheses and warrants further investigation into the histological and vascular variations recognised with this syndrome. In an effort to elucidate the pathogenesis of hypoplastic left heart syndrome, this review will focus on its unique myocardial and coronary pathology as well as evaluate the association of hypoplastic left heart syndrome with the endocardial fibroblastosis reaction.

Keywords: Hypoplastic left heart syndrome; endocardial fibroblastosis; coronary arteries; myocardial morphology; CHD

Received: 7 April 2015; Accepted: 31 May 2015; First published online: 8 July 2015

Methods

We performed a detailed review of PubMed for articles pertaining to hypoplastic left heart syndrome, which produced over 2700 articles that were then narrowed down to 67 articles based on relevance to histology, pathology, valve, myocardium, coronaries, endocardial fibroblastosis, and pathogenesis. We searched from 1940 until the present. We needed to evaluate articles dating back to 1940 because it was during that time period that much of the histological analysis for hypoplastic left heart syndrome was being performed. We also wanted to know the histologic appearance of specimens before any intervention. We included all articles we could locate pertaining to the pathogenesis of hypoplastic left heart syndrome.

Variable anatomy of hypoplastic left heart syndrome

Hypoplastic left heart syndrome is a severe and devastating heart defect that affects ~1 in 5–10,000

children born each year and accounts for up to 25% of all neonatal deaths from CHD.¹ Hypoplastic left heart syndrome is characterised by a diverse spectrum of malformations distinguished by underdevelopment of the left ventricle and its components, rendering it unable to support systemic circulation.^{2,3} The presence of anatomical variations within the classification of hypoplastic left heart syndrome yields a continuum of phenotypic heterogeneity that can be divided into broad subgroups.^{3,4} These variations are dependent upon the presence or absence of the following: an inlet to the left ventricle, a patent outflow tract, a ventricular septal defect, and/or any other associated cardiac defects. Each subtype can be associated with any atrial arrangement, with *situs solitus* being the most common.³ It is necessary to analyse these subgroups individually because of their differing histological characteristics and the possibility for differing inciting events. Sedrak et al provide an excellent breakdown of hypoplastic left heart syndrome subtypes, which appear in Table 1.

First, hearts with combined mitral and aortic atresia present with a thin-walled, slit-like left ventricle.⁴ The ascending aorta and arch are extremely hypoplastic, and flow is retrograde.⁵ Systemic output is ductal dependent. In the setting of

Correspondence to: P. Eghtesady M.D., PhD, Division of Pediatric Cardiothoracic Surgery, Washington University School of Medicine, Campus Box 8234, St. Louis, MO 63110, United States of America. Tel: +314 454 6165; Fax: +314 454 2381; Email: eghtesady@wudosis.wustl.edu

Table 1. Morphologic subgroups of hypoplastic left heart syndrome.³

- 1 Mitral and aortic atresia
- 2 Mitral atresia with a patent aortic root and ventricular septal defect
- 3 Aortic atresia with patent mitral valve
- 4 Aortic stenosis and dysplasia with patent mitral valve
- 5 Left ventricular hypoplasia with coarctation of the aortic arch

combined mitral and aortic atresia, if a ventricular septal defect is present, a larger left ventricular cavity will develop and the wall of the left ventricle will be proportionately thicker. The larger the ventricular septal defect, the more closely the left ventricle will approach normal dimensions.³ The observation that left ventricular dimensions are proportional to the size of the inlet suggests that myocardial development is dependent on adequate blood flow during development. Second, mitral atresia can also occur with a stenotic aortic valve, patent aortic root, and ventricular septal defect. These cases are characterised by a hypoplastic ascending aorta with a widely patent ductus arteriosus.⁴ In the setting of a large ventricular septal defect, irrespective of mitral valve dimensions, the aortic valve is rarely atretic, which suggests that valvular development is also dependent to some degree on adequate blood flow during development. Third, hearts with isolated aortic atresia and a patent mitral valve without a ventricular septal defect demonstrate thickening of the left ventricular free wall, ventricular septum, and endocardium. As in combined mitral and aortic atresia, the ascending aorta and arch are hypoplastic, and all systemic output is ductal dependent.⁵ Fourth, hypoplastic left heart syndrome hearts with aortic valve stenosis and a patent mitral valve maintain antegrade flow through the aortic valve. The degree of ascending aortic and arch hypoplasia is less than observed with aortic atresia.⁵ Fifth, in general, the degree of aortic arch hypoplasia correlates with the dimensions of the aortic root, but there is one subset of hypoplastic left heart syndrome hearts with severe aortic arch hypoplasia that is out of proportion to the degree of aortic root hypoplasia.^{3,6} This subtype of hypoplastic left heart syndrome can have multiple combinations of mitral and/or aortic valve abnormalities.

Lev et al described some common qualities between subtypes of hypoplastic left heart syndrome. A detailed analysis of 230 hypoplastic left heart syndrome hearts revealed that specimens had an enlarged heart with an apex forming the right ventricle. In all cases the right atrium was hypertrophied and enlarged, the left atrial appendage was small, and the tricuspid orifice was enlarged and impinged upon the infundibular region. The right ventricle was hypertrophied and enlarged.⁷ Although

right ventricular abnormalities are found in hypoplastic left heart syndrome, many of these abnormalities may be the result of the right ventricle compensating for an increased workload secondary to the primary left ventricular malformation.

Myocardial histopathology of normal hearts versus hypoplastic left heart syndrome hearts

Cardiac muscle cell orientation in non-diseased hearts exhibits a similar pattern in the embryo, fetus, child, and adult.⁸ Normally, the major portion of both ventricular walls and the mid portion of the interventricular septum have an orderly parallel arrangement of cells. Myocardial fibre disarray, defined as a lack of orderly parallel arrangement of myocytes, has been described in hypoplastic left heart syndrome. Myocardial fibre disarray can present in one of the following patterns: fibres branching at sharp angles to one another, groups of fibres cut longitudinally interspersed with fibres cut transversely, or fibres forming concentric whorls. These patterns may be found alone or in combination within a given heart.⁹ Myocardial fibre disarray is not unique to hypoplastic left heart syndrome but can be found to some degree in normal hearts, and more extensively in hearts with conditions such as hypertrophic cardiomyopathy,⁹ pulmonary atresia,⁸ and tetralogy of Fallot.¹⁰

Histopathology of left ventricle in hypoplastic left heart syndrome

Certain subtypes of hypoplastic left heart syndrome present with organised myocardial architecture at birth. Autopsy specimens of hearts with combined mitral and aortic atresia collected before the availability of effective surgical palliation demonstrate a normally arranged cell pattern without evidence of myocardial disarray in the left ventricular myocardium and rudimentary septum.⁸ In the event of mitral atresia with a large ventricular septal defect and a fully developed left ventricle chamber, myocardial fibre orientation is normal.⁸ A common finding in both of these cases is that they likely do not have increased left ventricular cavity haemodynamic pressure burden during development. The normal myocardial architecture in these cases also suggests that the aetiology of these cases is likely valvular.

Hypoplastic left heart syndrome specimens that are exposed to increased left ventricular haemodynamic pressure in utero (patent inflow and obstructed outflow) display myocardial disarray as newborns. Hearts with aortic atresia and a patent mitral valve demonstrate marked myocardial fibre disarray of both the ventricular septum and the left ventricular free wall. Connoret al observed that the distribution of

diseases is generally in the inner two-thirds of the myocardium and does not involve the outer compact myocardium or right ventricle.¹¹ Areas of focal calcification or scarring may also involve the subendocardial region of the left ventricle, interventricular septum, and papillary muscles.¹¹ Patients with hypoplastic left heart syndrome and coarctation of the aorta will develop myocardial pathology that correlates with valvular pathology. Postnatally, this subgroup can develop hypertrophy and fibrosis of the left ventricle.³ Characteristics of each subgroup are summarised in Table 2.

A recent histologic analysis of the myocardium of hypoplastic left heart syndrome hearts found large areas of randomly oriented, disorganized bundles of myocytes with variable myocyte size found between bundles in all left ventricle and right ventricle samples (Fig 1).¹⁶ Although subgroup classifications were not documented in this report, all cases presented with myocyte disarray.¹⁶ Previously reported histological studies observed organised myocardium in the subgroups with combined aortic atresia/mitral atresia or mitral atresia with a ventricular septal defect.¹⁷ The obvious difference between the study groups was the age of the patients and the advancements in palliative techniques. In early publications, the patients' mean lifespan was typically 1 week or less versus them our own study group in which the lifespan was 6 days to 10 months.¹⁶ The more modern patients were also exposed to inotropic support. This observation suggests that increased exposure to abnormal blood flow in the neonate will lead to myocardial disorganisation and scar formation in some hearts that perhaps had a histologically normal myocardium at birth. The differing myocyte architecture observed between subtypes of hypoplastic left heart syndrome during development highlights the potential for differing inciting events between subtypes. The cases of hypoplastic left heart syndrome with abnormal myocyte architecture at birth may have a primary myocardial pathogenesis that alters blood flow patterns and results in abnormal development of the immature valve structures. Recent studies have highlighted the possibility that some cases of hypoplastic left heart syndrome may result from abnormal cardiac myocyte proliferation during development.^{18,19}

Endocardial fibroelastosis

In an eloquent review, Lurie²⁰ describes endocardial fibroelastosis as a reaction of the endocardium, not a disease state. Grossly, the reaction is identified as a pearly or opaque white appearance of the endocardium, especially of the ventricles.²⁰ The normal endocardium is transparent and only around 10 µm

in thickness.²¹ Endocardial fibroelastosis is defined by thickening of the endocardium by layers of collagenous and elastic fibres to >20 µm.²⁰ Lurie²⁰ described the endocardial fibroelastosis reaction as a chronologic sequence of smooth muscle hyperplasia followed by their transformation and translocation from the inner, sub-endothelial layers to the outer, juxtamycardial layers (Fig 2). In the foetus, the fibroelastic lining is highly cellular, containing numerous smooth muscle cells and fibroblasts. Antepartum, the cellular characteristics disappear, leaving layers of collagen and elastic fibres.³ The endocardial fibroelastosis reaction is most active during foetal life and during periods of active growth.

Left ventricular endocardial fibroelastosis in hypoplastic left heart syndrome is often found in the setting of a patent mitral valve and severely stenotic or atretic aortic valve. Hearts that develop the endocardial fibroelastosis reaction are always under stress, either from pressure overload due to mechanical obstruction or from volume overload due to cardiac muscle disease.^{20,22} However, not every heart under stress will develop endocardial fibroelastosis.^{4,20} In addition, endocardial fibroelastosis is not seen within the left ventricle in the setting of combined mitral and aortic atresia when blood flow is absent from the left ventricle²³ but may affect the left atrium and mitral valve in the presence of mitral valve obstruction.²³

There is strong evidence that prenatal infection could account for some cases of left ventricular hypoplasia with endocardial fibroelastosis.²⁴ Coxsackie virus and murine virus were recognised as aetiological agents of endocardial fibroelastosis in the 1960s and 1970s.²⁰ These findings were further verified with more recent work identifying murine genome in the majority of hearts with endocardial fibroelastosis that were preserved from an era when murine was a widespread infection.²⁰ The incidence of endocardial fibroelastosis sharply declined with the administration of the murine vaccine.²⁰

Infants with foetal ventricular hypertrophy lack the endocardial fibroelastosis reaction²⁰ even though they are exposed to similar mechanical stressors seen with certain subtypes of hypoplastic left heart syndrome (patient in whom with obstructed outflow) that may present with endocardial fibroelastosis. Animal studies have been performed creating a severe left ventricular outflow tract obstruction in lamb fetuses. All foetuses had elevated end-diastolic pressure, but none developed endocardial fibroelastosis.²⁵ This finding suggests that mechanical stressors alone do not always cause endocardial fibroelastosis. Lurie hypothesised that in the conditions that increase contractility, such as hypertrophic cardiomyopathy, the endocardium may inhibit the endocardial fibroelastosis reaction. It is also possible that in

Table 2. Characteristics of hypoplastic left heart syndrome subtypes in newborns.

Subtype	Frequency	Gross anatomy	Histopathology of left ventricle	Endocardial fibroelastosis	Aortic valve pathology	Coronary artery pathology
Mitral and aortic atresia	36–46% *	Thin-walled, slit-like left ventricle. Right ventricle from cardiac apex	Normal ally arranged cell pattern	Absent	Atritic	Coronary abnormalities rarely documented. Histologically normal coronary arteries
Mitral atresia with patent aortic root and ventricular septal defect	N/A	The larger the ventricular septal defect, the more closely the left ventricle will approach normal dimensions	In the event of a large ventricular septal defect and fully developed ventricular chamber, myocardial fibre orientation will be normal	Absent	Thin friable valve leaflets. Few ventriculo-coronary stules. Usually abnormal with pathology ranging from valve leaflet fusion to severe stenosis	Ventriculo-coronary stules. Left coronary arteries have increased tortuosity, but preserved lumen diameter
Aortic atresia with a patent mitral valve	20–29% *	Thickened left ventricular free wall and thickened septum	Marked myocardial fibre disarray of the left ventricular free wall and septum. Myocyte disarray generally in the inner two-thirds of the myocardium that does not involve the outer compact myocardium. Areas of focal calcification and scarring	Present	Atritic	Common ventriculo-coronary stules, ranging from 32 to 100%. Diseased coronary arteries limited to the left ventricular free wall and septum. Diseased coronary arteries displayed thickening and tortuosity in the epicardial and intramycardial branches. Histologic findings consistent with circumferential medial muscular hypertrophy, reduplicated elastic bays, and focal intimal proliferation without significant luminal narrowing
Aortic valvular stenosis and dysplasia with patent mitral valve	23–26% *	Thickened left ventricular free wall and thickened septum	Marked myocardial fibre disarray of the left ventricular free wall and septum. Areas of calcification and fibrosis	Noular; thickened, and dysplastic valve leaflets. Fusion of valve leaflets	Common ventriculo-coronary stules, ranging from 32 to 100%. Diseased coronary arteries limited to the left ventricular free wall and septum. Diseased coronary arteries displayed thickening and tortuosity in the epicardial and intramycardial branches. Histologic findings consistent with circumferential medial muscular hypertrophy, reduplicated elastic bays, and focal intimal proliferation without significant luminal narrowing	Common ventriculo-coronary stules, ranging from 32 to 100%. Diseased coronary arteries limited to the left ventricular free wall and septum. Diseased coronary arteries displayed thickening and tortuosity in the epicardial and intramycardial branches. Histologic findings consistent with circumferential medial muscular hypertrophy, reduplicated elastic bays, and focal intimal proliferation without significant luminal narrowing
Left ventricular hypoplasia with coarctation of the aorta	24–80% of hypoplastic left heart syndrome cases with coarctation**	Proportional hypoplasia of left ventricle in relation to hypoplasia of valvular structures	Fibrosis may develop	Absent	Fusion of valve leaflets. Valves are not thickened or dysplastic	Proportional coronary abnormalities in relation to hypoplasia of valvular structures

*Frequencies documented before Norwood procedure^{12,13}**Frequencies unrelated to Norwood procedure^{6,14,15}

genetically susceptible patients contributions from both mechanical and immunologic factors are necessary for pathology. In hypoplastic left heart syndrome e, a two-hit phenomenon may contribute to the expression and/or severity of the endocardial fibroelastosis reaction.

In the past, endocardial fibroelastosis was thought to be a primary disease that could potentially lead to left

ventricular hypoplasia, termed the "contracted type" of primary endocardial fibroelastosis.²⁰ The current understanding is that endocardial fibroelastosis does not cause hypoplastic left heart syndrome e,²⁰ but its presence and severity worsen the prognosis.²⁶ M cElhinney et al²⁶ recently reported that the severity of endocardial fibroelastosis, as determined by prenatal echocardiography in patients with aortic stenosis and evolving hypoplastic left heart syndrome e, had an association with postnatal outcome following in utero balloon valvoplasty. Patients with more severe endocardial fibroelastosis had a lower probability of postnatal biventricular outcome. They also reported that from pre-intervention to late gestation the time-indexed change in left ventricular end-diastolic volume was significantly greater in fetuses with mild endocardial fibroelastosis compared with those with severe endocardial fibroelastosis.²⁶ Although our understanding of endocardial fibroelastosis has improved greatly over the last half century, many questions remain regarding the fascinating relationship between hypoplastic left heart syndrome e and endocardial fibroelastosis.




Figure 1. Hypoplastic left heart syndrome. Disorganized bundles of myocytes, large areas of fibrosis, and abnormal vascular formations (left ventricle from 3-month-old hypoplastic left heart syndrome patient, trichrome stain, original magnification 25×). Courtesy of Bahlmeyer.¹⁶

Congenital coronary artery fistulas

We now shift focus to the coronary artery histopathology found in hypoplastic left heart syndrome e and its contribution to the pathogenesis of the syndrome. We will first evaluate coronary artery




Figure 2. (a) Histologic stained section of endocardium (3420), showing dense fibers of collagen and elastin typical of endocardial fibroelastosis. Reproduced from Am J Pathol 1972; 66: 483–496, Figure 3; with permission from The American Society for Investigative Pathology and kind cooperation of G rover M . Hutchins, MD 13. (b) In contrast is the parallel diagram derived from electron microscopy showing cellular activity that generates the fibroelastosis. 1. Dark smooth muscle cell; 2. light smooth muscle cell; 3. light smooth muscle cell with loss of basement membrane; 4. leimyoid cell; and 5. fibroblast. Adapted from the Archives of Pathology Laboratory Medicine 1979; 103: 218, Figure 8; Copyright 1979 American Medical Association. All rights reserved. Courtesy of Luria.²⁰

stulas, followed by macro coronary artery disease or malformations, and will usually evaluate the microvasculature observed in hypoplastic left heart syndrome. A congenital coronary artery stula is an abnormal direct communication between any part of the coronary system and a cardiac chamber or great vessel, having bypassed the myocardial capillary bed.²⁷ Microscopic evaluation of hypoplastic left heart syndrome specimen ens with ventriculo-coronary connections demonstrate that sinusoids from the endocardial surface may extend into the myocardium.¹¹ Blake categorised these ventriculo-coronary connections into three subtypes:

- Arterio-luminal subtype - connects the left ventricular chamber directly to the vertically penetrating branches of the coronary arteries.
- Arterio-sinusoidal subtype - connects the left ventricular chamber indirectly to sinusoids from the endocardial surface of the coronary arteries.
- Arterio-capillary subtype - connects the left ventricular chamber indirectly to a rich network of thin-walled, capillary-sized vessels.

O'Connor et al¹¹ found a variable combination of ventriculo-coronary connections in all cases of hypoplastic left heart syndrome with patent coronary and obstructed outflow. The largest number of connections was found in subgroups with either stenosis or complete closure of the foramen ovale. No ventriculo-coronary connections were identified in the subgroup with mitral and aortic atresia.¹¹ Baffa et al²⁸ found ventriculo-coronary connections in 27 of 89 specimens with a patent mitral valve combined with aortic atresia and in 2 out of 52 specimens with mitral and aortic atresia. Most of the connections in hypoplastic left heart syndrome are the arterio-sinusoidal subtype rather than the direct arterio-luminal subtype.^{28,29} The arterio-sinusoidal subtype frequently coexists with endocardial fibroelastosis of the left ventricle.²⁸ The regional distribution of myocardial disarray supports the concept that vascularisation parallels myocardial organisation in the developing heart.³⁰ It has been postulated that elevated left ventricular intracavitory pressure incites the persistence of embryonic microvasculature, which leads to the creation of complex arterio-luminal connections to allow for the egress of blood.³¹

Coronary artery pathology

Physiologic coronary perfusion occurs in a cyclical pattern, with the major contribution during diastole. In the setting of an anatomically obstructed left ventricle, the coronary perfusion occurs during systole, as demonstrated by selective aortic root angiography in prior studies of hypoplastic left

heart syndrome.³² The high-pressure blood flow during ventricular systole may lead to wall thickening and tortuosity in the epicardial coronary arteries and their intramural myocardial branches^{11,29} (Fig 3). O'Connor et al observed that diseased vessels were limited to the left ventricular free wall and septum. The histological findings consisted of circumferential medial muscular hypertrophy, reduplicated elastic fibres, and focal intimal fibrous proliferation without significant luminal narrowing.¹¹ Sauer et al²⁹ found that specimens with mitral and aortic atresia did not demonstrate histological coronary artery abnormalities.

Coronary artery histopathologic findings are most prominent in arteries that communicate directly with the lumen (arterio-luminal subtype) and less prominently in those that communicate indirectly with the lumen (arterio-sinusoidal and arterio-capillary subtypes).¹¹ Baffa et al²⁸ noted increased tortuosity of the left coronary artery in mitral stenosis/aortic atresia specimens, but observed on histological examination that the ratio of coronary wall thickness relative to lumen diameter in the left anterior descending artery and circumflex artery revealed no significant differences from control hearts. Lloyd et al³³ also observed that the luminal diameters of the coronary arteries and ostia in hypoplastic left heart syndrome were not different from control specimen.

Anomalous coronary artery origins and coronary artery hypoplasia have rarely been reported in hypoplastic left heart syndrome.³⁴ Infrequently, case reports have described an anomalous origin of the left coronary³⁵⁻³⁸ or circumflex^{39,40} artery from the right pulmonary artery in the setting of hypoplastic left heart syndrome. Ito et al⁴¹ reported a case of hypoplastic left heart syndrome with a single coronary artery originating from the pulmonary artery. Saroli et al³⁷ reported three cases of hypoplastic left heart syndrome with superior origin of the left coronary artery. The origin of the right coronary artery from the descending thoracic aorta has been documented in one case of hypoplastic left heart syndrome.⁴² De Rose et al³⁴ described a case of hypoplasia of the left anterior descending artery that resulted in marked left ventricular ischaemia during a Norwood procedure, ultimately leading to the patient's demise.

Anomalous coronary artery stulas have also been described. Raghbir et al⁴³ described a case of hypoplastic left heart syndrome due to aortic atresia with a stula present between the left circumflex coronary artery and the coronary sinus. A recent case report documented a left ventricle-left coronary artery-pulmonary artery stula associated with combined mitral and aortic atresia and a ventricular septal defect.⁴⁴ The authors concluded that the coronary stula was the primary cause of hypoplastic left heart

syndrome in this case because of the existence of a ventricular septal defect that nullified the intraventricular pressure gradient, which is typical of

hypoplastic ventricles and thought to be responsible for the secondary development of stenosis in other cases.⁴⁴

In an autopsy report of 122 patients who died after a Norwood procedure, Batizal et al³⁹ found impairment of coronary artery perfusion to be the most frequent cause of death (33 patients, 27%). However, the cause of stenosis in the vast majority of these patients (31 of 33) was secondary to either intimal stenosis at the anastomosis or external kinking of the graft.^{34,39} The poor coronary perfusion in these cases was therefore the result of the surgical technique and was not the underlying coronary disease. Surgical ligation of an aberrant left circumflex coronary artery from the right pulmonary artery occurred in one case, resulting in biventricular infarction.³⁹ Generalised coronary artery hypoplasia was also found in one patient.³⁹ Despite the rare occurrence of coronary anomalies in hypoplastic left heart syndrome, detailed coronary artery assessment is recommended as part of the routine echocardiographic evaluation of hypoplastic left heart syndrome before surgical intervention.³⁷ The coronary arteries in hypoplastic left heart syndrome may become thickened or tortuous in cases with high intraventricular pressure, but have a preserved lumen, and with rare exception are likely not primary to the pathogenesis of disease.

Microvasculature

Focusing on the microvasculature of hypoplastic left heart syndrome, Salih et al⁴⁵ described an interesting finding that unoperated hearts with hypoplastic left heart syndrome have a higher mean and maximal diffusion distance from any arbitrary point to the nearest capillary than do normal age-matched control hearts. No differences were noted between left and right ventricles or between subtypes. The authors believe the reduction in capillarisation may be an inherent abnormality of hypoplastic left heart syndrome that may have implications for ventricular development. Rakusan et al⁴⁶ noted that congenital




Figure 3.
Coronary artery wall thickening observed in cases of hypoplastic left heart syndrome with patent inflow and obstructed outflow.
(a) Photomicrograph of a branching posterior descending artery in the epicardial groove. Magnification 14 \times . (b) Vessel outlined in (a) shows a prominent muscolastic zone (M). Eccentric intimal thickening (I) is present but does not appear to significantly narrow the calibre of the lumen. Magnification 120 \times . (c) Low-power view of interventricular septum with prominent muscularized arteries (arrows). Magnification 10 \times . All elastic stain. Courtesy of O'Connor.¹¹ LV = left ventricle; PD = patent ductus; RV = right ventricle.

aortic stenosis and coarctation of the aorta are characterised by an increase in capillary supply proportional to myocyte volume, maintaining capillary density similar to control hearts. They observed that pressure-overload left ventricular hypertrophy in children demonstrates proportional capillary angiogenesis, whereas in adults hypertrophy appears to be associated with failure of compensatory angiogenesis.

Could the pathogenesis of hypoplastic left heart syndrome stem in part from premature failure of compensatory angiogenesis? Jacobs suggested that the decreased capillarisation observed by Salih et al might not be an unalterable inherent abnormality of hearts with hypoplastic left heart syndrome, but rather a snapshot of the supply-demand mismatch encountered with unaltered hypoplastic left heart syndrome anatomy. As stated previously, perfusion of the myocardium in unoperated hearts with hypoplastic left heart syndrome occurs primarily in systole, rather than in diastole, as there is considerable diastolic runoff into the pulmonary circulation. Coronary perfusion is further limited by excessive myocardial wall tension.⁴⁷ Elimination of the diastolic runoff, by means of closure of the systemic to pulmonary artery shunt at the time of the second-stage surgery, results in the restoration of more normal diastolic coronary perfusion as well as reduction in the volume of work of the ventricle.⁴⁷ Jacobs⁴⁷ proposes that this higher diastolic blood pressure may result in improved capillarisation of the myocardium early in life. Further research is necessary to confirm or refute this theory, but it is clear that there are primary myocardial differences in capillary density in hypoplastic left heart syndrome compared with other left-sided CHDs, which further suggests that alterations in the myocardial development of hypoplastic left heart syndrome could contribute to the observed phenotype.

Inappropriate expression of platelet-endothelial cell adhesion molecule-1, also known as CD 31, has been identified in hearts samples with hypoplastic left heart syndrome¹⁶ and may give some insight into myocardial differences in capillary density, or capillary angiogenesis, observed in hypoplastic left heart syndrome. CD 31 is a member of the cell adhesion molecule family, which has a primary role in the regulation of tissue morphogenesis.¹⁶ Before a recent study, cardiac myocytes have never been shown to express CD 31 at any stage of development or in any disease state.¹⁶ The authors suggest that inappropriate expression of CD 31 or a related gene under the same regulatory control may be responsible for the disorganisation of hypoplastic left heart syndrome cardiac myocytes and potentially the higher-order cardiac structural abnormalities associated with this disease.¹⁶

CD 31 has also been associated with cell migration and cancer angiogenesis.⁴⁸ Recent work suggests that CD 31 has a novel role in arteriogenesis and collateral remodelling.⁴⁹ CD 31 has also been identified as the first molecule that determines pre-existing collateral diameter.⁴⁹ These findings highlight the possibility that abnormal CD 31 expression might be an attempt by the myocardium to induce angiogenesis secondary to the decreased capillarisation seen in hypoplastic left heart syndrome. Alternatively, the atypical capillarisation in hypoplastic left heart syndrome could be the product of abnormal CD 31 expression within the myocardium.

Pathogenesis of hypoplastic left heart syndrome

The aetiological mechanisms leading to hypoplastic left heart syndrome are largely unknown. About one-fourth of hypoplastic left heart syndrome cases occur in the context of recognised genetic disorders including, but not limited to, Turner, Jacobsen, Noonan, and Holt-Oram syndromes. However, studies involving non-syndromic families members have suggested that heritability is complex. No single disease-causing pathway has yet been identified.^{50,51} Clinical observations indicate that obstruction of blood flow through the left ventricle in an otherwise normal four-chambered fetal heart, caused by either aortic and/or mitral stenosis, leads to the development of left ventricular hypoplasia.⁵² Prenatal diagnosis of hypoplastic left heart syndrome can be made as early as 14 weeks' gestation,⁵³ but numerous cases of hypoplastic left heart syndrome have presented with normal or even dilated left ventricular cavities on routine fetal ultrasound at >19 weeks' gestation.^{52,54} The unifying aetiological explanation is that the growth and development of vascular structures are dependent to some degree on the relative quantity and quality of blood flow during development.³ Recent studies have shown that hypoplastic left heart syndrome myocytes are well differentiated, but have prolonged expression of foetal or 'heart failure' genes.¹⁶ These findings suggest that intrauterine insult to the foetus may occur after embryogenesis and highlights the possibility that foetal development of hypoplastic left heart syndrome may occur later in gestation.^{55,56} Recent hypotheses also question whether immunologic,⁵⁷ infectious,^{3,58} or autoimmune insults to genetically susceptible hosts may contribute to left ventricular hypoplasia from either direct myocardial injury or secondary to reduced left ventricular blood flow through damaged valves.

Genetic factors that alter valve development have been proposed as the aetiology in hypoplastic left heart syndrome.^{59,60} Mutations in the signalling and transcription regulator, NOTCH 1, have caused early

aortic valve defects in animal models.⁵ Genes involved in downstream NOTCH signalling and cardiac gene expression have been implicated in familial forms of hypoplastic left heart syndrome.⁶¹ A recent study showed a shared genetic linkage with hypoplastic left heart syndrome and bicuspid aortic valve.⁶² Bicuspid aortic valve is very common, affecting 1% of the general population, and is a known risk factor for aortic valve disease. Is it possible that a small subset of patients with most restrictive bicuspid aortic valves progress to hypoplastic left heart syndrome? Isolated aortic valve atresia and congenital aortic stenosis do not always progress to hypoplastic left heart syndrome, suggesting that aortic valve defects and hypoplastic left heart syndrome have distinct genetic susceptibilities.⁵⁰ Genetic factors are clearly present, but epigenetic mechanisms and/or environmental influences might be necessary for the phenotypic expression of hypoplastic left heart syndrome.⁵⁰

The complex heritability of hypoplastic left heart syndrome suggests the potential for environmental contributions to the observed phenotype. Maternal upper respiratory infection during the first trimester has been documented as a significant risk factor for hypoplastic left heart syndrome.⁶³ Population-based studies in the Baltimore-Washington region have identified environmental risk factors for hypoplastic left heart syndrome, including maternal exposure to organic solvents.⁶⁴ A recent study showed a seasonal pattern in the presentation of hypoplastic left heart syndrome with preponderance in summer months in contrast to the random pattern observed in other left-sided heart diseases.⁶⁵ These findings support a role for an environmental component in the aetiology of hypoplastic left heart syndrome.

An immune-mediated mechanism for the pathogenesis of hypoplastic left heart syndrome has been proposed where antibodies cross the placenta and create disease in genetically susceptible hosts.⁵⁷ In rheumatic heart disease, antibodies have already been shown to "cross-react" with human valvular and myocardial antigens through a mechanism known as molecular mimicry.⁶⁶ Our lab recently demonstrated that transplacental transfer of maternal anti-cardiac myosin autoantibodies leads to structural congenital cardiac defects in affected progeny that included diminished left ventricular cavity dimensions.⁶⁷ Foetuses that developed hypoplastic left heart syndrome-like phenotype had elevated serum titres of anti- β adrenergic receptor as well as increased protein kinase A activity, suggesting a potential mechanism for the observed pathological changes.⁶⁷

There is evidence that prenatal viral infection could account for some cases of left ventricular hypoplasia with endocardial fibroelastosis.^{3,24,58} Viral insult is a

well-accepted cause of cardiomopathy. Studies have been reported that viral RNA is present in myocardial samples from foetal hearts with aortic valvar stenosis.³

Alternative mechanisms for the pathogenesis of hypoplastic left heart syndrome have been suggested in which abnormal cardiac myocyte proliferation may be the primary defect. In this case, left ventricular hypoplasia could result from abnormal cellular signal transduction, which is critical for normal myocyte division and left ventricular growth.¹⁹ Conversely, the thickening of the left ventricular myocardium in subtypes of hypoplastic left heart syndrome with a patent ductus and obstructed outflow tract is thought to result from cardiac myocyte hyperplasia during development.^{3,67} The abnormal thickening of the left ventricular myocardium could lead to reduced left ventricular cavity dimensions, and valvular abnormalities as a result of altered haemodynamics. Many primary myocardial abnormalities have been identified that could contribute to the pathogenesis of hypoplastic left heart syndrome.

Conclusion

Because of the variability in subtypes and presentation of hypoplastic left heart syndrome, a common aetiology for all subtypes is unlikely. Genetic and environmental factors are likely working in concert to create the spectrum of phenotype observed. Cases of hypoplastic left heart syndrome with mitral atresia and aortic atresia or with mitral atresia, a ventricular septal defect, and an intact aortic root have been shown to have normal myocardial architecture at birth and normal coronary vasculature, which suggests that these subtypes are likely due to primary valvular pathogenesis. Future research efforts in these subtypes should look for genetic and environmental causes of valvular agenesis. Subtypes of hypoplastic left heart syndrome with patent ductus and obstructed outflow present with disparity of the myocardial architecture and thickening of the myocardium. These hearts are also more prone to vascular abnormalities, such as coronary artery stulas. As stated previously, the regional distribution of myofoils supports the concept that vascularisation parallels myocardial organisation in the developing heart. Future studies should evaluate for primary myocardial disorders or dysregulation of cardiac myocyte proliferation during development for these subtypes. In addition, some phenotypes of hypoplastic left heart syndrome, most likely the subtypes with an atretic valve and thickened ventricle, may require a combined insult to both the myocardium and valve structures. Furthermore, continued investigation into the reduction in capillarisation in hypoplastic left heart syndrome and the potential contribution of CD 31 in

this abnormality is warranted. If the pathogenesis of this disease is uncovered, the possibility for more effective treatment or perhaps even the prevention of certain subtypes of hypoplastic left heart syndrome may one day become reality.

Acknowledgment

The authors would like to thank Anoop Brar PhD and R. Scott Baker for their contribution in editing this review.

Financial Support

Financial support was provided by NIH R21-HL104391, NIH F32-HL103054 and Saving Tiny Hearts Society.

Conflicts of Interest

The authors have no relationships with industry or financial associations that might pose a conflict of interest with the submitted article.

References

1. Rosamond W, Flegal K, Furukawa T, et al. Heart disease and stroke statistics—2008 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2008; 117:e25–e146.
2. Barron DJ, Kirby MD, Davies B, Wright JS, Jones TJ, Brown WJ. Hypoplastic left heart syndrome. *Lancet* 2009; 374: 551–564.
3. Seward J, Cook AC, Shiralgi M, McQuinn TC. Current issues and perspectives in hypoplasia of the left heart. *Cardiol Young* 2005; 15:56–72.
4. Sinha SN, Rusnak SL, Sommers H, Cole RB, Muster AJ, Paul MH. Hypoplastic left ventricle syndrome. Analysis of thirty autopsy cases in infants with surgical considerations. *Am J Cardiol* 1968; 21:166–173.
5. Tweddell JS, Hoffman GM, et al. Hypoplastic left heart syndrome: morphological study of 21 autopsy cases. In: Allen HD, Driscoll DJ, Shaddy BE, Roberts RJ, Fetterley F (eds). *Modern Advances in Infants, Children and Adolescents*. Lippincott William and Wilkins, October 16, 2007: 1005–1038.
6. Szabolcs J, Skarzynski D, Swatek J, Karbowicz E. Hypoplastic left heart syndrome: morphological study of 21 autopsy cases. *Ann Univ Mariae Curie Skłodowska Med* 2002; 57:278–284.
7. Bhattacharyya M, Lev M. The surgical anatomy of hypoplasia of aortic tract complex. *J Thorac Cardiovasc Surg* 1984; 88:97–101.
8. Bulkley BH, Weissfeldt ML, Hutchins GM. Asymmetric septal hypertrophy and myocardial ber disease. Features of normal developing, and malformed hearts. *Circulation* 1977; 56: 292–298.
9. Bulkley BH, Damico B, Taylor AL. Extensive myocardial ber disease in aortic and pulmonary atresia. Relevance to hypertrophic cardiomyopathy. *Circulation* 1983; 67:191–198.
10. Jones M, Frazee VJ, Monrow AG, Roberts W C U. Infrastructure of crista supraventricularis muscle in patients with congenital heart diseases associated with right ventricular outflow tract obstruction. *Circulation* 1975; 51:39–67.
11. O'Connor WN, Cash JB, Cottrill CM, Johnson GL, Nicanor JA. Ventriculoauricular connections in hypoplastic left heart: an autopsy/micropscopic study. *Circulation* 1982; 66:1078–1086.
12. Mahle WT, Spritz TL, Weintraub G, Gaynor JW, Clark BJ 3rd. Survival after reconstructive surgery for hypoplastic left heart syndrome: a 15-year experience from a single institution. *Circulation* 2000; 102: III36–III41.
13. Tweddell JS, Hoffman GM, Muzatko KA, et al. Improved survival of patients undergoing palliation of hypoplastic left heart syndrome: lessons learned from 115 consecutive patients. *Circulation* 2002; 106: 182–189.
14. Hawkins JA, Clark EB, Doty DB. Morphologic characteristics of aortic atresia: implications for fetal hemodynamics. *Int J Cardiol* 1986; 10:127–132.
15. Von Rueden TJ, Knight L, Mueller JH, Edwards JE. Coarctation of the aorta associated with aortic valvular atresia. *Circulation* 1975; 52: 951–954.
16. Buhmeyer TJ, Helmke S, Ge S, et al. Hypoplastic left heart syndrome: myocytes are differentiated but possess a unique phenotype. *Cardiovasc Pathol* 2003; 12:23–31.
17. Bulkley BH, Weissfeldt ML, Hutchins GM. Isometric cardiac contraction. A possible cause of the disorganized myocardial pattern of idiopathic hypertrophic subaortic stenosis. *N Engl J Med* 1977; 296:135–139.
18. de la Sierra A, McQuinn T, Sedmera D. Increased ventricular preload is compensated by myocyte proliferation in normal and hypoplastic fetal chick left ventricle. *Circ Res* 2007; 100: 1363–1370.
19. Grossfeld P. Hypoplastic left heart syndrome: new insights. *Circ Res* 2007; 100:1246–1248.
20. Lurie PR. Changing concepts of endocardial fibroelastosis. *Cardiology* 2010; 20:115–123.
21. Neustein HB, Lurie PR, Fugita M. Endocardial fibroelastosis found on transmural endomyocardial biopsy in children. *Arch Pathol Lab Med* 1979; 103:214–219.
22. Black-Schaffer B. Infantile endocardial fibroelastosis; a suggested etiology. *Am J Arch Pathol* 1957; 63:281–306.
23. Nicanor JA, Nadas AS. The hypoplastic left heart syndrome: an analysis of 101 cases. *Pediatr Clin North Am* 1958; 5:1029–1056.
24. Ni J, Bowles NE, Kim YH, et al. Viral infection of the myocardium in endocardial fibroelastosis. Molecular evidence for the role of mumps virus as an etiologic agent. *Circulation* 1997; 95: 133–139.
25. Eighty P, Michaelson E, Altaye M, Ballard E, Hirsh R, Beekman RH 3rd. Revisiting animal models of aortic stenosis in the early gestation fetus. *Ann Thorac Surg* 2007; 83:631–639.
26. McElhinney DB, Vogel M, Benson CB, et al. Assessment of left ventricular endocardial fibroelastosis in fetuses with aortic stenosis and evolving hypoplastic left heart syndrome. *Am J Cardiol* 2010; 106:1792–1797.
27. Mavridis C, Becker CL, Roachini AP, Muster AJ, Gevirtz M. Coronary artery stulas in infants and children: a surgical review and discussion of coil embolization. *Ann Thorac Surg* 1997; 63: 1235–1242.
28. Baffa JM, Chen SL, Guttenberg ME, Nicasio W, Weinberg PM. Coronary artery abnormalities and right ventricular histology in hypoplastic left heart syndrome. *J Am Coll Cardiol* 1992; 20: 350–358.
29. Sauer U, Geitzenberger-de Groot AC, Geishauser M, Babic R, Buhmeyer K. Coronary arteries in the hypoplastic left heart syndrome. Histopathologic and histomorphological studies and implications for surgery. *Circulation* 1989; 80:1168–1176.
30. O'Connor WN, Stahr BJ, Cottrill CM, Todd EP, Nicanor JA. Ventriculoauricular connections in hypoplastic right heart syndrome: autopsy serial section study of six cases. *J Am Coll Cardiol* 1988; 11:1061–1072.

31. Beckman CB, Moller JH, Edwards JE. A rare pathway to pulmonary venous flow in left-sided obstructive anomalies. *Circulation* 1975;52:509-516.
32. Freedman RM, Culham JA, Mees CA, Harrington DP. Selective aortic root angiography in the hypoplastic left heart syndrome e. *Eur J Cardiol* 1976;4:25-29.
33. Lloyd TR, Evans TC, Marvin W Jr. Morphologic determinants of coronary blood flow in the hypoplastic left heart syndrome e. *Am Heart J* 1986;112:666-671.
34. DeRoege JJ Jr, Corliss R, Dirscholl M R, Eleazar J, Mosa RS. Isolated left ventricular ischaemia after the Norwood procedure. *Ann Thorac Surg* 2002;73:657-659.
35. Malek E, Mroczek T, Pajak J, Zajc A, Kolcz J, Januszewska K. Hypoplastic left heart syndrome with an anomalous origin of the left coronary artery. *Ann Thorac Surg* 2001;72:2129-2130.
36. Sandusky GE, Dunnam-Welch JJ, Ebaid M R, Letson LA, Mee RB. Anomalous origin of left coronary from right pulmonary artery in hypoplastic left heart syndrome e. *Ann Thorac Surg* 1997;64:836-838.
37. Sarlik T, Gelehrter S, Gammie Fifer CA, van der Velde M E, Bove EL, Ensing GJ. Anomalous origin of left coronary artery affecting surgical repair of hypoplastic left heart syndrome e and shone complex. *Echocardiography* 2008;25:727-731.
38. Nevels M, Ommer IC, Ponriban R. Hypoplastic left heart syndrome with anomalous origin of left coronary artery from the right pulmonary artery: successful surgical treatment in a neonate. *Eur J Cardiothorac Surg* 2005;28:497-498.
39. Bartizal U, Grunenfelder J, Van Praagh R. Causes of death after the modified Norwood procedure: a study of 122 postoperative cases. *Ann Thorac Surg* 1997;64:1795-1802.
40. Villa E, Biancaccio G, Caputo A, Francalanci P, Di Donato RM. Circumflex coronary artery from right pulmonary artery in hypoplastic left heart syndrome e. *Ann Thorac Surg* 2005;80:1919-1920.
41. Ito T, Nitobe M, Ishikawa J, et al. Hypoplastic left heart syndrome with a single coronary artery originating from the pulmonary artery. *Acta Paediatr Jpn* 1995;37:61-63.
42. Cheetham JP, Ruyledge NA, McDonald BM, Gunn G E. Origin of the right coronary artery from the descending thoracic aorta: angiographic diagnosis and unique coronary artery anatomy at autopsy. *Cathet Cardiovasc Diagn* 1987;13:321-324.
43. Raghib G, Blomendal RD, Kanjilal VI, Edwards JE. Aortic atresia and premature closure of foramen ovale. Myocardial sinusoids and coronary arteriovenous sulae serving as flow channel. *Am Heart J* 1965;70:476-480.
44. Paladini D, Camuso G, Sgavaro G, Lepidula C, Parisi G, Maitinelli P. Hypoplastic left heart syndrome associated with left ventricle-coronary artery-pulmonary artery stula: prenatal diagnosis and histological correlation. *Ultrasound Obstet Gynecol* 2005;26:78-80.
45. Salih C, Sheppard MN, Ho SY. Morphometry of coronary capillaries in hypoplastic left heart syndrome e. *Ann Thorac Surg* 2004;77:903-907; discussion 907.
46. Rakusan K, Flanagan MF, Gerva T, Southern J, Van Praagh R. Morphometry of human coronary capillaries during normal growth and the effect of age in left ventricular pressure-overload hypertrophy. *Circulation* 1992;86:38-46.
47. Jacobs M L. Invited commentary. *Ann Thorac Surg* 2004;77:907.
48. Delisser HM, Christodoulou M, Stricker RM, et al. Involvement of endothelial PECAM-1/CD31 in angiogenesis. *Am J Pathol* 1997;151:671-677.
49. Chen Z, Rubin J, Tzien R. Role of PECAM-1 in arteriogenesis and sprouting of preexisting collaterals. *Circ Res* 2010;107:1355-1363.
50. Grossfeld P, Ye M, Harvey R. Hypoplastic left heart syndrome e: new genetic insights. *J Am Coll Cardiol* 2009;53:1072-1074.
51. Grossfeld PD. Hypoplastic left heart syndrome e: it is all in the genes. *J Am Coll Cardiol* 2007;50:1596-1597.
52. Allen LD, Sharland G, Tynan M J. The natural history of the hypoplastic left heart syndrome e. *Int J Cardiol* 1989;25:341-343.
53. Carvalho JS. Early prenatal diagnosis of major congenital heart defects. *Curr Opin Obstet Gynecol* 2001;13:155-159.
54. Anderson NG, Brown J. Normal size left ventricle on antenatal scan in lethal hypoplastic left heart syndrome e. *Pediatr Radiol* 1991;21:436-437.
55. Makikallio K, McElhinney DB, Levine JC, et al. Patent aortic valve stenosis and the evolution of hypoplastic left heart syndrome e: patient selection for fetal intervention. *Circulation* 2006;113:1401-1405.
56. Tworetzky W, Wilkins-Hughes L, Jennings RW, et al. Balloon dilation of severe aortic stenosis in the fetus: potential for prevention of hypoplastic left heart syndrome e: candidate selection, technique, and results of successful intervention. *Circulation* 2004;110:2125-2131.
57. Eighty P. Hypoplastic left heart syndrome e: rheumatic heart disease of the fetus? *Mol H Hypotheses* 2006;66:554-565.
58. Oyer CE, Ongpin EH, Ni J, Bowles NE, Towbin JA. Fatal intrauterine adenosivvalvular endocarditis with aortic and pulmonary valve stenosis: diagnosis by polymerase chain reaction. *Hum Pathol* 2000;31:1433-1435.
59. Hor KN, Border W L, Cripe LH, Benson DW, Hinton RB. The presence of bicuspid aortic valve does not predict ventricular septal defect type. *Am J Med Genet A* 2008;146A:3202-3205.
60. Hinton RB Jr, Martin LJ, Tabangin M E, Mawim L, Cripe LH, Benson DW. Hypoplastic left heart syndrome e is heritable. *J Am Coll Cardiol* 2007;50:1590-1595.
61. Dafys S, Cserjesi P, Makhani BE, Molkentin JD. The transcription factors GATA4 and dHAND physically interact to synergistically activate cardiac gene expression through a p300-dependent mechanism. *J Biol Chem* 2002;277:24390-24398.
62. Hinton RB, Martin LJ, Ramagowda S, Tabangin M E, Cripe LH, Benson DW. Hypoplastic left heart syndrome e links to chromosomal 6q and 6qq isogenetically related to bicuspid aortic valve. *J Am Coll Cardiol* 2009;53:1065-1071.
63. Tikkkanen J, Heinonen OP. Risk factors for hypoplastic left heart syndrome e. *Teratology* 1994;50:112-117.
64. Kuehl KS, Loffredo CA. A cluster of hypoplastic left heart malformations in Baldwin County, Maryland. *Pediatr Cardiol* 2006;27:25-31.
65. Eighty P, Biar A, Hall M. Seasonality of hypoplastic left heart syndrome e in the United States: a 10-year time-series analysis. *J Thorac Cardiovasc Surg* 2011;141:432-438.
66. Cunningham MW. Autoimmunity and molecular mimicry in the pathogenesis of post-streptococcal heart disease. *Front Biosci* 2003;8:s533-s543.
67. Cole CR, Yutzy KE, Biar AK, et al. Congenital heart disease linked to maternal autoimmunity against cardiac myosin. *J Immunol* 2014;192:4074-4082.