

2016

Open source drug discovery with the Malaria Box compound collection for neglected diseases and beyond

Na Le Dang

Washington University School of Medicine in St. Louis

Tyler B. Hughes

Washington University School of Medicine in St. Louis

Donatella Taramelli

Washington University School of Medicine in St. Louis

et al.

Follow this and additional works at: http://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Dang, Na Le; Hughes, Tyler B.; Taramelli, Donatella; and et al., "Open source drug discovery with the Malaria Box compound collection for neglected diseases and beyond." *PLoS Pathogens*. 12,7. e1005763. (2016).
http://digitalcommons.wustl.edu/open_access_pubs/5248

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

RESEARCH ARTICLE

Open Source Drug Discovery with the Malaria Box Compound Collection for Neglected Diseases and Beyond

Wesley C. Van Voorhis^{1*}, John H. Adams², Roberto Adelfio^{3,4}, Vida Ahyong⁵, Myles H. Akabas⁶, Pietro Alano⁷, Aintzane Alday⁸, Yesmalie Alemán Resto⁹, Aishah Alsibae¹⁰, Ainhoa Alzualde⁸, Katherine T. Andrews^{11,12}, Simon V. Avery¹³, Vicky M. Avery¹¹, Lawrence Ayong¹⁴, Mark Baker¹⁵, Stephen Baker^{16,17,18}, Choukri Ben Mamoun¹⁹, Sangeeta Bhatia²⁰, Quentin Bickle²¹, Lotfi Bounaadja²², Tana Bowling²³, Jürgen Bosch²⁴, Lauren E. Boucher²⁴, Fabrice F. Boyom²⁵, Jose Brea²⁶, Marian Brennan¹⁰, Audrey Burton²³, Conor R. Caffrey²⁷, Grazia Camarda⁷, Manuela Carrasquilla^{28ma}, Dee Carter²⁹, Maria Belen Cassera³⁰, Ken Chih-Chien Cheng³¹, Worathad Chindaudomsate³², Anthony Chubb¹⁰, Beatrice L. Colon³³, Daisy D. Colón-López²⁴, Yolanda Corbett³⁴, Gregory J. Crowther¹, Noemi Cowan^{3,4}, Sarah D'Alessandro³⁴, Na Le Dang³⁵, Michael Delves³⁶, Joseph L. DeRisi⁵, Alan Y. Du³⁷, Sandra Duffy¹¹, Shama Abd El-Salam El-Sayed^{38,39}, Michael T. Ferdig⁴⁰, José A. Fernández Robledo⁹, David A. Fidock⁴¹, Isabelle Florent²², Patrick V. T. Fokou²⁵, Ani Galstian⁴², Francisco Javier Gamo⁴³, Suzanne Gokool⁴⁴, Ben Gold⁴⁵, Todd Golub⁴², Gregory M. Goldgor⁴⁶, Rajarshi Guha³¹, W. Armand Guiguemde⁴⁷, Nil Gural²⁰, R. Kiplin Guy⁴⁷, Michael A. E. Hansen¹⁴, Kirsten K. Hanson^{48,49}, Andrew Hemphill⁵⁰, Rob Hooft van Huijsduijnen⁵¹, Takaaki Horii⁵², Paul Horrocks⁵³, Tyler B. Hughes³⁵, Christopher Huston⁵⁴, Ikuo Igarashi³⁸, Katrin Ingram-Sieber^{3,4}, Maurice A. Itoe⁴⁹, Ajit Jadhav³¹, Amornrat Naranuntarat Jensen⁵⁵, Laran T. Jensen³², Rays H. Y. Jiang², Annette Kaiser⁵⁶, Jennifer Keiser^{3,4}, Thomas Ketas⁴⁵, Sebastien Kicka⁵⁷, Sunyoung Kim⁵⁸, Kieran Kirk⁵⁹, Vidya P. Kumar¹⁹, Dennis E. Kyle², Maria Jose Lafuente⁴³, Scott Landfear⁶⁰, Nathan Lee⁵¹, Sukjun Lee¹⁴, Adele M. Lehane⁵⁹, Fengwu Li⁶⁰, David Little⁴⁵, Liqiong Liu⁵⁸, Manuel Llinás²⁸, Maria I. Loza²⁶, Aristeo Lubar⁶¹, Leonardo Lucantoni¹¹, Isabelle Lucet⁶², Louis Maes⁶³, Dalu Mancama⁶⁴, Nuha R. Mansour²¹, Sandra March²⁰, Sheena McGowan⁶⁵, Iset Medina Vera⁴⁹, Stephan Meister³⁷, Luke Mercer²³, Jordi Mestres⁶⁶, Alvine N. Mfopa²⁵, Raj N. Misra⁶⁷, Seunghyun Moon¹⁴, John P. Moore⁴⁵, Francielly Morais Rodrigues da Costa⁶⁸, Joachim Müller⁵⁰, Arantza Muriana⁸, Stephen Nakazawa Hewitt¹, Bakela Nare²³, Carl Nathan⁴⁵, Nathalie Narraidoo¹³, Sujeevi Nawaratna^{11,12}, Kayode K. Ojo¹, Diana Ortiz⁶⁰, Gordana Panic^{3,4}, George Papadatos⁶⁹, Silvia Parapini³⁴, Kailash Patra⁶¹, Ngoc Pham¹¹, Sarah Prats⁴³, David M. Plouffe⁷⁰, Sally-Ann Poulsen¹¹, Anupam Pradhan², Celia Quevedo⁸, Ronald J. Quinn¹¹, Christopher A. Rice², Mohamed Abdo Rizk^{38,71}, Andrea Ruecker³⁶, Robert St. Onge⁷², Rafaela Salgado Ferreira⁷³, Jasmeet Samra²⁸, Natalie G. Robinett^{24,74}, Ulrich Schlecht⁷², Marjorie Schmitt⁷⁴, Filipe Silva Villela⁷³, Francesco Silvestrini⁷, Robert Sinden⁷⁵, Dennis A. Smith⁷⁶, Thierry Soldati⁵⁷, Andreas Spitzmüller⁶⁶, Serge Maximilian Stamm²⁴, David J. Sullivan⁷⁷, William Sullivan⁷⁸, Sundari Suresh⁷³, Brian M. Suzuki²⁷, Yo Suzuki⁷⁹, S. Joshua Swamidass³⁶, Donatella Taramelli³⁵, Lauve R. Y. Tchokouaha²⁵, Anjo Theron⁶⁴, David Thomas⁴², Kathryn F. Tonissen^{11,80}, Simon Townson⁴⁴, Abhai K. Tripathi⁷⁷, Valentin Trofimov⁵⁷, Kenneth O. Udenze², Imran Ullah⁵³, Cindy Vallieres¹³, Edgar Vigil³⁷, Joseph M. Vinetz⁶¹, Phat Voong Vinh¹⁶, Hoan Vu¹¹, Nao-aki Watanabe⁵², Kate Weatherby²⁹, Pamela M. White⁷⁸, Andrew F. Wilks^{81,82}, Elizabeth A. Winzeler³⁷, Edward Wojcik⁵⁸, Melanie Wree³⁷, Wesley Wu⁵, Naoaki Yokoyama³⁸, Paul H. A. Zollo²⁵, Nada Ablal⁵¹, Benjamin Blasco⁵¹, Jeremy Burrows⁵¹, Benoît Laleu⁵¹, Didier Leroy⁵¹, Thomas Spangenberg^{51ab}, Timothy Wells⁵¹, Paul A. Willis⁵¹



OPEN ACCESS

Citation: Van Voorhis WC, Adams JH, Adelfio R, Ahyong V, Akabas MH, Alano P, et al. (2016) Open Source Drug Discovery with the Malaria Box Compound Collection for Neglected Diseases and Beyond. *PLoS Pathog* 12(7): e1005763. doi:10.1371/journal.ppat.1005763

Editor: Margaret A Phillips, U Tex SouthWestern, UNITED STATES

Received: April 1, 2016

Accepted: June 21, 2016

Published: July 28, 2016

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files. Much of the data also appears on ChemBL <https://www.ebi.ac.uk/chembl/>

Funding: Thanks to the UK DFID and the Bill and Melinda Gates Foundation Grand Challenges Explorations for providing funding for testing of the Malaria Box and funding the support of individual groups including: Medicines for Malaria Venture MMV Challenge Grant, Grant Numbers MMV 12/0048 and MMV 12/0076 (to JHA), the Australian Research Council (FT10100185 to SAP; FT0991213 to KTA

1 Departments of Medicine, Microbiology, and Global Health, Center for Emerging and Re-emerging Infectious Diseases (CERID) University of Washington, Seattle, Washington, United States of America, 2 Center for Global Health and Infectious Diseases Research, Department of Global Health, University of South Florida, Tampa, Florida, United States of America, 3 Medical Parasitology and Infection Biology,

and LP120200557 awarded to VMA), Bill & Melinda Gates Foundation Grant OPP1040394 to PA, OPP1040399 to DAF and VMA and OPP1086189 to KKH, OPP1069393 and OPP1119049 to ML, OPP1024029 to CN, the Bloomberg Family Foundation (JBr), JHMRI for a predoctoral fellowship, the US NIH for the CBI training grant T32GM080189 (to LEB), R01GM104486 (to PAW & WS), R01AI117017 (to JHA) the National Science Foundation Graduate Research Fellowship Program Grant No.DGE-1232825 (DDCL), the South African Medical Research Council Strategic Health Innovation Partnerships (grant V6YBT51 to DM) and the Council for Scientific and Industrial Research (grant V1YTB95, to DM), and the French ANR program Mammamia (ANR-12-BS07-0020-01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The following commercial organizations employ or employed some of the authors, which might be considered a conflict of interest by some readers: BBD BioPhenix SL—BIOBIDE: Aintzane Alday PhD, Ainhoa Alzualde PhD, and Arantza Muriana; Celia Quevedo PhD; SCYNEXIS, Inc.: Tana Bowling, Audrey Burton, Luke Mercer, and Bakela Nare PhD; GlaxoSmithKline: Francisco Javier Gamo, PhD, Maria Jose Lafuente, PhD, and Sarah Prats; Eisai Co., Ltd.: Takaaki Horii Ph.D. and Nao-aki Watanabe Ph.D.; Novartis Inc.: Mark Baker PhD MSc (med), and David M. Plouffe; Definiens AG: Andreas Spitzmüller PhD; and, Merck Serono Inc: Thomas Spangenberg PhD.

Swiss Tropical and Public Health Institute, Basel, Switzerland, **4** University of Basel, Basel, Switzerland, **5** Howard Hughes Medical Institute, Department of Biochemistry and Biophysics, University of California, San Francisco, California, United States of America, **6** Departments of Physiology & Biophysics, Neuroscience and Medicine, Albert Einstein College of Medicine, New York, New York, United States of America, **7** Dipartimento Malattie Infettive, Parassitarie ed Immunomediate Istituto Superiore di Sanità, Roma, Italia, **8** BBD BioPhenix SL—BIOBIDE, Donostia, Gipuzkoa, Spain, **9** Bigelow Laboratory for Ocean Sciences, East Boothbay, Maine, United States of America, **10** Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, Dublin, Ireland, **11** Eskitis Institute for Drug Discovery, Griffith University, Nathan, QLD, Australia, **12** QIMR Berghofer Medical Research Institute Herston, Brisbane, Australia, **13** School of Life Sciences, University of Nottingham, Nottingham, Nottinghamshire, England, United Kingdom, **14** Institut Pasteur Korea, Pangyo Techno-Valley, Gyeonggi Province, Korea, **15** Clinical Pharmacology, Novartis Consumer Health, Nyon, Switzerland, **16** Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, The Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam, **17** Nuffield Department of Clinical Medicine, Centre for Tropical Medicine, Oxford University, Oxford, England, United Kingdom, **18** The London School of Hygiene and Tropical Medicine, London, England, United Kingdom, **19** Internal Medicine, Yale University, New Haven, Connecticut, United States of America, **20** Health Sciences and Technology/Institute for Medical Engineering and Science, Massachusetts Institute of Technology, Cambridge, Massachusetts, United States of America, **21** Department of Immunology & Infection, London School of Hygiene and Tropical Medicine, London, England, United Kingdom, **22** Museum of National History, Sorbonne Universities, Paris, France, **23** SCYNEXIS, Inc., Durham, North Carolina, United States of America, **24** Department of Biochemistry and Molecular Biology and Johns Hopkins Malaria Research Institute, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, United States of America, **25** Department of Biochemistry, University of Yaoundé, Yaoundé, Cameroon, **26** CIMUS Research Centre, University of Santiago de Compostela, Santiago de Compostela, A Coruña, Spain, **27** Center for Discovery and Innovation in Parasitic Diseases, Department of Pathology, University of California San Francisco, San Francisco, California, United States of America, **28** Department of Biochemistry and Molecular Biology and Huck Center for Malaria Research, Pennsylvania State University, University Park, Pennsylvania, United States of America, **29** School of Life and Environmental Sciences, University of Sydney, Darlington New South Wales, Australia, **30** Department of Biochemistry and Virginia Tech Center for Drug Discovery, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, United States of America, **31** National Center of Advancing Translational Sciences, NIH, Bethesda, Maryland, United States of America, **32** Department of Biochemistry, Mahidol University, Bangkok, Thailand, **33** Department of Molecular Medicine, Morsani College of Medicine, University of South Florida, Tampa, Florida, United States of America, **34** Department of Pharmacological and Biomolecular Sciences, Università degli Studi di Milano, Milano, Italy, **35** Department of Pathology and Immunology, Washington University in St Louis, St. Louis, Missouri, United States of America, **36** Department of Life Sciences, Imperial College London, London, England, United Kingdom, **37** Division of Pharmacology and Drug Discovery, Department of Pediatrics, School of Medicine University of California San Diego, La Jolla, California, United States of America, **38** National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido, Japan, **39** Department of Biochemistry and Chemistry of nutrition, Mansoura University, Mansoura City, Egypt, **40** Eck Institute for Global Health, Department of Biological Sciences, University of Notre Dame, Notre Dame Indiana, United States of America, **41** Department of Microbiology & Immunology and Division of Infectious Diseases, Department of Medicine, Columbia University Medical Center, New York, New York, United States of America, **42** Broad Institute, Cambridge, Massachusetts, United States of America, **43** Biochemistry and Parasitology Department, Malaria DPU, Diseases of the Developing World (DDW), GlaxoSmithKline R&D, Tres Cantos, Madrid, Spain, **44** Tropical Parasitic Diseases Unit, Northwick Park Institute for Medical Research, Harrow, Middlesex, England, United Kingdom, **45** Department of Microbiology and Immunology, Weill Cornell Medical College, New York, New York, United States of America, **46** Medical Scientist Training Program, University of California, San Diego, San Diego, California, United States of America, **47** Department of Chemical Biology & Therapeutics, St. Jude Children's Research Hospital, Memphis, Tennessee, United States of America, **48** Dept. of Biology and South Texas Center for Emerging Infectious Diseases, University of Texas, San Antonio, San Antonio, Texas, United States of America, **49** Instituto de Medicina Molecular, Lisboa, Portugal, **50** Institute of Parasitology, University of Berne, Bern, Switzerland, **51** Medicines for Malaria Venture, Geneva, Switzerland, **52** Global Health Research Section, hhc Data Creation Center, Eisai Co., Ltd, Tsukuba-shi, Ibaraki, Japan, **53** Institute for Science and Technology in Medicine, Keele University, Keele, Staffordshire, United Kingdom, **54** Department of Medicine, College of Medicine, University of Vermont, Burlington, Vermont, United States of America, **55** Department of Pathobiology, Faculty of Science, Mahidol University, Bangkok Thailand, **56** Medical Research Centre, Institute for Pharmacogenetics, Essen, Germany, **57** Department of Biochemistry, University of Geneva, Geneva, Switzerland, **58** Department of Biochemistry and Molecular Biology, LSU Health Sciences Center, New Orleans, Louisiana, United States of America, **59** Research School of Biology, Australian National University, Canberra, Australian Capital Territory, Australia, **60** Department of Molecular Microbiology & Immunology, Oregon Health & Science University,

Portland, Oregon, United States of America, **61** Department of Medicine, University of California San Diego, San Diego, California, United States of America, **62** Department of Biochemistry and Molecular Biology, Monash University, Clayton, Australia, **63** University of Antwerp, Department of Biomedical Sciences, Antwerp, Belgium, **64** Biosciences Unit, Council for Scientific and Industrial Research, Pretoria, South Africa, **65** Department of Microbiology, Monash University, Clayton, Australia, **66** Chemotargets S.L. and Research Group on Systems Pharmacology, Research Program on Biomedical Informatics (GRIB), IMIM Hospital del Mar Institute of Medical Research and University Pompeu Fabra, Barcelona, Catalonia, Spain, **67** Division of Cancer Therapeutics and Diagnosis, Drug Synthesis and Chemistry Branch, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, United States of America, **68** Graduate Program in Bioinformatics, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **69** ChEMBL group, European Molecular Biology Laboratory—European Bioinformatics Institute (EMBL-EBI), Hinxton, Cambridgeshire, United Kingdom, **70** Genomics Institute of the Novartis Research Foundation, San Diego, California, United States of America, **71** Department of Internal Medicine and Infectious Diseases, Faculty of Veterinary Medicine, Mansoura University, Mansoura City, Egypt, **72** Department of Biochemistry and Stanford Genome Technology Center, Stanford University, Palo Alto, California, United States of America, **73** Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, **74** Laboratoire de Chimie Moléculaire, CNRS,—UMR 7509, COB-IRJBD, Mulhouse Cedex, France, **75** The Jenner Institute, University of Oxford, Oxford, England, United Kingdom, **76** Department of Chemistry, University of Capetown, Capetown, South Africa, **77** H. Feinstone Department of Molecular Microbiology and Immunology, Johns Hopkins School of Public Health, Baltimore, Maryland, United States of America, **78** Molecular, Cell and Developmental Biology, University of California, Santa Cruz, Santa Cruz, California, United States of America, **79** Department of Synthetic Biology and Bioenergy, J. Craig Venter Institute, La Jolla, California, United States of America, **80** School of Natural Sciences, Griffith University, Nathan, Queensland, Australia, **81** Hudson Institute of Medical Research; Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, Australia, **82** SYNthesis Research, Parkville, Victoria, Australia

^{‡a} Current address: Wellcome Trust Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridgeshire, United Kingdom

^{‡b} Current address: Global Health, Merck, Coisins, Switzerland

* wesley@uw.edu.

Abstract

A major cause of the paucity of new starting points for drug discovery is the lack of interaction between academia and industry. Much of the global resource in biology is present in universities, whereas the focus of medicinal chemistry is still largely within industry. Open source drug discovery, with sharing of information, is clearly a first step towards overcoming this gap. But the interface could especially be bridged through a scale-up of open sharing of physical compounds, which would accelerate the finding of new starting points for drug discovery. The Medicines for Malaria Venture Malaria Box is a collection of over 400 compounds representing families of structures identified in phenotypic screens of pharmaceutical and academic libraries against the *Plasmodium falciparum* malaria parasite. The set has now been distributed to almost 200 research groups globally in the last two years, with the only stipulation that information from the screens is deposited in the public domain. This paper reports for the first time on 236 screens that have been carried out against the Malaria Box and compares these results with 55 assays that were previously published, in a format that allows a meta-analysis of the combined dataset. The combined biochemical and cellular assays presented here suggest mechanisms of action for 135 (34%) of the compounds active in killing multiple life-cycle stages of the malaria parasite, including asexual blood, liver, gametocyte, gametes and insect ookinete stages. In addition, many compounds demonstrated activity against other pathogens, showing hits in assays with 16 protozoa, 7 helminths, 9 bacterial and mycobacterial species, the dengue fever mosquito vector, and the

NCI60 human cancer cell line panel of 60 human tumor cell lines. Toxicological, pharmacokinetic and metabolic properties were collected on all the compounds, assisting in the selection of the most promising candidates for murine proof-of-concept experiments and medicinal chemistry programs. The data for all of these assays are presented and analyzed to show how outstanding leads for many indications can be selected. These results reveal the immense potential for translating the dispersed expertise in biological assays involving human pathogens into drug discovery starting points, by providing open access to new families of molecules, and emphasize how a small additional investment made to help acquire and distribute compounds, and sharing the data, can catalyze drug discovery for dozens of different indications. Another lesson is that when multiple screens from different groups are run on the same library, results can be integrated quickly to select the most valuable starting points for subsequent medicinal chemistry efforts.

Author Summary

Malaria leads to the loss of over 440,000 lives annually; accelerating research to discover new candidate drugs is a priority. Medicines for Malaria Venture (MMV) has distilled over 25,000 compounds that kill malaria parasites *in vitro* into a group of 400 representative compounds, called the "Malaria Box". These Malaria Box sets were distributed free-of-charge to research laboratories in 30 different countries that work on a wide variety of pathogens. Fifty-five groups compiled >290 assay results for this paper describing the many activities of the Malaria Box compounds. The collective results suggest a potential mechanism of action for over 130 compounds against malaria and illuminate the most promising compounds for further malaria drug development research. Excitingly some of these compounds also showed outstanding activity against other disease agents including fungi, bacteria, other single-cellular parasites, worms, and even human cancer cells. The results have ignited over 30 drug development programs for a variety of diseases. This open access effort was so successful that MMV has begun to distribute another set of compounds with initial activity against a wider range of infectious agents that are of public health concern, called the Pathogen Box, available now to scientific labs all over the world (www.PathogenBox.org).

Introduction

Preclinical development for drugs in neglected diseases remains a slow process due to a lack of access to compounds, and legal complications over intellectual property ownership. One way to accelerate drug discovery is to provide open access to bioactive molecules with public disclosure of the resulting biological data. The data from open access of bioactive molecules can help prioritize which compounds to investigate further through medicinal chemistry for the original indication and can also uncover other indications for compound development. It was in this spirit of providing open access of malaria-bioactive compounds, and disseminating the results in the public domain, that the Malaria Box project was initiated by the Medicines for Malaria Venture.

Origins of the 'Malaria Box' compound set

Since 2007, over 6 million compounds were screened against asexual-stage *Plasmodium falciparum*, at two pharmaceutical companies (GlaxoSmithKline [1] and Novartis [2]), and two

academic centers (St. Jude, Memphis [3], and Eskitis, Australia [4]), resulting in over 20,000 compounds active in the low- to sub-micromolar range. The structures of the 20,000 anti-malaria hits were made available in ChEMBL (www.ebi.ac.uk/chembl), but discussions with biology groups had underlined the importance of access to the compounds themselves for testing. Cluster analysis and commercial availability reduced this to a set of 400 representative compounds, the 'Malaria Box', which was distributed freely to researchers who provided a rationale for screening [5]. This paper presents a summary and analysis of the collected results of the Malaria Box screening from 55 groups who performed a wide variety of assays, the large majority of which are presented in this paper. The collective results are greater than the sum of the individual assays, because each compound can be queried for activity, pharmacokinetic, and safety data to gauge its suitability as a starting point for subsequent medicinal chemistry optimization efforts.

Results

The Heat Map (S1 Table) reports the data from over 290 assays run on the Malaria Box compounds; a snapshot is shown in Fig 1. The results are color coded, where the compounds with the highest activity are coded red and those with relative inactivity green. In the center of the box in S1 Table, the numerical value for the compound is given. It can be seen immediately that some compounds have activities in several biological assays across multiple species and these tend to have activity against mammalian cells as well, whereas other compounds have a rather limited spectrum of activity and are less toxic to mammalian cells.

The data demonstrated in S1 Table are provided by 55 groups who have performed 291 assays to screen the Malaria Box. The vast majority of the data are presented for the first time in this paper. In supplementary data S1 Table, note that columns with data presented for the first time in this paper, representing 236 assays, are colored pink on the top row; published /in press data columns, 55, are grey, with citations provided. Presenting the combined dataset provides insights into the hit rates in these various assays while allowing rapid access to the data by the wider scientific community.

The Heat Map (S1 Table) presents the Malaria Box chemicals grouped by chemical relatedness. Of the 400 compounds, over 100 are closely-related paired molecules so immediate structure-activity-relationships (SAR) can often be seen from hits with these pairs. The Heat Map identified obvious correlations in chemistry and biology between compounds (both Mechanism-of-Action and phenotypic activity). Some biological assays are relatively similar; for example, there were a large number of different *P. falciparum* gametocyte assays (S1 Table, columns AV-CB), which also cluster, although not perfectly. As such, the aggregate screening data help overcome inter-laboratory bias and identify outstanding activities. For example, compounds that were active in multiple gametocyte assays represent more solid positives than a compound that was active in only one screening assay. However, the gametocyte assays were often performed using different techniques and screening concentrations (see S1 Methods and Results, for details) and one assay may be preferred over another to select compounds with gametocyte activity. Thus having the aggregate data presented together with the individual protocols is more valuable than just having each individual data set to look at sequentially.

Malaria Box safety and pharmacokinetic data

Early safety data were obtained by testing all compounds against 73 human cell lines at 10 μM or above, and developing zebrafish embryos were exposed at 5 μM , providing further clues on potential safety issues. A frequent cardiotoxicity safety concern is QTc prolongation, and all compounds were screened for hERG inhibition [6], which is a proxy for this risk (S1 Table




Fig 1. Malaria Box Heatmap. Shown are selected data from the HeatMap (S1 Table) for the 400 Malaria Box compounds. Each column represents an assay (grouped by category), compounds are represented in rows. The red-green gradient represents higher to lower activity. Favorable PK activities are

scored green. *Pf*: *Plasmodium falciparum*, *Pb*: *Plasmodium berghei*, PK: pharmacokinetics, sol.: solubility, hERG: human ether-a-go-go channel inhibition, DDI: drug-drug interactions (predicted).

doi:10.1371/journal.ppat.1005763.g001

column GI). The efficacy and safety of anti-malarial compounds could be altered in endemic regions when administered to patients who are also treated for HIV (Human Immunodeficiency Virus) or TB (tuberculosis), due to drug-drug interactions in the liver. To flag such interactions, we employed two recent breakthrough models: a bioengineered microscale human liver in a high-throughput assay format that accurately captures human drug-drug interactions not detectable in animals or cell lines [7] and a custom-made, robotic high-throughput Luminex bead-based method for profiling the expression of 83 human liver drug-metabolizing enzymes [8]. Combining these tools, we profiled the Malaria Box compounds for induction or inhibition of drug-metabolizing pathways (S1 Table, columns GL-HA) and thereby ranked compounds for potential for drug interactions with existing HIV and TB regimens, to enhance selection of compounds with the lowest safety risks. We also scored the Malaria Box compounds for acute hepatotoxicity by monitoring morphology and daily albumin and urea secretion from hepatocytes (S1 Table, columns FQ-FS).

G protein-coupled receptors (GPCRs) represent the largest human drug target class [9]; they affect neurological and cardiovascular physiology and are included in routine safety pharmacology panels [10]. Therefore, *in vitro* affinity determinations on 23 selected human off-target GPCRs were performed on a subset (10%) of MMV compounds (S1 Table, columns HC-HZ). One of the most severe GPCR-related adverse effects is cardiac valvulopathy linked to 5-HT_{2B} activation [11, 12]. Therefore, some of the MMV compounds with significant binding affinity for the 5-HT_{2B} receptor were also tested on the corresponding functional assay to determine a potential agonistic effect. In addition, predictions of compound glutathione reactivity and epoxidation potential were calculated for each of the Malaria Box compounds (S1 Table, columns IB-IC). These combined safety results alert us to compounds with issues that hopefully can be resolved in subsequent medicinal chemistry programs.

Prior to *in vivo* pharmacology evaluation it is important to know that an effective plasma concentration can be reached; this exposure was measured in rodents for all compounds, from a single high oral dose (140 $\mu\text{mol/kg}$). Around one third of the compounds generated high plasma C_{max} (>1 $\mu\text{g/ml}$) and/or high overall exposure (S1 Table, columns GD-GE). This is a higher than expected percentage of compounds with measureable oral bioavailability than if compounds were randomly selected, and probably reflects the large number of drug-like leads selected for the Malaria Box. The combination of *in vitro* potency and bioavailability provides a rough dosing estimate, informing subsequent decision-making around selection of development leads.

The combined analysis of all of these safety and pharmacokinetic data allows selection of the most promising compounds to advance to medicinal chemistry, and which parameters should be monitored and improved during a medicinal chemistry program.

New insights into malaria

The activity of Malaria Box compounds against the asexual, erythrocytic stage of *P. falciparum* was confirmed by five laboratories on seven different *P. falciparum* strains. There were sometimes 5-10-fold differences in the effective concentration that caused a 50% reduction in growth (EC_{50}) in each assay, and these may have been due to variations in the readouts for the screening assays (LDH release, MitoTracker or Sybr Green dye incorporation, hypoxanthine incorporation, DAPI imaging assay), variations in the protein concentration in the assay medium (affecting the free compound concentration), the time the compound incubated time,

Table 1. Malaria Box compounds with activity in biological assays (malaria, helminths, *Wolbachia*, and cancer cells) and lacking toxicity at therapeutic levels. Selectivity Index, SI, is toxicity level/activity level; p, probe-like; d, drug-like.

Antimalarial positives		Anthelmintic positives			Anti- <i>Wolbachia</i> positives	Anticancer positives		Notes
All drug-like (d) <i>P. falciparum</i> early ring stage compounds (EC50 <200 nM, SI >10) with gametocyte activity	<i>P. falciparum</i> asexual, liver, & gametocyte activity	<i>Brugia malayi</i>	<i>Trichuris trichuria</i>	<i>Ancylostoma ceylanicum</i>	<i>Wolbachia</i>	MMV Number	MGLI50 = Mean Growth Log I50	
MMV000248 ^d	MMV006913 ^d	MMV007907 ^d	MMV008294 ^p	MMV666607 ^p	MMV000642 ^p	MMV007384 ^p	-7.34	Colon, differential, potent
MMV006087 ^d	MMV007116 ^d	MMV019241 ^p	MMV666601 ^p		MMV008138 ^d	MMV019074 ^d	-4.91	Specific lines sensitive
MMV006455 ^d	MMV007199 ^p	MMV665831 ^p			MMV396664 ^p	MMV665803 ^d	-5.01	
MMV006706 ^d	MMV007907 ^d	MMV666054 ^p			MMV665824 ^p	MMV665796 ^d	-5	
MMV011567 ^d	MMV020700 ^d				MMV665841 ^p	MMV020275 ^d	-4.93	
MMV011795 ^d	MMV665843 ^d				MMV665890 ^d	MMV000760 ^d	-6.02	Differential
MMV020505 ^d	MMV665977 ^p				MMV665897 ^d	MMV666020 ^p	-4.9	
MMV020660 ^d	MMV666095 ^p				MMV665948 ^d	MMV665969 ^p	-6.29	
MMV396749 ^d					MMV666075 ^d	MMV666597 ^p	-5.57	Differential
MMV396794 ^d					MMV666597 ^p	MMV006962 ^p	-5.61	Differential, CNS
MMV665805 ^d					MMV666601 ^p			
MMV665878 ^d					MMV666607 ^p			
MMV665915 ^d								

doi:10.1371/journal.ppat.1005763.t001

or other differences. However, usually the results were consistent and strain-independent. We have documented which sub-stage of the asexual lifecycle the compounds acted upon ([S1 Table](#), columns AA-AE). This information is important in identifying compounds that may overcome existing resistance against artemisinin and other antimalarials. For instance, compounds that target early ring stage intra-erythrocytic parasites and have fast-killing dynamics are sought after because, like artemisinins, they kill parasites rapidly and may reduce patient mortality. [Table 1](#) shows compounds that also target liver stages of the parasite's life cycle.

Targeting disease-relevant malaria stages

P. berghei liver stage (LS) inhibition, using parasite-encoded luciferase activity as a readout of infection in HepG2 cells, was independently determined by two groups at very different screening concentrations (Hanson: 5 μM, Winzeler: 50 μM). Forty-three compounds, roughly 10% of the compound library, inhibited infection by at least 50% at 5 μM and 90% at 50 μM (referred to as LS double actives). HepG2 cell toxicity, (50% or greater reduction in HepG2 abundance based on direct or indirect readouts) was observed with 63% of Malaria Box compounds at 50 μM, while only 10% were toxic at the 5 μM concentration. After excluding those that showed significant toxicity in HepG2 cells at both 5 and 50 μM, Malaria Box compounds were stratified by potential mode-of-action annotation ([S1 Table](#), column M). Five potential modes of action stood out as enriched in LS double actives: (i) cysteine protease inhibitors (cruzain, rhodesain): 1.8% of all Malaria Box compounds (7/400) and 4.7% (2/43) of LS double actives; (ii) possible respiratory-dependent targets (Δ IC₅₀ in low oxygen vs. normal oxygen): 0.8% (3/

400) of all Malaria Box compounds and 4.7% (2/43); (iii) targeting yeast respiration: 3.5% of all Malaria Box compounds (14/400) and 9.3% of LS double actives (4/43); (iv) suspected or known *PfDHODH* (dihydroorotate dehydrogenase) inhibitors: 2.5% of all Malaria Box compounds (10/400) and 9.3% of LS double active (4/43); and (v) suspected or known cytochrome bc1 inhibitors: 4.3% of all Malaria Box compounds (17/200) and 16.3% of LS double actives (7/43). Compounds with activity against *PfATP4*, now the most common intra-erythrocytic asexual target seen in phenotypic screens, were not found amongst the LS double actives.

There is a great need for antimalarials that kill *dormant*, liver-stage *P. vivax* (hypnozoites), but there is a lack of assays that measure this activity. Only nine compounds (Table 1) show simultaneous activity against gametocytes, liver, and asexual stages, whilst lacking evidence of toxicity in zebrafish and broad cytotoxicity to mammalian cells. These would be compounds to prioritize for *in vitro* and *in vivo* screening against *P. vivax* hypnozoites and would benefit from additional MoA studies.

Gametocytocidal drugs would block transmission from the human to the mosquito and break the parasite's life cycle. The data shown in Table 1 include series with activities on both gametocyte and liver stages, and some of the data intriguingly challenges existing assumptions. For instance, MMV007116 in this category is a mitochondrial (bc1) inhibitor (S1 Table, column M, line 168) and has activity in a number of gametocytocidal assays, but other bc1 inhibitors are not generally gametocytocidal, suggesting another MoA for this compound. We also see 4-aminoquinolines as inhibitors of some gametocyte assays, although the parent 4-aminoquinoline compound chloroquine is known not to be gametocytocidal for *P. falciparum*. Again, this may imply a different MoA for some 4-aminoquinoline compounds or perhaps multiple modes of action for certain compounds. These findings re-emphasize the strength of looking at assay data in a wider context in Open Source drug discovery.

Mechanism-of-action screening

Data from one-hundred-nineteen MoA assays for compounds from the Malaria Box are included, identifying potential targets for 135 of them (S1 Table and S1 Methods and Results). The MoA assay data are presented in Column M of S1 Table, and further information about the screens and their results are given in S1 Methods and Results. These screens included biochemical screens for enzyme inhibition, protein-protein interactions, behavior by altered yeast or malaria organisms, and a variety of other screens. Some associations are strong and have been followed up with additional experimentation (e.g. MMV008138 and its target *Pf-IspD* [13–15]), but most target associations are still tentative. Indeed, some listed MoA activities occur only at higher concentrations than activity in cell-based screens and therefore are unlikely to explain that compound's activity against a pathogen or tumor cell. In addition, many MoAs have been inferred for malaria, but are less likely to apply to the diverse groups of organisms screened with the Malaria Box compounds.

Surface plasmon resonance (SPR) was used to identify nine compounds which inhibit four sets of protein-protein interactions (PPI), without overlap between sets (S1 Methods and Results), suggesting that molecules were identified that specifically target these protein-protein interfaces. Compounds inhibiting *P. falciparum* (autophagy-related proteins) Atg8-Atg3 PPI were MMV007907, MMV001246 and MMV665909 (S1 Table, column M). They had a pronounced effect on all stages of gametocyte development, which supports the idea of *PfAtg8-Atg3* being involved in remodeling and vesicular trafficking in gametocyte development. Six compounds inhibited *in vitro* translation in *P. falciparum* lysates by more than 60% at a concentration of 1 μ M (S1 Table, column L; [16]). One of these protein translation-inhibiting

compounds, MMV007907, is interesting in that it had activity against both liver and gametocyte stages as well as a broad range of other pathogens, and has low toxicity to human cell lines. Twenty-six compounds either inhibited the mitochondrial electron transport chain (bc1, 11 compounds) or DHODH (15 compounds). Since both the bc1 and DHODH pathways converge on pyrimidine biosynthesis, it is interesting that almost all bc1 inhibitors had anti-liver stage and anti-male gametocyte activity, while the anti-male gametocyte property was generally lacking in most DHODH inhibitors [17–19].

PfATP4 is a *P. falciparum* plasma membrane protein with genetic variants that confer resistance to several new clinical and preclinical antimalarials [20–24]. *PfATP4* has been proposed to function as a $\text{Na}^+:\text{H}^+$ pump, effluxing Na^+ from (and importing H^+ into) the malaria parasite [21]. Parasites exposed to 28 MMV Malaria Box compounds have shown ion-homeostasis changes similar to those observed with likely *PfATP4* inhibitors (indicated in column K, S1 Table) [25], and thus are inferred to be *PfATP4* inhibitors. Analysis of the 281 assays' results with these compounds, reported here, allows detailed conclusions about the potential effects of ATP4 inhibition in *Plasmodium* as well as other organisms. From the Malaria Box data summarized here, it is evident that the 28 *PfATP4*-associated hits tended to be inactive against the variety of non-Apicomplexan protozoa, helminths, insects, yeast and bacteria that were tested. An exception was *Trypanosoma cruzi*, that was growth-inhibited by almost 40% of the *PfATP4* inhibitors (11/28), compared to an overall hit rate of 20%. It should be noted that the non-*Plasmodium* Apicomplexan parasites against which the majority of the compounds were tested—*Cryptosporidium parvum*, *Toxoplasma gondii*, *Theileria equi* and three species of *Babesia*—were not, in general, particularly susceptible to the *PfATP4*-associated hits. There is not, to our knowledge, any evidence that the other Apicomplexan parasites against which the Malaria Box was tested are exposed to a high- Na^+ environment within their host cells, and this may explain the lower sensitivity to inhibition of a Na^+ efflux mechanism. In contrast, infection of an erythrocyte by *Plasmodium* is followed by an increase in the Na^+ concentration in the erythrocyte cytosol as a result of the induction of broad-specificity (Na^+ -permeable) 'New Permeability Pathways' in the host erythrocyte membrane [26–28]. This suggests that perturbation of Na^+ efflux through inhibition of *PfATP4* is uniquely, highly detrimental to intra-erythrocytic malaria parasites.

There is prior evidence that *PfATP4*-associated compounds are active against gametocyte stages of *P. falciparum* [5, 22–24, 29–32]. Twenty-five of the 28 *PfATP4*-associated hits (89%) caused some inhibition of male gamete formation at 1 μM (i.e. had positive % inhibition values; S1 Table). It should be noted, however, that approximately half of the *PfATP4*-associated hits have IC_{50} values for the killing of asexual parasites that are similar to or higher than the 1 μM concentration used in the gamete formation assay. Only 65% of the *PfATP4* non-hits tested had positive values for inhibition of male gamete formation at 1 μM . An increase in extracellular pH is known to trigger the exflagellation of male *P. falciparum* gametes, raising the possibility that an increase in intracellular pH in male gametocytes or gametes, resulting from *PfATP4* inhibition, triggers premature exflagellation, leading to parasite death. Thus, it is possible that an increase in intracellular pH in male gametocytes or gametes resulting from *PfATP4* inhibition triggers premature exflagellation leading to their death.

Malaria box compounds were also screened against asexual stages using metabolomic and chemogenomic profiling (Fig 2). Using metabolomic profiling to examine the metabolic responses to the 80 compounds in plate A, six of seven compounds believed to target *PfATP4* [25] showed a distinct metabolic response characterized by an accumulation of dNTPs, and a decrease in hemoglobin-derived peptides (Fig 2A, S2 Table). Twenty-one compounds clustered with atovaquone, an inhibitor of the bc1 complex of the electron transport chain, exhibiting an atovaquone-like signature characterized by the dysregulation of pyrimidine synthesis. Of these





Fig 2. Metabolomic and chemogenomic profiling. (A) Metabolic profiling: Heat map showing metabolic fingerprints of 80 Malaria Box compounds and atovaquone control. Parasite extracts were analyzed by LC-MS, and changes in metabolite pools were calculated for drug-treated parasites as compared to untreated controls. Hierarchical clustering was performed on ²log-fold changes in metabolites (data in [S2 Table](#)), scaled from -3 to +3. Six of seven compounds (indicated in red) reported to target *Pf*ATP4 [25] showed a distinct metabolic response characterized by the accumulation of dNTPs and a decrease in hemoglobin-derived peptides. A large cluster of compounds (indicated in blue) clustered with the atovaquone control (indicated in orange), and exhibit an atovaquone-like signature characterized by dysregulation of pyrimidine biosynthesis, and showed a distinct metabolic response characterized by the accumulation of dNTPs and a decrease in hemoglobin-derived peptides. (B) Chemogenomic profiling: A collection of 35 *P. falciparum* single insertion *piggyBac* mutants were profiled with 53 MMV compounds and 3 artemisinin (ART) compounds [Artesunate (AS), Arteminic acid (AL) and Artemether (AM)] for changes in IC₅₀ relative to the wild-type parent NF54 (data in [S3 Table](#), genes queried in [S4 Table](#)). Clone PB58 carried a *piggyBac* insertion in the promoter region of the K13 gene and has an increased sensitivity to ART compounds as do PB54 and PB55 [33]. Drug-drug relationships based on similarities in IC₅₀ deviations of compounds generated with *piggyBac* mutants created chemogenomic profiles used to define drug-drug relationships. The significance of similarity in MoA between Malaria Box compounds and ART was evaluated by Pearson's correlation calculations from pairwise comparisons. The X axis shows the chemogenomic profile correlation between a Malaria Box compound and AS, the Y axis with AM; the color gradient indicates the average correlation with all ART derivatives tested. Five Malaria Box compounds (MMV006087, MMV006427, MMV020492, MMV665876, MMV396797) were identified as having similar drug-drug chemogenomic profiles to the ART sensitivity cluster.

doi:10.1371/journal.ppat.1005763.g002

21 atovaquone-like compounds, 17 were also identified by other groups as targeting the electron transport chain or pyrimidine synthesis. For chemogenomic profiling, a collection of 35 *P. falciparum* single insertion *piggyBac* [33] mutants were profiled with 53 MMV compounds and three artemisinin (ART) compounds [Artesunate (AS), Arteminic acid (AL) and Artemether (AM)] for changes in IC₅₀ relative to the wild-type parent NF54 ([Fig 2B](#), [S3 Table](#), [S4 Table](#)). Five Malaria Box compounds (MMV006087, MMV006427, MMV020492, MMV665876 and MMV396797) were identified as having similar drug-drug chemogenomic profiles to the ART-sensitivity cluster ([Fig 2B](#)). These compounds may be rapid killers, like artemisinin, and should be explored further for confirmation, and whether they can overcome artemisinin-resistance for ring-stage killing.

Screening on yeast to suggest MoAs

Four groups carried out screens on *S. cerevisiae* strains engineered to help elucidate the MoA of test compounds. One screen established that 35 Malaria Box compounds were active on a multiple ABC-transporter deficient strain (also known as the 'monster strain') *S. cerevisiae* [34]. Since yeasts are generally resistant to compound inhibition due to transporters, this monster strain can now be analyzed for MoA of inhibition by these 35 compounds. A second study measured selective growth inhibition of *S. cerevisiae* using different carbon sources. Growth was measured in three different growth media: rich or minimal media using dextrose as a carbon source, or minimal media using ethanol and glycerol as carbon sources. Compounds affecting growth in a media-specific manner may represent inhibitors of key metabolic pathways. A third group used a yeast strain expressing the *Pf* phosphoethanolamine methyltransferase (*Pf*PMT) to screen for phosphocholine (PC) synthesis inhibitors. This screen relies on the incapability of this yeast strain to synthesize PC in the absence of exogenous choline, and thus depends on the malaria *Pf*PMT for survival. Screening the Malaria Box compounds, and a variety of controls including wild-type PMT and choline supplemented media, led to the identification of MMV007384, MMV007041 MMV396736, MMV396723, MMV000304, MMV000570, MMV000704, MMV666071, MMV000445, MMV667491, and MMV666080 as possible *Pf*PMT inhibitors. Finally, a fourth group screened *S. cerevisiae* grown either on ethanol-containing media requiring respiration or glucose-fermentative media not requiring respiration, and identified 12 compounds that gave superior inhibition on ethanol media suggesting

that these compounds inhibit a respiratory target. Seven of these were not associated with any other targets; the others were potential inhibitors of DHODH (3, 49), bc1, and IspD.

Activity against protozoa other than Plasmodium

The Malaria Box was screened against 16 additional protozoa, all of which are of medical or veterinary significance. Compounds with activity against three or more protozoa were usually toxic for the zebrafish or non-cancer mammalian cell lines, underlining the need to limit the concentrations used in assays, to avoid meaningless positives. [Table 2](#) lists compounds with activity against protozoa that were nontoxic to zebrafish and most mammalian cells. In the *Cryptosporidium parvum* assay there were numerous active compounds, but none were completely devoid of toxicity for zebrafish and mammalian cell lines. MMV665917 had a >20-fold Selectivity Index (SI) for *C. parvum* over mammalian cells. *Trypanosoma cruzi* actives were non-overlapping between groups, and are listed separately, but *T. brucei* actives overlapped extensively with other screens and are presented together. There were seven non-toxic hits that were active against extracellular amastigotes of *Leishmania infantum*, but no non-toxic compounds were active on intracellular macrophage growth of *L. infantum*. There were five non-toxic Malaria Box compounds active against *T. gondii* (MMV666095, MMV007363, MMV007791, MMV007881 and MMV006704). Many of the compounds that were active on *Neospora caninum* raised no toxicity flags on the accompanying host cell fibroblast screen, but many were toxic at 10 μ M or below for mammalian cells and zebrafish. The remaining non-toxic *N. caninum* actives that bear further investigation include: MMV019670, MMV000911 and MMV006309. Most compounds active against *Entamoeba histolytica*, *Naegleria fowleri*, or exflagellation of *Chromera velia* were toxic. An exception was MMV665979, an outstanding hit for *Naegleria fowleri*, with limited toxicity elsewhere in the dataset. With respect to screening *Babesia* and *Theileria*, ten novel anti-*Babesia* and anti-*Theileria* hits with nanomolar IC₅₀s were identified: MMV666093, MMV396794, MMV006706, MMV665941, MMV085203, MMV396693, MMV006787, MMV073843, MMV007092 and MMV665875. The most interesting hits were MMV396693, MMV073843, MMV666093, MMV665875 and MMV006706 with mean SIs greater than 230 and IC₅₀s ranging from 43 to 750 nM for both bovine *Babesia* and equine *Babesia* and *Theileria* parasites. Additionally, 64, 45 and 49 Malaria Box compounds exhibited IC₅₀s lower than those of diminazene aceturate (the most widely used antibabesial drug) against the *in vitro* growth of *B. bovis*, *B. bigemina* and *T. equi*, respectively.

In vitro screening of Open Access Malaria Box compounds against *Babesia bovis*, *B. bigemina*, *Theileria equi* and *B. caballi* has led to the discovery of 10 novel potent anti-babesial hits exhibiting submicromolar potency against both bovine *Babesia* and equine *Babesia* and *Theileria*. *In vitro* follow up of the many of the hits identified in this study for *B. bovis*, *B. bigemina*, *B. caballi*, and *T. equi* parasites, revealed IC₅₀s lower than that obtained with the previously described drug-leads luteolin, pyronaridine, nimbolide, gedunin and enoxacin [35]. The ten potent hits for bovine *Babesia* and equine *Babesia* and *Theileria* identified in this study exhibited IC₅₀s lower than that obtained with the apicoplast-targeting antibacterials (ciprofloxacin, thiostrepton, and rifampin), miltefosine, fusidic acid or allicin [36–39].

Activity on helminths, mycobacteria, and bacteria

Many Malaria Box compounds were active on helminths at 10 μ M, but most of these were also toxic for mammalian cells or zebrafish. The remaining non-toxic compounds had activity against *Brugia malayi* (lymphatic filariasis) and *Ancylostoma ceylanicum* (hookworm; [Table 1](#)). But no non-toxic compounds were found with consistent activity against *Schistosoma mansoni*, *Strongyloides stercoralis*, *Trichuris muris*, *Haemonchus contortus*, or *Onchocerca linealis*.

Table 2. Antiprotozoal Malaria Box compounds with activity in biological assays and lacking toxicity at therapeutic levels. Selectivity Index, SI, is toxicity level/activity level; p, probe-like; d, drug-like.

Antiprotozoal positives										
<i>Cryptosporidium parvum</i> (SI>20)	<i>Trypanosoma brucei</i> (positive in all three screens, and SI >10)	<i>T. cruzi</i> (Scynexis screen)	<i>T. cruzi</i> (Maes screen)	<i>Leishmania donovani</i> amastigotes axenic (extracellular):	<i>Toxoplasma gondii</i>	<i>Babesia spp. (3) AND Theileria equi</i>	<i>Babesia spp. (3) OR Theileria equi</i> (excluding active on all 4 spp.)	<i>Naegleria fowleri</i>	<i>Neospora caninum</i>	<i>Perkinsus marinus</i>
MMV665917 ^p	MMV020505 ^d	MMV006558 ^p	MMV001230 ^d	MMV000356 ^d	MMV007363 ^d	MMV006706 ^d	MMV000620 ^{Bc}	MMV665979 ^d	MMV000911 ^d	MMV666081 ^d
	MMV020548 ^d	MMV006706 ^d	MMV007907 ^d	MMV007907 ^d	MMV007791 ^d	MMV666093 ^d	MMV000653 ^{Bbi+Bbo}		MMV006309 ^p	
		MMV006764 ^p		MMV011438 ^p	MMV666095 ^p	MMV396693 ^p	MMV000911 ^{Bc}		MMV019670 ^d	
		MMV007571 ^d		MMV019127 ^d	MMV007881 ^d	MMV073843 ^p	MMV006764 ^{Bbo+Bbi}			
		MMV011256 ^d		MMV665979 ^d	MMV006704 ^d	MMV665875 ^p	MMV007374 ^{Bc}			
		MMV019199 ^p		MMV665987 ^p			MMV007557 ^{Bc+Bbi}			
		MMV498479 ^p		MMV666686 ^p			MMV007808 ^{Bc}			
		MMV665841 ^p					MMV007906 ^{Bc}			
		MMV665843 ^d					MMV008173 ^{Bbo+Bbi}			
		MMV665878 ^d					MMV008455 ^{Bbi}			
		MMV665890 ^d					MMV011438 ^{Bbo+Bbi}			
		MMV665899 ^d					MMV018984 ^{Bc+Bbo}			
		MMV665908 ^p					MMV019199 ^{Bbo+Bbi}			
							MMV019202 ^{Bc}			
							MMV019670 ^{Bc}			
							MMV019995 ^{Bbi}			
							MMV396681 ^{Bc}			
							MMV665878 ^{Bbi}			
							MMV665879 ^{Bbo+Bbi}			
							MMV665890 ^{Bbo+Bbi}			
							MMV665908 ^{Bbi+Bbo+Te}			
							MMV665939 ^{Bbo+Bbi}			
							MMV665946 ^{Te}			
							MMV666009 ^{Bc+Bbi}			
							MMV666025 ^{Bbo}			
							MMV666095 ^{Bc}			
							MMV666116 ^{Te+Bbi}			
							MMV666689 ^{Bbo+Bbi}			

doi:10.1371/journal.ppat.1005763.t002

There remains the possibility that some of the toxic hits against these species can be addressed by medicinal chemistry.

With respect to activity against mycobacteria and bacteria, although every screen delivered actives, the majority were again discarded because of a toxicity signal against zebrafish and/or mammalian cells. The exceptions were non-toxic Malaria Box compounds that were active against *Wolbachia* (Table 1). *Wolbachia* bacteria are targeted as anti-filarials in order to deprive nematodes causing river blindness and elephantiasis from essential nutrients provided by this bacterium [40].

Activity on cancer cells

The US National Cancer Institute has screened 59 human tumor cell lines ('NCI60') against the Malaria Box compounds at 10 μ M (S1 Table and S1 Methods and Results). Among the 133 compounds further evaluated for dose-responses, and the ten of these then tested in confirmatory assays (S1 Text), MMV007384 was selected for potency and focused activity against colon cancer cells, and has been advanced to an *in vivo* proof-of-concept experiment.

Discussion

Academic drug discovery is highly fragmented. Many biology groups, especially those in disease-endemic countries, excel in developing highly disease-relevant pathogen models suitable for low- to medium-high throughput screening, but suffer from lack of access to innovative compounds. If they do have access to compounds, then they may fail to share the results, or lack drug development skills. The Malaria Box Project demonstrates how an open source approach allows effective data sharing: this publication serves as much to share the data among the 180+ co-authors as with the wider scientific community. By publishing in concert this ensures early publication and also sharing of ideas and expertise in drug discovery. New insights and series have been obtained for malaria (nine pan-stage active molecules which had not been previously prioritized). Moreover, screening against pathogens for additional neglected diseases has been catalyzed and hits found. The sharing of data from safety screens flags compounds that probably work through a general toxicity mechanism, and those compounds can be down-prioritized at an early stage. This is key to prioritizing compounds for medicinal chemistry, since the paucity of good starting points against some parasites has encouraged groups to screen at what may be inappropriately high drug concentrations. Another advantage of having a standardized, publically available library and dataset is that this allows benchmarking assay sensitivity, setting compound concentrations for expanded screens and deciding on acceptable hit criteria [41].

We saw some discrepancies in the values obtained for the same compounds in similar assays that were carried out by multiple groups, such as activity against asexual or gametocyte forms of *P. falciparum*, *Trypanosoma spp.*, and mammalian cells. In this sense, compounds that were positive in more than one assay would clearly be more likely to represent a true positive than compounds that were positive in only one screen. Some of these apparent discrepancies were probably due to variations in the techniques used for the screens. For instance, many methods used to measure gametocytocidal activity measure a specific metabolic activity. Because the metabolism will be affected by many factors that will lead to differences in output, including media composition (albumax versus serum), how old the media used was, purity of the gametocytes (how much asexual contamination and cell debris is present). In addition, the tested compounds varied widely in their propensity to bind to protein in the assay medium, and large differences in the protein content in two assays could lead to differences in unbound compound. Only the free compound would likely be available for activity in biological assays. Some

assays had extensive follow up, and if a compound was tested and activity confirmed with a dose-response, it is more likely to be a true positive than a compound flagged as positive from a single screening run. This complex dataset highlights the need to consider integrating more standardized criteria, such as similar (free) compound concentrations, assay media, or compound exposure duration, into future screening initiatives of this nature. This could potentially reduce inter-assay differences, and facilitate more direct data comparison across the different platforms. However it is clear in the case of gametocyte screens, that different assays that interrogate different biological processes do not necessarily achieve the same result for a given compound, even when the assay conditions have been standardized [42]. And trying to standardize assays may be counterproductive with the goal of convincing multiple groups to run their assays on a given set of compounds.

The MoAs associated with compounds (S1 Table, Column M) vary from very strong associations such as chemical-genetic evidence, to relatively weak associations, such as activity in a single biochemical screen at relatively high compound concentration. Thus most of the associations should not be taken as definitive MoA of the compounds for their biological activities. All associations were presented because not only could they be hypothesis-building for the discovery of a compound's disease-relevant MoA, but also because the Malaria Box compounds now represent a rich source of bioactive compound tools.

With its outcomes continually evolving, the Malaria Box has already made an impact by stimulating medicinal chemistry for many diseases. We are aware of such new medicinal chemistry programs against pathogens such as *Plasmodium* [43–45], *Babesia*, *Toxoplasma* [46], *Trypanosoma* [47–49], *Cryptosporidium* [31], *Schistosoma* [50], filaria, *Echinococcus*, helminths, bacteria, cancer and other diseases [30]. Ensuring that data becomes freely available is a challenge, and this paper represents the first such summary of over 290 screens against the compound collection, highlighting new activities and new MoAs. For the future, three goals are important. First, to track these compound series to ascertain whether any of these hits do become leads of drug development candidates. Second, data must be rapidly published, even with follow-up incomplete. Finally and most importantly, this model can be taken further. A second collection of 400 compounds, the Pathogen Box, (www.PathogenBox.org) based on compounds known to be active in phenotypic screens against an expanded set of pathogens responsible for neglected and tropical diseases has now become available from the Medicines for Malaria Venture. It is hoped this can be the start of equally fruitful collaborative networks.

Methods

See [S1 Methods and Results](#) for further details.

The Malaria Box is a set of 400 compounds that were previously shown to be active against asexual stages of *P. falciparum* *in vitro*. The process for Malaria Box compound selection was published previously [5], with 200 drug-like compounds as starting points for oral drug discovery and development and 200 diverse probe-like compounds for use as bioactive tools research. The selection was made to represent the broadest cross-section of structural diversity and, in the case of the drug-like compounds, properties commensurate with excellent oral absorption and the minimum presence of known toxicophores. One limiting factor was that compounds had to be commercially available; this limited the chemical space displayed in the original set of 20,000 malaria bioactives.

The Malaria box was shipped to 193 different research groups in 29 different countries as frozen 96-well plates with the compounds dissolved at 10 mM in 20 μ l DMSO (dimethylsulphoxide). Two years after shipping the first Malaria Box, the 193 groups were re-contacted and asked if they wanted to participate in a group publication disseminating and comparing the

results from the Malaria Box screens. Forty-seven of these groups did not reply to our multiple requests. Fifty-nine groups had not yet initiated screening, but 26 of these had only received the Malaria Box in the preceding three months. Thirty-one groups had publications in preparation and 39 papers have already been published [5, 14, 16, 25, 30–32, 42–46, 48–75]. Fifty-five groups agreed to contribute data and participate in this paper and provided data from 291 assays.

The compounds were then screened in biochemical and biological screens as documented in detail in [S1 Methods and Results](#). More detailed methods are provided for screens presented in this paper than for those whose results have already published. In addition, [S1 Methods and Results](#) provides data for both positive and negative controls obtained for each assay. In most assays, a single-concentration screen was run first and bioactives were identified. Some work was stopped after the primary screen, but most groups went on to perform confirmatory assays, and many provided hit concentrations that achieve 50% activity ([S1 Table](#)). The assays included a variety of cell-based pathogen screens covering multiple taxonomic groups, including *Plasmodium* (multiple life-stages), other protozoa, bacteria, mycobacteria, HIV, and also multicellular-organism screens such as helminths and a mosquito (See [Fig 1](#) and [S1 Table](#)).

Supporting Information

S1 Methods and Results. Supplementary Methods and Results.

(DOCX)

S1 Table. Malaria Box HeatMap. (Reference numbers refer to [S1 Methods and Results](#) references; Pink Headers signify data presented first in this paper data; Grey headers are published, submitted, or in press). Red shading means active and green means inactive and values are provided in each square. Favorable PK activities are scored green.

(XLSX)

S2 Table. Metabolomic data. The file is ordered as shown in [Fig 2A](#) and shows log₂ fold changes compared to an untreated control. Positive and negative values indicate increase and decrease, respectively, as compared to the control.

(XLSX)

S3 Table. Chemigenomic data.

(XLSX)

S4 Table. Genes queried in Chemogenomic approach.

(XLSX)

S1 Text. Supplementary Dose-Response data on selected cancer cell lines.

(PDF)

Acknowledgments

We thank Scynexis and Evotec for plating and distribution, Syngene for synthesis of the Malaria Box compounds, Essen Bioscience (particularly Del Trezise, Tim Dale, and Srinivasan Kanumilli) for determination of hERG inhibition, WuXi App Tec (particularly Genfu Chen, Hongshi Yu, Yusong Zhu, Ying Huang, and Wen Hua) for the PK/DMPK determinations, and ChEMBL for data maintenance and particularly Louisa Bellis and Anna Gaulton of the European Bioinformatics Institute, EBI. The purified HSP90 was a gift from Prof. Juan Pizzaro. We acknowledge the technical assistance of An Matheeußen, Jessica Richard of Louisiana State University, LSU, and Koji Sagane, Ph.D., Eisai Co., Ltd. for the construction of transgenic yeast

clones. We wish to thank Suresh Solapure PhD for discussions about PK determinations, Dr. Jean-Robert Ioset of Drugs for Neglected Diseases *initiative* (DNDi) for his many suggestions and support for kinetoplastid testing, Dr Nicoletta Basilico for invaluable advice, and Maria Mota (Instituto de Medicina Molecular, Lisboa) for her support. We thank the staff of the NCI Developmental Therapeutics Program for providing NCI60 tumor cell line data. We thank Compounds Australia (compoundsaustralia.com.au) for assistance with preparation of assay ready plates of Malaria Box compounds. Thanks to the Australian Red Cross Blood Service and AVIS (Associazione Volontari Italiani Sangue) Milano for provision of human blood and/or sera. We wish to thank all the departments of MMV that worked behind the scenes to make the Malaria Box work, including Finance, Legal, Administration, Human Resources, and Discovery.

Author Contributions

Conceived and designed the experiments: WCVV JHA MHA PA AAlz KTA SVA VMA LA MBa SBa CBM SBh QB LB JBo LEB FFB MBr CRC DC MBC GJC SD'A MD JLD SDu MTF JAFR DAF IF FJG BG TG RKG KKH AH TH PH CH II LTJ AK JK SKic KK DEK SLa ML LMa DM NRM SMa SMc SMe JMe RNM JPM BN CN KKO SAP RJQ RSO RSF NGR FS RSi DAS TSo WS YS SJS DTa DTh KFT ST AKT VT JMV AFW EAW EW JBu TSp TW PAW. Performed the experiments: RA VA PA AAlz YAR AAls AAlz KTA LA QB TB JBr AB CRC GC MC DC MBC KCCC WC AC BLC DDCL YC GJC NC SD'A MD AYD SDu SAESES IF PVTf AG SG BG GMG RG WAG NG KKH PH TBH KIS MAI AJ ANJ AK TK SKic SKim VPK MJL SLe AML FL DLi LLi MIL AL LLu IL DM NRM SMa SMc IMV SMe LMe ANM FMRdC JMü SNH NN SN KKO DO GPan SPa KP NP SPPr DMP SAP AP CQ CAR MAR AR JS NGR MS FSV FS AS SMS SS BMS YS LRYT AT ST AKT VT KOU IU CV EV PVV HV NW KW PMW MW WW NY. Analyzed the data: WCVV JHA RA VA MHA PA AAlz YAR AAls AAlz KTA SVA VMA LA MBa SBa CBM SBh QB LB TB JBo LEB FFB JBr MBr AB CRC GC MC DC MBC KCCC WC AC BLC DDCL YC GJC NC SD'A NLD MD JLD AYD SDu SAESES MTF JAFR DAF IF PVTf AG FJG SG BG TG GMG RG WAG NG RKG MAEH KKH AH RHvH TH PH TBH CH II KIS MAI AJ ANJ LTJ RHYJ AK JK TK SKic SKim KK VPK DEK MJL SLa NL SLe AML FL DLi LLi ML MIL AL LLu IL LMa DM NRM SMa SMc IMV SMe LMe JMe ANM RNM FMRdC SMo JPM JMü AM SNH BN CN NN SN KKO DO GPan GPap SPa KP NP SPPr DMP SAP AP CQ RJQ CAR MAR AR RSO RSF JS NGR US MS FSV FS RSi DAS TSo AS SMS DJS WS SS BMS YS SJS DTa LRYT AT DTh KFT ST AKT VT KOU IU CV EV JMV PVV HV NW KW PMW AFW EAW EW MW WW NY PHAZ NA BB JBu BL DLe TSp TW PAW. Contributed reagents/materials/analysis tools: AYD. Wrote the paper: WCVV JHA RA VA MHA PA AAlz YAR AAls AAlz KTA SVA VMA LA MBa SBa CBM SBh QB LB TB JBo LEB FFB JBr MBr AB CRC GC MC DC MBC KCCC WC AC BLC DDCL YC GJC NC SD'A NLD MD JLD AYD SDu SAESES MTF JAFR DAF IF PVTf AG FJG SG BG TG GMG RG WAG NG RKG MAEH KKH AH RHvH TH PH TBH CH II KIS MAI AJ ANJ LTJ RHYJ AK JK TK SKic SKim KK VPK DEK MJL SLa NL SLe AML FL DLi LLi ML MIL AL LLu IL LMa DM NRM SMa SMc IMV SMe LMe JMe ANM RNM FMRdC SMo JPM JMü AM SNH BN CN NN SN KKO DO GPan GPap SPa KP NP SPPr DMP SAP AP CQ RJQ CAR MAR AR RSO RSF JS NGR US MS FSV FS RSi DAS TSo AS SMS DJS WS SS BMS YS SJS DTa LRYT AT DTh KFT ST AKT VT KOU IU CV EV JMV PVV HV NW KW PMW AFW EAW EW MW WW NY PHAZ NA BB JBu BL DLe TSp TW PAW.

References

1. Gamo FJ, Sanz LM, Vidal J, de Cozar C, Alvarez E, Lavandera JL, et al. Thousands of chemical starting points for antimalarial lead identification. *Nature*. 2010; 465(7296):305–10. doi: [10.1038/nature09107](https://doi.org/10.1038/nature09107) PMID: [20485427](https://pubmed.ncbi.nlm.nih.gov/20485427/).

2. Meister S, Plouffe DM, Kuhen KL, Bonamy GM, Wu T, Barnes SW, et al. Imaging of *Plasmodium* liver stages to drive next-generation antimalarial drug discovery. *Science*. 2011; 334(6061):1372–7. doi: [10.1126/science.1211936](https://doi.org/10.1126/science.1211936) PMID: [22096101](https://pubmed.ncbi.nlm.nih.gov/22096101/); PubMed Central PMCID: PMC3473092.
3. Guiguemde WA, Shelat AA, Bouck D, Duffy S, Crowther GJ, Davis PH, et al. Chemical genetics of *Plasmodium falciparum*. *Nature*. 2010; 465(7296):311–5. doi: [10.1038/nature09099](https://doi.org/10.1038/nature09099) PMID: [20485428](https://pubmed.ncbi.nlm.nih.gov/20485428/); PubMed Central PMCID: PMC2874979.
4. Avery VM, Bashyam S, Burrows JN, Duffy S, Papadatos G, Puthukkuti S, et al. Screening and hit evaluation of a chemical library against blood-stage *Plasmodium falciparum*. *Malar J*. 2014; 13(1):190. Epub 2014/06/03. doi: [10.1186/1475-2875-13-190](https://doi.org/10.1186/1475-2875-13-190) PMID: [24886460](https://pubmed.ncbi.nlm.nih.gov/24886460/).
5. Spangenberg T, Burrows JN, Kowalczyk P, McDonald S, Wells TN, Willis P. The open access malaria box: a drug discovery catalyst for neglected diseases. *PLoS One*. 2013; 8(6):e62906. doi: [10.1371/journal.pone.0062906](https://doi.org/10.1371/journal.pone.0062906) PMID: [23798988](https://pubmed.ncbi.nlm.nih.gov/23798988/); PubMed Central PMCID: PMC3684613.
6. Priest BT, Bell IM, Garcia ML. Role of hERG potassium channel assays in drug development. *Channels (Austin)*. 2008; 2(2):87–93. PMID: [18849661](https://pubmed.ncbi.nlm.nih.gov/18849661/).
7. Khetani SR, Bhatia SN. Microscale culture of human liver cells for drug development. *Nat Biotechnol*. 2008; 26(1):120–6. doi: [10.1038/nbt1361](https://doi.org/10.1038/nbt1361) PMID: [18026090](https://pubmed.ncbi.nlm.nih.gov/18026090/).
8. Chan TS, Yu H, Moore A, Khetani SR, Tweedie D. Meeting the challenge of predicting hepatic clearance of compounds slowly metabolized by cytochrome P450 using a novel hepatocyte model, Hepato-Pac. *Drug metabolism and disposition: the biological fate of chemicals*. 2013; 41(12):2024–32. doi: [10.1124/dmd.113.053397](https://doi.org/10.1124/dmd.113.053397) PMID: [23959596](https://pubmed.ncbi.nlm.nih.gov/23959596/).
9. Overington JP, Al-Lazikani B, Hopkins AL. How many drug targets are there? *Nat Rev Drug Discov*. 2006; 5(12):993–6. doi: [10.1038/nrd2199](https://doi.org/10.1038/nrd2199) PMID: [17139284](https://pubmed.ncbi.nlm.nih.gov/17139284/).
10. Whitebread S, Hamon J, Bojanic D, Urban L. Keynote review: in vitro safety pharmacology profiling: an essential tool for successful drug development. *Drug Discov Today*. 2005; 10(21):1421–33. doi: [10.1016/S1359-6446\(05\)03632-9](https://doi.org/10.1016/S1359-6446(05)03632-9) PMID: [16243262](https://pubmed.ncbi.nlm.nih.gov/16243262/).
11. Fitzgerald LW, Burn TC, Brown BS, Patterson JP, Corjay MH, Valentine PA, et al. Possible role of valvular serotonin 5-HT(2B) receptors in the cardiopathy associated with fenfluramine. *Mol Pharmacol*. 2000; 57(1):75–81. PMID: [10617681](https://pubmed.ncbi.nlm.nih.gov/10617681/).
12. Roth BL. Drugs and valvular heart disease. *N Engl J Med*. 2007; 356(1):6–9. doi: [10.1056/NEJMp068265](https://doi.org/10.1056/NEJMp068265) PMID: [17202450](https://pubmed.ncbi.nlm.nih.gov/17202450/).
13. Yao ZK, Krai PM, Merino EF, Simpson ME, Slebodnick C, Cassera MB, et al. Determination of the active stereoisomer of the MEP pathway-targeting antimalarial agent MMV008138, and initial structure-activity studies. *Bioorg Med Chem Lett*. 2015. Epub 2015/03/11. doi: [10.1016/j.bmcl.2015.02.020](https://doi.org/10.1016/j.bmcl.2015.02.020) PMID: [25754494](https://pubmed.ncbi.nlm.nih.gov/25754494/).
14. Wu W, Herrera Z, Ebert D, Baska K, Cho SH, DeRisi JL, et al. A chemical rescue screen identifies a *Plasmodium falciparum* apicoplast inhibitor targeting MEP isoprenoid precursor biosynthesis. *Antimicrob Agents Chemother*. 2015; 59(1):356–64. Epub 2014/11/05. doi: [10.1128/aac.03342-14](https://doi.org/10.1128/aac.03342-14) PMID: [25367906](https://pubmed.ncbi.nlm.nih.gov/25367906/); PubMed Central PMCID: PMCPmc4291372.
15. Imlay LS, Armstrong CM, Masters MC, Li T, Price KE, Edwards RL, et al. *Plasmodium* IspD (2-C-methyl-D-erythritol 4-phosphate cytidyltransferase), an essential and druggable antimalarial target. *ACS Infectious Diseases*. 2015; 1(4):157–67. PMID: [26783558](https://pubmed.ncbi.nlm.nih.gov/26783558/)
16. Ah Yong V, Sheridan CM, Leon KE, Witchley JN, Diep J, DeRisi JL. Identification of *Plasmodium falciparum* specific translation inhibitors from the MMV Malaria Box using a high throughput in vitro translation screen. *Malar J*. 2016; 15(1):173. Epub 2016/03/19. doi: [10.1186/s12936-016-1231-8](https://doi.org/10.1186/s12936-016-1231-8) PMID: [26987601](https://pubmed.ncbi.nlm.nih.gov/26987601/); PubMed Central PMCID: PMCPmc4794828.
17. Painter HJ, Morrissey JM, Mather MW, Vaidya AB. Specific role of mitochondrial electron transport in blood-stage *Plasmodium falciparum*. *Nature*. 2007; 446(7131):88–91. doi: [10.1038/nature05572](https://doi.org/10.1038/nature05572) PMID: [17330044](https://pubmed.ncbi.nlm.nih.gov/17330044/).
18. Phillips MA, Rathod PK. *Plasmodium* dihydroorotate dehydrogenase: a promising target for novel anti-malarial chemotherapy. *Infectious disorders drug targets*. 2010; 10(3):226–39. Epub 2010/03/26. PMID: [20334617](https://pubmed.ncbi.nlm.nih.gov/20334617/); PubMed Central PMCID: PMCPMC2883174.
19. Phillips MA, Lotharius J, Marsh K, White J, Dayan A, White KL, et al. A long-duration dihydroorotate dehydrogenase inhibitor (DSM265) for prevention and treatment of malaria. *Science Translational Medicine*. 2015; 7(296):296ra111. doi: [10.1126/scitranslmed.aaa6645](https://doi.org/10.1126/scitranslmed.aaa6645) PMID: [26180101](https://pubmed.ncbi.nlm.nih.gov/26180101/)
20. Rottmann M, McNamara C, Yeung BK, Lee MC, Zou B, Russell B, et al. Spiroindolones, a potent compound class for the treatment of malaria. *Science*. 2010; 329(5996):1175–80. doi: [10.1126/science.1193225](https://doi.org/10.1126/science.1193225) PMID: [20813948](https://pubmed.ncbi.nlm.nih.gov/20813948/); PubMed Central PMCID: PMC3050001.
21. Spillman NJ, Allen RJ, McNamara CW, Yeung BK, Winzeler EA, Diagana TT, et al. Na(+) regulation in the malaria parasite *Plasmodium falciparum* involves the cation ATPase PfATP4 and is a target of the

- spiroindolone antimalarials. *Cell host & microbe*. 2013; 13(2):227–37. Epub 2013/02/19. doi: [10.1016/j.chom.2012.12.006](https://doi.org/10.1016/j.chom.2012.12.006) PMID: [23414762](https://pubmed.ncbi.nlm.nih.gov/23414762/); PubMed Central PMCID: [PMCPMC3574224](https://pubmed.ncbi.nlm.nih.gov/PMC3574224/).
22. Vaidya AB, Morrissey JM, Zhang Z, Das S, Daly TM, Otto TD, et al. Pyrazoleamide compounds are potent antimalarials that target Na⁺ homeostasis in intraerythrocytic *Plasmodium falciparum*. *Nature communications*. 2014; 5:5521. Epub 2014/11/26. doi: [10.1038/ncomms6521](https://doi.org/10.1038/ncomms6521) PMID: [25422853](https://pubmed.ncbi.nlm.nih.gov/25422853/).
 23. Jimenez-Diaz MB, Ebert D, Salinas Y, Pradhan A, Lehane AM, Myrand-Lapierre ME, et al. (+)-SJ733, a clinical candidate for malaria that acts through ATP4 to induce rapid host-mediated clearance of *Plasmodium*. *Proc Natl Acad Sci U S A*. 2014. Epub 2014/12/03. doi: [10.1073/pnas.1414221111](https://doi.org/10.1073/pnas.1414221111) PMID: [25453091](https://pubmed.ncbi.nlm.nih.gov/25453091/).
 24. van Pelt-Koops JC, Pett HE, Graumans W, van der Vegte-Bolmer M, van Gemert GJ, Rottmann M, et al. The spiroindolone drug candidate NITD609 potently inhibits gametocytogenesis and blocks *Plasmodium falciparum* transmission to anopheles mosquito vector. *Antimicrob Agents Chemother*. 2012; 56(7):3544–8. Epub 04/18. AAC.06377-11 [pii] doi: [10.1128/AAC.06377-11](https://doi.org/10.1128/AAC.06377-11) PMID: [22508309](https://pubmed.ncbi.nlm.nih.gov/22508309/).
 25. Lehane AM, Ridgway MC, Baker E, Kirk K. Diverse chemotypes disrupt ion homeostasis in the malaria parasite. *Mol Microbiol*. 2014; 94(2):327–39. doi: [10.1111/mmi.12765](https://doi.org/10.1111/mmi.12765) PMID: [25145582](https://pubmed.ncbi.nlm.nih.gov/25145582/).
 26. Ginsburg H, Kutner S, Krugliak M, Cabantchik ZI. Characterization of permeation pathways appearing in the host membrane of *Plasmodium falciparum* infected red blood cells. *Mol Biochem Parasitol*. 1985; 14(3):313–22. Epub 1985/03/01. PMID: [3887158](https://pubmed.ncbi.nlm.nih.gov/3887158/).
 27. Kirk K, Horner HA, Elford BC, Ellory JC, Newbold CI. Transport of diverse substrates into malaria-infected erythrocytes via a pathway showing functional characteristics of a chloride channel. *J Biol Chem*. 1994; 269(5):3339–47. Epub 1994/02/04. PMID: [8106373](https://pubmed.ncbi.nlm.nih.gov/8106373/).
 28. Staines HM, Ellory JC, Kirk K. Perturbation of the pump-leak balance for Na(+) and K(+) in malaria-infected erythrocytes. *Am J Physiol Cell Physiol*. 2001; 280(6):C1576–87. Epub 2001/05/15. PMID: [11350753](https://pubmed.ncbi.nlm.nih.gov/11350753/).
 29. Flannery EL, McNamara CW, Kim SW, Kato TS, Li F, Teng CH, et al. Mutations in the P-type cation-transporter ATPase 4, PfATP4, mediate resistance to both aminopyrazole and spiroindolone antimalarials. *ACS Chem Biol*. 2015; 10(2):413–20. Epub 2014/10/17. doi: [10.1021/cb500616x](https://doi.org/10.1021/cb500616x) PMID: [25322084](https://pubmed.ncbi.nlm.nih.gov/25322084/); PubMed Central PMCID: [PMCPmc4340351](https://pubmed.ncbi.nlm.nih.gov/PMC4340351/).
 30. Alemán Resto Y, Fernández Robledo JA. Identification of MMV Malaria Box inhibitors of *Perkinsus marinus* using an ATP-based bioluminescence assay. *PLoS One*. 2014; 9(10):e111051. doi: [10.1371/journal.pone.0111051](https://doi.org/10.1371/journal.pone.0111051) PMID: [25337810](https://pubmed.ncbi.nlm.nih.gov/25337810/); PubMed Central PMCID: [PMCPMC4206467](https://pubmed.ncbi.nlm.nih.gov/PMC4206467/).
 31. Bessoff K, Spangenberg T, Foderaro JE, Jumani RS, Ward GE, Huston CD. Identification of *Cryptosporidium parvum* Active Chemical Series by Repurposing the Open Access Malaria Box. *Antimicrob Agents Chemother*. 2014; 58(5):2731–9. Epub 2014/02/26. doi: [10.1128/aac.02641-13](https://doi.org/10.1128/aac.02641-13) PMID: [24566188](https://pubmed.ncbi.nlm.nih.gov/24566188/).
 32. Duffy S, Avery VM. Identification of inhibitors of *Plasmodium falciparum* gametocyte development. *Malar J*. 2013; 12(1):408. doi: [10.1186/1475-2875-12-408](https://doi.org/10.1186/1475-2875-12-408) PMID: [24206914](https://pubmed.ncbi.nlm.nih.gov/24206914/); PubMed Central PMCID: [PMC3842684](https://pubmed.ncbi.nlm.nih.gov/PMC3842684/).
 33. Di Matteo M, Matrai J, Belay E, Firdissa T, Vandendriessche T, Chuah MK. PiggyBac toolbox. *Methods Mol Biol*. 2012; 859:241–54. doi: [10.1007/978-1-61779-603-6_14](https://doi.org/10.1007/978-1-61779-603-6_14) PMID: [22367876](https://pubmed.ncbi.nlm.nih.gov/22367876/).
 34. Suzuki Y, St Onge RP, Mani R, King OD, Heilbut A, Labunskyy VM, et al. Knocking out multigene redundancies via cycles of sexual assortment and fluorescence selection. *Nat Methods*. 2011; 8(2):159–64. Epub 2011/01/11. doi: [10.1038/nmeth.1550](https://doi.org/10.1038/nmeth.1550) PMID: [21217751](https://pubmed.ncbi.nlm.nih.gov/21217751/); PubMed Central PMCID: [PMCPmc3076670](https://pubmed.ncbi.nlm.nih.gov/PMC3076670/).
 35. Rizk MA, El-Sayed SA, Terkawi MA, Youssef MA, El Said el Sel S, Elsayed G, et al. Optimization of a Fluorescence-Based Assay for Large-Scale Drug Screening against *Babesia* and *Theileria* Parasites. *PLoS One*. 2015; 10(4):e0125276. doi: [10.1371/journal.pone.0125276](https://doi.org/10.1371/journal.pone.0125276) PMID: [25915529](https://pubmed.ncbi.nlm.nih.gov/25915529/); PubMed Central PMCID: [PMCPMC4411034](https://pubmed.ncbi.nlm.nih.gov/PMC4411034/).
 36. AbouLaila M, Batadoj D, Salama A, Munkhjargal T, Ichikawa-Seki M, MAT, et al. Evaluation of the inhibitory effects of miltefosine on the growth of *Babesia* and *Theileria* parasites. *Veterinary parasitology*. 2014; 204(3–4):104–10. doi: [10.1016/j.vetpar.2014.05.023](https://doi.org/10.1016/j.vetpar.2014.05.023) PMID: [24938827](https://pubmed.ncbi.nlm.nih.gov/24938827/).
 37. Aboulaila M, Munkhjargal T, Sivakumar T, Ueno A, Nakano Y, Yokoyama M, et al. Apicoplast-targeting antibacterials inhibit the growth of *Babesia* parasites. *Antimicrob Agents Chemother*. 2012; 56(6):3196–206. doi: [10.1128/AAC.05488-11](https://doi.org/10.1128/AAC.05488-11) PMID: [22391527](https://pubmed.ncbi.nlm.nih.gov/22391527/); PubMed Central PMCID: [PMCPMC3370714](https://pubmed.ncbi.nlm.nih.gov/PMC3370714/).
 38. Salama AA, Aboulaila M, Moussa AA, Nayel MA, El-Sify A, Terkawi MA, et al. Evaluation of in vitro and in vivo inhibitory effects of fusidic acid on *Babesia* and *Theileria* parasites. *Veterinary parasitology*. 2013; 191(1–2):1–10. Epub 2012/09/19. doi: [10.1016/j.vetpar.2012.08.022](https://doi.org/10.1016/j.vetpar.2012.08.022) PMID: [22985928](https://pubmed.ncbi.nlm.nih.gov/22985928/).

39. Salama AA, AbouLaila M, Terkawi MA, Mousa A, El-Sify A, Allaam M, et al. Inhibitory effect of allicin on the growth of *Babesia* and *Theileria equi* parasites. *Parasitol Res.* 2014; 113(1):275–83. doi: [10.1007/s00436-013-3654-2](https://doi.org/10.1007/s00436-013-3654-2) PMID: [24173810](https://pubmed.ncbi.nlm.nih.gov/24173810/).
40. Taylor MJ, Hoerauf A, Townson S, Slatko BE, Ward SA. Anti-Wolbachia drug discovery and development: safe macrofilaricides for onchocerciasis and lymphatic filariasis. *Parasitology.* 2014; 141(1):119–27. doi: [10.1017/S0031182013001108](https://doi.org/10.1017/S0031182013001108) PMID: [23866958](https://pubmed.ncbi.nlm.nih.gov/23866958/); PubMed Central PMCID: [PMCPMC3884836](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3884836/).
41. Katsuno K, Burrows JN, Duncan K, Hooft van Huijsduijnen R, Kaneko T, Kita K, et al. Hit and lead criteria in drug discovery for infectious diseases of the developing world. *Nat Rev Drug Discov.* 2015; 14:751–8. Epub 2015/10/06. doi: [10.1038/nrd4683](https://doi.org/10.1038/nrd4683) PMID: [26435527](https://pubmed.ncbi.nlm.nih.gov/26435527/).
42. Reader J, Botha M, Theron A, Lauterbach SB, Rossouw C, Engelbrecht D, et al. Nowhere to hide: interrogating different metabolic parameters of *Plasmodium falciparum* gametocytes in a transmission blocking drug discovery pipeline towards malaria elimination. *Malar J.* 2015; 14(1):213. Epub 2015/05/23. doi: [10.1186/s12936-015-0718-z](https://doi.org/10.1186/s12936-015-0718-z) PMID: [25994518](https://pubmed.ncbi.nlm.nih.gov/25994518/).
43. Paiardini A, Bamert RS, Kannan-Sivaraman K, Drinkwater N, Mistry SN, Scammells PJ, et al. Screening the Medicines for Malaria Venture "Malaria Box" against the *Plasmodium falciparum* Aminopeptidases, M1, M17 and M18. *PLoS One.* 2015; 10(2):e0115859. Epub 2015/02/24. doi: [10.1371/journal.pone.0115859](https://doi.org/10.1371/journal.pone.0115859) PMID: [25700165](https://pubmed.ncbi.nlm.nih.gov/25700165/); PubMed Central PMCID: [PMCPMC4336144](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC4336144/).
44. Ruecker A, Mathias DK, Straschil U, Churcher TS, Dinglasan RR, Leroy D, et al. A male and female gametocyte functional viability assay to identify biologically relevant malaria transmission-blocking drugs. *Antimicrob Agents Chemother.* 2014; 58(12):7292–302. doi: [10.1128/AAC.03666-14](https://doi.org/10.1128/AAC.03666-14) PMID: [25267664](https://pubmed.ncbi.nlm.nih.gov/25267664/); PubMed Central PMCID: [PMCPMC4249523](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC4249523/).
45. Fong KY, Sandlin RD, Wright DW. Identification of β -hematin inhibitors in the MMV Malaria Box. *International Journal for Parasitology: Drugs and Drug Resistance.* 2015; 5(6):84–91. doi: [10.1016/j.ijpddr.2015.05.003](https://doi.org/10.1016/j.ijpddr.2015.05.003)
46. Boyom FF, Fokou PV, Tchoukouaha LR, Spangenberg T, Mfopa AN, Kouipou RM, et al. Repurposing the open access malaria box to discover potent inhibitors of *Toxoplasma gondii* and *Entamoeba histolytica*. *Antimicrob Agents Chemother.* 2014; 58(10):5848–54. doi: [10.1128/AAC.02541-14](https://doi.org/10.1128/AAC.02541-14) PMID: [25049259](https://pubmed.ncbi.nlm.nih.gov/25049259/); PubMed Central PMCID: [PMCPMC4187973](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC4187973/).
47. Jefferson T, McShan D, Warfield J, Ogungbe IV. Screening and Identification of Inhibitors of *Trypanosoma brucei* Cathepsin L with Antitrypanosomal Activity. *Chem Biol Drug Des.* 2015. doi: [10.1111/cbdd.12628](https://doi.org/10.1111/cbdd.12628) PMID: [26242248](https://pubmed.ncbi.nlm.nih.gov/26242248/).
48. Kaiser M, Maes L, Tadoori LP, Spangenberg T, Ioset JR. Repurposing of the Open Access Malaria Box for Kinetoplastid Diseases Identifies Novel Active Scaffolds against Trypanosomatids. *J Biomol Screen.* 2015; 20(5):634–45. doi: [10.1177/1087057115569155](https://doi.org/10.1177/1087057115569155) PMID: [25690568](https://pubmed.ncbi.nlm.nih.gov/25690568/).
49. Fletcher S, Avery VM. A novel approach for the discovery of chemically diverse anti-malarial compounds targeting the *Plasmodium falciparum* Coenzyme A synthesis pathway. *Malar J.* 2014; 13:343. doi: [10.1186/1475-2875-13-343](https://doi.org/10.1186/1475-2875-13-343) PMID: [25174342](https://pubmed.ncbi.nlm.nih.gov/25174342/); PubMed Central PMCID: [PMCPMC4168161](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC4168161/).
50. Ingram-Sieber K, Cowan N, Panic G, Vargas M, Mansour NR, Bickle QD, et al. Orally active antischistosomal early leads identified from the open access malaria box. *PLoS Negl Trop Dis.* 2014; 8(1):e2610. Epub 2014/01/15. doi: [10.1371/journal.pntd.0002610](https://doi.org/10.1371/journal.pntd.0002610) PMID: [24416463](https://pubmed.ncbi.nlm.nih.gov/24416463/); PubMed Central PMCID: [PMCPMC3886923](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3886923/).
51. Bowman JD, Merino EF, Brooks CF, Striepen B, Carlier PR, Cassera MB. Antiapicoplast and gametocytocidal screening to identify the mechanisms of action of compounds within the malaria box. *Antimicrob Agents Chemother.* 2014; 58(2):811–9. Epub 2013/11/20. doi: [10.1128/aac.01500-13](https://doi.org/10.1128/aac.01500-13) PMID: [24247137](https://pubmed.ncbi.nlm.nih.gov/24247137/); PubMed Central PMCID: [PMCPMC3910863](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3910863/).
52. Celik H, Hong SH, Colon-Lopez DD, Han J, Kont YS, Minas TZ, et al. Identification of Novel Ezrin Inhibitors Targeting Metastatic Osteosarcoma by Screening Open Access Malaria Box. *Mol Cancer Ther.* 2015. doi: [10.1158/1535-7163.MCT-15-0511](https://doi.org/10.1158/1535-7163.MCT-15-0511) PMID: [26358752](https://pubmed.ncbi.nlm.nih.gov/26358752/).
53. Cowan N, Datwyler P, Ernst B, Wang C, Vennerstrom JL, Spangenberg T, et al. Activities of N,N'-Diarylurea MMV665852 analogs against *Schistosoma mansoni*. *Antimicrob Agents Chemother.* 2015; 59(4):1935–41. doi: [10.1128/AAC.04463-14](https://doi.org/10.1128/AAC.04463-14) PMID: [25583726](https://pubmed.ncbi.nlm.nih.gov/25583726/); PubMed Central PMCID: [PMCPMC4356791](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC4356791/).
54. Frame I, Deniskin R, Rinderspacher A, Katz F, Deng S, Moir RD, et al. Yeast-based High-Throughput Screen Identifies *Plasmodium falciparum* Equilibrative Nucleoside Transporter 1 Inhibitors That Kill Malaria Parasites. *ACS Chem Biol.* 2015. Epub 2015/01/21. doi: [10.1021/cb500981y](https://doi.org/10.1021/cb500981y) PMID: [25602169](https://pubmed.ncbi.nlm.nih.gov/25602169/).
55. Galusic S. MMV Malaria Box activity screening in dormant *Plasmodium falciparum* phenotypes. Thesis, University of South Florida. 2015.
56. Hain AU, Bartee D, Sanders NG, Miller AS, Sullivan DJ Jr, Levitskaya J, et al. Identification of an Atg8-Atg3 protein-protein interaction inhibitor from the Medicines for Malaria Venture Malaria Box active in

- blood and liver stage *Plasmodium falciparum* parasites. *J Med Chem*. 2014. Epub 2014/05/03. doi: [10.1021/jm401675a](https://doi.org/10.1021/jm401675a) PMID: [24786226](https://pubmed.ncbi.nlm.nih.gov/24786226/).
57. Kher SS, Penzo M, Fulle S, Ebejer JP, Finn PW, Blackman MJ, et al. Quinoxaline-Based Inhibitors of Malarial Protease PfSUB1. *Chemistry of Heterocyclic Compounds*. 2015; 50(10):1457–63.
 58. Linares M, Viera S, Crespo B, Franco V, Gomez-Lorenzo MG, Jimenez-Diaz MB, et al. Identifying rapidly parasiticidal anti-malarial drugs using a simple and reliable in vitro parasite viability fast assay. *Malar J*. 2015; 14(1):441. doi: [10.1186/s12936-015-0962-2](https://doi.org/10.1186/s12936-015-0962-2) PMID: [26542470](https://pubmed.ncbi.nlm.nih.gov/26542470/); PubMed Central PMCID: PMCPMC4635989.
 59. Liu L, Richard J, Kim S, Wojcik EJ. Small molecule screen for candidate antimalarials targeting Plasmodium Kinesin-5. *J Biol Chem*. 2014; 289(23):16601–14. Epub 2014/04/17. doi: [10.1074/jbc.M114.551408](https://doi.org/10.1074/jbc.M114.551408) PMID: [24737313](https://pubmed.ncbi.nlm.nih.gov/24737313/); PubMed Central PMCID: PMCPMC4047425.
 60. Lucantoni L, Duffy S, Adjalley SH, Fidock DA, Avery VM. Identification of MMV malaria box inhibitors of plasmodium falciparum early-stage gametocytes using a luciferase-based high-throughput assay. *Antimicrob Agents Chemother*. 2013; 57(12):6050–62. doi: [10.1128/AAC.00870-13](https://doi.org/10.1128/AAC.00870-13) PMID: [24060871](https://pubmed.ncbi.nlm.nih.gov/24060871/); PubMed Central PMCID: PMCPMC3837862.
 61. Lucantoni L, Silvestrini F, Signore M, Siciliano G, Eldering M, Dechering KJ, et al. A simple and predictive phenotypic High Content Imaging assay for Plasmodium falciparum mature gametocytes to identify malaria transmission blocking compounds. *Scientific reports*. 2015; 5:16414. Epub 2015/11/11. doi: [10.1038/srep16414](https://doi.org/10.1038/srep16414) PMID: [26553647](https://pubmed.ncbi.nlm.nih.gov/26553647/).
 62. Njuguna JT, von Koschitzky I, Gerhardt H, Lammerhofer M, Choucry A, Pink M, et al. Target evaluation of deoxyhypusine synthase from *Theileria parva* the neglected animal parasite and its relationship to Plasmodium. *Bioorg Med Chem*. 2014. Epub 2014/06/10. doi: [10.1016/j.bmc.2014.05.007](https://doi.org/10.1016/j.bmc.2014.05.007) PMID: [24909679](https://pubmed.ncbi.nlm.nih.gov/24909679/).
 63. Ramsey NB, Andersen OS. Bilayer Effects of Antimalarial Compounds. *PLoS One*. 2015; 10(11):e0142401. Epub 2015/11/10. doi: [10.1371/journal.pone.0142401](https://doi.org/10.1371/journal.pone.0142401) PMID: [26551613](https://pubmed.ncbi.nlm.nih.gov/26551613/).
 64. Sanders NG, Sullivan DJ, Mlambo G, Dimopoulos G, Tripathi AK. Gametocytocidal Screen Identifies Novel Chemical Classes with Plasmodium falciparum Transmission Blocking Activity. *PLoS One*. 2014; 9(8):e105817. Epub 2014/08/27. doi: [10.1371/journal.pone.0105817](https://doi.org/10.1371/journal.pone.0105817) PMID: [25157792](https://pubmed.ncbi.nlm.nih.gov/25157792/); PubMed Central PMCID: PMCPMC4144897.
 65. Spitzmuller A, Mestres J. Prediction of the P. falciparum target space relevant to malaria drug discovery. *PLoS Comput Biol*. 2013; 9(10):e1003257. Epub 2013/10/23. doi: [10.1371/journal.pcbi.1003257](https://doi.org/10.1371/journal.pcbi.1003257) PMID: [24146604](https://pubmed.ncbi.nlm.nih.gov/24146604/); PubMed Central PMCID: PMCPMC3798273.
 66. Sun W, Tanaka TQ, Magle CT, Huang W, Southall N, Huang R, et al. Chemical signatures and new drug targets for gametocytocidal drug development. *Scientific reports*. 2014; 4:3743. doi: [10.1038/srep03743](https://doi.org/10.1038/srep03743). PMID: [24434750](https://pubmed.ncbi.nlm.nih.gov/24434750/); PubMed Central PMCID: PMC3894558.
 67. von Koschitzky I, Kaiser A. Chemical profiling of deoxyhypusine hydroxylase inhibitors for antimalarial therapy. *Amino Acids*. 2013; 45(5):1047–53. Epub 2013/08/15. doi: [10.1007/s00726-013-1575-0](https://doi.org/10.1007/s00726-013-1575-0) PMID: [23943044](https://pubmed.ncbi.nlm.nih.gov/23943044/).
 68. Turner H. Spiroindolone NITD609 is a novel antimalarial drug that targets the P-type ATPase PfATP4. *Future Med Chem*. 2016. Epub 2016/01/30. doi: [10.4155/fmc.15.177](https://doi.org/10.4155/fmc.15.177) PMID: [26824174](https://pubmed.ncbi.nlm.nih.gov/26824174/).
 69. Tiwari NK, Reynolds PJ, Calderon AI. Preliminary LC-MS Based Screening for Inhibitors of Plasmodium falciparum Thioredoxin Reductase (PfTRxR) among a Set of Antimalarials from the Malaria Box. *Molecules*. 2016; 21(4). Epub 2016/04/05. doi: [10.3390/molecules21040424](https://doi.org/10.3390/molecules21040424) PMID: [27043496](https://pubmed.ncbi.nlm.nih.gov/27043496/).
 70. Stadelmann B, Rufener R, Aeschbacher D, Spiliotis M, Gottstein B, Hemphill A. Screening of the Open Source Malaria Box Reveals an Early Lead Compound for the Treatment of Alveolar Echinococcosis. *PLoS Negl Trop Dis*. 2016; 10(3):e0004535. Epub 2016/03/12. doi: [10.1371/journal.pntd.0004535](https://doi.org/10.1371/journal.pntd.0004535) PMID: [26967740](https://pubmed.ncbi.nlm.nih.gov/26967740/).
 71. Hostettler I, Muller J, Hemphill A. In vitro screening of the open source MMV malaria box reveals novel compounds with profound activities against *Theileria annulata* schizonts. *Antimicrob Agents Chemother*. 2016. Epub 2016/03/16. doi: [10.1128/aac.02801-15](https://doi.org/10.1128/aac.02801-15) PMID: [26976863](https://pubmed.ncbi.nlm.nih.gov/26976863/).
 72. D'Alessandro S, Camarda G, Corbett Y, Siciliano G, Parapini S, Cevenini L, et al. A chemical susceptibility profile of the Plasmodium falciparum transmission stages by complementary cell-based gametocyte assays. *The Journal of antimicrobial chemotherapy*. 2016. Epub 2016/02/19. doi: [10.1093/jac/dkv493](https://doi.org/10.1093/jac/dkv493) PMID: [26888912](https://pubmed.ncbi.nlm.nih.gov/26888912/).
 73. Crowther GJ, Hillesland HK, Keyloun KR, Reid MC, Lafuente-Monasterio MJ, Ghidelli-Disse S, et al. Biochemical Screening of Five Protein Kinases from Plasmodium falciparum against 14,000 Cell-Active Compounds. *PLoS One*. 2016; 11(3):e0149996. Epub 2016/03/05. doi: [10.1371/journal.pone.0149996](https://doi.org/10.1371/journal.pone.0149996) PMID: [26934697](https://pubmed.ncbi.nlm.nih.gov/26934697/).

74. Bilsland E, Bean DM, Devaney E, Oliver SG. Yeast-Based High-Throughput Screens to Identify Novel Compounds Active against *Brugia malayi*. *PLoS Negl Trop Dis*. 2016; 10(1):e0004401. Epub 2016/01/27. doi: [10.1371/journal.pntd.0004401](https://doi.org/10.1371/journal.pntd.0004401) PMID: [26812604](https://pubmed.ncbi.nlm.nih.gov/26812604/).
75. Aroonsri A, Akinola O, Posayapisit N, Songsungthong W, Uthaipibull C, Kamchonwongpaisan S, et al. Identifying antimalarial compounds targeting dihydrofolate reductase-thymidylate synthase (DHFR-TS) by chemogenomic profiling. *Int J Parasitol*. 2016. Epub 2016/05/07. doi: [10.1016/j.ijpara.2016.04.002](https://doi.org/10.1016/j.ijpara.2016.04.002) PMID: [27150044](https://pubmed.ncbi.nlm.nih.gov/27150044/).