

Xavier, N. M; Silva, S.; Madeira, P. J. A; Florêncio, M. H.; Silva, F. V. M.¹ ; Justino, J.¹; Thiem, J.; Rauter, A. P. (2008) - Synthesis and biological evaluation of sugars containing α,β -Unsaturated γ -Lactones. *European journal of organic chemistry*, 36: 6134-6143. Retrieved from: <http://onlinelibrary.wiley.com/doi/10.1002/ejoc.200800763/pdf> DOI: 10.1002/ejoc.200800763

1 - Escola Superior Agrária de Santarém, Inst. Politécnico de Santarém, Apartado 310, 2001-904 Santarém, Portugal

Abstract

The stereocontrolled synthesis of new sugar derivatives carrying the α,β -unsaturated δ -lactone (butenolide) moiety is described. Sugar-fused or sugar-linked butenolides can be constructed by an efficient reaction sequence involving Wittig olefination of 3- or 5-keto sugars and intramolecular cyclization of the intermediate γ -hydroxy α,β -unsaturated esters. The antimicrobial activities of the products and that of a known sugar-derived pyranoid α,β -unsaturated δ -lactone were investigated against six pathogenic bacteria and six fungi. The pyranoid α,β -unsaturated δ -lactone 29 proved to be the most active compound in this series towards the plant pathogenic fungi *Colletotrichum coffeanum* (coffee berry disease) and *Pyricularia oryzae* (rice blast disease).

Keywords: furanosidic systems; scaffolds; Wittig reaction; Lactones; Carbohydrates