

Voskoglou, M. (2000). The Process of Learning Mathematics: A Fuzzy set Approach. *Millenium*, 17

---

## THE PROCESS OF LEARNING MATHEMATICS: A FUZZY SET APPROACH

MICHAEL G. VOSKOGLOU

Technological Educational Institute, Mesolonghi, Greece

The fuzzy sets theory is used for the description of the process of learning and, as an application, a classroom experiment is presented about learning mathematics.

### 1. Introduction.

There are often situations in real life in which definitions do not have clear boundaries; for example this happens when we speak about the "long rivers" or "high mountains" of a country, about the "young people" of a town, about the "tall pupils" of a school, e.t.c.. The fuzzy sets theory was created in response to the need to have a mathematical representation of such kind of situations.

Let  $U$  denote the universal set of the discourse. Then a fuzzy set  $A$  in  $U$ , initiated by Zadeh [7], is defined by means of the membership function  $m_A$ , which assigns to each element of  $U$  a real value from the interval  $[0,1]$ . More specifically  $A = \{(x, m_A(x)) : x \in U\}$ , where  $m_A : U \rightarrow [0,1]$ .

The value  $m_A(x)$ , called the membership degree (or grade) of  $x$  in  $A$ , expresses the degree to which  $x$  verifies the characteristic property of  $A$ . Thus, the nearer the value  $m_A(x)$  to 1, the higher the membership degree of  $x$  in  $A$ .

It is always said that the formal mathematical knowledge comes from the very exactness of the science of Mathematics and therefore there is no possible space in Mathematics for any lack of definition, or vagueness. But, the knowledge that students have about the various mathematical concepts is usually imperfect, characterized by a different degree of depth. This suggests that the

application of the fuzzy sets theory in mathematical education could be a very promising tool in order to represent such situations and get useful conclusions.

## **2. The model of Voss for the process of learning.**

The concept of learning has been fundamental to the study of human cognitive action. But, while everyone in general knows what learning is, the understanding of the nature of this process has proved to be complicated.

Voss [6] has developed an argument that learning is a specific case of the general class of the transfer of knowledge. In adopting this perspective he argues that any instance of acquisition of knowledge involves the use of existing knowledge. Basically, according to his argument, learning consists of successive problem-solving activities, in which the input information is represented of existing knowledge with the solution occurring when the input is appropriately interpreted. The whole process involves the following states:

a) representation of the input data, b) interpretation of these data; c) generalization of the new knowledge and d) categorization of this knowledge.

The representation of the stimulus input is relied upon the individual's ability to use content of his memory in order to find information that will facilitate a solution development.

Learning consists of developing an appropriate number of interpretations of the various input information and generalizing the respective interpretations to a variety of situations. When knowledge becomes substantial, much of the process involves categorization, i.e. the input information is interpreted in terms of the classes of the existing knowledge. Thus the individual becomes able to relate new information to his knowledge structures.

## **3. A fuzzy set representation of the process of learning.**

Let us consider a group  $G$  of  $n$  students during the process of learning in the classroom,  $n \in \mathbb{N}$ ,  $n \geq 2$ .  $\Sigma$

Obviously, from the point of view of the teacher, there exists an uncertainty about the degree of acquisition of each state of the process (as it has been described in the previous paragraph) from

his students, a fact which gave us the hint to introduce the fuzzy sets theory in order to achieve a mathematical representation of the process of learning in the classroom. For this, let us denote by  $A_i$ ,  $i=1, 2, 3$ , the state of interpretation, generalization and categorization respectively and by  $a, b, c, d, e$  the linguistic labels of negligible, low, intermediate, high and complete acquisition respectively of each of the  $A_i$ 's. Consider the set  $U = \{a, b, c, d, e\}$ , then we are going to represent the  $A_i$ 's as fuzzy sets in  $U$ .

In fact, if  $n_{ia}, n_{ib}, n_{ic}, n_{id}$  and  $n_{ie}$  denote the numbers of the students that have achieved negligible, low, intermediate, high and complete acquisition of the state  $A_i$  respectively,  $i=1, 2, 3$ , we can define the membership function  $m_{A_i}$  by  $m_{A_i}(x) = n_{ix}/n$ , for each  $x$  in  $U$  and therefore we can write  $A_i = \{(x, n_{ix}/n) : x \in U\}$ .

It becomes clear then that  $\sum m_{A_i}(x) = 1$ .

$x \in U, i=1, 2, 3$

At this point notice that a fuzzy relationship, like the classical ones, can be considered as a fuzzy set of tuples each one of which possesses a degree of membership included between 0 and 1.

Consider now the fuzzy relation  $R = \{(s, m_R(s)) : s = (x, y, z) \in U^3\}$ , where the membership function  $m_R$  is defined by  $m_R(s) = m_{A_1}(x)m_{A_2}(y)m_{A_3}(z)$ , for all  $s = (x, y, z)$  in  $U^3$ . This definition satisfies the axioms of aggregation operations in fuzzy sets and further we have that  $\sum m_R(s) = 1$  (see section 6.8 of [3], p.p. 282-286).

$s \in U^3$

The fuzzy relation  $R$  represents all the possible profiles (overall states) of the behaviour of a student during the learning process.

In the next, and in order to simplify our notation, we shall write  $m_s$  instead of  $m_R(s)$ .

Assume now that one wants to study the behaviour of  $k$  groups of students during the learning process of the same subject, or the behaviour of the same group of students during the learning process of  $k$  different subjects,  $k \in \mathbb{N}, k \geq 2$ .

In this case it becomes necessary to introduce the fuzzy variables  $A_i(t)$ , where  $i=1, 2, 3$  and  $t=1, 2, \dots, k$ .

Then the pseudofrequency  $f(s)$  of the overall state  $s(t)$  is

$k$

given by the sum  $\sum_{t=1}^k m_s(t)$ , while the probability of  $s(t)$  is

$t=1$

given by  $p(s) = f(s) / \sum_{s \in U} f(s)$ , where  $\sum_{s \in U} f(s)$  denotes the sum of

$s \in U$

all pseudofrequencies. But, since  $\sum_{s \in U} m_s = 1$ , it becomes

$\sum_{s \in U} f(s) = k$

clear that  $\sum_{s \in U} f(s) = k$  and therefore  $p(s) = f(s)/k$ .

$s \in U$

Finally the possibility of  $s(t)$  is given by  $r(s) = f(s) / \max_{s \in U} f(s)$ , where  $\max_{s \in U} f(s)$  denotes the maximal

$s \in U$

pseudofrequency. The possibility of  $s(t)$  measures the degree of evidence of combined results, i.e. in other words one may say that  $r(s)$  gives the "relative probability" of  $s(t)$  with respect to the other overall states

#### **4. An application of the fuzzy sets theory to the process of learning mathematics: A classroom experiment.**

The following experiment took place at the Technological Educational Institute (T.E.I.) of Mesolonghi (Greece), when I was teaching the definite integral to a group of 35 students of the School of Administration and Economics.

During my 3 hour lecture I used the method of rediscovery keeping in mind what Polya [5] says about active learning: "For an effective learning the learner discovers alone the biggest possible, under the circumstances, part of the new information".

Thus, in my short introduction, I presented the concept of the definite integral through the need of calculating an area under a curve, but I gave the fundamental theorem of integral calculus (connecting the definite with the indefinite integral of a continuous function in the interval  $[a, b]$  without proof. Then I left my students to work alone on their papers and I was inspecting their works and reactions, giving to them from time to time suitable hints or instructions. My aim was to help them to understand the basic methods of calculating the definite integrals, with respect to what they already knew about the indefinite integrals (state A1). I observed that 17,8 and 10 students achieved intermediate, high and complete understanding of the new subject respectively ( $n_{1a}=n_{1b}=0$ ,  $n_{1c}=17$ ,  $n_{1d}=8$  and  $n_{1e}=10$ ).

Thus  $A1 = \{(a,0), (b,0), (c,17/35), (d,8/35), (e,10/35)\}$ .

In the next step I gave for solution to the students a number of exercises and simple problems (involving the use of the rules and the basic techniques of integration for the case of definite integrals, calculation of improper integrals as limits of definite integrals, and calculation of the area under a curve or among curves), aiming to help them in generalizing the new information to a variety of situations (state A2).

In the same way I found that

$A2 = \{(a,6/35), (b,6/35), (c,14/35), (d,9/35), (e,0)\}$ .

At the final step I gave to the students for solution a number of composite problems (involving applications to economics, such as present value of cash flows, consumer's and producer's surplus, calculation of probability density functions e.t.c.; see [1], chapter 17) relating the new information to their existing knowledge structures (state A3).

In this case I found that

$A3 = \{(a,12/35), (b,10/35), (c,13/35), (d,0), (e,0)\}$ .

A few days later I gave the same lecture to a group of 30 students of another department of the School of Administration and Economics. This time I found that

$$A1 = \{ (a,0), (b,6/30), (c,15/30), (d,9/30), (e,0) \}$$

$$A2 = \{ (a,6/30), (b,8/30), (c,16/30), (d,0), (e,0) \}$$

$$\text{and } A3 = \{ (a,12/30), (b,9/30), (c,9/30), (d,0), (e,0) \}.$$

Looking at the  $A_i$ 's, it can be seen that, for both groups, the higher the  $i$ , the lower the second members of the pairs having as first members the higher degrees of acquisition  $d$  and  $e$  (in all cases except one). This means that the higher the state of learning, the lower the degree of acquisition of it.

Further, on comparing, in the same way, the degrees of acquisition of the states of learning from the two groups, it can be observed that the first group was better in general than the second one.

In table 1 (see below) the probabilities and possibilities are given of the profiles having non zero pseudofrequencies. It turns out that the profile  $s=(c, c, a)$  had the highest pseudofrequency for the two groups of students of our experiment ( $ms(1)=mA1(c)mA2(c)mA3(a)=(17/35) \cdot (14/35) \cdot (12/35)=0.067$ ,  $ms(2)=(15/30) \cdot (16/30) \cdot (12/30)=0.107$  and therefore  $f(s)=0.174$ ). Therefore it had also the highest probability of occurrence ( $p(s)=0.174/2=0.087$ , or 8.7%), while its possibility was 1.

**TABLE 1**

Probabilities and possibilities of profiles ( $f(s)=0$ ) of the classroom experiment (section 3)

A <sub>1</sub>	A <sub>2</sub>	A <sub>3</sub>	ms(1)	ms(2)	f(s)	p(s)	r(s)
b	b	b	0	0.016	0.016	0.008	0.092
b	a	b	0	0.012	0.012	0.006	0.069
b	c	b	0	0.032	0.032	0.016	0.184
b	b	a	0	0.021	0.021	0.010	0.121
b	b	c	0	0.016	0.016	0.008	0.092
b	a	a	0	0.016	0.016	0.008	0.092
b	a	c	0	0.012	0.012	0.006	0.069
b	c	a	0	0.042	0.042	0.021	0.241
b	c	c	0	0.032	0.032	0.016	0.184
c	c	c	0.072	0.080	0.152	0.076	0.874
c	a	c	0.082	0.030	0.112	0.056	0.644
c							
c	b	c	0.031	0.040	0.071	0.036	0.408
c	d	c	0.046	0	0.046	0.023	0.264
c	c	a	0.067	0.107	0.174	0.087	1
c	c	b	0.056	0.008	0.064	0.032	0.368
c	a	a	0.028	0.040	0.068	0.034	0.391
c	a	b	0.024	0.030	0.054	0.027	0.310
c	b	a	0.028	0.053	0.081	0.040	0.466
c	b	b	0.024	0.040	0.064	0.032	0.368
c	d	a	0.043	0	0.043	0.022	0.247
c	d	b	0.036	0	0.036	0.018	0.207
d	d	a	0.020	0	0.020	0.010	0.115
d	d	b	0.017	0	0.017	0.008	0.098
d							
d	d	c	0.022	0	0.022	0.011	0.126
d	a	a	0.013	0.024	0.037	0.018	0.213
d	a	b	0.011	0.018	0.029	0.014	0.167
d	a	c	0.015	0.018	0.033	0.016	0.190
d	b	a	0.013	0.032	0.045	0.022	0.259
d	b	b	0.011	0.024	0.035	0.018	0.201
d	b	c	0.014	0.024	0.038	0.019	0.218
d	c	a	0.031	0.064	0.095	0.048	0.546
d	c	b	0.026	0.048	0.074	0.037	0.425
d	c	c	0.034	0.048	0.082	0.041	0.471
e	a	a	0.017	0	0.017	0.008	0.098
e	a	b	0.014	0	0.014	0.007	0.080
e	a	c	0.018	0	0.018	0.009	0.103
e	b	a	0.017	0	0.017	0.008	0.098
e	b	b	0.014	0	0.014	0.007	0.080
e	b	c	0.018	0	0.018	0.009	0.103
e	c	a	0.039	0	0.039	0.020	0.224
e	c	b	0.033	0	0.033	0.016	0.190
e	c	c	0.042	0	0.042	0.021	0.241
e	d	a	0.025	0	0.025	0.012	0.144
e	d	b	0.021	0	0.021	0.010	0.121
e	d	c	0.027	0	0.027	0.014	0.155

NOTE : In Table 1 we should have  $\sum ms(i)=1$  ( $i=1,2$ ),

$S \in U3$

$\sum f(s)=2, \sum p(s)=1$  and  $\sum r(s)=(\sum f(s))/\max f(s)=2/0.174 =$

$s \in U3 \ s \in U3 \ s \in U3 \ s \in U3 \ s \in U3$

$=11.494$ , but, because of the roundoff errors appeared, we get slightly different values for the sums above.

### **5. Remarks and conclusions.**

From the experiment of the previous paragraph it becomes evident that the use of the fuzzy sets theory for the mathematical representation of the process of learning, apart from its theoretical interest, leads to useful quantitative conclusions which can give an effective help to the teacher of mathematics in getting a concentrating view of his students abilities and skills and therefore to readapt, according to each case, the process, the rate and the method of his tuition.

It becomes also evident that the same method, with the proper each time modifications, could be used in the classroom during the learning process, not only of mathematics but of many other cognitive subjects.

We note that an analogous application of the fuzzy sets theory has been attempted on the van Hiele model for the teaching of the Euclidian Geometry [4].

Finally we must note that analogous efforts, but with different methodologies of development, have been attempted and by other investigators in the area of mathematical education (e.g. see [2] and its references).



## REFERENCES

- [1] DOWLING E. T. , Mathematics for Economists, Schaum's Outime Series, Mc Graw - Hill, New York, 1980.
- [2] ESPIN E. A. - OLIVERAS C. M. L. , Introduction to the use of the fuzzy logic in the assessment of mathematics teachers' Proceedings 1st Mediterranean Conf. Math., 107-113 , Cyprus,1997.
- [3] KLIR G. J. - FOLGER T. A. , Fuzzy sets ' Uncertainty and Information, Prentice - Hall Int., London, 1988.
- [4] PERDIKARIS 5., Mathematizing the van Hiele levels: a fuzzy set approach, Int. J. Math. Educ. Sci. Technol., 27, 41-47, 1996.
- [5] POLYA G., On learning , teaching and learning teaching American Math. Monthly, 70, 605-619, 1963.
- [6] VOSS J. F., Learning and transfer in subject-matter learning: A problem solving model, Int. J. Educ. Research,11, 607-622, 1987.
- [7] ZADEH, L. A. Information and Control, 8,. 338-353, 1965.