

The North Atlantic is the primary deep ventilator of the oceans playing a key role in the thermohaline circulation through the Atlantic Meridional Overturning Circulation whose strength can be affected by regional or global climatic changes. During abrupt climate change events, e.g. Heinrich Events, the southward migration of the Polar Front, induced by the discharges of icebergs and melt-water pulses, led to abrupt decreases in sea surface temperatures and salinity thereby affecting primary productivity. Paleoproductivity signals in the North Atlantic are controversial and not straightforward (Villanueva et al., 2001). Coccolithophores are primary producers and the most important calcifying organisms in the ocean, playing a key role not only in the ecosystem but also in the global carbon cycle. The coccolith Sr/Ca ratio is linked to productivity with a higher ratio indicating higher growth rates and thus coccolith productivity.

This study infers paleoproductivity variability based on coccolith Sr/Ca ratios at IODP Site U1313 (41°N, 33°W) between 676 and 355 ka ago. MIS 12 is considered the most extreme glacial of the last 500 ka and experienced Heinrich-type ice-rafting events while MIS 14 was a relative mild glacial. For the MIS 12-10 interval, for which data corrections and interpretation have already been completed, the Sr/ Ca data clearly shows that coccolithophore productivity was substantially higher during glacial than interglacial or interstadial periods. Heinrich events, conversely, greatly diminished paleo-productivity.

**Palavras chave:** circulação termohalina, coccolitóforos, produtividade, Sr/Ca, evento Heinrich-type.

**Keywords:** *thermohaline circulation, coccolithophores, productivity, Sr/Ca, Heinrich-type event.*

## Proxy calibration: What are coccolithophores of Galician waters and surface sediments telling us?

Calibração de proxy: O que nos dizem os coccolitóforos e coccolitos encontrados nas águas da Galiza e sedimentos superficiais?

**C. Cavaleiro (1,2,3), E. Salgueiro (1,2,3), A. Silva (4), A. Narciso (5), J. Abel (6), B. Arbones (7), D. Zúñiga (7), F. de la Granda (7), N. Villaceros-Robineau (7), F. Alonso (7), C. G. Castro (7)**

- (1) IPMA – Instituto Português do Mar e da Atmosfera. [catarina.cavaleiro@lneg.pt](mailto:catarina.cavaleiro@lneg.pt)
- (2) Unidade de Geologia Marinha, Laboratório Nacional de Energia e Geologia, Lisboa, Portugal.
- (3) CIMAR - Centro de Investigação Marinha e Ambiental, Portugal
- (4) IPMA – Instituto Português do Mar e da Atmosfera (former INRB/MADRP), Portugal.
- (5) Departamento de Geologia, Faculdade de Ciências de Lisboa, Portugal.
- (6) Grupo de Geociencias Oceanicas, Universidade de Salamanca, Espanha.

- (7) Grupo de Oceanologia do Instituto de Investigações Mariñas (CSIC), Vigo, Espanha.

Calibration of nowadays proxies, namely micro and nanofossils, validate their use in paleoenvironmental reconstructions. CALIBERIA project, under which this study has been developed, aims a multi-proxy calibration from the water column into the surface sediments off Galician margin.

Coccolithophores are one of the major groups of marine phytoplankton and among the most important pelagic calcifying organisms. They affect the climate system and global carbon cycle through biological (sink of organic carbon) and alkalinity pumps (sink of inorganic carbon). Coccoliths, little platelets that constitute their calcareous exoskeleton, can be used as tracers for paleoceanographic reconstructions. Coccolithophores are thus a paleoclimatic tracer due to their stable calcite skeleton and wide geographic distribution under direct climatic control.

This work aims to describe and understand the distribution of coccolithophores off Galician margin based on (1) water column samples collected at two stations; one on the continental shelf (75 m depth) and the other on the shelf-break (350 m depth) at aprox. 42°N, (2) sediment trap samples from a subsurface deploy at the inshore station and (3) sediment surface samples collected between 41.5° - 42°N and 9 - 10°W. The integration of these three data sets will, for the first time, to allow us to improve our coccolithophore proxy for this area. Additionally, hydrographic (temperature and salinity from CTD profiler) and biogeochemical data of discrete water column samples, such as oxygen, inorganic nutrients and stable isotopes, will be used to study the ecology of coccolithophores in this coastal upwelling region.

**Palavras chave:** calibração, paleoceanografia, cocólito, abundâncias, ecologia.

**Keywords:** *calibration, paleoceanography, coccolithophores, Galician margin.*

## A Preliminary Environmental Sensitivity Index (ESI) Map to assist oil spill management in the Aveiro coastal and estuarine region

Mapa Preliminar de Índices de Sensibilidade Ambiental (ESI) para apoio à gestão de derrames petrolíferos na região costeira e estuarina de Aveiro

**C. R. Lemos (1), A. Luís (2), L. M. Pinheiro (1), E. Levine (3), C. Coelho (4)**

- (1) CESAM & Departamento de Geociências, Universidade de Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal. [catarinalemos@ua.pt](mailto:catarinalemos@ua.pt)
- (2) Departamento de Biologia, Universidade de Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal.