

Isotopic Zn and Cd multitracing study of metal transfer processes in soils and lichens from an old mining area (São Domingos Mine, Portugal)

Mattielli Nadine ^{1*}, Africano Fatima ², Branquinho Christina ³, Delahaye Arno, Vogel-Mota Réjane, Couder Eléonore, Matos Joao X. ⁴

- 1 : G-Time, Département des Sciences de la Terre et de l'Environnement, Université Libre de Bruxelles (ULB)
CP 160/02, 50 Av. FD Roosevelt, 1050 Brussels
- 2 : Centro de Geologia, Faculdade de Ciências, Universidade de Lisboa
Campo Grande 1749-016 Lisbon
- 3 : Centro de Biologia Ambiental, Faculdade de Ciências, Universidade de Lisboa
Campo Grande 1749-016 Lisbon
- 4 : Portuguese Geological Survey (LNEG)
R. Frei Amador, 39 rc. Ap. 104, 7801-902 Beja
- * : Corresponding author

Isotopic multitracing approach has been conducted in an old mining area for a better understanding of the metal transfer processes in a semi-arid environment.

Belonging to the Iberian Pyrite Belt (Alentejo, Portugal), the São Domingos Mine (SDM) accounted for one of the most important copper-pyrite mining and refinery area of the Iberian Peninsula until its closure in 1966. In the neighbourhood of the São Domingos village, about 5x10⁶ metric ton of waste deposits and an acid mine drainage system (AMD) are observed.

Elemental and multi-isotope Zn-Cd analyses were performed on lichen samples, topsoils, unprocessed and processed ores and AMD salts collected within a 16 km² surface area.

As, Cd, Pb, Zn, Cu and Mn are identified as main atmospheric metal pollutants in soils and lichens, with maximum values recorded inside the village (As from 78 to 1500mg/kg). Lichens display higher enrichments over a wider area relative to soils.

Elemental and Zn isotopic compositions of the soils samples clearly distinguish highly contaminated soils ($d_{66}\text{Zn} = -0.69 \pm 0.03$ to $+0.07 \pm 0.05$? (2SD); Zn from 300 to 1054mg/kg; Pb from 1500 to 3745mg/kg) from geogenic soils ($d_{66}\text{Zn} = +0.10 \pm 0.05$ to $+0.22 \pm 0.04$? ; Zn from 23 to 150mg/kg; Pb from 29 to 100mg/kg). The lowest $d_{66}\text{Zn}$ values are observed in the village.

$d_{66}\text{Zn}$ values of the lichens are only consistent with the soil isotopic signatures at the geogenic soil sites. At the highly contaminated spots, lichens significantly differ from the soils because either (1) in the village, lichen $d_{66}\text{Zn}$ values ($+0.05 \pm 0.01$ to $+0.11 \pm 0.02$?) reflect re-suspension of unprocessed Cu-pyrite tailings, or (2) in the valley, the lichen $d_{66}\text{Zn}$ values (-0.07 ± 0.03 to -0.05 ± 0.04 ?) reflect Zn fractionation by interaction with AMD salts (jarosite).

Preliminary $d_{114}\text{Cd}$ data in soils show enrichment in light Cd isotopes, which suggests a strong effect of Cd adsorption on Fe-Mn oxi-hydroxides in regosols.

Zn-Cd isotopic multitracers applied to soils and lichens constitute a sensitive approach to discriminate processes that might simultaneously occur in complex metal polluted environments.