

Journal of Minerals and Materials Characterization and Engineering, 2014, 2, 135-143  
Published Online March 2014 in SciRes. <http://www.scirp.org/journal/jimmce>  
<http://dx.doi.org/10.4236/jimmce.2014.22018>



# Towards the Recovery of By-Product Metals from Mine Wastes: An X-Ray Absorption Spectroscopy Study on the Binding State of Rhenium in Debris from a Centennial Iberian Pyrite Belt Mine

Maria-Ondina Figueiredo<sup>1,2\*</sup>, Teresa Pereira da Silva<sup>2</sup>, João Pedro Veiga<sup>1</sup>, Daniel de Oliveira<sup>2</sup>, Maria João Batista<sup>2</sup>

<sup>1</sup>CENIMAT/I3N, Faculty of Sciences and Technology, New University of Lisbon, Campus da Caparica, Lisbon, Portugal

<sup>2</sup>Unit of Mineral Resources and Geophysics, National Laboratory for Energy and Geology, Amadora, Portugal

Email: \*[ondina.figueiredo@lneg.pt](mailto:ondina.figueiredo@lneg.pt)

Received 3 December 2013; revised 3 January 2014; accepted 20 January 2014

Copyright © 2014 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

## Abstract

Rhenium is a very scarce element, occurring in the Earth's crust mainly carried by molybdenite ( $\text{MoS}_2$ ). Due to a very low availability comparative to actual industrial demand, rhenium is nowadays one of the most expensive mineral commodities and an increased interest is focused on exploring residues resulting from a long-term mining, particularly of sulphide ore deposits. It is therefore noteworthy to assign the presence of rhenium (in a concentration up to 3 ppm) in the waste materials from the old sulphur factory at the abandoned mine of São Domingos (Iberian Pyrite Belt, Southeast Portugal), exploited since the Roman occupation of Iberia. Aiming at a potential sustainable recovery of rhenium as a by-product, X-ray near-edge absorption spectroscopy (XANES) was applied to clarify the Re-binding and mode of occurrence by comparing Re  $L_3$ -edge XANES spectra obtained from mine waste samples (previously fully characterized by X-ray laboratory techniques) with similar spectra collected from Re-rich molybdenites ( $\text{Mo}_{1-x}\text{Re}_x\text{S}_2$ ) and from Re-O model compounds configuring various valences and coordination environments of rhenium ions. Obtained results are commented, ruling out a possible Re-S binding and rather conforming with the binding of rhenium to oxygen in the analysed mine waste materials.

## Keywords

Mine Waste Materials By-Products; Iberian Pyrite Belt; Rhenium; XANES; Re  $L_3$ -Edge

\*Corresponding author.

**How to cite this paper:** Figueiredo, M.O., Silva, T.P., Veiga, J.P., de Oliveira, D. and Batista, M.J. (2014) Towards the Recovery of By-Product Metals from Mine Wastes: An X-Ray Absorption Spectroscopy Study on the Binding State of Rhenium in Debris from a Centennial Iberian Pyrite Belt Mine. *Journal of Minerals and Materials Characterization and Engineering*, 2, 135-143. <http://dx.doi.org/10.4236/jimmce.2014.22018>