#### INTRODUCTION

Traditional Madeira wines are "baked" and aged. One of the aging volatiles is sotolon (4,5-dimethyl-3-hydroxy-2 (5H)-furanone), which imparts an aroma usually described as caramel, burnt sugar, "madeirised", "rancio", nutty and curry-like, and which increases in concentration with age [1]. The sensory effect of sotolon on the "perceived age" of Madeiras was evaluated by comparing unbaked wines with added sotolon to traditional baked wines. Six traditional Madeiras made with Bual grapes (vintage 1961, 1968, 1980, 1993, 1996, and 1999), and 3 young, unbaked Bual wines with 0 ppb, 10 ppb and 19 ppb added sotolon, respectively, were compared by sensory descriptive analysis.

## GOAL:

Evaluate the sensory impact of sotolon on the "perceived age" of Madeira Wine



## 1st STEP: Establish the sensory profile of Madeiras



#### 2nd STEP: Training Sessions

Descriptive Analysis Panel Training and Selection. Twelve University of California, Davis, graduate students were selected from a group of 32 persons on the basis of their sensory performances [2].




#### 3rd STEP: Descriptive Analysis

The wines used in this study were 6 traditional Madeiras made with Bual grapes that were 5, 8, 11, 24, 38 and 43 years-old (vintages 2004-1999, 1996, 1993, 1980, 1968, 1961) and 3 young, unbaked Bual wines, from 2003, with 0 ppb, 10 ppb and 19 ppb added sotolon (Table.1). All wines were donated by The Madeira Wine Company, Funchal, Madeira, Portugal.



Sotolon was donated by International Flavors & Fragrances, Inc.(150 Docks Corner Road, Dayton,

## 5th STEP: Colour Analysis. Differences in colour between wines



# 6th STEP: Data Analysis

Data Analysis: All statistical analyzes were performed using SAS (SAS Institutes, Cary, NC). The results of all the descriptive analysis sessions were analyzed by analysis of variance and multivariate analysis of variance (proc glm, with judges as a random factor), to test if the 19 flavor attributes generated by the trained panel, discriminated significantly among the wines. Calculations of the least significant difference (LSD) [3] were applied for validation of the comparison of the different wines for each significant attribute. Principal component analysis was applied to disclose the interrelationship among the wines.

A plot produced from the frequency data associated with each wine and color chips by correspondence analysis (proc corresp) graphically showed the relationship among the colors of the different wines.

# **MADEIRA WINES:**



# SENSORY DESCRIPTIVE ANALYSIS OF TRADITIONAL WINES

## AND WINES WITH ADDED SOTOLON

# Beatriz Machado\*, A. C. Silva Ferreira\*, Hildegarde Heymann\*\*

\*Escola Superior de Biotecnologia, Universidade Católica Portuguesa Rua Dr. Ántónio Bernardino de Almeida 4200-072 Porto, Portugal
\*\*Department of Viticulture and Enology, University of California, Davis, One Shields Avenue, Davis, 95616 CA, USA Hildegarde Heymann, Ph.D. Telephone: 001-530-754-4816; Fax 001-530-752-0882; E-mail address: hheymann@ucdavis.edu

### RESULTS


### ·ANOVA:




significantly different across 16 of the 19 flavor attributes generated by the trained panel.

significantly among the wines using for error the wine\*judge

interactions.



### **•COLOURS: VINTAGES ONLY**



# CONCLUSION

The ANOVA results showed that the wines were significantly different across 16 of the 19 flavor attributes generated by the trained panel.

PCA, results (Fig 1) showed that the baked and the unbaked wines were located on opposite ends of PC1. The unbaked wines were higher in musty, mushroom and dried fruit flavor-by-mouth, while the baked wines had more complexity in flavor and were higher in multiple attributes. The older vintages from were characterized by higher intensities of cherry, nutty, coffee, astringency and cocoa flavors-by-mouth. The younger vintages were higher in vanilla, caramel, oak, brown sugar rum, baked and citrus flavors-by-mouth.

Correspondence analysis was used to analyze the color data for all the wines. Using only the traditional Madeira wines results showed that the first 2 axes accounted for 89% of the variance. In Fig 2, the results show a progression of increasing brownness with increasing age up to about 23 years of aging.

Although the two unbaked wines with added sotolon seemed to be more complex in flavor than the one without, this compound alone did not make them similar

The baking process is important to the flavors of traditional Madeiras. A future modification of this study would be to add the sotolon after baking but before aging and to use a trained panel of connoisseurs of aged dessert wines, because the majority of persons in the panel were not familiar with the characteristics of old



POCTI
This work was financially supported by Fundação para a Ciência e a Tecnologia (FCT), as part of the Projects - POCTI / AGRI / 41'92 / 2002.
- "IMPACT" POCTI, Agencia de Inovação SA and FEDER)