



CATÓLICA
UNIVERSIDADE CATÓLICA PORTUGUESA | PORTO
Escola Superior de Biotecnologia

***THE INFLUENCE OF THE GRAPE VARIETY ON THE
PRODUCTION OF VOLATILE PHENOLS IN
PORTUGUESE WINES***

Master Thesis presented to *Escola Superior de Biotecnologia* of the *Universidade Católica Portuguesa* as part of the European Master of Science Degree in Food Science, Technology and Nutrition

By Sophie Weiss

Under the supervision of Francisco Campos and José António Couto

March 2014

ABSTRACT

*The main purpose of this work is to determine whether there is a correlation between the susceptibility to volatile phenol production of wines and grape variety. Therefore, 11 single varietal red wines from Portugal were heat sterilized, contaminated with *Dekkera bruxellensis* PYCC 4801 and incubated for at least 10 days at 30°C. Since yeasts did not grow in pure wines due to inhibition by ethanol, the experiments were conducted in diluted samples with similar initial pH and ethanol levels. Yeast growth was monitored and produced levels of 4-ethylguaiacol, 4-ethylphenol and 4-vinylphenol were measured using gas chromatography in order to compare values between varieties.*

*This approach led to the production of 4-ethylphenol and 4-ethylguaiacol in all samples; 4-vinylphenol however, could not be detected in any sample. All values reached or exceeded perception threshold levels established in literature. This suggests that all examined wines are at risk of developing phenolic off-flavour once contaminated with *Brettanomyces/Dekkera* yeasts, especially considering that samples were diluted and thus contained lower amounts of precursors than are potentially present in pure wines.*

4-ethylguaiacol values were generally lower and varied less between samples and varieties compared to 4-ethylphenol. Tinta Roriz and Touriga Franca showed the highest potential in regard to volatile phenol production from the natural precursors available in the wines while Sousão wines appeared to be the least prone and Touriga Nacional wines exhibited intermediate volatile phenol values. Furthermore, 4-ethylphenol:4-ethylguaiacol ratios were calculated and show differences between and similarities within varieties.

An attempt was made to estimate the precursor quantities originally present in the wines by comparing the results of spiked and unspiked wines but the obtained results were inconclusive.

From the results it can be suggested that the grape variety may have an influence on the production of volatile phenols by predetermining the availability of precursors. However, several other factors including vinification and wine aging methods may have an impact on the production of volatile phenols.

ACKNOWLEDGEMENTS

Foremost, I would like to thank Escola Superior de Biotecnologia da Universidade Católica Portuguesa (ESBUCP) for accepting me for this Master's thesis project and providing all the instrumental support necessary for conducting the research.

In particular, I would like to express my great appreciation to my supervisors Francisco Campos and José Antonio Couto for the continuous support at all times, motivation and help throughout the project.

I would also like to thank the winemakers from the Douro and Alentejo region who have provided me with samples for the work.

Finally, I would like to dedicate this project to my parents who have been supporting me throughout my studies and all the friends I have made during this master program.

ABBREVIATIONS

CFU	Colony Forming Units
FID	Flame Ionisation Detector
HCA	Hydroxycinnamic Acids
LAB	Lactic Acid Bacteria
log	Logarithm to the base 10
PAD	Phenolic Acid Decarboxylase
S1-S4	Sousão
SY	Syrah
TF	Touriga Franca
TN1-TN4	Touriga Nacional
TR	Tinta Roriz
v	Volume
4EC	4-Ethylcatechol
4EG	4-Ethylguaiacol
4EP	4-Ethylphenol
4VC	4-Vinylcatechol
4VG	4-Vinylguaiacol
4VP	4-Vinylphenol

ABBREVIATIONS OF GENUS NAMES

<i>B.</i>	<i>Brettanomyces</i>
<i>C.</i>	<i>Candida</i>
<i>D.</i>	<i>Dekkera</i>
<i>L.</i>	<i>Lactobacillus</i>
<i>P.</i>	<i>Pichia</i>
<i>Ped.</i>	<i>Pediococcus</i>
<i>S.</i>	<i>Saccharomyces</i>

LIST OF TABLES

Table 1: Most common volatile phenols found in wine, their synthesis pathway and odour ..	12
Table 2: Concentration ranges of volatile phenols in red wine	13
Table 3: Variety, origin and analytical parameters of the wine samples.....	24
Table 4: Sample numbers and their composition.....	25
Table 5: Range of retention times for calibrated volatile phenols and internal standard	27
Table 6: Cell concentration of inocula	28
Table 7: Mean potential of varieties to produce 4-ethylphenol in diluted wines in ppm	39
Table 8: Molar conversion rates (%) of p-coumaric acid to 4-ethylphenol by <i>D. bruxellensis</i> in different varietal wines fortified with the precursor	40
Table 9: Molar conversion rates (%) of ferulic acid to 4-ethylguaiacol by <i>D. bruxellensis</i> in different varietal wines fortified with the precursor	41

LIST OF FIGURES

Figure 1: Phenolic acids found in grapes and wine	14
Figure 2: Average contents of phenolic acids in wines from continental Portugal and Azores	15
Figure 3: Formation pathway of volatile phenols via the decarboxylation of hydroxycinnamic acids	16
Figure 4: Calibration curves for 4-ethylguaiacol, 4-ethylphenol and 4-vinylphenol including trendline equations	27
Figure 5: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Touriga Franca (TF) at 30°C.....	29
Figure 6: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Tinta Roriz (TR) at 30°C	29
Figure 7: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Syrah (SY) at 30°C	30
Figure 8: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Touriga Nacional (TN1) at 30°C.	30
Figure 9: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Touriga Nacional (TN2) at 30°C.	30
Figure 10: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Touriga Nacional (TN3) at 30°C	31
Figure 11: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Touriga Nacional (TN4) at 30°C	31
Figure 12: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Sousão (S1) at 30°C.....	31
Figure 13: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Sousão (S2) at 30°C.....	32
Figure 14: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Sousão (S3) at 30°C.....	32
Figure 15: Growth of <i>D. bruxellensis</i> PYCC 4801 in diluted Sousão (S4) at 30°C.....	33
Figure 16: Volatile phenol concentration in examined red wines	33
Figure 18: Initial volatile phenol concentration in Tinta Roriz (TR) and Touriga Franca and amount present in diluted wine after contamination with <i>D. bruxellensis</i> PYCC 4801	35
Figure 19: Initial volatile phenol concentration in Syrah (SY) and the amount present in diluted wine after contamination with <i>D. bruxellensis</i> PYCC 4801	35
Figure 20: Initial volatile phenol concentration in Touriga Nacional wines and the amount present in diluted wines after contamination with <i>D. bruxellensis</i> PYCC 4801	36
Figure 21: Initial volatile phenol concentration in Sousão wines compared to the amount present in diluted wines after contamination with <i>D. bruxellensis</i> PYCC 4801	37
Figure 22: Concentration of 4-ethylguaiacol produced by <i>D. bruxellensis</i> in diluted varietal red wines.....	38
Figure 23: Concentration of 4-ethylphenol produced by <i>D. bruxellensis</i> in diluted varietal red wines.....	39

OUTLINE

ABSTRACT.....	2
ACKNOWLEDGEMENTS	3
ABBREVIATIONS	4
ABBREVIATIONS OF GENUS NAMES.....	5
LIST OF TABLES.....	6
LIST OF FIGURES	7
1 INTRODUCTION.....	10
2 LITERATURE REVIEW	11
2.1 Phenolic compounds in wine	11
2.2 Volatile phenols in wine.....	11
2.2.1 Concentration ranges.....	12
2.2.2 Perception thresholds	13
2.2.3 Precursors.....	13
2.2.4 Synthesis	15
2.3 Microbial origin	16
2.3.1 <i>Dekkera/Brettanomyces</i>	18
2.3.2 Sources of contamination.....	19
2.4 Grape variety and volatile phenol production by <i>Brettanomyces bruxellensis</i>	20
2.4.1 Portuguese wines.....	21
2.5 Other factors influencing volatile phenol production	22
3 MATERIALS AND METHODS	24
3.1 Wine samples.....	24
3.2 Yeast strain.....	25
3.3 Preparation of phenolic compound solutions	25
3.4 Sample preparation	25
3.5 Determination of yeast growth.....	26
3.6 Extraction of volatile phenols	26
3.7 Gas chromatography	26
3.8 Calibration curves	27

4	RESULTS.....	28
4.1	Inoculum size	28
4.2	Yeast growth	28
4.3	Volatile phenols in varietal wines.....	33
4.4	Volatile Phenols production by <i>D. bruxellensis</i> PYCC 4801 in diluted wines.....	34
4.4.1	Estimation of the potential for volatile phenol production in diluted wines	38
4.4.2	Estimation of volatile phenol precursor quantities in diluted wines.....	39
5	DISCUSSION.....	42
5.2	Volatile phenols production.....	43
5.3	Potential of volatile phenol production	44
5.4	Estimation of precursor quantities	45
6	CONCLUSIONS	46
	REFERENCES.....	47

1 INTRODUCTION

Volatile phenols such as 4-ethylphenol, 4-ethylguaiacol, 4-vinylguaiacol and 4-vinylphenol impart the so-called phenolic off-flavour that is described as inducing barnyard, horse sweat or leather-like defects in wine aroma. It can also be referred as to "Brett character" since volatile phenols are formed by *Brettanomyces/Dekkera* yeasts utilizing phenolic precursors present in grapes and wine. This alteration, mainly occurring in red wines is considered detrimental to wine quality leading to rejection of those concerned wines. Therefore, since the accumulation of volatile phenols carries a high risk of financial losses for the wine industry, the interest in control and prevention measures of *Brettanomyces/Dekkera* spoilage has been rising in the past few years.

While several factors influencing the formation of volatile phenols in wine have been widely discussed, little research has been focused on the impact of the grape variety. Empirical evidence obtained by Portuguese winemakers suggests that some grape varieties may be more susceptible to *Brettanomyces/Dekkera* spoilage and production of volatile phenols than others. In spite of Portugal having a substantial diversity of native grape varieties, those have not yet been studied in as much detail as varieties from other origins, especially with regard to production of volatile phenols. Therefore, this project aims to evaluate whether there are recognizable differences in the susceptibility of different Portuguese varieties to produce volatile phenols. In order to attain this purpose, the following objectives have to be achieved.

- Determining the ability of chosen single varietal wines to support the growth of a particular *Dekkera* strain, namely *Dekkera bruxellensis* PYCC 4801.
- Determining the ability of *Dekkera bruxellensis* PYCC 4801 to produce volatile phenols in this environment.
- Determining a potential correlation between grape variety and volatile phenol production.
- Determining the suitability of contamination with *D. bruxellensis* PYCC 4801 and subsequent determination of volatile phenols as method to obtain information on the wines potential to produce volatile phenols.

The theoretical part of this work gives an overview over phenolic compounds in wine, focusing on those acting as precursors, the synthesis of volatile phenols and their microbial origin. Furthermore, previous works evaluating volatile phenol occurrence in several varietal wines as well as some from different origins is discussed. In the experimental part different native varietal wines from Portugal were contaminated with *Dekkera bruxellensis* in order to determine and evaluate the possible effect of the grape variety on the production of volatile phenols. Eventually it is discussed if and to what extent the results indicate a correlation between variety and volatile phenol production.

This study is meant to give a first impression of this potential correlation and to indicate a direction for further research.

2 LITERATURE REVIEW

2.1 Phenolic compounds in wine

Grapes and wine contain various phenolic compounds that are related to both the colour and flavour attributes of wine and may also have antioxidant properties that may be beneficial to health (De Beer, et al., 2002; Ribéreau-Gayon, et al., 2006). These compounds are mainly extracted from grape skins, particularly the epidermal cells and the seeds during winemaking but smaller amounts also originate from wood, introduced during wine aging (Ribéreau-Gayon, et al., 2006). The phenolic composition of wines is greatly influenced by the grape variety, location of the vineyard, climate, soil type, system and practices of cultivation and harvest as well as the process of winemaking including all process steps such as pressing, maceration, aging, etc. (Rodríguez-Delgado, et al., 2002)

Based on their carbon skeleton, phenolic compounds can basically be divided into two groups: flavonoids and non-flavonoids. Flavonoids present in wine are anthocyanins, flavan-3-ol monomers, polymers and flavonols (Castillo-Sánchez, et al., 2008). Their structure is a composition of 15 carbon atoms, including two benzene rings that are connected by a linear three-carbon chain (Kheir, et al., 2013). Non-flavonoids including hydroxycinnamic acids and their derivatives, hydroxybenzoic acids and volatile phenols by contrast possess a simpler structure consisting of one primary aromatic ring bond to either one or three carbons (Castillo-Sánchez, et al., 2008; Kheir, et al., 2013; Basha, et al., 2004). Apart from these compounds, a third group of phenolic compounds in wine has recently been mentioned by Basha et al. (2004): phenolic-protein-polysaccharides.

2.2 Volatile phenols in wine

Certain non-flavonoids, particularly phenolic acids (hydroxycinnamic) in their free forms, act as precursors to the production of odorous volatile phenols. Volatile phenols associated with wine include ethylphenols (such as 4-ethylphenol and 4-ethylguaiacol) that are quantitatively more significant in red wines as well as vinylphenols (such as 4-vinylphenol and 4-vinylguaiacol) which are mostly found in white wines (Chatonnet, et al., 1997). The low concentrations of vinylphenols in red wine were thought to be due to the subsequent conversion of these intermediates to their corresponding ethyl-derivates by lactic acid bacteria during malolactic fermentation. It has been shown however that it is rather related to the inhibition of conversion enzymes by certain grape phenols (Ribéreau-Gayon, et al., 2006). Recently, 4-ethylcatechol has been found to contribute to phenolic off-flavour in wine as well. However, due to difficulties in its detection more attention has been paid to the four above mentioned volatile phenols (Hesford, et al., 2004).

Although commonly occurring in very low concentrations, volatile phenols have great influence on organoleptic characteristics, especially when sensory thresholds are exceeded (Kheir, et al., 2013; Rentzsch, et al., 2009). Excessive volatile phenol production is regarded as spoilage, imparting phenolic, animal or even stable odours that are also referred as to phenolic character (Chatonnet, et al., 1992). Specifically, 4-ethylphenol possesses woody, smoky, leather and animal flavours while 4-vinyl and 4-ethylguaiacol impart spicy clove-like notes (Edlin, et al., 1995).

Table 1 shows the most commonly occurring volatile phenols, summarizes their pathway of formation and sensory descriptors. These off-flavours can appear in red wines at various stages during fermentation and ageing (Chatonnet, et al., 1997). However, below certain threshold levels their presence may be considered as contributing to the complexity of wine aroma generating notes of leather, smoke or spices (Kheir, et al., 2013).

Table 1: Most common volatile phenols found in wine, their synthesis pathway and odour

Volatile phenol	Synthesis	Odour	Reference
4-ethylphenol	Conversion of 4-vinylphenol by vinylphenol reductase	stable, barnyard, phenol, woody	Chatonnet et al., 1992 Heresztyn et al., 1986 Edlin et al.,
4-ethylguaiacol	Conversion of 4-vinylguaiacol by vinylphenol reductase	spicy, clove-like, medicinal	Nelson, 2008 Edlin et al., 1995
4-ethylcatechol	Conversion of caffeic acid by hydrocinnamate decarboxylase	horse-like, animal, sweat, disinfectant	Hesford et al., 2004 Nelson, 2008
4-vinylphenol	Conversion of <i>p</i> -coumaric acid by hydrocinnamate decarboxylase	medicinal	Dias et al., 2003 Nelson, 2008 Oelofse et al., 2008
4-vinylguaiacol	Conversion of ferulic acid by hydrocinnamate decarboxylase	flowery, spicy, clove-like	Jackson, 2008 Edlin et al., 1995 Oelofse et al., 2008

2.2.1 Concentration ranges

Volatile phenol quantities found in wines strongly vary depending on variety, vintage and the wine's origin. Pollnitz et al. (2000) examined 61 bottled commercial red wines of various vintages using gaschromatography-mass spectrometry and found 4-ethylphenol concentrations ranging from 2 µg/l in Merlot up to 2660 µg/l in Shiraz with a mean value of 795 µg/l. 4-ethylguaiacol was detected in a range from 1 µg/l in Pinot Noir to 437 µg/l in Merlot with a mean concentration of 99 µg/l. Oelofse et al. (2008) as well summarized quantity rates commonly found in red wines showing ranges similar to those Pollnitz et al. (2000) discovered. Additionally, Table 2 shows concentration ranges for ethyl and vinylphenols occurring in red wines.

Table 2: Concentration ranges of volatile phenols in red wine (reproduced from Oelofse et al, 2008)

Volatile phenol	Concentration range (µg/l)
4-ethylphenol	118-3696
4-ethylguaiacol	1-432
4-ethylcatechol	27-427
4-vinylphenol	4.3-8.8
4-vinylguaiacol	0.2-15

2.2.2 Perception thresholds

Chatonnet et al. (1992) established perception thresholds for ethyl-derivatives in water as well as in red wine. Due to the aromatic complexity of wine, threshold values in water are much lower. It was shown that 4-ethylguaiacol was more easily perceived compared to 4-ethylphenol with threshold values of 140 µg/l and 620 µg/l in wine, respectively. The rough proportion in which both these compounds are normally present in red wine is a 10:1 average ratio for the mixture 4-ethylphenol : 4-ethylguaiacol. If the combined threshold value of 426 µg/l is exceeded, the phenolic character may occur (Chatonnet, et al., 1992). However, these threshold values may not be equally applicable in all wines. It has been shown that certain compounds possess a masking effect on the detection of ethyl-derivatives and therefore sensory descriptors do not always correlate with the actual ethylphenol contents. In fact, it was proven that the presence of isobutyric and isovaleric acids, synthesized during volatile phenol production by *Brettanomyces bruxellensis*, raise the detection threshold of ethylphenols about up to three times compared to the aforementioned value (Romano, et al., 2008).

2.2.3 Precursors

Both grape juice and wine contain phenolic acids including hydroxycinnamic and hydroxybenzoic acids. They can be present in *cis* and *trans* configurations while the latter are more stable and hence more prevalent (Rentzsch, et al., 2009). According to Ribéreau-Gayon et al. (2006), the combined concentration of benzoic and cinnamic acids in red wine ranges from 100 to 200 mg/l and from 10 to 20 mg/l in white wines. They may vary among different grape varieties, ripening conditions and be dependent on vinification practices (Ribéreau-Gayon, et al., 2006). For example, lower occurrence of precursors in white wine may be explained by the fact that they are extracted from the skins and seeds during maceration which is usually avoided in the production of white wine (Kheir, et al., 2013).

Figure 1 shows the chemical structures of benzoic and cinnamic acids present in grapes and wine. In grapes, benzoic acids are mainly bound to glycosides or esters and released during winemaking by the hydrolysis of these combinations or other breakdown reactions of

anthocyanins (Ribéreau-Gayon, et al., 2006). Therefore in wine, benzoic acids are more prevalent in their free forms while cinnamic acids are found rather in combinations such as esterified with tartaric acid. (Kheir, et al., 2013; Rentzsch, et al., 2009; Ribéreau-Gayon, et al., 2006). In wine, fertaric acid and *p*-coutaric acid, tartaric esters of ferulic and *p*-coumaric acids, respectively, which are the most prevalent precursors in volatile phenol production are present at values around 16.0 mg/l and 55.0 mg/l, respectively (Rentzsch, et al., 2009).




Figure 1: Phenolic acids found in grapes and wine (reproduced from Ribéreau-Gayon, et al., 2006)

Research conducted on both red and white wines from Portugal found total phenolic acid contents close to the range mentioned by Ribéreau-Gayon et al. (2006), 98.7 mg/l were present in red and 54.4 mg/l in white wine (Ribeiro de Lima, et al., 1998). Moreover, average concentrations of some specific phenolic acids were determined using HPLC with diode array detector procedures. The results, presented in Figure 2, show average contents of phenolic acids obtained from examining a total of 72 wines. Furthermore, they confirm that apart from gentistic acid, white wines show lower values for phenolic acids (Ribeiro de Lima, et al., 1998).




Figure 2: Average contents of phenolic acids in wines from continental Portugal and Azores (Data compiled from Ribeiro de Lima et al., 1998)

2.2.4 Synthesis

The production of volatile phenols is related to the enzymatic conversion of free hydroxycinnamic acid precursors, particularly *p*-coumaric acid, ferulic acid and caffeic acid (Kheir, et al., 2013; Ribéreau-Gayon, et al., 2006). Two sequential conversion steps facilitated by cinnamate decarboxylase and vinylphenol reductase are necessary in order to form volatile phenols (Chatonnet, et al., 1997). A cinnamate carboxylase enzyme or phenolic acid decarboxylase (PAD) decarboxylates hydroxycinnamic acids to intermediate hydroxystyrenes, namely 4-vinylphenol, 4-vinylguaiacol and 4-vinylcatechol by cleaving off the C3 carbon from the side chain and releasing CO₂ (Kheir, et al., 2013; Chatonnet, et al., 1997). Research conducted by Barthelmebs et al. (2000) suggests that in some bacteria the synthesis of PAD enzymes, specifically *p*-coumaric acid decarboxylase may be a stress response to high concentrations of phenolic acids inducing the conversion of *p*-coumaric acid into a less toxic derivate. The second step of volatile phenol formation involves the reduction of vinyl-derivativee double bond by vinylphenol reductase (VPR) in order to form respective ethyl-derivatives (4-ethylphenol, 4-ethylguaiacol and 4-ethylcatechol) (Kheir, et al., 2013). The conversion pathway is shown in Figure 3.




Figure 3: Formation pathway of volatile phenols via the decarboxylation of hydroxycinnamic acids (reproduced from Oelofse et al., 2008)

An alternative pathway was proposed in some lactic acid bacteria by Whiting and Carr (1959) where the phenolic acids are first reduced to their substituted phenyl propionoc acids by *p*-coumaric and ferulic acid reductase activities (PAR) and subsequently decarboxylated to generate their 4-ethyl-derivates (Barthelmebs, et al., 2000; Ota, et al., 2001; Whiting, et al., 1959). This pathway was found in *Lactobacillus plantarum* and *Lactobacillus paracollinoides* (Barthelmebs, et al., 2000).

2.3 Microbial origin

The microbial origin of volatile phenols in wine has been widely discussed. Initially, their occurrence was associated with lactic acid bacteria (LAB) and their action during malolactic fermentation. Growing cultures of *Lactobacillus brevis*, *L. plantarum* and *Pediococcus* have been identified to be capable of decarboxylating phenolic acids, especially ferulic and *p*-coumaric acid (Couto, et al., 2006; Chatonnet, et al., 1995; Cavin, et al., 1993). Couto et al. (2006) proposed that LAB's capacity to produce volatile phenols is higher for *p*-coumaric than for ferulic acid. Although some LAB (*Lactobacillus brevis*, *Ped. pentosaceus*, *Leuconostoc oenos*) were able to produce vinylphenols in ranges from a few hundred up to thousands of µg/l in growth media, only insignificant amounts below 10.0 µg/l of ethylphenols were found to be released in wines (Chatonnet, et al., 1995; Cavin, et al., 1993). Chatonnet et al. (1995) suggested *L. plantarum* to be the only LAB capable of reducing vinyl-derivates into significant amounts of ethyl-derivates, 4-ethylphenol being the most predominant. These findings were confirmed by Couto et al. (2006) who found that 37% of 35 strains (20 species) exhibited cinnamate decarboxylase activity but only 9% were capable of producing ethylphenol. These

results were confirmed by a later study which identified, using a molecular screening approach, the presence of the *pdh* gene (which codifies for PAD) in certain LAB species (De las Rivas, et al., 2009). However, in previous studies conducted with wines, the released concentrations of volatile phenols by LAB did not exceed the perception threshold and therefore had no detrimental effect on wine aroma (Chatonnet, et al., 1995). Therefore, excessive volatile phenol production could not be linked to LAB or malolactic fermentation despite the fact that some LAB do possess enzymatic activity for transforming phenolic precursors into their corresponding volatile phenols.

Other microorganisms considered to be more relevant in regard to volatile phenol production are certain yeasts. *Saccharomyces cerevisiae*, the main yeast responsible for alcoholic fermentation in wines exhibits a cinnamate decarboxylase activity and is therefore capable of forming vinylguaiacol from hydroxycinnamic precursors, however does not reduce them to their corresponding ethyl-derivatives (Chatonnet, et al., 1993; Heresztyn, 1986). Dias et al. (2003) have shown that several yeast species associated with wine production or wine-related environment including *Dekkera bruxellensis*, *D. anomala*, *Candida cantarelli*, *Candida wickerhamii*, *Debaryomyces hansenii*, *Kluyveromyces lactis* and *Pichia guilliermondii* additionally exhibit the capability of both decarboxylating *p*-coumaric and reducing the arising 4-vinylphenol into its respective ethylphenol. The highest molar conversion rates were detected in *D. bruxellensis* and *D. anomala* with an efficiency rate of about 90%. Similar results were obtained for efficiency of *P. guilliermondii* although conversion rates were variable within strains. The remaining showed themselves to be rather weak ethylphenol producers (Dias, et al., 2003b). Chatonnet et al. (1995) suggested that *D. intermedia* produces 4-ethylphenol in amounts up to 16 times the perception threshold value while LAB barely exceeded half the threshold value. Other yeast species and strains, not related to wine but other fermented foods were found by Chatonnet et al. (1992) and confirmed to form ethylphenols by Dias et al. (2003). These include *C. halophila*, *C. mannifaciens* and *C. versailis* that are associated with soy sauce production. It can be concluded that the fermenting yeast *S. cerevisiae* or species other than *Brettanomyces/Dekkera bruxellensis*, *intermedia* and *anomala* or LAB are considered rather weak ethylphenol producers under wine conditions that may not influence wine aroma. Furthermore, *P. guilliermondii* which produces significant amounts of ethylphenol was indeed recovered from grapes and grape juice but not from wine (Dias, et al., 2003b). Therefore, certain *Dekkera/Brettanomyces* strains remain the most significant agents in creating phenolic off-flavours in wine. The species most prevalent in wine is *B. bruxellensis* (Chatonnet, et al., 1992).

2.3.1 *Dekkera/Brettanomyces*

The name *Brettanomyces* is most probably derived from "British brewing fungus" since it was isolated from old English stock beer and imparts typical flavours. These flavours were suggested to be linked to a secondary fermentation by the non-spore forming non-*Saccharomyces* yeast *Brettanomyces* that was first described in 1904 by Claussen (Oelofse, et al., 2008; Van der Walt, et al., 1958). Later on, *Brettanomyces* strains were found to play a role in various fermented beverages such as wine, cider, kombucha as well as other types of beer but also in fermented food including olives or cheeses (Nelson, 2008; Suárez, et al., 2007). The first association with wine was not made until 1940 when a research by M.T.J. Custers showed that of 17 strains isolated from beverages, one originated from a French wine whereas most of them were found in beer (Oelofse, et al., 2008).

According to Van der Walt et al. (1958) *Brettanomyces* species are characterized by their slow growth, the production of pseudomycelium, the formation of 'ogive' cells, the presence of fermentative metabolism, the production of acetic acid under anaerobic conditions and the absence of spore formation. Currently, of the genera *Brettanomyces* five species are recognized: *B. bruxellensis*, *B. anomalus*, *B. naardenensis*, *B. nanus* and *B. custersianus* (Oelofse, et al., 2008; Arvik, et al., 2002). A few years after the first characterization of *Brettanomyces* however, spore forming sexual strains were discovered and therefore the genus revised. Two species were assigned to the new genus *Dekkera*: *D. bruxellensis* and *D. intermedia*, of which the former represents the type strain (Van der Walt, 1964).

Brettanomyces/Dekkera yeasts are ubiquitous in regions associated with production of wine. *Brettanomyces* spp. has been isolated from wines and winery environment in Germany, France, Portugal, South Africa, Italy, New Zealand, Spain, Great Britain and the United States (Nelson, 2008). Moreover, in 31 winemaking regions in Australia, 8 different genotypes of *D. bruxellensis* were found in 244 *D. bruxellensis* isolates obtained from wineries and wine (Curtin, et al., 2007).

The yeast species *Brettanomyces/Dekkera bruxellensis* is considered responsible for spoilage of wine and other fermented beverages. Other names such as *D. intermedius*, *B. intermedius*, *B. lambicus* or *B. custersii* which enologists have referred to as the spoilage organism in the past, are nowadays considered synonyms for *Brettanomyces/Dekkera bruxellensis* (Loureiro, et al., 2003).

Regarding the growth phase in which the major production of volatile phenols by *Brettanomyces/Dekkera* occurs, Dias et al. (2003) observed the onset of 4-ethylphenol

production at about the mid-exponential phase lasting until the beginning of the stationary phase. In contrast Chatonnet et al. (1992) described an earlier onset of 4-ethylphenol production starting already from the latency phase and keeping up until the end of exponential phase. 4-vinylphenol decreased in both studies with the increase of 4-ethylphenol production. It was suggested this may be due to subsequent conversion to 4-ethylphenol (Chatonnet, et al., 1992; Dias, et al., 2003b)

Apart from volatile phenols, *Dekkera/Brettanomyces* exhibits the ability to produce other volatile compounds inducing off-flavours. These include tetrahydropyridines, responsible for mousy off-flavour as well as 3-methylbutyric acid (isovaleric acid) imparting sweaty and leathery notes (Nikfardjam, et al., 2009). Other effects of *B. bruxellensis* contamination are haziness or turbidity (Van der Walt, et al., 1958) and the formation of high levels of acetic acid (Nelson, 2008).

2.3.2 Sources of contamination

The primary source of contamination has been contentiously discussed and still remains uncertain. In 1958, Van der Walt et al. suggested a source to be winery environment rather than the yeast flora related to grapes and must. While *Brettanomyces* could be recovered from cellars and cellar equipment, they were not present on husks, pomaces or grapes. Therefore it was concluded that the contamination may arise only from within the winery. These findings were confirmed a couple of years later by the same authors with a method more appropriate for the recovery of *Brettanomyces* (Van der Walt, et al., 1960). More recent research using precise PCR techniques however, has shown the presence of *Brettanomyces/Dekkera* yeasts on grapes and in grape processing lines as well (Loureiro, et al., 2003). Generally, the rare detection on grapes in the past has been attributed to their low cell numbers in the diverse microbial environment of the grape skin in which other microorganisms or species may dominate. With the development of more specific media, *B. bruxellensis* could be detected on grapes at different stages of berry development (Oelofse, et al., 2008).

The occurrence of *Brettanomyces/Dekkera* has also been linked to grapes damaged by sour rot (Oelofse, et al., 2008). Furthermore, *Brettanomyces* was detected in air samples from crush, tank, barrel and bottling rooms of a winery. Thus, it could be shown that these spoilage yeasts can be present in the air (Connell, et al., 2002). However, the most critical stage in wine production for *Brettanomyces/Dekkera* spoilage to occur, is recognized to be the ageing in barrels after alcoholic fermentation has been completed, when the numbers of competitive microorganisms have decreased and the presence of residual sugars allows

proliferation (Chatonnet, et al., 1992). Barrels with their porous wood structure present an ecological niche for *Brettanomyces/Dekkera* which allows a little influx of oxygen and therefore promotes growth. Moreover, improper barrel sanitation and sulphite utilization might result in an incomplete elimination of these yeasts and thus favor the contamination of wines transferred to these barrels (Oelofse, et al., 2008; Suárez, et al., 2007; Loureiro, et al., 2003).

2.4 Grape variety and volatile phenol production by *Brettanomyces bruxellensis*

There are several factors affecting the formation of volatile phenols by *Brettanomyces/Dekkera*. Firstly, the availability of phenolic precursors greatly influences the formation of volatile products (Kheir, et al., 2013; Suárez, et al., 2007). This aspect has been widely related to both geographical origin and variety. In several studies from various countries and wine producing regions, it was suggested that phenolic compound concentrations vary between different grape varieties. For distinct differentiation of wines regarding grape variety however, the content of polyphenols has proven to be more useful than data on volatile compounds or phenolic acids. While some colored phenols such as the flavonoid quercetin show large variations between varieties, hydroxycinnamic acids have not been proven to differ significantly, although trends are recognizable (Villiers, et al., 2005). Gambelli et al. (2004) compared the phenolic compound contents of Italian red wines from different geographical origins. The findings indicate that concentrations of phenolic acids including *p*-coumaric and ferulic acid are more dependent on cultivars than on geographical origin (Gambelli, et al., 2004). Similar results were obtained by Pena-Neira et al. (1999) who observed quantitative differences in phenolic compounds in Spanish red wines. It was suggested they may be related to grape variety but are also influenced by vinification method and employed process of wine aging (Pena-Neira, et al., 2000). White wines generally show lower amounts of phenolic compounds due to shorter extraction times from skins and seeds. In some samples however, ferulic acid contents were similar to those found in red wine (Villiers, et al., 2005).

Pollnitz et al. (2000) describes research conducted by Goldberg et al. (1998) who analyzed *p*-coumaric acid concentrations of 547 commercial red wines from different wine producing regions and countries using high performance liquid chromatography. Although values seemed to vary slightly within varieties, the lowest values were observed in Pinot Noir compared to Shiraz and Cabernet Sauvignon. These results from Australian wines were confirmed by the analysis of Californian and South African wines which showed the lowest *p*-coumaric acid concentrations in Pinot Noir as well (Villiers, et al., 2005; Goldberg, et al., 1998). The highest concentrations were detected in Cabernet Sauvignon and Merlot wines

from France, Australia, California, South Africa and Canada (Goldberg, et al., 1998; Soleas, et al., 1997).

Pollnitz et al. (2000) analyzed Australian red wines regarding varietal influences on the presence of volatile phenols (4-ethylphenol and 4-ethylguaiacol). Again, within varieties wide ranges of concentrations could be observed. The general trend however showed that the mean concentration of 4-ethylphenol was lowest in Pinot (with an average of 338.0 µg/l), with the authors suggesting that a relation might exist to the low precursor occurrence in this grape variety detected in previous research (Pollnitz, et al., 2000). Pinot Noir was followed by Shiraz with an average concentration of 605.0 µg/l and Cabernet Sauvignon with the highest average 4-ethylphenol concentration of 1250.0 µg/l. Varietal differences in 4-ethylguaiacol were found to be insignificant (Pollnitz, et al., 2000). Similar results were obtained by a study conducted on German wines from the Württemberg region. Varieties that are known to be rich in hydroxycinnamic precursors such as Acolon, Cabernet Cubin, Dornfelder or Portugieser, have shown to possess high ethylphenol contents that partly exceed the sensory threshold value established by Chatonnet et al. (1992) (Nikfardjam, et al., 2009). These results suggest that depending on their contents of phenolic precursors, varieties may be more or less prone to formation of volatile phenols.

There are however certain factors that may affect the precursor concentration so that even within varieties, values may change. Grapes grown in hot climates for example are richer in hydroxycinnamic acids. Moreover, the maturity of grapes as well as technological aspects during vinification play important roles (Ribéreau-Gayon, et al., 2006). For instance, the extraction process including temperature profiles during maceration impact precursor availability. High temperatures, especially at the end of maceration have been shown to promote extraction processes by affecting the permeability of cells and membranes of grape skins and thus increasing formation of volatile phenols (Gerbeaux, et al., 2002).

2.4.1 Portuguese wines

Only a few studies have been conducted on Portuguese varieties and none of them has tried to find a relation between precursor availability and the potential to form volatile phenols but rather evaluated the possibility to differentiate varieties on the basis of their phenolic profiles. Ribeiro de Lima et al. (1998) have proven varietal differences in *p*-coumaric acid content, with Touriga Francesa (also referred as to Touriga Franca) (38.3 mg/l) exhibiting highest values in red and Verdelho (5.0 mg/l) in white wines. Other authors that evaluated anthocyanin profiles rather than non-coloured phenolics have shown that anthocyanins may be a good marker to differentiate varieties (Dopico-García, et al., 2008; Mateus, et al., 2002).

While Dopico-García et al. (2008) examined 10 different varieties of red Vinho Verde grapes and could observe clear differences in their specific anthocyanin profiles, Mateus et al. (2002) only evaluated two varieties, namely Touriga Nacional and Touriga Franca but obtained similar results. Observations regarding the influence of grape origin and harvest conditions on non-coloured phenolic composition and anthocyanin profile have shown that the impact of these factors is higher in non-coloured phenolics rather than anthocyanins (Dopico-García, et al., 2008). Although no correlation to volatile phenols was specifically examined, the results suggest that the grape variety highly impacts their phenolic profile and therewith the potential precursor composition of grapes.

2.5 Other factors influencing volatile phenol production

Besides factors affecting the phenolic composition, microbiological factors may also influence volatile phenol production. For instance, the capacity to form those compounds from their precursors varies depending on the strain responsible for spoilage (Suárez, et al., 2007; Valentao, et al., 2007; Shinohara, et al., 2000). This may be explained by differences in enzymatic specificity, activity and the metabolism of phenolic acids between species (Edlin, et al., 1995). Furthermore, a high correlation between *Brettanomyces/Dekkera* population size and conversion time of *p*-coumaric acid into volatile phenols was found by Benito et al. (2009) suggesting that the amount of odorous product formed is greatly influenced by the extent of contamination by the spoilage organism (Benito, et al., 2008). A decrease of volatile phenols formed may also be due to a decrease of phenolic precursors caused by their adsorption onto yeast cell walls (Salameh, et al., 2008). Both Salameh et al. (2008) and Cabrita et al. (2012) have found the initial concentration of *p*-coumaric as well as ferulic and caffeic acid to be decreased immediately after inoculation with *Brettanomyces/Dekkera* yeast in both wine and synthetic medium and attributed the major part of this loss to cell adsorption. Besides adsorption to cell walls smaller amounts of acids may be lost due to esterification with ethanol or instability at high temperatures (Cabrita, et al., 2012; Salameh, et al., 2008). Although this and other chemical reactions can decrease the quantity of *p*-coumaric acid available for bioconversion, other reactions can cause its increase. This effect was described by Dugelay et al. (1993) who observed the hydrolysis of *p*-coumaroyltartaric acid facilitated by hydrolytic enzymes including pectinase, (hemi-)cellulase and cinnamate esterase releasing *p*-coumaric acid in must. Such enzyme preparations may be used for clarification improvement purposes, colour extraction or flavour enhancement (Dugelay, et al., 1993).

Dias et al. (2003) examined the effects of temperature, ethanol and different carbon sources on ethylphenol production in synthetic media and grape juice. It was reported that an

increase in temperature (30°C compared to 16°C) raised the production rate of ethylphenol although it did not impact the eventual yield. Moreover, it was shown that glucose or ethanol are necessary carbon sources in order to obtain conversion rates up to 90%. It should be noted however, that ethanol concentrations above 13 %(v/v) have been restricting both the growth of *Brettanomyces/Dekkera bruxellensis* and the formation of ethylphenols.

In conclusion, several authors have shown that the grape variety indeed impacts the phenolic composition of grapes and wine and thus the availability of precursors. Due to various other factors related to the vinification process and contaminating yeasts however a concrete correlation between variety and formation of volatile phenols has not been found yet.

3 MATERIALS AND METHODS

The experimental work described in this dissertation involved analyzing Portuguese wines for the occurrence of volatile phenols. Single varietal wines were selected in order to identify the influence of the grape variety on volatile phenol production in purposely contaminated wines. A total of 11 samples were collected from different wineries. Subsequently, the wines were modified regarding pH- and alcohol content in order to serve as base media for yeast growth and/or volatile phenol metabolism. Freshly prepared cultures of a *Dekkera bruxellensis* strain were used to inoculate the wines and samples were incubated for 10-14 days at 30°C. Yeast growth was monitored by viable yeast counts three times a week. Once stationary phase was reached volatile compounds were extracted (using a liquid-liquid extraction protocol) in order to prepare samples for subsequent chromatographic screening. The samples were analysed for the presence of three particular volatile phenols: 4-Ethylphenol (4EP), 4-Ethylguacaol (4EG) and 4-Vinylphenol (4VP).

3.1 Wine samples

A total of 11 single variety wines were provided by different wine producers from two major wine regions in Portugal; the Douro Valley and Alentejo region. Table 3 summarizes the preliminary information available for some chemical parameters of all wines used in this work.

In order to eliminate wine organisms that might interfere with the experiments, all samples were heat treated prior to actual sample preparation. Pasteurization was performed at 50°C for 3 min. using a heat plate with incorporated magnetic stirrer to facilitate even heat distribution throughout the sample. If not immediately used, pasteurized samples were stored at 7°C.

Table 3: Variety, origin and analytical parameters of the wine samples

Sample code	Variety	Vintage	Region	Vol. %	pH
TF	Touriga Franca	2011	Douro	14.7	3.74
SY	Syrah	2012	Alentejo	15.3	3.48
TR	Tinta Roriz	2012	Douro	15.6	3.86
TN1	Touriga Nacional	2012	Douro	13.6	3.77
TN2	Touriga Nacional	2012	Douro	14,0	3,63
TN3	Touriga Nacional	n.a.*	Douro	14.8	3.81
TN4	Touriga Nacional	2012	Douro	13.7	3.89
S1	Sousão	2012	Douro	12.0	3.48
S2	Sousão	2012	Douro	14,2	3,33
S3	Sousão	2013	Douro	12,9	3,45
S4	Sousão	2013	Douro	12,4	3.36

*n.a.: not available

3.2 Yeast strain

For the experimental work the reference strain *Dekkera bruxellensis* PYCC 4801, from the Portuguese Yeast Culture Collection (Caparica, Portugal) was chosen due to its relevance as a volatile phenol producing spoilage organism in the wine industry. The stock culture was prepared in liquid YM broth (Difco™, Becton, Dickinson and Company, France) and grown to late exponential phase for about 3-5 days at 30°C. In order to preserve the stock culture it was subsequently inoculated in a YM agar (2.0% w/w) medium slant, grown at 30°C and eventually stored at 7°C for later use. The fresh culture used for inoculating the samples was prepared from the slant using sterile YM broth and in the same manner as the stock. Prior to inoculation, the fresh culture was centrifuged (3000 g, 5 min), washed and resuspended in sterile deionized water in order to avoid growth medium being transferred into the sample.

3.3 Preparation of phenolic compound solutions

Individual concentrated solutions (10 g/l) of volatile phenol precursors were prepared by weighing in about 0.1 g of both ferulic acid and *p*-coumaric acid and dissolving each in 10.0 ml of ethanol (99.5%, Aga, Portugal). The solutions were properly mixed using a vortex and stored in the freezer at -18°C. All commercial phenolic compounds had a purity of at least 99% and were obtained from Sigma-Aldrich (Steinheim, Germany).

3.4 Sample preparation

Considering the high alcohol content and low pH of the samples, dilutions were prepared of each wine in order to avoid conditions that may cause stress to the yeasts. In sterile Schott glasses, the wines were mixed with either sterile YM broth or sterile deionized water, in a 50:50 (v/v) ratio constituting a total of 80.0 ml per sample. In order to determine the influence of certain precursors in volatile phenol production, 400 µl of each ferulic and *p*-coumaric acid solutions were added to selected samples up to a final concentration of 5.0 ppm or 50.0 ppm. The exact composition of the samples is shown in Table 4. To guarantee identical conditions for the yeasts, additionally, pH and ethanol content were adjusted to 3.8 and 7.5% vol., respectively, in all wines. Each sample was prepared as duplicate.

Table 4: Sample numbers and their composition

Sample	Wine	YM broth	Water	<i>p</i> -coumaric acid solution	ferulic acid solution
W+YM	40 ml	40 ml	-	-	-
W+YM/A	40 ml	40 ml	-	0.400 ml	0.400 ml
W+W	40 ml	-	40 ml	-	-
W+W/A	40 ml	-	40 ml	0.400 ml	0.400 ml

Subsequently, each sample was inoculated with 800 µl of freshly prepared *Dekkera bruxellensis* PYCC 4801 culture to give an approximate initial cell density of 10⁶ cells/ml.

3.5 Determination of yeast growth

Yeast growth in samples was determined using the viable plate count technique as described by Miles and Misra (1938). Prior to plating, decimal dilutions were made in 9.0 ml of sterile Ringer solution (Biokar Diagnostics, France) until a 10^{-5} dilution was obtained. After thoroughly mixing using a rotative vortex, 20 μ l drops of each dilution were plated in YM agar (2.0% w/w) plates equally divided into sectors and labeled with their respective dilution. Each sample was plated in duplicates in order to obtain average values. Following incubation at 30°C for 3-4 days, colonies were counted and the number of colony forming units (CFU) per ml in the original sample was calculated.

3.6 Extraction of volatile phenols

50.00 ml of each sample was transferred into volumetric flasks. In order to allow for the appropriate comparison of results, 50.0 μ l of 3-octanol were added as internal standard. Afterwards, 4.0 ml of a 50:50 (v/v) solution of n-hexan (Merck, Germany) and diethyl-ether (Panreac, Spain) were added to each volumetric flask and subsequently stirred for 5 minutes. Next, the mixtures were transferred to separation funnels where aqueous and organic phases were separated. While the latter were collected in new vials, the aqueous phases were poured back into respective volumetric flasks for further extraction. The extraction procedure was repeated two more times with 2.0 ml of solvent in order to achieve maximum extraction yields. Eventually, organic phases were collected in vials and reduced under nitrogen flow to about one third of their original volume.

3.7 Gas chromatography

Subsequent to extraction, the samples were analysed using a gas chromatograph from Hewlett Packard (5890A) equipped with a flame ionisation detector (GC-FID). The column employed was a FFAP type (BP1) with dimensions of 50 m x 0.22 mm x 0.25 μ m (SGE, Austin, Texas). Hydrogen was used as carrier gas and the flow adjusted to 1.0 ml/min. Once the split flow was adjusted to around the optimum of 30.0 ml/min and the injector had heated up to about 250°C, a sample volume of 2.0 μ l was injected. The oven temperature program applied was as following: starting off with 40°C held for 1 min and followed by an increase at a rate of 2°C per minute reaching up to 220°C where it was held for 20 minutes.

3.8 Calibration curves

In order to identify volatile phenols in the samples a calibration curve was established by preparing standard solutions containing 4-ethylphenol (99%, Sigma-Aldrich, Germany), 4-ethylguaiacol (98%, Sigma-Aldrich, Germany) and 4-vinylphenol (10% solution in propylene glycol, SAFC, United Kingdom). The range of consequent retention times for given GC parameters are shown in Table 5.

Table 5: Range of retention times for calibrated volatile phenols and internal standard

Compounds	Average retention time in min
3-octanol	29.56
4-ethylguaiacol	67.42
4-ethylphenol	74.69
4-vinylphenol	85.01

Moreover, all peaks were well separated and as can be seen in Figure 4 R^2 values show that there is a good correlation between the different data points of each compound.




Figure 4: Calibration curves for 4-ethylguaiacol, 4-ethylphenol and 4-vinylphenol including trendline equations

4 RESULTS

4.1 Inoculum size

In all experiments, the initial population of yeasts was inoculated at a minimum of 10^6 cells/ml. As can be seen in Table 6, all values ranged between 1.5 and 85×10^6 .

Table 6: Cell concentration of inocula

Sample code	Inoculum (cells/ml)
TF	85×10^6
TN3	45×10^6
TN4, TR	1.5×10^6
TN1, S1, SY	70×10^6
TN2, S2, S3, S4	45×10^6

(TF: Touriga Franca; TR: Tinta Roriz; TN1, TN3, TN4: Touriga Nacional; S1, S2, S3: Sousão)

4.2 Yeast growth

In test trials, pasteurized non-diluted wine was inoculated with *D. bruxellensis* in order to follow yeast growth and the production of volatile phenols. However, no yeast growth, but rather a decline in cell numbers could be observed (results not shown). Therefore, yeast growth was documented in wines diluted with either YM broth or water in a 50:50 ratio for a period of at least 10 days. The resulting growth curves of *D. bruxellensis* are presented from Figure 5 to Figure 15.

It can be seen that the initial cell concentration in all samples ranges from approximately 5.0 log CFU/ml to 6.0 log CFU/ml, except for the samples TN4 (Figure 11) and TR (Figure 6) which, in accordance with the inoculum size, contained lower counts at the day of inoculation. No significant differences in growth could be observed comparing samples diluted with water or YM although a slight trend towards higher final counts in YM diluted is visible in most of the samples. However, the opposite trend can be seen in Touriga Franca (TF) and one Touriga Nacional (TN3), presented in Figure 5 and Figure 10, respectively. Moreover, the supplementation with 5mg/l of *p*-coumaric and ferulic acids such as in TF (Figure 5) and TN3 (Figure 10), did not affect the yeast growth, while with addition of concentrations as high as 50 ppm (remaining figures) one can observe slightly lower cell numbers in supplemented samples. Therefore, it can be suggested that the addition of phenolic precursors in high quantities (such as 50 ppm) might be inhibitory to yeast growth.

Apart from three wines made from the Sousão variety (namely, S2, S3 and S4), the growth curves in different media do not show any significant differences in their behavior regarding growth phases within one sample. The end of exponential phase is attained after a minimum of 3 days, as to be seen for SY (Figure 7), TN1 (Figure 8) and TN2 (Figure 9). The latest onset of stationary phase occurred at around day 7 and 8 in TF (Figure 5), TN3 (Figure 10), S2 (Figure 13) and S4 (Figure 15).




Figure 5: Growth of *D. bruxellensis* PYCC 4801 in diluted Touriga Franca (TF) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 6: Growth of *D. bruxellensis* PYCC 4801 in diluted Tinta Roriz (TR) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 7: Growth of *D. bruxellensis* PYCC 4801 in diluted Syrah (SY) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 8: Growth of *D. bruxellensis* PYCC 4801 in diluted Touriga Nacional (TN1) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 9: Growth of *D. bruxellensis* PYCC 4801 in diluted Touriga Nacional (TN2) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 10: Growth of *D. bruxellensis* PYCC 4801 in diluted Touriga Nacional (TN3) at 30°C; values obtained from two determinations (W: wine; W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 11: Growth of *D. bruxellensis* PYCC 4801 in diluted Touriga Nacional (TN4) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 12: Growth of *D. bruxellensis* PYCC 4801 in diluted Sousão (S1) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)

As mentioned previously and as can be seen in Figure 13, Figure 14 and Figure 15, the Sousão samples do not show the same patterns when comparing the curves shape both within one sample as well as between the samples. The growth phases are not clearly visible. Furthermore, the maximum of cell concentrations obtained were (especially in wine diluted with water) much lower when compared to other samples, meaning that there was less cell growth. In some cases (Figure 14) numbers declined even below the original inoculation amount of cells. However, that they were detectable with viable plate count technique at any stage shows that they were viable and thus able to produce volatile phenols at any time.




Figure 13: Growth of *D. bruxellensis* PYCC 4801 in diluted Sousão (S2) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 14: Growth of *D. bruxellensis* PYCC 4801 in diluted Sousão (S3) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)




Figure 15: Growth of *D. bruxellensis* PYCC 4801 in diluted Sousão (S4) at 30°C; values obtained from two determinations (W+YM: wine diluted with YM medium; W+W: wine diluted with water; /A: with added phenolic acids)

4.3 Volatile phenols in varietal wines

As indicated by Figure 16, some of the original wines showed presence of 4-ethylphenol and 4-ethylguaiaicol, while 4-vinylphenol could not be detected.

It can also be seen that in the non-contaminated wines 4-ethylguaiaicol occurred quite often compared to 4-ethylphenol. While 4-ethylguaiaicol was found in 7 wines, 4-ethylphenol was detected in only 4 wines at considerably low ranges from 0.02 ppm up to 0.09 ppm and from 0.03 ppm to 0.07 ppm, respectively. The values however are extremely low and do not exceed quantities that are actually considered as detectable by GC. Hence, all values found are far below the perception threshold level established by Chatonnet et al. (1992) and therefore not considered detrimental to wine aroma. Three samples, the Syrah (SY) and each one of the Touriga Nacional and Sousão wines, namely, TN2 and S4 did not show any volatile phenols.




Figure 16: Volatile phenol concentration in examined red wines (TF: Touriga Franca; TR: Tinta Roriz; TN1, TN3, TN4: Touriga Nacional; S1, S2, S3: Sousão)

4.4 Volatile Phenols production by *D. bruxellensis* PYCC 4801 in diluted wines

The amount of volatile phenols in the diluted wines was measured using gas chromatography after contamination with *D. bruxellensis* and subsequent incubation for at least 10 days at 30°C. All the artificially contaminated samples showed presence of 4-ethylphenol and 4-ethylguaiacol but not of 4-vinylphenol.

When considering the concentration of volatile phenols observed in the diluted wines, it must be considered that potentially, the values could be twice as high as in the original wine since the samples were diluted into a 50:50 ratio and therefore in theory only contained half the amount of volatile phenol precursors compared to non-diluted wine. However, if non-diluted wine was directly contaminated, some factors such as inhibition of yeast growth by higher alcohol levels could have possibly impacted product outcome and therefore it cannot be presumed that levels were actually precisely twice as high. Hence, in all following results the volatile phenol levels presented, are the ones determined in diluted wines.

Figure 17 to Figure 20 present 4-ethylphenol and 4-ethylguaiacol concentrations detected in the original wines and in the same wines diluted with water or YM medium and artificially contaminated with *D. bruxellensis* PYCC 4801. As appears in the figures, with one exception (TN2), all samples show a trend to slightly higher volatile phenol production in wines diluted with YM compared to water. This may be attributed to an enhanced yeast growth in the presence of the growth medium which could be related to increased nutrient availability.

As indicated by Figure 17 *D. bruxellensis* PYCC 4801 has formed considerable amounts of volatile phenols in Touriga Franca and Tinta Roriz wines. The released quantities of 4-ethylphenol and 4-ethylguaiacol in both diluted wines exceed the perception thresholds of 0.64 ppm and 0.14 ppm (Chatonnet, et al., 1992), respectively and thus it can be assumed the wines should exhibit a phenolic off-flavour. Moreover, the highest 4-ethylphenol quantity in this entire experiment was detected in Tinta Roriz (diluted with YM) with 1.59 ppm. Touriga Franca on the other hand showed the highest value for 4-ethylguaiacol with 0.25 ppm in the YM diluted sample.

Furthermore, the 4EG:4EP ratios of 1:7 (in YM diluted TF) and 1:6 (in water diluted TF) are close to the ones suggested in literature (Chatonnet, et al., 1992). While the 1:10 ratio in Tinta Roriz diluted with YM equals references in literature, the ratio in water diluted TR is a little different at 1:16.




Figure 17: Initial volatile phenol concentration in Tinta Roriz (TR) and Touriga Franca and amount present in diluted wine after contamination with *D. bruxellensis* PYCC 4801; the error bars represent the standard deviation of two determinations (W: wine; W+YM: wine diluted with YM medium; W+W: wine diluted with water)

Compared to the two previous samples (Touriga Franca and Tinta Roriz), the Syrah wine has shown a smaller difference in volatile phenol production between water and YM diluted wine. Furthermore, as can be seen in Figure 18, the perception thresholds for 4-ethylphenol and 4-ethylguaiacol as defined by Chatonnet et al. (1992) are not exceeded. From that however, it cannot be concluded that there is no phenolic off flavour since also the combined threshold of 0.42 ppm (Chatonnet, et al., 1992) and other factors have to be considered. Regarding the ratio 4EG:4EP which is about 1:6 in YM diluted and 1:5 in water diluted wine, it can be observed that the production of 4-ethylphenol in relation to 4-ethylguaiacol was smaller than suggested in literature (Chatonnet, et al., 1992).




Figure 18: Initial volatile phenol concentration in Syrah (SY) and the amount present in diluted wine after contamination with *D. bruxellensis* PYCC 4801; the error bars represent the standard deviation of two determinations (W: wine; W+YM: wine diluted with YM medium; W+W: wine diluted with water)

The results for all four Touriga Nacional wines that were analysed are presented below in Figure 19. In two of the samples, namely TN2 und TN4, the 4-ethylphenol perception threshold as defined by Chatonnet et al. (1992) was, at least in the YM diluted sample, exceeded. Apart from TN4, the 4-ethylguaiacol concentrations either exceeded or were really close to perceivable quantities. As mentioned before, if both compounds are present in wine, in combination they are more easily perceivable and thus the off-flavour may occur even if individual threshold values are not exceeded (Chatonnet, et al., 1992). Furthermore, 4EG:4EP ratios are quite similar in all Touriga Nacional samples. They comprise ratios of 1:3, 1:4 and 1:5.




Figure 19: Initial volatile phenol concentration in Touriga Nacional wines and the amount present in diluted wines after contamination with *D. bruxellensis* PYCC 4801; the error bars represent the standard deviation of two determinations (W: wine; W+YM: wine diluted with YM medium; W+W: wine diluted with water)

As can be seen by comparing Figure 20 to all previous wines, in the Sousão wines the 4-ethylguaiacol quantities are much higher in relation to 4-ethylphenol. While those levels are really close or even exceeding the perception threshold, ethylphenol quantities only reach half of the amounts that are considered perceivable. However, it should be considered that these amounts of volatile phenols were obtained in (50:50) diluted wines, which may indicate that in undiluted wines higher amounts of these compounds may be produced by *Dekkera/Brettanomyces* strains.




Figure 20: Initial volatile phenol concentration in Sousão wines compared to the amount present in diluted wines after contamination with *D. bruxellensis* PYCC 4801; the error bars represent the standard deviation of two determinations (W: wine; W+YM: wine diluted with YM medium; W+W: wine diluted with water)

4.4.1 Estimation of the potential for volatile phenol production in diluted wines

The estimation of the "potential" for volatile phenol production in varietal wines following *D. bruxellensis* PYCC 4801 contamination was determined by subtracting the quantity of 4-ethylphenol and 4-ethylguaiacol in the diluted wines after contamination from the initial amount of the same compounds in the original wines. Thus, this "potential" represents the amounts of volatile phenols that were produced by the action of the spoilage yeasts only. As mentioned previously, the results were calculated from values obtained from diluted wines although theoretically quantities could presumably be twice as high in the full concentration wines.

As can be seen when comparing Figure 21 and Figure 22, generally all wines exhibit a higher potential to produce 4-ethylphenol than 4-ethylguaiacol. Moreover, the differences in potential for 4-ethylguaiacol production between varieties seems rather low and therefore no significant statement can be made (Figure 21).




Figure 21: Concentration of 4-ethylguaiacol produced by *D. bruxellensis* in diluted varietal red wines (TF: Touriga Franca; TR: Tinta Roriz; SY: Syrah; TN1-TN4: Touriga Nacional; S1-S4: Sousão)

In regard to 4-ethylphenol however, a clear trend is to be seen comparing Touriga Franca and Tinta Roriz to the remaining wines. Both of them exhibit a potential at least three times higher than other varieties, in both wine diluted with water and YM medium. Furthermore, the values of the four examined Touriga Nacional samples fluctuate between 0.38 ppm and 0.72 ppm (4-ethylphenol in wine diluted with YM) but can be clearly differentiated from Touriga Franca and Tinta Roriz samples. However, the potential of the examined Syrah appears to lie in the same range and can therefore not be differentiated. From the data presented below, it can be suggested moreover that Sousão is a variety with a rather low potential. All four samples show similar quantities of 4-ethylphenol in ranges rarely exceeding 0.2 ppm.




Figure 22: Concentration of 4-ethylphenol produced by *D. bruxellensis* in diluted varietal red wines (TF: Touriga Franca; TR: Tinta Roriz; SY: Syrah; TN1-TN4: Touriga Nacional; S1-S4: Sousão)

Calculating the mean value of produced 4-ethylphenol in varieties of which more than one wine was examined, it can be observed that the potential of Touriga Nacional indeed is twice as high as compared to Sousão. It has to be considered that the results for Sousão are more consistent since the standard deviation is smaller. Furthermore, the mean value is slightly higher in YM medium. In order to compare with other varieties and obtain representative results, more varietal wines have to be selected and examined.

Table 7: Mean potential of varieties to produce 4-ethylphenol in diluted wines in ppm (+/- standard deviation)

Variety	W+YM	W+W
Touriga Nacional	0,56 +/-0,16	0,48 +/-0,22
Sousão	0,22 +/-0,05	0,20 +/-0,09

(W+YM: wine diluted with YM medium; W+W: wine diluted with water)

4.4.2 Estimation of volatile phenol precursor quantities in diluted wines

In an attempt to estimate the quantity of volatile phenol precursors naturally present in the wines, diluted wines were supplemented with either 5.0 or 50.0 ppm of *p*-coumaric and ferulic acid and inoculated with *D. bruxellensis* PYCC 4801. The production of volatile phenols in the supplemented and artificially contaminated wines was monitored and compared to the levels of the original wines. If the conversion rates of precursors to volatile phenols in the supplemented wines were close to 100%, a “standard addition method” approach could be used for each sample in order to estimate the precursor amounts in wines. In order to calculate the molar conversion rate, the amount of volatile phenols that were produced only from the added precursors were calculated by subtracting the 4-ethylphenol or 4-ethylguaiacol content formed in non-supplemented samples, from the quantity produced in supplemented samples. The result was put in relation to 3.7 or 37.0

ppm (for 4-ethylphenol) and 3.7 or 37.0 ppm (for 4-ethylguaiacol) accounting for a 100% conversion rate at a precursor addition of 5.0 and 50.0 ppm, respectively.

As can be seen in Table 8, the molar conversion ratios of *p*-coumaric acid to 4-ethylphenol were variable and did not exceed 56%. In case of the ferulic acid to 4-ethylguaiacol conversion (Table 9), the molar conversion rates were also very different between samples and no specific pattern for any of the wines that could be correlated to the grape variety factor, can be observed. However, in most samples, a slight trend towards higher conversion rates in YM diluted samples, for both 4-ethylphenol and 4-ethylguaiacol production was observed (Tables 8 and 9).

Table 8: Molar conversion rates (%) of *p*-coumaric acid to 4-ethylphenol by *D. bruxellensis* in different varietal wines fortified with the precursor

Sample	<i>P</i>-coumaric acid added (ppm)	Molar conversion rate in wine+water (%)	Molar conversion rate in wine+YM (%)
TF	5	56	32
TR	50	28	43
SY	50	20	34
TN1	50	37	40
TN2	50	6	30
TN3	5	36	39
TN4	50	27	21
S1	50	34	45
S2	50	1	37
S3	50	6	29
S4	50	1	0

(**TF**: Touriga Franca; **TR**: Tinta Roriz; **SY**: Syrah; **TN1-TN4**: Touriga Nacional; **S1-S4**: Sousão)

Table 9: Molar conversion rates (%) of ferrulic acid to 4-ethylguaiacol by *D. bruxellensis* in different varietal wines fortified with the precursor

Sample	Ferrulic acid added (ppm)	Molar conversion rate in wine+water (%)	Molar conversion rate in wine+YM (%)
TF	5	64	42
TR	50	40	45
SY	50	28	41
TN1	50	42	42
TN2	50	18	30
TN3	5	37	43
TN4	50	26	12
S1	50	37	48
S2	50	18	30
S3	50	16	38
S4	50	57	74

(**TF**: Touriga Franca; **TR**: Tinta Roriz; **SY**: Syrah; **TN1-TN4**: Touriga Nacional; **S1-S4**: Sousão)

5 DISCUSSION

In this project, 11 different Portuguese single varietal wines were analyzed for their potential to produce volatile phenols if contaminated by a *Dekkera bruxellensis* strain. In order to obtain representative results for varietal wines from Portugal however, more samples of each variety have to be selected and analyzed. Additionally, other regions apart from the Douro valley have to be represented. Hence, all observations discussed here shall serve only as preliminary results which can lead to further research being conducted.

5.1 Yeast growth

Test trials showed that certain factors, such as alcohol content or pH value influence yeast growth. The lack of growth and decline in cell numbers when *D. bruxellensis* PYCC 4801 is inoculated in pure wines but an opposite behavior when grown in diluted wine confirms the results of Dias et al. (2003a) who had observed reduced growth and production of volatile phenols with high ethanol concentrations. Especially in alcohol concentrations equal to or higher than 13% (as present in most of the wines used in these experiments), growth is very unlikely (Dias, et al., 2003a). In the present experiments, dilution with YM has shown that there may be a growth enhancing influence of added substrate that is not very significant in terms of volatile phenol production however. Dias et al. (2003a) confirms that supplementation with sugar and thus energy may be beneficial for the growth of *Dekkera/Brettanomyces*.

From the results presented above it can be seen that *D. bruxellensis* PYCC 4801 is able to grow in diluted wine at 7.5 % vol. of ethanol and pH 3.8. In most of the samples, the phases characteristic for yeast growth can be differentiated, with the exception of lag that is not quite visible. In three Sousão samples the phases are not easily recognizable and lower final cell numbers occur. This may be due to some LAB contamination in the culture that was observed in these samples. However the volatile phenol levels produced by the yeasts were similar to non-contaminated experiments. Due to time constraints it was not possible to repeat the experiments with these wines.

Furthermore, some of the results suggest a slight inhibition of growth when *p*-coumaric and ferulic acid are present in high quantities such as 50 ppm. Grbin (2008) has shown that in growth medium, ferulic acid can inhibit growth at concentrations of about 19.4 ppm and even completely suppress *D. bruxellensis* strains at 194.2 ppm. A synergistic effect has been observed in the presence of ethanol and other hydroxycinnamic acids such as *p*-coumaric acid. The latter alone however did not show an inhibitory effect as strong as ferulic acid

(Grbin, P.R., 2008). Contrary to observations of Grbin (2008), in the present experiments no delay in the onset of the stationary phase could be noticed in supplemented samples.

5.2 Volatile phenols production

First of all, the results show that *D. bruxellensis* PYCC 4801 is able to convert different amounts of precursors in wines to their corresponding ethyl derivatives. Since the original wines contained quantities far below the threshold and detection levels, the volatile phenols present after incubation of inoculated wines were present due to yeast metabolic activity. In all diluted red wine samples, both 4-ethylphenol and 4-ethylguaiacol were detected. The results show, in agreement with literature, that the 4-ethylphenol values reached, were higher than 4-ethylguaiacol and both lie within the ranges detected by other authors (Table 2). Moreover, the 4-ethylguaiacol values reached, did not vary as much between varieties, while 4-ethylphenol quantities did, confirming Pollnitz et al. (2000), who stated that there are no significant differences in 4-ethylguaiacol between varieties. Furthermore, no 4-vinylphenol was detected in any of the samples which indicates that *D. bruxellensis* PYCC 4801 was converting effectively, producing more end product (4-ethylphenol) than intermediate product (4-vinylphenol). Another reason for the absence of 4-vinylphenol may be its reaction with anthocyanins forming stable pyranoanthocyanins normally leading to lower amounts of this compound in red wine (Benito, et al., 2009).

Since, in most cases, volatile phenol concentrations exceed the perception threshold established by Chatonnet et al. (1992) it shows that, generally, all these wines are at risk of developing off-flavours if contaminated by *Dekkera/Brettanomyces bruxellensis*. However, that the examined Sousão wines, for instance, did not in any case reach those thresholds, does not imply that the development of phenolic off-flavour is not possible. Due to the synergistic effects between 4-ethylguaiacol and 4-ethylphenol, their intensity is not directly proportional to the concentration present. Furthermore, the sensory value of wines is such a very complex matter that threshold values may not be applicable if certain other compounds mask or enhance the sensorial effect of volatile phenols. It has been reported that wines with 4-ethylphenol values below the threshold levels exhibited phenolic character while wines strongly exceeding those levels were recognized as off-flavour free in sensorial trials (Benito, et al., 2008). In regard to perception thresholds, it also has to be considered that the wine potentially has twice as many precursors available for conversion to ethyl derivatives as the diluted samples used in these experiments.

Furthermore, since samples of the same varieties showed similar 4-ethylguaiacol to 4-ethylphenol ratios, it can be suggested from the present results that varieties show distinct

ratios. The highest ratio was found in water diluted Tinta Roriz with 1:16 and lowest in diluted Sousão wines exhibiting ratios between 1:1 and 1:2. These findings are confirmed by Pollnitz et al. (2000) who also found that some varieties may be differentiated based on the 4EG:4EP ratio.

5.3 Potential of volatile phenol production

The results indicate that wines may be grouped regarding their potential to produce volatile phenols after being artificially contaminated with *D. bruxellensis* PYCC 4801. This potential considers the quantity of *p*-coumaric acid naturally present in the wines that was converted into 4-ethylphenol by the action of *D. bruxellensis* PYCC 4801. As compared to a chemical analysis of only free *p*-coumaric acid, this biological approach considers precursors in different forms such as an esterified form and therefore may be more accurate in order to estimate the volatile phenol potential of a wine.

The highest potential was observed in the Touriga Franca and Tinta Roriz wines. This is in accordance with results obtained by Ribeiro de Lima et al. (1998) who found Touriga Franca to be the variety with the highest quantities of *p*-coumaric acid of all examined wines. In the present experiment Sousão wines produced the lowest amounts of volatile phenols while the Touriga Nacional and Syrah wines are in the middle. Regarding the quantities of *p*-coumaric acid present in these varieties, no references for comparison are available in literature currently.

As can be seen in the results, all examined Sousão wines show similarly small potential while the Touriga Nacional wines possess higher ones that are as well in an analogous range within the variety. Therefore, it can be suggested that wines made from one variety can be grouped together regarding the potential of volatile phenol production. However, some of the examined wines show similar values such as Tinta Roriz and Touriga Franca amongst each other or Syrah compared to the Touriga Nacional wines and can therefore not be differentiated by this characteristic. Certainly, especially for the last mentioned varieties a higher number of samples have to be examined in the future in order to find out whether there are similarities in potential within them.

When estimating the varietal influence on volatile phenol production, it has to be noted however, that the present approach only considers the availability of precursors to the yeast, not the actual susceptibility of the undiluted wine to contamination and growth of *D. bruxellensis* as well as any other external factors.

5.4 Estimation of precursor quantities

In order to determine indirectly the quantities of precursors of volatile phenols originally present in the wines, the molar conversion rates of *p*-coumaric acid and ferulic acid to 4-ethylphenol and 4-ethylguaiacol, respectively, were calculated for the samples supplemented with the precursors. Since the amounts of added precursors were relatively high and the molar conversion ratios (calculated by dividing the molar concentration of volatile phenols by the initial molar concentration of phenolic acids) were lower than 100%, it was not possible to use the proposed “standard addition” approach for estimating the precursor levels in the wines and then comparing them between varieties. Furthermore, it is known that besides the free phenolic acids present in wines, there are other potential precursors of volatile phenols such as esters of phenolic acids with ethanol, tartaric acid and anthocyanins (Ribéreau-Gayon, et al., 2006). Therefore, it is difficult to estimate how many of the volatile phenols produced from the wine precursors are related to free and how many to combined phenolic acids. Moreover, it has been found that *Dekkera* yeasts can decrease the amount of free *p*-coumaric acid available for conversion by adsorbing them to their cell walls (Salameh, et al., 2008) and thereby interfere with the calculation of exact conversion rates. In conclusion, further experiments are necessary to study the effect of the type and concentration level of precursors in the production of volatile phenols in wines by *Dekkera/Brettanomyces*.

6 CONCLUSIONS

- All the mono varietal wines used in this work (diluted 50:50) were found to be able to support the growth of *D. bruxellensis* PYCC 4801. In most cases, the addition of YM medium to the wines did not lead to a significant improvement of yeast growth.
- A correlation between grape variety and the production of volatile phenols can be suggested. Touriga Franca and Tinta Roriz wines produced higher concentrations of volatile phenols while the Sousão wines clearly remained below the other wines. However, several other factors such as the winemaking practices have to be considered as well.
- A small variability in the production of volatile phenols was found within the Touriga Nacional wines (from 0.4 to 0.8 ppm of 4-ethylphenol).
- The present study confirms that *D. bruxellensis* PYCC 4801 produces 4-ethylphenol and 4-ethylguaiacol from phenolic acids while the intermediary volatile phenols were not detected.
- The contamination with *D. bruxellensis* PYCC 4801 and subsequent determination of volatile phenols in wines may be used as a method to obtain information on the wines potential to produce volatile phenols.

Further work should focus on the selection of more samples of the same variety and compare different vintages in order to obtain representative results. Furthermore, other influencing factors such as the susceptibility to contamination of different wines should be considered and included. Experiments on undiluted wines as well as long-term studies and the evaluation of naturally spoiled wines could also reveal interesting outcomes.

REFERENCES

- Arvik T.J. and Henick-Kling T.** *Brettanomyces bruxellensis* occurrence, growth and effect on wine flavour [Article] // Practical Winery & Vineyard. - 2002. - Vol. 24. - pp. 48-56.
- Barthelmebs L., Divies C. and Cavin J.-F.** Knockout of the p-Coumarate Decarboxylase Gene from *Lactobacillus* Other Inducible Enzymatic Activities Involved in Phenolic Acid Metabolism [Article] // Applied and Environmental Microbiology. - 2000. - Vol. 66, No. 8. - pp. 3368-3375.
- Basha S.M., Musingo M. and Colova V.S.** Compositional differences in the phenolics compounds of muscadine and bunch grape wines [Article] // African Journal of Biotechnology. - October 2004. - pp. 523-528.
- Benito S. [et al.]** A method for estimating *Dekkera/Brettanomyces* populations in wines [Article] // Journal of Applied Microbiology. - 2008. - Vol. 106. - pp. 1743-1751.
- Benito S. [et al.]** Minimization of ethylphenol precursors in red wines via the formation of pyranoanthocyanins by selected yeasts [Article] // International Journal of Microbiology. - 2009. - Vol. 132. - pp. 145-152.
- Cabrita M.J. [et al.]** Conversion of hydroxycinnamic acids into volatile phenols in a synthetic medium and in red wine by *Dekkera bruxellensis* [Article] // Ciência e Tecnologia de Alimentos. - 2012. - Vol. 32. - pp. 106-111.
- Campos F.M., Couto J.A. and Hogg T.A.** Influence of phenolic acids on growth and inactivation of *Oenococcus oeni* and *Lactobacillus hilgardii* [Article] // Journal of Applied Microbiology. - 2003. - Vol. 94. - pp. 167-174.
- Castillo-Sánchez J.X. [et al.]** Phenolic compounds and colour stability of Vinha~o wines: Influence of wine-making protocol and fining agents [Article] // Food Chemistry. - 2008. - 106. - pp. 16-26.
- Cavin J.F. [et al.]** Ability of Wine Lactic Acid Bacteria to Metabolize Phenol Carboxylic Acids [Article] // American Journal of Enology and Viticulture. - 1993. - Vol. 44, Nr. 1. - pp. 76-80.
- Chatonnet P. [et al.]** Synthesis of Volatile Phenols by *Saccharomyces cerevisiae* in Wines [Article] // Journal of the Science of Food and Agriculture. - 1993. - Vo. 62. - pp. 191-202.
- Chatonnet P., Dubourdieu D. and Boidron J.N.** The Influence of *Brettanomyces/Dekkera* sp. Yeasts and Lactic Acid Bacteria on the Ethylphenol Content of Red Wines [Article] // American Journal of Enology and Viticulture. - 1995. - Vol. 46, No. 4. - pp. 463-468.
- Chatonnet P., Dubourdieu D. and Boidron J.-N.** The Origin of Ethylphenols in Wines [Article] // Journal of Food Science and Agriculture. - 1992. - Vol. 60. - pp. 165-178.
- Chatonnet P., Viala C. and Dubourdieu D.** Influence of Polyphenolic Components of Red Wines on the Microbial Synthesis of Volatile Phenols [Article] // American Journal of Enology and Viticulture. - 1997. - Vol. 48, No. 4. - pp. 443-448.
- Connell L., Stender H. and Edwards C.G.** Rapid Detection and Identification of *Brettanomyces* from Winery Air Samples Based on Peptide Nucleic Acid Analysis [Article] // American Journal of Enology and Viticulture. - 2002. - Vol. 53, No. 5. - pp. 322-324.

- Couto J.A. [et al.]** Ability of Lactic Acid Bacteria to Produce Volatile Phenols [Article] // American Journal of Enology and Viticulture. - 2006. - Vol. 57, No. 2. - pp. 166-171.
- Curtin C.D. [et al.]** Genetic diversity of *Dekkera bruxellensis* yeasts isolated from Australian Wines [Article] // FEMS Yeast Research. - 2007. - Vol. 7. - pp. 471-481.
- De Beer D. [et al.]** Phenolic Compounds: A Review of Their Possible Role as In Vivo Antioxidants of Wine [Article] // South African Journal for Enology and Viticulture. - South Africa : [s.n.], February 2002. - Vol. 23, No. 2. - pp. 48-61.
- De las Rivas B. and Rodríguez H.** Molecular Screening of Wine Lactic Acid Bacteria Degrading Hydroxycinnamic Acids [Article] // Journal of Agricultural and Food Chemistry. - 2009. - Vol. 57. - pp. 490-494.
- Dias L. [et al.]** Factors affecting the production of 4-ethylphenol by the yeast *Dekkera Bruxellensis* in enological conditions [Article] // Food Microbiology. - 2003a. - Vol. 20. - pp. 377-384.
- Dias L. [et al.]** Identification of yeasts isolated from wine-related environments and capable of producing 4-ethylphenol [Article] // Food Microbiology. - 2003b. - Vol. 20. - pp. 567-574.
- Dopico-García M.S. [et al.]** Principal components of phenolics to characterize red Vinho Verde grapes: Anthocyanins or non-coloured compounds? [Article] // Talanta. - 2008. - Vol. 75. - pp. 1190-1202.
- Dugelay I. [et al.]** Role of cinnamoyl esterase activities from enzyme preparations on the formation of volatile phenols during winemaking [Article] // Journal of Agricultural and Food Chemistry. - 1993. - Vol. 41. - pp. 2092-2096.
- Edlin D.A.N. [et al.]** The biotransformation of simple phenolic compounds by *Brettanomyces anomalus* [Article] // FEMS Microbiology Letters. - 1995. - Vol. 125. - pp. 311-316.
- Gambelli L. and Santaroni G.P.** Polyphenols content in some Italian red wines of different geographical origins [Article] // Journal of Food Composition and Analysis. - 2004. - Vol. 17. - pp. 613-618.
- Gerbeaux V., Vincent B. and Bertrand A.** Influence of maceration temperature and enzymes on the content of volatile phenols in Pinot noir wines [Article] // American Journal of Enology and Viticulture. - 2002. - Vol. 53. - pp. 131-137.
- Goldberg D.M [et al.]** Quercetin and p-coumaric acid concentrations in commercial wines [Article] // American Journal of Enology and Viticulture. - 1998. - Vol. 49. - pp. 142-151.
- Grbin, P.R.** Grape and Wine Research and Development Organisation [Online] // Growth and physiology of *Brettanomyces/Dekkera* yeast and the production of flavour compounds in wine. - 2008. - March 2014. - <http://www.gwrdc.com.au/wp-content/uploads/2012/09/UA-03-01.pdf>.
- Heresztyn T.** Metabolism of volatile phenolic compounds from hydroxycinnamic acids by *Brettanomyces* yeast [Article] // Archives of Microbiology. - 1986. - Vol. 146. - pp. 96-98.
- Hesford F. and Schneider K.** Die Entdeckung eines dritten Ethylphenols als Mitversursacher des *Brettanomyces*-Fehltons [Article] // Schweizer Obst-Weinbau. - 2004. - Vol. 13. - pp. 11-13.

Jackson R.S. Chemical Constituents of Grapes and Wine [Book Section] = Phenols and Related Phenol (Phenyl) // Wine Science - Principles and Applications. - San Diego, California : Elsevier, 2008. - 3..

Kheir J. [et al.] Impact of volatile phenols and their precursors on wine quality and control measures of *Brettanomyces/Dekkera* yeasts [Article] // European Food Research and Technology. - 2013.

Loureiro V. and Malfeito-Ferreira M. Spoilage yeasts in the wine industry [Article] // International Journal of Food Microbiology. - 2003. - Vol. 86. - pp. 23-50.

Mateus N., Machado J.M. and De Freitas V. Development changes of anthocyanins in *Vitis vinifera* grapes grown in the Douro Valley and concentration in respective wines [Article] // Journal of the Science of Food and Agriculture. - 2002. - Vol. 82. - pp. 1689-1695.

Nelson L. The production of volatile phenols by wine microorganisms [Article] // Master thesis at Stellenbosch University. - 2008.

Nikfardjam M.P., May B. and Tschiersch C. 4-Ethylphenol and 4-ethylguaiacol contents in bottled wines from the German 'Württemberg' region [Article] // European Food Research and Technology. - 2009. - pp. 333-341.

Oelofse A., Pretorius I.S. and du Toit M. Significance of *Brettanomyces* and *Dekkera* during Winemaking: A Synoptic Review [Article] // South African Journal of Enology and Viticulture. - 2008. - Vol. 29, No. 2. - pp. 128-144.

Ota A. [et al.] Interactions of p-coumaric, caffeic and ferulic acids and their styrenes with model lipid membranes [Article] // Food Chemistry. - 2001. - Vol. 125. - pp. 1256-1261.

Pena-Neira A. [et al.] A survey of phenolic compounds in Spanish wines of different geographical origin [Article] // European Food Research and Technology. - 2000. - pp. 445-448.

Pollnitz A.P., Pardon K.H. and Sefton M.A. Quantitative analysis of 4-ethylphenol and 4-ethylguaiacol in red wine [Article] // Journal of Chromatographie A. - 2000. - Vol. 874. - pp. 101-109.

Rentzsch M., Wilkens A. and Winterhalter P. Non-flavonoid Phenolic Compounds [Book Section] // Wine Chemistry and Biochemistry / ed. Moreno-Arribas M.V. and Polo M.C.. - [s.l.] : Springer, 2009.

Ribeiro de Lima M.T. and Cabanis J.C. The phenolic acid content of different varieties and vintages of wines in Continental Portugal and the Azores Islands [Article] // Bulletin de l'OIV. - 1998. - pp. 1012-1027.

Ribéreau-Gayon P. [et al.] Chemical Nature, Origin and Consequences of the Main Organoleptic Defects [Book Section] // Handbook of Enology Volume 2: The Chemistry of Wine Stabilization and Treatments. - England : John Wiley & Sons. Ltd, 2006.

Ribéreau-Gayon P. [et al.] Phenolic Compounds [Book Section] // Handbook of Enology Volume 2: The Chemistry of Wine Stabilization and Treatments. - England : John Wiley & Sons, Ltd, 2006. - 2..

Rodríguez-Delgado M.-A. [et al.] Principal component analysis of the polyphenol content in young red wines [Article] // Food Chemistry. - 2002. - Vol. 78. - pp. 523-532.

Romano A. [et al.] <https://home.zhaw.ch> [Online] // Sensorial Aspects of "Brett-Character": Re-evaluation of the Olfactory Perception Threshold of Volatile Phenols in Red Wine. - 2008. - November 16., 2013. - <https://home.zhaw.ch/~yere/pdf/Teil63%20-%20Expression%20of%20Multidisciplinary.pdf>.

Salameh D. [et al.] Highlight on the problems generated by p-coumaric acid analysis in wine fermentations [Article] // Food Chemistry. - 2008. - Vol. 107. - pp. 1661-1667.

Shinohara T., Kubodera S. and Yanagida F. Distribution of Phenolic Yeasts and Production of Phenolic Off-Flavors in Wine Fermentation [Article] // Journal of Bioscience and Bioengineering. - 2000. - Vol. 90, No. 1. - pp. 90-97.

Soleas G.J. [et al.] Toward the Fingerprinting of Wines: Cultivar-Related Patterns of Polyphenolic Constituents in Ontario Wines [Article] // Journal of Agriculture and Food Chemistry. - 1997. - Vol. 45. - pp. 3871-3880.

Suárez R. [et al.] The production of ethylphenols in wine by yeasts of the genera *Brettanomyces* and *Dekkera*: A review [Article] // Food Chemistry. - 2007. - Vol. 102. - pp. 10-21.

Valentao P. [et al.] Influence of *Dekkera bruxellensis* on the contents of anthocyanins, organic acids and volatile phenols of Dao red wine [Article] // Food Chemistry. - 2007. - Vol. 100. - pp. 64-70.

Van der Walt J.P. and Van der Kerken A.E. The Wine Yeasts of the Cape Part I. - A Taxonomical Survey of the Yeasts Causing Turbidity in South African Table Wines [Article] // Antonie van Leeuwenhoek. - 1958. - Vol. 27. - pp. 239-252.

Van der Walt J.P. and Van der Kerken A.E. The Wine Yeasts of the Cape Part V. - Studies in the Occurrence of *Brettanomyces Intermedius* and *Brettanomyces Schanderlii* [Article] // Antonie van Leeuwenhoek. - 1960. - Vol. 24. - pp. 81-90.

Van der Walt J.P. *Dekkera*, a new genus of the Saccharomycetaceae [Article] // Antonie van Leeuwenhoek. - 1964. - Vol. 30. - pp. 273-280.

Villiers A. [et al.] Classification of South African red and white wines according to grape variety based on the non-coloured phenolic content [Article] // European Food Research and Technology. - 2005. - Vol. 221. - pp. 520-528.

Whiting G.C. and Carr J.G. Metabolism of Cinnamic Acid and Hydroxy-Cinnamic Acids by *Lactobacillus pastorianus* var. *quinicus* [Article] // Nature. - 1959. - Vol. 184. - pp. 1427-1428.