

4-1-1999

Unevening the Odds of "Even Up"

Arthur T. Benjamin
Harvey Mudd College

Jennifer J. Quinn
Occidental College

Recommended Citation

Benjamin, Arthur T. and Jennifer J. Quinn. "Unevening the Odds of 'Even Up.'" *Mathematics Magazine*, Vol 72, No. 2, pp. 149-150, April 1999.

This Article is brought to you for free and open access by the HMC Faculty Scholarship at Scholarship @ Claremont. It has been accepted for inclusion in All HMC Faculty Publications and Research by an authorized administrator of Scholarship @ Claremont. For more information, please contact scholarship@cuc.claremont.edu.



Unevening the Odds of "Even Up"

Author(s): Arthur T. Benjamin and Jennifer J. Quinn

Source: *Mathematics Magazine*, Vol. 72, No. 2 (Apr., 1999), pp. 145-146

Published by: [Mathematical Association of America](#)

Stable URL: <http://www.jstor.org/stable/2690601>

Accessed: 11/06/2013 17:11

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Mathematical Association of America is collaborating with JSTOR to digitize, preserve and extend access to *Mathematics Magazine*.

<http://www.jstor.org>

Acknowledgment This paper was written when the first author was visiting the Department of Mathematics at Wilfrid Laurier University, December 1996–August 1997. The hospitality of WLU is greatly appreciated. The authors would like to thank the referee for many constructive suggestions, which substantially improved this paper.

REFERENCES

1. Benkart, G. M., et. al., Problem #600, *College Math. Journal* 28:2 (1997), 146.
2. Rosen, Kenneth H., *Elementary Number Theory and Its Applications*, 3rd ed., Addison Wesley, Reading, MA, 1993.
3. Shan, Zun and Wang, Edward T. H., Solution to Problem #600 (Unit Multiples), *College Math. Journal*, 29:2 (1998), 156–157.
4. Sierpinski, W., *Elementary Theory of Numbers*, North-Holland, Amsterdam, The Netherlands, 1988.

Unevening the Odds of “Even Up”

ARTHUR T. BENJAMIN
Harvey Mudd College
Claremont, CA 91711

JENNIFER J. QUINN
Occidental College
Los Angeles, CA 90041

The Disclaimer The authors take no responsibility for any gambling hustles or scams based on applications of the principles contained in this note.

The Game “Even Up” is a game of solitaire played with 40 cards from a standard deck that has its jacks, queens, and kings removed. The cards are shuffled and dealt in a row. If a consecutive pair of cards adds to an even number, then that pair can be removed. The object of the game is to remove all of the cards.

More generally, we can play Even Up with $2n$ cards, x of them being odd and $2n - x$ being even. We require the number of cards to be even since the game cannot be won with an odd number of cards. In fact, the game cannot be won when x is odd since odd valued cards are removed in pairs. Harkleroad [1] showed that the game involves no skill, in that the outcome is predetermined by the original order of the $2n$ cards, and that the probability of winning is $p(2n, x) = \binom{n}{x/2}^2 / \binom{2n}{x}$. Thus the probability of winning the original game is $p(40, 20) = 0.248$.

A few remarks about $p(2n, x)$ are called for. Clearly $p(2n, 0) = 1 = p(2n, 2n)$. By comparing $p(2n, x)$ with $p(2n, x - 2)$, one sees that for fixed n the probability of winning is minimized when $x = n$. When n is large, we can use Stirling’s formula ($n! \approx (n/e)^n \cdot \sqrt{2\pi n}$) to obtain $p(2n, n) \approx 2/\sqrt{\pi n}$.

For our purposes, any arrangement of $2n$ cards can be represented as the product of a ’s and b ’s with a ’s denoting odd cards and b ’s denoting even cards. The rules of Even Up reduce to the two multiplications $a^2 = 1$ and $b^2 = 1$. Every game simplifies to exactly one string of the form $(ab)^z$, where $-n \leq z \leq n$ and $(ab)^{-z} = (ba)^z$. Winning games occur when $z = 0$. Letting $f(2n, x, z)$ denote the number of arrange-

ments of x a 's and $2n - x$ b 's that reduce to $(ab)^z$, Harkleroad gives two proofs that

$$f(2n, x, z) = \binom{n}{(x+z)/2} \binom{n}{(x-z)/2}. \quad (1)$$

One proof used induction and the other used a complicated summation. Here we provide a direct combinatorial explanation of equation (1).

The Proof Consider a string of $2n$ symbols as n ordered pairs, where each pair is either (a, a) , (b, b) , (a, b) , or (b, a) . Such a string will reduce to $(a, b)^z$, where z is the number of (a, b) pairs minus the number of (b, a) pairs. If we let the four pairs have scores of 0, 0, 1, and -1 respectively, then z denotes the total score. Another way to calculate the score is to assign a value to each symbol as follows: a beginning a in an ordered pair gets a score of 1, an ending a in an ordered pair gets a score of -1 , and all b 's get a score of 0. Hence the total score is equal to the number of beginning a 's minus the number of ending a 's. If x' denotes the number of beginning a 's ($0 \leq x' \leq x$), then the score $z = x' - (x - x') = 2x' - x$. Thus to achieve a score of z , we must choose $x' = (x + z)/2$ of our n pairs to begin with an a and $x - x' = (x - z)/2$ of our n pairs to end with an a . This completes the proof of equation (1).

The probability that a game with x a 's and $2n - x$ b 's reduces to $(ab)^z$ is $f(2n, x, z) / \binom{2n}{x}$. Notice that $z = 2x' - x$ will always have the same parity as x . (This is reflected in (1) since $\binom{n}{k/2}$ is 0 when k is odd.) Thus, we see again that a game with an *even* number of cards but *odd* number of odd cards is impossible to win.

The Scam After explaining the game of Even Up to your mark, challenge him to a duel. Wager that you can win the game in strictly fewer attempts than he can. He can shuffle your cards before every deal. You play the game until you win. Say it takes you 4 attempts. Now it's his turn to play. How can you be sure that it will take him at least 5 attempts? When *you* play the game, use 10 even cards and 10 odd cards. Each attempt has a 34% chance of success. After you win, shuffle the cards, but secretly add (or remove, if you prefer) one even and one odd card to his deck. This will make his winning probability 0.

REFERENCE

1. Leon Harkleroad, Lowering the odds for "Even up", this MAGAZINE 63 (1990), 160–163.