

Journal section: *Oral Medicine and Pathology*

doi:10.4317/medoral.18245

Publication Types: *Review*<http://dx.doi.org/doi:10.4317/medoral.18345>

Intralesional corticosteroid injections in the treatment of central giant cell lesions of the jaws: A meta-analytic study

Rafael-Lima-Verde Osterne ¹, Phelype-Maia Araújo ², Abrahao-Cavalcante-Gomes de Souza-Carvalho ³, Roberta-Barroso Cavalcante ⁴, Eduardo Sant'Ana ⁵, Renato-Luiz-Maia Nogueira ⁶

¹ DDS, MSc, Assistant Professor, Department of Pathology, Fortaleza University School of Medicine, Fortaleza, Brazil

² DDS, MSc student, Federal University of Ceará School of Dentistry, Fortaleza, Brazil

³ DDS, MSc, PhD, Assistant Professor, Faculdade Católica Rainha do Sertão, School of Dentistry, Quixadá, Brazil

⁴ DDS, MSc, PhD, Associate Professor, Department of Oral and Maxillofacial Pathology, Fortaleza University School of Dentistry, Fortaleza, Brazil

⁵ DDS, MSc, PhD, Associate Professor of Oral and Maxillofacial Surgery, Bauru School of Dentistry, University of São Paulo, Bauru, Brazil

⁶ DDS, MSc, PhD, Associate Professor, Department of Dental Clinic, Discipline of Oral and Maxillofacial Surgery and Stomatology, Federal University of Ceara School of Dentistry, Fortaleza, Brazil. Oral and Maxillofacial Surgeon, Department of Oral and Maxillofacial Surgery, Memorial Batista Hospital, Fortaleza, Brazil

Correspondence:

Universidade de Fortaleza

Fortaleza University/ Universidad de Fortaleza

Av. Washington Soares

1321, Edson Queiroz

60811-905 Fortaleza

Ceará, Brasil P.O. Box 1258

rlimaverde@unifor.br

Osterne RLV, Araújo PM, ACG de Souza-Carvalho, Cavalcante RB, Sant'Ana E, Nogueira RLM. Intralesional corticosteroid injections in the treatment of central giant cell lesions of the jaws: A meta-analytic study. *Med Oral Patol Oral Cir Bucal*. 2013 Mar 1;18 (2):e226-32. <http://www.medicinaoral.com/medoralfree01/v18i2/medoralv18i2p226.pdf>

Received: 13/03/2012

Accepted: 25/08/2012

Article Number: 18345 <http://www.medicinaoral.com/>
© Medicina Oral S. L. C.I.F. B 96689336 - pISSN 1698-4447 - eISSN: 1698-6946
eMail: medicina@medicinaoral.com

Indexed in:

Science Citation Index Expanded
Journal Citation Reports
Index Medicus, MEDLINE, PubMed
Scopus, Embase and Emcare
Indice Médico Español

Abstract

Objective: The aim of this study was to evaluate the response of treatment of central giant cell lesion to intralesional corticosteroid injections.

Study Design: Review of articles indexed in PubMed on the topic between the years 1988 and 2011, and development of a descriptive meta-analysis of the results.

Results: Sample of 41 patients primarily treated with intralesional corticosteroid injections was obtained, with a male female ratio of 1:0.95, being 23 aggressive and 18 non-aggressive central giant cell lesions. Triamcinolone acetonide and triamcinolone hexacetonide were the drugs used, and 78.0% cases were considered as good result, 14.6% were considered as moderate response and 7.3% were considered as negative result to treatment. Considering the aggressiveness, 88.9% of non-aggressive lesions presented a good response to treatment, in aggressive central giant cell lesions, 69.6% presented a good response to intralesional corticosteroid injections.

Conclusion: In view of the results analyzed, intralesional corticosteroid injections could be considered as first treatment option for central giant cell lesion.

Key words: *Central giant cell lesion, corticosteroids injections, triamcinolone hexacetonide, triamcinolone acetonide.*

Introduction

Central giant cell lesion (CGCL) is an uncommon type of benign jaw lesion that can be classified as aggressive or non-aggressive (1,2). CGCL is more common in females (3-6), with a female/male ratio of 1.3:1 (7). This lesion can occur at all ages, but most cases are diagnosed in the second and third decades of life (3,7). The mandible is usually more affected than the maxilla, with a mandible/maxilla incidence ratio of 2:1 (7). In radiographic analyses, CGCL may range from small apical lesions to large destructive multilocular radiolucencies involving large areas of the jaws (7).

Chuong et al. (2) first described aggressive and non-aggressive CGCL. The former is characterized by one or more of the following: pain, paresthesia, root resorption, rapid growth, cortical perforation, and a high recurrence rate. Non-aggressive lesions present with slower growth and without cortical perforation or tooth resorption. Aggressive lesions are usually larger and more frequently produce swelling (2). The pathogenesis of CGCL has yet to be elucidated.

Surgery is currently the most common proposed treatment for CGCL in the literature (5,6,8,9); surgical treatment methods range from simple curettage to aggressive en-bloc resection (4-6,10), which can lead to significant facial disfigurement. Intralesional corticosteroid injections are increasingly being used in the clinic, and some reports have shown excellent results. Intralesional corticosteroid injections can avoid extensive mutilating surgeries and successfully manage CGCL; the injections can be used alone or in combination with other treatment options, such as calcitonin or surgery (11). As most of the published articles on intralesional corticosteroid injections are case reports, the literature lacks data about this treatment modality. The aim of this study was to perform a meta-analytic study of intralesional corticosteroid injections for the treatment of CGCL.

Material and Methods

The articles referenced in the bibliography were collected through a search of PubMed, using the following keywords: central giant cell granuloma, central giant cell lesion, and intralesional corticosteroid. Study articles and case report articles were selected. Case reports were included, as only one research article has been published on this topic. The time parameters of the search were set between 1988 and 2011. Additionally,

the report by Terry and Jacoway (12) was included in this review, as this was the first report to document intralesional corticosteroid treatment for CGCL. The data were grouped into tables 1,2,3.

The inclusion criteria were as follows: articles published between the years 1988 and 2011 and cases using intralesional corticosteroid injections as the first choice for treatment of CGCL. The following exclusion criteria were used: studies that included reports on peripheral giant cell lesion and those that used a combination treatment of intralesional corticosteroids with other treatment methods, such as calcitonin or surgery, as the first therapeutic choice. Fourteen articles that met the inclusion criteria were selected. Of the articles selected, one was a research article, and thirteen were case reports.

The data obtained were analyzed for the following variables: number of cases, gender, mean age, location, aggressiveness of CGCL, drug and protocol used, whether any additional procedures were necessary, result of the treatment and follow-up. The aggressiveness of CGCL was defined as proposed by Chuong et al. (2) using data available from the articles. Non-aggressive lesions were those that presented as slow growing and without symptoms, cortical perforation, or root resorption. Aggressive lesions were those associated with pain, rapid growth, cortical perforation, root resorption, or a large size. Treatment results were analyzed as proposed by Nogueira et al. (13) using a four-item scoring system: A score of 1 indicated stabilization or regression in lesion size, as evaluated by the clinical aspect of the lesion and follow-up radiographs. A score of 2 represented the absence of symptoms. A score of 3 indicated an increased radio-opacity in the radiographs, representing peripheral and/or central calcification of the lesion. A score of 4 indicated an increasing difficulty in a solution diffusing into the lesion upon multiple applications. When a case was positive for all four scores, the response was classified as good, two or three scores as moderate, and one or zero scores implied a negative response to the treatment. If a case report did not indicate that the lesion was increasing in size or that the symptoms had not been controlled, these items were considered to have not happened, and the scores 1 and 2 were given to the report.

Results

The search resulted in a total of 14 articles, with 13 case or series reports (12,14-25) and one research article (13). A sample of 41 patients was obtained (20 males

Table 1. Characteristics of the central giant cell lesion studied.

Authors	Number of cases	Gender	Age (years)	Aggressiveness	Site	Drug and protocol	Additional treatment	Results	Follow-up
Terry and Jacoway (12) 1994	4	3 male; 1 female	8, 9, 12 and 19	3 aggressive; 1 non-aggressive	4 in mandible	Triamcinolone acetamide 10 mg/mL and marcaine 0.5%, 1:1; 2 mL per 2 cm of the lesion, once a week for 6 weeks	Second session of treatment and curettage in one case	3 good responses; 1 moderate response	16-36 months
Kermer et al. (14) 1994	1	1 Male	40	Non-aggressive	Mandible	Triamcinolone acetamide and lidocaine 1:1; once a week for 6 weeks	New CGCL or recurrence after 14 months, treated with the same protocol	Good response	3 years
Rajeevan and Soumithran (15) 1998	1	1 Female	17	Aggressive	Mandible	Triamcinolone acetamide 10 mg/mL and lidocaine 1:1; once a week for 6 weeks	None	Good response	10 months
Khafif et al. (16) 2001	1	Female	36	Aggressive	Maxilla	Triamcinolone acetamide 40 mg/mL and marcaine 0.5%, 1:1; once a week for 6 weeks	None	Good response	24 months
Kurtz et al. (17) 2001	1	Female	10	Aggressive	Mandible	Triamcinolone acetamide 10 mg/mL and Marcaine 0.5%, 1:1; 2 mL per 2 cm of the lesion, once a week for 6 weeks.	Second session	Good response after two treatment session	19 months
Adornato and Paticoff (18) 2001	1	Female	10	Non-aggressive	Mandible	Triamcinolone acetamide 10 mg/mL and marcaine 0.5%, 1:1; 2 mL per 2 cm of the lesion, once a week for 6 weeks	None	Good response	7 months
Carlos and Sedano (19) 2002	4	3 male; 1 Female	2, 6, 31 and 34	2 aggressive; 2 non-aggressive	2 maxilla; 2 mandible	Triamcinolone acetamide 10 mg/mL and marcaine 0.5%, 1:1; 6 mL, variable number of injections (3-20).	Curettage in one case	Good response in three cases; moderate response in one case	2-7 years
Abdo et al. (20) 2005	1	Female	14	Non-aggressive	Mandible	Weekly; 3 weeks 5-ml injection of Triamcinolone acetamide 10 mg/ml, and lidocaine solution 2% with epinephrine 1:200,000. 1:1; once a week for 6 weeks	None	Good response	18 months
Sezer et al. (21) 2005	1	Male	11	Aggressive	Mandible	Triamcinolone acetamide 10 mg/ml, and lidocaine solution 2% with epinephrine 1:200,000. 1:1; once a week for 6 weeks	None	Good response	3 Years

Table 1. Continue.

Mohanty and Jhamb (22) 2009	2	1 male; 1 Female	10 and 20	2 Aggressive	Mandible	Triamcinolone 10 mg/ml and lidocaine, 1:1, weekly for 5 weeks in one case; 9 injections in the other case	None	Good response	13-18 months				
Wendt et al. (23) 2009	1	Female	8	aggressive	Maxilla	Triamcinolone acetamide 10 mg/ml and 0.5% bupivacaine, 1:1; weekly, 11 weeks	None	Good response	6 Years				
Nogueira et al. (13) 2010	21	11 male; 10 female	Average: 16	10 aggressive; 11 non-aggressive	8 in maxilla; 13 in mandible	Triamcinolone hexacetamide 20 mg/ml, and 2% lidocaine/epinephrine 1:200,000, 1:1; biweekly; 6 injections	Osteoplasty in 8 cases; curettage in 3 cases; surgical resection in 2 cases	15 good responses; 4 moderate responses; 2 negative responses	4-7 years				
Ferretti et al. (24) 2011	1	Female	16	Non-aggressive	Mandible	Triamcinolone acetamide 40 mg/ml and 0.5% bupivacaine; biweekly; 3 injections	None	Good response	4 years				
Shirani et al. (25) 2011	1	Female	13	Aggressive	Mandible	40 mg/ml Triamcinolone acetamide mixed with 5 cc Lidocaine 1% with 1/200,000 epinephrine; once a week for 6 weeks	Surgical resection	Negative	2 years				
Total	41	20 male; 21 female	Average: 15.9	23 aggressive; 18 non-aggressive	12 in maxilla; 29 in mandible		Osteoplasty in 9 cases; curettage in 5 cases; surgical resections in 3 cases; new injections in 3 cases	32 good responses; 6 moderate responses; 3 negative responses	7 months-7 years				

and 21 females), with a female/male ratio of 1:0.95. The average age was 15.9 years; for aggressive lesions, the average age was 13.9 years, and for non-aggressive lesions, the average age was 18.3 years old. Twelve lesions were in the maxilla and 29 in the mandible, with a maxilla:mandible ratio of 1:2.4. According to the criteria defined by Chuong et al.(2) 23 lesions were classified as aggressive CGCL and 18 lesions as non-aggressive. Triamcinolone acetamide (10 mg/ml or 40 mg/ml) and triamcinolone hexacetamide (20 mg/ml) were the adopted drugs, but one case report did not indicate the type of corticosteroid used (20). The drugs were always diluted with an anesthetic solution of marcaine, lidocaine or bupivacaine in equal parts. A 2-ml dose of this solution for every 2 cm of radiolucency was the most cited dosage, but a dosage of 1 ml for every 1 cm³ was also reported. The most frequently used protocol was a regimen of 6 weekly injections, but a biweekly protocol was also described, and in one patient, 20 injections were given. According to the criteria previously defined by Nogueira et al. (13) in 2010, 32 (78.0%) cases were considered good results, 6 (14.6%) were considered moderate responses and 3 (7.3%) showed negative results to treatment (Data shown in tables 1,2,3).

Discussion

Surgery is the most common treatment of choice for CGCLs, and the extent of surgery ranges from curettage with or without adjuvant therapy, such as cryosurgery, peripheral ostectomy and carnoy solution, to aggressive en bloc resection (6,10), resulting in varying degrees of deformity (4). Conservative curettage and enucleation can lead to high recurrence rates, whereas en bloc resection may sacrifice adjacent structures, thus requiring surgical reconstructive procedures to recover satisfactory functional and esthetic results (2). As some cases of CGCL affect children and young adults, defects in the developing dentition and jaws are of great concern (13). Reconstructive and rehabilitative procedures, which usually require multiple hospitalizations, and the use of dental implants and prosthetic devices are very costly. Several non-surgical methods have been proposed to treat CGCLs, including radiotherapy (26), systemic calcitonin (27-29), intralesional injection with corticosteroids (12-25) and systemic α interferon (30,31). Among these non-surgical treatment methods, intralesional corticosteroid injection has shown promising results and can lead to a complete resolution of the lesion or a significant reduction in size, allowing a more conservative surgery (12,13). Body et al. (32) first reported the use of corticosteroids in the treatment of CGCL. In their report, the authors used dexamethasone to treat an aggressive recurrent case of CGCL, achieving a significant reduction in lesion size; however, complications resulted because a systemic

Table 2. Characteristics of aggressive central giant cell lesion studied.

Authors	Number of cases	Gender	Age (years)	Site	Additional treatment	Results
Terry and Jacoway (12) 1994	3	2 male; 1 female	8, 9 and 19	3 in mandible	None	3 good response
Rajeevan and Soumithran, (15) 1998	1	1 Female	17	Mandible	None	Good response
Khafif et al. (16) 2001	1	Female	36	Maxilla	None	Good response
Kurtz et al. (17) 2001	1	Female	10	Mandible	Second session	Good response after two treatment session
Carlos and Sedano (19) 2002	2	1 male; 1 Female	2 and 6	1 maxilla; 1 mandible	Curettage in one case	Good response in one cases; moderate response in one case
Sezer et al. (21) 2005	1	Male	11	Mandible	None	Good response
Mohanty and Jhamb (22) 2009	2	1 male; 1 Female	10 and 20	Mandible	None	Good response
Wendt et al. (23) 2009	1	Female	8	Maxilla	None	Good response
Nogueira et al. (13) 2010	10	5 male; 5 female	Average: 15.2	5 in Maxilla; 5 in Mandible	4 osteoplasty; 2 surgical resection; 1 curettage	5 good response; 3 moderate response; 2 negative response
Shirani et al. (25) 2011	1	Female	13	Mandible	Surgical resection	Negative
Total	23 cases	10 male; 13 female	average: 13.9	9 in maxilla; 14 in mandible	4 cases underwent osteoplasty; 2 cases underwent curettage; 3 cases underwent surgical resection	16 good response; 4 moderate response; 3 negative response

Table 3. Characteristics of non-aggressive central giant cell lesions studied.

Authors	Number of cases	Gender	Age (years)	Site	Additional treatment	Results
Terry and Jacoway (12) 1994	1	1 male	12	1 in mandible	Second session of treatment and curettage	Moderate response
Kermer et al. (14) 1994	1	1 Male	40	Mandible	New CGCL or recidive? after 14months treated with same protocol	Good response
Adornato and Paticoff (18) 2001	1	Female	10	Mandible	None	Good response
Carlos and Sedano (19) 2002	2	2 male	31 and 34	1 maxilla; 1 mandible	None	Good response
Abdo et al. (20) 2005	1	Female	14	Mandible	None	Good response
Nogueira et al. (13) 2010	11	6 male; 5 female	Average 15.8	3 in Maxilla; 8 in Mandible	osteoplasty in 4 cases; curettage in 2 cases	10 good response; 1 moderate response
Ferretti et al. (24) 2011	1	Female	16	Mandible	None	Good response
Total	18 cases	10 male; 8 female	Average 18.3	4 in maxilla; 14 in mandible	Osteoplasty in 4 cases; curettage in 3 cases; new injections in 2 cases	16 good response; 2 moderate response

corticosteroid was used. Consequently, the dose was reduced, and the lesion re-grew. Terry and Jacoway (12) first reported the use of intralesional corticosteroid injections in the treatment of CGCL. Intralesional injections are preferable because they can achieve an elevated and localized concentration in the tissue (12). In addition, the complications associated with systemic corticosteroid administration usually do not appear, as none of the articles included in this meta-analysis reported complications related to the corticosteroids.

The drugs used for treatment included triamcinolone acetonide (10 mg/ml or 40 mg/ml) and triamcinolone hexacetonide (20 mg/ml); both showed similar efficacies. The injections were administered weekly or bi-weekly. Nogueira et al. (13) proposed that using a more concentrated drug, such as triamcinolone hexacetonide 20 mg/ml, may allow for a biweekly interval, facilitating a greater reactivity and radiographic perception of the increasing radio-opacity. Six injections were the most common treatment regimen, but cases with up to 20 intralesional injections were reported (19). Triamcinolone is always diluted in equal parts with an anesthetic solution, and a dose of 2 ml for each 2 cm of radiolucency or 1 ml for each cm³ of lesion is injected.

Although CGCL is more frequently diagnosed in female patients (3-6), data from the present study found an almost equal frequency; 51.22% of cases occurred in female and 48.8% in male patients. Similar to other reports (3-7), in young patients, CGCL occurred more often in the mandible. In imaging studies, CGCL appeared as a uniloculated or multiloculated expansive osteolytic lesion and was frequently associated with tooth displacement (12,13,17,23-25). Root resorption was seen in some cases (12,13), as was cortical perforation (13,21,22,25); the later is better seen in computerized tomography (CT) scans. Microscopically, CGCL appeared as multinucleated giant cells in a cellular background composed of mononucleated stromal cells with an ovoid or spindle-shape, with hemorrhage foci (13,19,20,23,25). Two reports performed microscopic exams after the treatment and revealed a reduced number or, in some cases, the complete absence of CGCL, surrounded by markedly fibrocollagenous stroma and showing reduced vascularization (13,19).

According to the criteria defined by Chuong et al. (2), 23 lesions were classified as aggressive and 18 as non-aggressive. Aggressive lesions tended to affect younger patients compared with non-aggressive lesions, with an average age of 13.9 years for aggressive CGCL and 18.3 years for non-aggressive lesions. According to the criteria previously defined by Nogueira et al. (13), a good response to intralesional corticosteroid injections was seen in 78.0% of CGCL patients, and only 7.3% showed a negative result. Considering only aggressive lesions, 69.6% of cases presented a good response to treatment,

and 13.0% showed a negative result. In non-aggressive CGCL, an even better result was found, as 88.9% these patients presented a good response to treatment, and none presented a negative result. These are excellent results, and it must not be forgotten that in cases showing a negative response to treatment, others treatment options are available. In fact, surgical resection was performed in all cases with a negative result (13,25). It has been previously described that CGCL contain glucocorticoid receptors in multinucleated giant cells and mononucleated stromal cells (33); this may be the reason why CGCL regresses upon corticosteroid treatment. More recently, Nogueira et al. (34) showed that in cases with good response to intralesional corticosteroid injections, a higher expression of glucocorticoid receptors were seen in multinucleated giant cells. Although in most cases intralesional injections were used alone, additional treatment may be necessary. Esthetic osteoplasty was performed in 9 cases in the present study. Curettage was performed in 5 cases, mainly to remove residual lesion tissue.

Although the authors are aware of the limitations of meta-analytic studies, these data suggest that a non-surgical, intralesional corticosteroid injection approach for CGCL could be provided as a first-line treatment option. This treatment can lead to complete resolution of the lesion, or it may lead to a significant reduction in lesion size, allowing for a more conservative surgical approach. Further controlled prospective studies should be incentivized to corroborate these results.

References

1. de Lange J, van den Akker HP, van den Berg H, Richel DJ, Gortzak RA. Limited regression of central giant cell granuloma by interferon alpha after failed calcitonin therapy: a report of 2 cases. *Int J Oral Maxillofac Surg.* 2006;35:865-9.
2. Chuong R, Kaban LB, Kozakewich H, Perez-Atayde A. Central giant cell lesions of the jaws: a clinicopathologic study. *J Oral Maxillofac Surg.* 1986;44:708-13.
3. Kruse-Lösler B, Diallo R, Gaertner C, Mischke KL, Joos U, Kleinheinz J. Central giant cell granuloma of the jaws: a clinical, radiologic, and histopathologic study of 26 cases. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006;101:346-54.
4. Rawashdeh MA, Bataineh AB, Al-Khateeb T. Long-term clinical and radiological outcomes of surgical management of central giant cell granuloma of the maxilla. *Int J Oral Maxillofac Surg.* 2006;35:60-6.
5. Motamedi MH, Talesh KT, Jafari SM, Khalifeh S. Peripheral and central giant cell granulomas of the jaws: a retrospective study and surgical management. *Gen Dent.* 2010;58:e246-51.
6. Tosco P, Tanteri G, Iaquina C, Fasolis M, Rocca F, Berrone S, et al. Surgical treatment and reconstruction for central giant cell granuloma of the jaws: a review of 18 cases. *J Craniomaxillofac Surg.* 2009;37:380-7.
7. de Lange J, van den Akker HP. Clinical and radiological features of central giant-cell lesions of the jaw. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005;99:464-70.
8. Roberts J, Shores C, Rose AS. Surgical treatment is warranted in aggressive central giant cell granuloma: a report of 2 cases. *Ear Nose Throat J.* 2009;88:E8-E13.
9. Nicolai G, Lorè B, Mariani G, Bollero P, De Marinis L, Cala-

- brese L. Central giant cell granuloma of the jaws. *J Craniofac Surg.* 2010;21:383-6.
10. Bataineh AB, Al-Khateeb T, Rawashdeh MA. The surgical treatment of central giant cell granuloma of the mandible. *J Oral Maxillofac Surg.* 2002; 60:756-61.
 11. Rachmiel A, Emodi O, Sabo E, Aizenbud D, Peled M. Combined treatment of aggressive central giant cell granuloma in the lower jaw. *J Craniomaxillofac Surg.* 2012;40:292-7.
 12. Terry BC, Jacoway JR. Management of central giant cell lesions – an alternative to surgical therapy. *Oral Maxillofac Surg Clin North Am.* 1994;6:579-600.
 13. Nogueira RL, Teixeira RC, Cavalcante RB, Ribeiro RA, Rabenhorst SH. Intralesional injection of triamcinolone hexacetonide as an alternative treatment for central giant-cell granuloma in 21 cases. *Int J Oral Maxillofac Surg.* 2010;39:1204–10.
 14. Kermer C, Millesi W, Watzke IM. Local injection of corticosteroids for central giant cell granuloma. A case report. *Int J Oral Maxillofac Surg.* 1994;23:366-8.
 15. Rajeevan NS, Soumithran CS. Intralesional corticosteroid injection for central giant cell granuloma. A case report. *Int J Oral Maxillofac Surg.* 1998;27:303-4.
 16. Khafif A, Krempl G, Medina JE. Treatment of giant cell granuloma of the maxilla with intralesional injection of steroids. *Head Neck.* 2000;22:822-5.
 17. Kurtz M, Mesa M, Alberto P. Treatment of a central giant cell lesion of the mandible with intralesional glucocorticosteroids. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2001;91:636-7.
 18. Adornato MC, Patcoff KA. Intralesional corticosteroid injection for treatment of central giant-cell granuloma. *J Am Dent Assoc.* 2001;132:186-90.
 19. Carlos R, Sedano HO. Intralesional corticosteroids as an alternative treatment for central giant cell granuloma. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2002;93:161-6.
 20. Abdo EN, Alves LC, Rodrigues AS, Mesquita RA, Gomez RS. Treatment of a central giant cell granuloma with intralesional corticosteroid. *Br J Oral Maxillofac Surg.* 2005;43:74-6.
 21. Sezer B, Koyuncu B, Gomel M, Günbay T. Intralesional corticosteroid injection for central giant cell granuloma: a case report and review of the literature. *Turk J Pediatr.* 2005;47:75-81.
 22. Mohanty S, Jhamb A. Central giant cell lesion of mandible managed by intralesional triamcinolone injections. A report of two cases and literature review. *Med Oral Patol Oral Cir Bucal.* 2009;14:e98-102.
 23. Wendt FP, Torriani MA, Gomes AP, de Araujo LM, Torriani DD. Intralesional corticosteroid injection for central giant cell granuloma: an alternative treatment for children. *J Dent Child (Chic).* 2009;76:229-32.
 24. Ferretti C, Muthray E. Management of central giant cell granuloma of mandible using intralesional corticosteroids: case report and review of literature. *J Oral Maxillofac Surg.* 2011;69:2824-9.
 25. Shirani G, Abbasi AJ, Mohebbi SZ, Shirinbak I. Management of a locally invasive central giant cell granuloma (CGCG) of mandible: report of an extraordinary large case. *J Craniomaxillofac Surg.* 2011;39:530-3.
 26. Eisenbud L, Stern M, Rothberg M, Sachs SA. Central giant cell granuloma of the jaws: experiences in the management of thirty-seven cases. *J Oral Maxillofac Surg.* 1988;46:376–84.
 27. de Lange J, Rosenberg AJ, Van Den Akker HP, Koole R, Wirds JJ, Van Den Berg H. Treatment of central giant cell granuloma of the jaw with calcitonin. *Int J Oral Maxillofac Surg.* 1999;28:372- 6.
 28. Borges HO, Machado RA, Vidor MM, Beltrao RG, Heitz C, Filho MS. Calcitonin: a non-invasive giant cells therapy. *Int J Pediatr Otorhinolaryngol.* 2008;72:959-63.
 29. Sadiq Z, Goodger NM. Calcitonin-induced osteoplastic reaction in the mandible. *Br J Oral Maxillofac Surg.* 2011;49:578-9.
 30. Baker SB, Parikh PM, Rhodes DN, Abu-Ghosh A, Shad AT. Aggressive central giant cell lesion of the maxilla: surgical management and the use of adjuvant interferon alfa-2a. *Plast Reconstr Surg.* 2008;122:77e-9e.
 31. de Lange J, van Rijn RR, van den Berg H, van den Akker HP. Regression of central giant cell granuloma by a combination of imatinib and interferon: a case report. *Br J Oral Maxillofac Surg.* 2009;47:59-61.
 32. Body JJ, Jortay AM, de Jager R, Ardichvili D. Treatment with steroids of a giant cell granuloma of the maxilla. *J Surg Oncol.* 1981;16:7-13.
 33. Vered M, Buchner A, Dayan D. Immunohistochemical expression of glucocorticoid and calcitonin receptors as a tool for selecting therapeutic approach in central giant cell granuloma of the jawbones. *Int J Oral Maxillofac Surg.* 2006;35:756–60.
 34. Nogueira RL, Faria MH, Osterne RL, Cavalcante RB, Ribeiro RA, Rabenhorst SH. Glucocorticoid and calcitonin receptor expression in central giant cell lesions: implications for therapy. *Int J Oral Maxillofac Surg.* 2012;41:994-1000. 2012 Feb 23 [Epub ahead of print].