

Spacetime correlators of perturbations in slow-roll de Sitter inflation

Adrian del Rio and Jose Navarro-Salas

Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC. Facultad de Física, Universidad de Valencia, Burjassot-46100, Valencia, Spain

(Dated: March 13, 2014)

Two-point correlators and self-correlators of primordial perturbations in quasi-de Sitter spacetime backgrounds are considered. For large separations two-point correlators exhibit nearly scale invariance, while for short distances self-correlators need standard renormalization. We study the deformation of two-point correlators to smoothly match the self-correlators at coincidence. The corresponding angular power spectrum is evaluated in the Sachs-Wolfe regime of low multipoles. Scale invariance is maintained, but the amplitude of C_ℓ could change in a nontrivial way.

Key Words: quantum field theory in curved spacetime, correlators, asymptotic behaviour, renormalization, cosmology.

PACS numbers: 04.62.+ , 98.80.Cq

I. INTRODUCTION

In recent years de Sitter space has received considerable attention. Astronomical observations [1] are pointing out that our universe now has a very tiny positive cosmological constant, which, however, embodies around three quarters of the energy of the observable universe. Moreover, according to inflationary cosmology the very early universe underwent a period of very rapid expansion powered by a large effective cosmological constant. The discovery of the anisotropies in the cosmic microwave background (CMB) [2] constitutes a very sensitive probe of the primordial density perturbations and its quantum mechanical origin [3]. The comparison of observations [4] with theoretical predictions is currently a sharp tool to test inflation and the theory of quantized fields in curved backgrounds [5, 6]. Therefore, a precise understanding of the quantum properties of fields in de Sitter space is fundamental for both the very early and the late-time universe.

In this article we will focus on the quantum treatment of primordial perturbations, which will be regarded as quantum fields ϕ living in a curved (quasi-de Sitter) spacetime. The two-point correlation function $\langle \phi(t, \vec{x}) \phi(t, \vec{x}') \rangle$ exhibits scale invariance at large separations $|\vec{x} - \vec{x}'|H \gg 1$ or, equivalently, at late-time $Ht \gg 1$. On the other hand, the amplitude of the perturbation at a given spacetime point can be quantified by the self-correlator $\langle \phi^2(t, \vec{x}) \rangle$, which requires one to get rid off the corresponding ultraviolet (UV) divergences and renormalize the expectation value.

In the first part of this work we will compute and analyze the above quantities in a slow-roll de Sitter background. We will also project the large-distance behavior of the correlator of scalar perturbations on a sphere of fixed radius. This sphere is linked by time evolution to the last scattering surface, where the cosmic microwave background and its anisotropies are formed, and the angular power spectrum can easily be obtained within this spacetime picture. In the second part we will redo the

calculation for the angular power spectrum by using a deformed two-point correlator. The new correlator is defined in such a way that it matches the self-correlator at coincidence. To this end we naturally use methods of renormalization in homogeneous backgrounds [5, 6]. The revised angular power spectrum maintains the nearly scale invariance, but the amplitude of the multipole coefficients C_ℓ may be altered in a non-trivial way. We will focus on the lower multipoles, where the Sachs-Wolfe effect dominates.

II. SPACETIME CORRELATORS IN SLOW-ROLL INFLATION

A. Correlator of tensorial perturbations in a slow-roll scenario

Tensorial perturbations can be described by two independent, massless scalar fields propagating in the unperturbed quasi de Sitter background. These two scalar fields represent the two independent polarization components of the fluctuation tensorial modes \mathcal{D}_{ij} in the inflationary universe: $ds^2 = dt^2 - a^2(t)(\delta_{ij} + \mathcal{D}_{ij})dx^i dx^j$. Expanding the fluctuating fields \mathcal{D}_{ij} in plane wave modes $\mathcal{D}_k(t)e_{ij}e^{i\vec{k}\vec{x}}$, where e_{ij} is a constant polarization tensor obeying the conditions $e_{ij} = e_{ji}$, $e_{ii} = 0$ and $k_i e_{ij} = 0$, one obtains the equation $\ddot{\mathcal{D}}_k + 3H\dot{\mathcal{D}}_k + \frac{k^2}{a^2}\mathcal{D}_k = 0$, with $k \equiv |\vec{k}|$ and $H = \dot{a}/a$. The conditions for the polarization tensor imply that the perturbation field \mathcal{D}_{ij} can be decomposed into two polarization states described by a couple of massless scalar fields $\mathcal{D}_{ij} = \mathcal{D}_+ e_{ij}^+ + \mathcal{D}_\times e_{ij}^\times$, where $e_{ij}^s e_{ij}^{s'} = 2\delta_{ss'}$ ($s = +, \times$ stands for the two independent polarizations), both obeying the above wave equation (see, for instance, [7]). For simplicity we omit the subindex $+$ or \times .

In the slow-roll approximation one assumes that the Hubble parameter $H(t)$ changes very gradually, and the change is parametrized by a slow-roll parameter $\epsilon \equiv$

$-\dot{H}/H^2 \ll 1$. Within this approximation it is possible to solve the wave equation in a closed form in terms of the conformal time $\eta \equiv \int dt/a(t)$. Taking into account that $(1-\epsilon)\eta = -\frac{1}{a\dot{H}}$, the wave equation for \mathcal{D}_k turns out to be of the form

$$\frac{d^2\mathcal{D}_k}{d\eta^2} - \frac{2}{\eta(1-\epsilon)} \frac{d\mathcal{D}_k}{d\eta} + k^2\mathcal{D}_k = 0. \quad (1)$$

Treating now the parameter ϵ as a constant, one can univocally solve the above equation with the requirement of recovering, for $\epsilon \rightarrow 0$, the Bunch-Davies vacuum [8]. The properly normalized solutions for the modes are

$$\mathcal{D}_k(t) = \frac{\sqrt{16\pi G}}{\sqrt{2(2\pi)^3 a^3}} (-\eta a \pi/2)^{1/2} H_\nu^{(1)}(-k\eta), \quad (2)$$

where G is the Newton constant and the index of the Hankel function is exactly $\nu = \frac{3}{2} + \frac{\epsilon}{1-\epsilon}$. Having the explicit form of the modes, we can now compute the two-point function. At equal times $t = t'$ we find

$$\begin{aligned} \langle \mathcal{D}(t, \vec{x}) \mathcal{D}(t, \vec{x}') \rangle &= \frac{G}{\pi a^2 \eta^2} \Gamma\left(\frac{3}{2} + \nu\right) \Gamma\left(\frac{3}{2} - \nu\right) \\ &\times {}_2F_1\left(\frac{3}{2} + \nu, \frac{3}{2} - \nu; 2; 1 - \frac{(\Delta x)^2}{4\eta^2}\right) \end{aligned} \quad (3)$$

where $\Delta x \equiv |\vec{x} - \vec{x}'|$. For $\nu = 3/2$ ($\epsilon = 0$) we have the unavoidable infrared divergence of the Bunch-Davies vacuum [9].

For large separations $a\Delta x \gg H^{-1}$ one obtains

$$\langle \mathcal{D}(t, \vec{x}) \mathcal{D}(t, \vec{x}') \rangle \sim \frac{4G\Gamma(3/2 - \nu)\Gamma(\nu)}{\pi^{3/2} a^2 \eta^2} \left(\frac{\Delta x}{-\eta}\right)^{2(\nu-3/2)}. \quad (4)$$

One can immediately observe that the amplitude above is nearly scale invariant. Moreover, the term $(-\eta)^{1-2\nu}/a^2$ is time independent, which allows us to evaluate it at the most convenient time. In fact, the correlator can be rewritten as

$$\langle \mathcal{D}(t, \vec{x}) \mathcal{D}(t, \vec{x}') \rangle \sim -\frac{16\pi G}{2\epsilon} \left(\frac{H(t_{\Delta x})}{2\pi}\right)^2, \quad (5)$$

where the time $t_{\Delta x}$ is defined as $a(t_{\Delta x})\Delta x = H^{-1}(t_{\Delta x})$. Note that there is an implicit Δx dependence on $H(t_{\Delta x})$, given by the one in (4).

B. Correlator of scalar perturbations in a slow-roll scenario

Scalar perturbations can be studied through the gauge-invariant field \mathcal{R} (the comoving curvature perturbation; see, for instance, [7]). For single-field inflation, the modes of the scalar perturbation are given by

$$\mathcal{R}_k(t) = (-\pi\eta/4(2\pi)^3 z^2)^{1/2} H_\nu^{(1)}(-\eta k), \quad (6)$$

where now $\nu = 3/2 + (2\epsilon + \delta)/(1-\epsilon)$ and $\delta \equiv \ddot{H}/2H\dot{H}$ is a second slow-roll parameter. Moreover, $z \equiv a\dot{\phi}_0/H$, where $\phi_0(t)$ is the homogeneous part of the inflaton field. These modes determine the vacuum state of scalar perturbations. Such a state can also be regarded as the natural extension of the Bunch-Davies vacuum of de Sitter space. The corresponding two-point function $\langle \mathcal{R}(t, \vec{x}), \mathcal{R}(t, \vec{x}') \rangle$ is given by

$$\begin{aligned} \langle \mathcal{R}(t, \vec{x}) \mathcal{R}(t, \vec{x}') \rangle &= \frac{1}{16\pi^2 z^2 \eta^2} \Gamma\left(\frac{3}{2} + \nu\right) \Gamma\left(\frac{3}{2} - \nu\right) \\ &\times {}_2F_1\left(\frac{3}{2} + \nu, \frac{3}{2} - \nu; 2; 1 - \frac{(\Delta x)^2}{4\eta^2}\right). \end{aligned} \quad (7)$$

For separations larger than the Hubble radius $a|\vec{x} - \vec{x}'| \gg H^{-1}$ we get

$$\langle \mathcal{R}(t, \vec{x}) \mathcal{R}(t, \vec{x}') \rangle \sim \frac{\Gamma(\frac{3}{2} - \nu)\Gamma(\nu)}{4\pi^2 z^2 \eta^2} \frac{1}{\sqrt{\pi}} \left(\frac{\Delta x}{-\eta}\right)^{2(\nu-3/2)}. \quad (8)$$

This expression can be rewritten, assuming $\nu - 3/2 \equiv (1-n)/2 \approx 0$ (n is the scalar spectral index), as

$$\langle \mathcal{R}(t, \vec{x}) \mathcal{R}(t, \vec{x}') \rangle \sim -\frac{4\pi G}{(1-n)\epsilon} \left(\frac{H(t_{\Delta x})}{2\pi}\right)^2. \quad (9)$$

C. Angular power spectrum

Restricting the two-point function of scalar perturbations to points such that $|\vec{x}| = |\vec{x}'|$, we can further obtain $\Delta x^{1-n} = 2^{\frac{1-n}{2}} |\vec{x}|^{(1-n)} (1 - \cos\theta)^{(1-n)/2}$ where θ is the angle formed by $\vec{n} = \vec{x}/|\vec{x}|$ and $\vec{n}' = \vec{x}'/|\vec{x}'|$. Then, taking $|\vec{x}| = r_L$, where r_L is the comoving radial coordinate of the last scattering surface

$$r_L = H(t_0)^{-1} a(t_0)^{-1} \int_{\frac{1}{1+z_L}}^1 \frac{dx}{\sqrt{\Omega_\Lambda x^4 + \Omega_M x + \Omega_R}}, \quad (10)$$

with the standard cosmological values for z_L , Ω_Λ , Ω_R , and Ω_M [4], the correlator of scalar perturbations for large separations (8) shows exactly

$$\begin{aligned} \langle \mathcal{R}(t, \vec{x}) \mathcal{R}(t, \vec{x}') \rangle &\sim \frac{4\pi G}{\epsilon} \frac{H^2(1-\epsilon)^2}{16\pi^2} \frac{4\Gamma(2 - \frac{n}{2})}{\sqrt{\pi}} \Gamma\left(\frac{n-1}{2}\right) \\ &\times 2^{\frac{1-n}{2}} \bar{r}_L^{1-n} (1 - \cos\theta)^{\frac{1-n}{2}} \end{aligned} \quad (11)$$

where we have defined the dimensionless quantity $\bar{r}_L(t) \equiv H(1-\epsilon)ar_L$. This two-point function can be related to the temperature fluctuations in the CMB,

$$\langle \Delta T(\vec{n}) \Delta T(\vec{n}') \rangle = \sum_\ell C_\ell \frac{2\ell+1}{4\pi} P_\ell(\cos\theta), \quad (12)$$

where P_ℓ is the Legendre polynomial, via the Sachs-Wolfe effect (see, e.g., [7]) $\langle \Delta T(\vec{n}) \Delta T(\vec{n}') \rangle_{SW} = \frac{T_0^2}{25} \langle \mathcal{R}(r_L \vec{n}) \mathcal{R}(r_L \vec{n}') \rangle$. The coefficients C_ℓ are obtained by Legendre transformation of (12),

$$C_\ell^{SW} = \frac{2\pi T_0^2}{25} \int_{-1}^1 d\cos\theta P_\ell(\cos\theta) \langle \mathcal{R}(r_L \vec{n}) \mathcal{R}(r_L \vec{n}') \rangle \quad (13)$$

Therefore, the low multipole coefficients, dominated by the Sachs-Wolfe effect, are proportional to the integral

$$C_\ell^{SW} \propto \int_{-1}^1 dy (1-y)^{\frac{1-n}{2}} P_\ell(y), \quad (14)$$

with $y \equiv \cos \theta$. The above integral can be computed analytically [10, 11], and we finally find

$$C_\ell^{SW} = \frac{8\pi T_0^2}{25} \frac{4\pi G}{\epsilon} \frac{H^2(1-\epsilon)^2}{16\pi^2} \frac{\Gamma(3-n)\Gamma(\ell + \frac{n-1}{2})}{\Gamma(\ell + 2 - \frac{n-1}{2})} \bar{r}_L^{1-n}, \quad (15)$$

in exact agreement with the result obtained with the momentum-space power spectrum $P_s(k) = |N|^2 k^{n-1}$ [7],

$$C_{\ell, SW} = \frac{16\pi^2 T_0^2}{25} \int_0^\infty \frac{dk}{k} P_s(k) j_\ell^2(kr_L), \quad (16)$$

where the amplitude $|N|^2$ is given by $|N|^2 = \frac{4\pi G}{\epsilon} \frac{H^2(1-\epsilon)^2}{16\pi^2} \frac{2^{3-n}}{\pi^2} \Gamma(2 - \frac{n}{2})^2 \left(\frac{\bar{r}_L}{r_L}\right)^{1-n} \sim \frac{8\pi G H^2}{32\pi^3 \epsilon}$. For completeness, taking approximately $n \approx 1$ in (15) and using the standard assumption $\bar{r}_L^{(1-n)} \approx O(1)$ [7], the estimated order of magnitude for the amplitude of C_ℓ^{SW}

is

$$\ell(\ell+1)C_\ell^{SW} \approx \frac{2GH^2T_0^2}{25\epsilon} \bar{r}_L^{1-n} \sim \frac{2GH^2T_0^2}{25\epsilon}. \quad (17)$$

We will go back to this point at the end of Sec. IV.

We note that if the coefficients C_ℓ^{SW} in (13) were actually evaluated using the exact expression (7) for the two-point function $\langle \mathcal{R}(t, \vec{x}) \mathcal{R}(t, \vec{x}') \rangle$, the integral (13) would have been divergent [due to the UV divergences of (7) as points \vec{x} and \vec{x}' merge]. The use of the large distance behavior (8 and 11) everywhere in the integral (13) bypasses the UV divergences and makes the integral convergent. We will see in Sec. IV how the use of a renormalized form of the two-point correlator $\langle \mathcal{R}(t, \vec{x}) \mathcal{R}(t, \vec{x}') \rangle$ does the same job, but with a slightly different final result for the integral. In a certain limit both results eventually agree, but in general we find a difference that could be potentially probed by observations.

III. AMPLITUDE OF QUANTUM PERTURBATIONS

We shall use ϕ to denote both scalar and tensorial fluctuations, and ν to represent the corresponding Hankel index. The two-point function $\langle \phi(t, \vec{x}), \phi(t, \vec{x}') \rangle$ can be expanded at short distances as

$$\langle \phi(t, \vec{x}) \phi(t, \vec{x}') \rangle = \frac{H^2(1-\epsilon)^2}{16\pi^2} \left\{ \frac{4}{\Delta \bar{x}^2} + \left(\frac{1}{4} - \nu^2\right) \left(-1 + 2\gamma + \psi(3/2 - \nu) + \psi(3/2 + \nu) + \log \frac{\Delta \bar{x}^2}{4}\right) + O(\Delta \bar{x}^2) \right\}, \quad (18)$$

where we have introduced the dimensionless quantity $\Delta \bar{x} \equiv H(1-\epsilon) a \Delta x$. An additional prefactor, $4\pi G/\epsilon$ or $16\pi G$, needs to be included in considering scalar or tensorial perturbations, respectively. As expected, one encounters the typical quadratic and logarithmic UV divergences of a quantum field in a curved background. Since we are now interested in evaluating the mean square fluctuation $\langle \phi^2(t, \vec{x}) \rangle$ at a given spacetime point, we have to remove these divergences by standard renormalization in curved spacetime [5, 6]. Different methods can be used to this end. A preferred method for our purposes is the point-splitting version of the adiabatic regularization scheme [5, 12, 13]. In short, the two-point function $\langle \phi(t, \vec{x}) \phi(t, \vec{x}') \rangle$ at coincidence can be naturally renormalized by subtracting the second-order adiabatic terms $G_{Ad}^{(2)}((t, \vec{x}), (t, \vec{x}'))$

$$\langle \phi(t, \vec{x}) \phi(t, \vec{x}') \rangle_{ren} \equiv \langle \phi(t, \vec{x}) \phi(t, \vec{x}') \rangle - G_{Ad}^{(2)}((t, \vec{x}), (t, \vec{x}')) \quad (19)$$

and taking the limit $\vec{x}' \rightarrow \vec{x}$. The method determines univocally the subtraction terms, which are found to be

$$G_{Ad}^{(2)}((t, \vec{x}), (t, \vec{x}')) = \frac{H^2(1-\epsilon)^2}{16\pi^2} \left\{ \frac{4}{\Delta \bar{x}^2} + \left(\frac{1}{4} - \nu^2\right) \log \frac{\Delta \bar{x}^2}{4} + \frac{2-\epsilon}{3(1-\epsilon)^2} + \left(\frac{1}{4} - \nu^2\right) \left(2\gamma + \log \frac{\mu^2}{H^2(1-\epsilon)^2}\right) \right\}, \quad (20)$$

where μ is a renormalization scale and the corresponding prefactor mentioned above for scalar or tensorial perturbations must be considered [14]. We observe immediately that the UV divergences cancel exactly and we are left with

$$\langle \phi^2(t, \vec{x}) \rangle_{ren} = \frac{H^2(1-\epsilon)^2}{16\pi^2} \left\{ \left(\frac{1}{4} - \nu^2\right) \left(-1 + \psi(3/2 - \nu) + \psi(3/2 + \nu) - \log \frac{\mu^2}{H^2(1-\epsilon)^2}\right) - \frac{2-\epsilon}{3(1-\epsilon)^2} \right\}. \quad (21)$$

The above self-correlators quantify the amplitude of perturbations at a given spacetime point.

IV. MODIFIED CORRELATORS AND ANGULAR POWER SPECTRUM

In previous sections we have studied the correlator $\langle \phi(t, \vec{x}) \phi(t, \vec{x}') \rangle$ and self-correlator $\langle \phi^2(t, \vec{x}) \rangle$ of tensorial

and scalar perturbations in slow-roll inflation. For an ordinary quantum mechanical system, with a finite number N of degrees of freedom, expectation values of the form $\langle \phi(i)\phi(j) \rangle$ and $\langle \phi^2(i) \rangle$ match when $j = i$ [for instance, in a chain of spins with $\phi(i) \equiv S_z(i)$ and $i = 1, \dots, N$]. However, we are facing here a field theory (with an infinite number of degrees of freedom), and the above matching is not *a priori* guaranteed. This is so because the self-correlator requires renormalization. We may either assume this discontinuity or modify the two-point correlation function to force it to match $\langle \phi^2(t, \vec{x}) \rangle_{ren}$ in the coincidence limit $\vec{x}' \rightarrow \vec{x}$ [5]. This second possibility was indirectly explored in [16] by analyzing the power spectrum of perturbations in momentum space. It has been somewhat debated in the literature and properly reviewed in [17]. One could naturally argue that the point-separated correlator has a well-defined definition in the distributional sense and there is not a mathematical need for any regularization [17, 18]. However, as the spatial points approach each other, the two-point correlator will grow without bound and diverge as the points merge. Therefore, from the physical point of view it seems reasonable to use a regularized form of the two-point correlator to consistently match the self-correlator at coincidence [5, 19]. In the conventional approach the expectation value of the self-correlator $\langle \phi^2(t, \vec{x}) \rangle$ plays almost no role. We assume here that the (renormalized) self-correlator is actually playing a physical role (as in the Casimir effect). As we will see shortly, the regu-

larized form of the two-point correlator makes the integral (13) UV convergent. The consequences of this merit to be explored. Therefore, we further analyze here this possibility taking advantage of the spacetime viewpoint sketched above.

We shall modify the correlators by adding the subtraction terms prescribed by renormalization and according to (19). We note that a distinguishing characteristic of adiabatic renormalization is that the subtraction terms $G_{Ad}^{(2)}((t, \vec{x}), (t, \vec{x}'))$ are well-defined for arbitrary point separation. In general this is not possible for an arbitrary spacetime, but for the homogeneous spaces relevant in cosmology the adiabatic subtraction terms extend to arbitrary large distances. With this in mind, we will finally compute the angular power spectrum for primordial perturbations using the modified spacetime correlators.

As a previous step we will compute the two-point function at leading order in slow-roll.

A. Two-point function at leading order in slow-roll

The procedure is similar for scalar and tensorial fluctuations, so we will do a general treatment. First, we start off splitting Eqs. (3) and (7) as a combination of two hypergeometric functions. To this end we use the transformation properties of hypergeometric functions [11]

$$F\left(\frac{3}{2} + \nu, \frac{3}{2} - \nu, 2, 1 - Z\right) = \frac{Z^{-\frac{3}{2}-\nu}\Gamma(-2\nu)}{\Gamma(\frac{3}{2}-\nu)\Gamma(\frac{1}{2}-\nu)} \operatorname{Re} \left\{ F\left(\frac{3}{2} + \nu, \frac{1}{2} + \nu, 1 + 2\nu, \frac{1}{Z}\right) \right\} \\ + \frac{Z^{-\frac{3}{2}+\nu}\Gamma(2\nu)}{\Gamma(\frac{3}{2}+\nu)\Gamma(\frac{1}{2}+\nu)} \operatorname{Re} \left\{ F\left(\frac{3}{2} - \nu, \frac{1}{2} - \nu, 1 - 2\nu, \frac{1}{Z}\right) \right\} \quad (22)$$

with $Z = \Delta\bar{x}^2/4 \geq 0$. We now expand expression (22) as a power series of the ‘‘slow-roll’’ parameter ν around $\nu = 3/2$, and stay at first order (for details see the Appendix). Grouping terms, we arrive at the following expression for the two-point function:

$$\langle \phi(x)\phi(x') \rangle \approx \frac{H^2(1-\epsilon)^2}{16\pi^2} \left\{ \frac{4}{\Delta\bar{x}^2} - 2 \log \frac{\Delta\bar{x}^2}{4} - 1 + \frac{2}{(3/2-\nu)} \left(\frac{\Delta\bar{x}^2}{4}\right)^{\nu-3/2} + 4 \operatorname{Re} \left[\log \left(\frac{\Delta\bar{x}}{2} + \sqrt{\frac{\Delta\bar{x}^2}{4} - 1} \right) \right] \right\} \quad (23)$$

Notice that the UV divergences are just the same as those found in (18), but now they are obtained at leading order in the slow-roll expansion. We recover exactly expression (18) taking the limit $\Delta\bar{x} \rightarrow 0$ and the slow-roll approximation.

B. Modified two-point function

We can now proceed to do the subtraction. The modified two-point function then reads

$$\langle \phi(x)\phi(x') \rangle_{ren} \approx \frac{H^2(1-\epsilon)^2}{16\pi^2} \left\{ \frac{2}{(3/2-\nu)} \left(\frac{\Delta\bar{x}^2}{4}\right)^{\nu-3/2} + 4 \operatorname{Re} \left[\log \left(\frac{\Delta\bar{x}}{2} + \sqrt{\frac{\Delta\bar{x}^2}{4} - 1} \right) \right] - \frac{5}{3} + 4\gamma + 2 \log \frac{\mu^2}{H^2} \right\} . \quad (24)$$

We remark that, at leading order in the slow-roll expansion, this is an expression valid for small and large

separations. For scales larger than the Hubble hori-

zon, $\Delta\bar{x} \gg 1$, we can further take the approximation, $4Re \left[\log \left(\frac{\Delta\bar{x}}{2} + \sqrt{\frac{\Delta\bar{x}^2}{4} - 1} \right) \right] \approx 2 \log(\Delta\bar{x}^2)$.

C. Angular power spectrum

We now compute the corresponding angular power spectrum from the modified two-point function for scalar perturbations and for low multipoles

$$C_\ell^{SW} = \frac{2\pi T_0^2}{25} \int_{-1}^1 \langle \mathcal{R}(x)\mathcal{R}(x') \rangle_{ren}(y) P_\ell(y) dy. \quad (25)$$

By construction this is a finite quantity, even without taking the large separation limit for the two-point function [as it was assumed in going from (13) to (14) and (15)]. To evaluate the logarithmic contributions of (24) to (25) we take into account that $\int_{-1}^1 dy \log(1-y) P_\ell(y) = -2/\ell(\ell+1)$, $\ell = 1, 2, \dots$. The final result for the angular power spectrum with the modified two-point function is very well approximated by the following analytical expression:

$$C_\ell^{SW} \approx \frac{4\pi G}{\epsilon} \frac{8\pi T_0^2}{25} \frac{H^2(1-\epsilon)^2 \bar{r}_L^{1-n}}{16\pi^2} \times \left\{ \frac{\Gamma(\ell + \frac{n-1}{2})}{\Gamma(\ell + 2 - \frac{n-1}{2})} - \frac{\bar{r}_L^{n-1}}{\ell(\ell+1)} \right\}, \quad (26)$$

where we have used $\nu - \frac{3}{2} = \frac{1-n}{2}$, and n represents the scalar index of inflation $n = 1 - 4\epsilon - 2\delta + O(\epsilon, \delta)^2$. Also notice that expression (26) is valid for $\ell \geq 1$, as for $\ell = 0$ there would be present all the constant contributions from the renormalized two-point function (24), including the one depending on the renormalization scale. In fact, the renormalization scale may be fixed by imposing the natural condition $C_0^{SW} = 0$.

Notice that the first term in (26) reproduces the standard result (15). The second one comes from the subtraction terms that we have added to the two-point correlator to continuously match the self-correlator at coincidence, but it shows scale invariance as well. Therefore, Eq. (26) is consistent with observations [4].

However, the two terms in (26) are competing, and the resulting amplitude for the coefficients C_ℓ^{SW} depends on the instant of time one evaluates \bar{r}_L . The first term is proportional to $H^2(t)\bar{r}_L^{(1-n)}(t)/\epsilon(t)$, and it is time independent. However, the second term depends slightly on time. The value of \bar{r}_L varies along the inflationary period, ranging from $\bar{r}_L \approx 1$, immediately after the instant of time t_i at which the scale r_L crosses the Hubble horizon [$a(t_i)r_L \approx H^{-1}(t_i)$], to $\bar{r}_L \approx e^{60}$, at the end of inflation (we have assumed that inflation lasts for around $N = 60$ e -foldings since the scale r_L exited the horizon at t_i). In the former case, $\bar{r}_L \approx 1$, the amplitude is severely reduced. In the latter situation, $\bar{r}_L^{(n-1)} \sim 10^{-1}$, where we have assumed that $n \approx 0.96$ [4], and the amplitude is then reduced at least 10%. The adequate value of

\bar{r}_L to properly evaluate the resulting amplitude in (26) is unclear. This question is closely related to the so-called “quantum-to-classical transition” [21], characterizing the period of time at which the primordial quantum perturbations behave as classical ones and define the initial conditions for the postinflationary evolution, along with its associated power spectrum. In momentum space (mode-by-mode picture) this process is thought to happen a few Hubble times after horizon exit [21], when the modes are frozen as classical perturbations. It seems natural to evaluate \bar{r}_L during this period, where quantum fluctuations are imprinted as classical perturbations. However, this quantum-to-classical mechanism is poorly understood, and it has not been rigorously established in the literature. Therefore one may regard $\bar{r}_L^{(n-1)} \equiv \alpha$ as a phenomenological parameter, varying in the range $1 > \alpha > 0$. Note that in the limiting case $\alpha \rightarrow 0$ one recovers the standard prediction, and this happens when the subtraction term is evaluated after inflation. This parameter has influence on the relative strength between multipole amplitudes

$$\frac{l_2(l_2+1)C_{l_2}^{SW}}{l_1(l_1+1)C_{l_1}^{SW}} = \frac{l_2(l_2+1) \frac{\Gamma(l_2 + \frac{n-1}{2})}{\Gamma(l_2 + 2 - \frac{n-1}{2})} - \alpha}{l_1(l_1+1) \frac{\Gamma(l_1 + \frac{n-1}{2})}{\Gamma(l_1 + 2 - \frac{n-1}{2})} - \alpha}. \quad (27)$$

Observations may properly fit the value of this parameter. It produces an observable effect for a significant range of values of α . As remarked above, the details of how this “quantum-to-classical transition” takes place are not well established in the literature, and further work is needed to fully understand this process. Within the present understanding of quantum gravity it is difficult to determine theoretically the value of α and hence the relative impact of the subtraction term in the observed angular power spectrum. However, as we showed above it can potentially be tested with observations.

V. CONCLUSIONS

We have analyzed two-point correlators and self-correlators of primordial perturbations in quasi-de Sitter spacetime backgrounds. For large separations two-point correlators exhibit nearly scale invariance in a very elegant way. We have deformed the two-point correlators to smoothly match the self-correlators at coincidence. To this end we have used renormalization methods in homogeneous backgrounds. We have studied the physical consequences for the angular power spectrum at low multipoles. Scale invariance is maintained, but the amplitude of C_ℓ could change significantly. If one accepts a mismatch between the standard two-point correlators and the self-correlators and keeps only the large-scale behavior, the conventional predictions remain unaltered.

We finally stress the importance of getting a better understanding of how to renormalize cosmological observables. The analysis carried out in the spacetime

framework for the tree-level power spectrum may offer a way to experimentally probe this issue.

Acknowledgements. J. N-S. would like to thank I. Agullo, A. Fabbri, G. J. Olmo and L. Parker for very useful discussions. This work is supported by the Spanish Grant No. FIS2011-29813-C02-02 and the Consolider Program CPANPHY-1205388.

APPENDIX

In this appendix we give the basic steps to obtain the result (23). We consider (22) first. Since the first prefactor is of order $O((\frac{3}{2}-\nu)^1)$, we only need the corresponding hypergeometric function to be of order $O((\frac{3}{2}-\nu)^0)$. One can see that

$$Z^{-\frac{3}{2}-\nu} \text{Re} \left\{ F \left(\frac{3}{2} + \nu, \frac{1}{2} + \nu, 1 + 2\nu, \frac{1}{Z} \right) \right\} \Big|_{\nu=3/2} = \quad (28)$$

$$6 \text{Re} \{ \log(Z-1) \} - 6 \log(Z) + \frac{3}{Z} - \frac{3}{(1-Z)}$$

On the other hand, the second prefactor of (22) is of order $O((\frac{3}{2}-\nu)^0)$, so it is necessary to evaluate the second hypergeometric function at first order in the slow-roll series. To this end we will employ the following relation

[11]:

$$F \left(\frac{3}{2} - \nu, \frac{1}{2} - \nu, 1 - 2\nu, \frac{1}{Z} \right) = \quad (29)$$

$$\left(1 - \frac{1}{Z} \right)^{-3/4} P_{1/2}^\nu \left[\frac{2Z-1}{2\sqrt{Z(Z-1)}} \right] 2^{-2\nu} \Gamma(1-\nu) Z^{-\nu},$$

together with

$$P_{1/2}^\nu(Z) = \left(\frac{Z+1}{Z-1} \right)^{\nu/2} \frac{F(-\frac{1}{2}, \frac{3}{2}, 1-\nu, \frac{1-Z}{2})}{\Gamma(1-\nu)}. \quad (30)$$

At this point one can expand

$$F \left(-\frac{1}{2}, \frac{3}{2}, 1-\nu, \frac{1-Z}{2} \right) \approx F \left(-\frac{1}{2}, \frac{3}{2}, -\frac{1}{2}, \frac{1-Z}{2} \right) \quad (31)$$

$$+ \left(\nu - \frac{3}{2} \right) \frac{dF \left(-\frac{1}{2}, \frac{3}{2}, 1-\nu, \frac{1-Z}{2} \right)}{d\nu} \Big|_{\nu=3/2},$$

where the derivative can be performed using the representation series of the hypergeometric function. Doing all the calculation properly one finally arrives at the following result:

$$\text{Re} \left\{ F \left(\frac{3}{2} - \nu, \frac{1}{2} - \nu, 1 - 2\nu, \frac{1}{Z} \right) \right\} \approx 1$$

$$+ \left(\frac{3}{2} - \nu \right) \left[\frac{1}{4Z} + \frac{1}{4(1-Z)} - \frac{1}{2} \text{Re} \{ \log(Z-1) \} - \frac{1}{2} \log(Z) \right.$$

$$\left. + 2 \text{Re} \left\{ \log \left(\sqrt{Z} + \sqrt{Z-1} \right) \right\} \right]. \quad (32)$$

Taking all these results together for $Z \equiv \Delta \bar{x}^2 / 4$ in (22) we can approximate the two-point function as in (23). We have also checked numerically that this expansion works well irrespectively of the value of Z .

-
- [1] A.G. Riess *et al.*, *Astron. J.* **116**, 1009 (1998); S. Perlmutter *et al.* *Astron. J.* **517**, 565 (1999).
- [2] G. Smoot, *et al.* *Astrophys. J. Lett.* **396**, L1 (1992).
- [3] V.F. Mukhanov and G.V. Chibisov, *JETP Lett.* **33**, 532(1981); S. W. Hawking, *Phys. Lett.* **115B**, 295 (1982); A. Guth and S.-Y. Pi, *Phys. Rev. Lett.* **49**, 1110 (1982); A. A. Starobinsky, *Phys. Lett.* **117B**, 175 (1982); J.M. Bardeen, P.J. Steinhardt, and M.S. Turner, *Phys. Rev. D* **28**, 679 (1983).
- [4] Planck Collaboration, *Planck 2013 Results. XV. CMB power spectra and likelihood*, arXiv:1303.5075 (2013).
- [5] L. Parker and D.J. Toms, *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity* (Cambridge University Press, Cambridge, UK, 2009).
- [6] N.D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, UK, 1982).
- [7] S. Weinberg, *Cosmology* (Cambridge University Press, Cambridge, UK, 2008).
- [8] T.S. Bunch and P.C.W. Davies, *Proc. R. Soc. A* **360**, 117 (1978).
- [9] L. H. Ford and L. Parker, *Phys. Rev. D* **16**, 245 (1977); A. Vilenkin and L. H. Ford, *Phys. Rev. D* **26**, 1231 (1982); B. Allen, *Phys. Rev. D* **32**, 3136 (1985); B. Allen and A. Folacci, *Phys. Rev. D* **35**, 3771 (1987).
- [10] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products* (Academic Press, New York, 1994).
- [11] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions* (National Bureau of Standards, New York, 1972).
- [12] L. Parker and S. A. Fulling, *Phys. Rev. D* **9**, 341 (1974).
- [13] A. Landete, J. Navarro-Salas, and F. Torrenti, *Phys. Rev. D* **89**, 044030 (2014); *Phys. Rev. D* **88**, 061501 (2013).
- [14] We note that for a free conformal-invariant field (like the electromagnetic one), the logarithmic subtraction term is absent and hence also the renormalization scale μ . The two-point correlator is diluted by inflation and only the self-correlator survives as a quantity with non-negligible amplitude [15].
- [15] I. Agullo and J. Navarro-Salas, *Conformal anomaly and primordial magnetic fields*, arXiv:1309.3435 [gr-qc].
- [16] L. Parker, *Amplitude of perturbations from inflation*, arXiv:hep-th/0702216; I. Agullo, J. Navarro-Salas, G. J.

- Olmo and L. Parker, *Phys. Rev. Lett.* **103**, 061301 (2009).
- [17] M. Bastero-Gil *et al.*, *Phys. Rev. D* **87**, 087302 (2013); R. Rangarajan, arXiv:1401.8082.
- [18] F. Finelli, G. Marozzi, G. P. Vacca, and G. Venturi *Phys. Rev. D* **76**, 103528 (2007).
- [19] Alternatively, one could also consider short-distance modifications of the two-point function due to quantum gravity effects. See, for instance, Ref. [20].
- [20] T. Padmanabhan, *Phys. Rev. Lett.* **60**, 2229 (1988).
- [21] A. R. Liddle and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure* (Cambridge University Press, Cambridge, UK, 2006).