

# Relating quarks and leptons without grand-unification

S. Morisi,<sup>1,\*</sup> E. Peinado,<sup>1,†</sup> Yusuke Shimizu,<sup>2,‡</sup> and J. W. F. Valle<sup>1,§</sup>

<sup>1</sup>*AHEP Group, Institut de Física Corpuscular – C.S.I.C./Universitat de València  
Edificio Institutos de Paterna, Apt 22085, E-46071 Valencia, Spain*

<sup>2</sup>*Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan  
(Dated: September 1, 2011)*

In combination with supersymmetry, flavor symmetry may relate quarks with leptons, even in the absence of a grand-unification group. We propose an  $SU(3) \times SU(2) \times U(1)$  model where both supersymmetry and the assumed  $A_4$  flavor symmetries are softly broken, reproducing well the observed fermion mass hierarchies and predicting: (i) a relation between down-type quarks and charged lepton masses, and (ii) a correlation between the Cabibbo angle in the quark sector, and the reactor angle  $\theta_{13}$  characterizing CP violation in neutrino oscillations.

PACS numbers: 11.30.Hv 14.60.-z 14.60.Pq 14.80.Cp 14.60.St 23.40.Bw

## INTRODUCTION

Understanding the observed pattern of quark and lepton masses and mixing [1, 2] constitutes one of the deepest challenges in particle physics. Flavor symmetries provide a very useful approach towards reducing the number of free parameters describing the fermion sector [3]. It has long been advocated that grand unification offers a suitable framework to describe flavor. In what follows we will adopt the alternative approach, assuming that flavor is implemented directly at the  $SU(3) \times SU(2) \times U(1)$  level. Typically this requires several  $SU(2)$  doublet scalars in order to break spontaneously the flavor symmetry so as to obtain an acceptable structure for the masses and mixing matrices. (One may alternatively introduce “flavons” instead of additional Higgs doublets, but in this case one would have to give up renormalizability).

In order to construct a “realistic” extension of the Standard Model (SM) with flavor symmetry one needs a suitable alignment of the scalar vacuum expectation values (vevs) in the theory [4–7]. There are several multi-doublet extensions of the SM with flavor in the market, but renormalizable supersymmetric extensions of the SM with a flavor symmetry are only a few [8], usually because the existence of additional Higgs doublets spoils the unification of the coupling constants.

Here we choose to renounce to this theoretical argument, noting that gauge coupling unification may happen in multi-doublet schemes due to other effects. What we now present is a supersymmetric extension of the SM based on the  $A_4$  group where all the matter fields as well as the Higgs doublets belong to the same  $A_4$  representation, namely, the triplet. This leads us to two theoretical predictions. The first a mass relation

$$\frac{m_\tau}{\sqrt{m_e m_\mu}} \approx \frac{m_b}{\sqrt{m_d m_s}}, \quad (1)$$

involving down-type quarks and charged lepton mass ratios. Such relation can be obtained by a suitable com-

ination of the three Georgi-Jarlskog (GJ) mass relations [9],

$$m_b = m_\tau, \quad m_s = 1/3 m_\mu, \quad m_d = 3 m_e, \quad (2)$$

which arise within a particular ansatz for the  $SU(5)$  model and hold at the unification scale. In contrast to eq. (2), our relation requires no unification group and holds at the electroweak scale. It would, in any case, be rather robust against renormalization effects as it involves only mass ratios.

The second prediction obtained in our flavor model is a correlation between the Cabibbo angle for the quarks and the so-called “reactor angle”  $\theta_{13}$  characterizing the strength of CP violation in neutrino oscillations [10, 11]. Within a reasonable approximation we find

$$\lambda_C \approx \frac{1}{\sqrt{2}} \frac{m_\mu m_b}{m_\tau m_s} \sqrt{\sin^2 2\theta_{13}} - \sqrt{\frac{m_u}{m_c}}. \quad (3)$$

which arises mainly from the down-type quark sector [12] with a correction coming from the up isospin diagonalization matrix. This is a very interesting relation, discussed below in more detail.

## THE MODEL

Here we propose a supersymmetric model based on an  $A_4$  flavor symmetry realized in an  $SU(3) \times SU(2) \times U(1)$  gauge framework. The field representation content is given in Table I. Note that all quarks and leptons transform as  $A_4$  triplets. Similarly the Higgs superfields with opposite hypercharge characteristic of the MSSM are now upgraded into two sets, also transforming as  $A_4$  triplets. Note that since all matter fields transform in the same way under the flavor symmetry one may in principle embed the model into a grand-unified scheme. However, given the large number of scalar doublets, gauge coupling unification must proceed differently, see, for example, Ref. [13].

fields	$\hat{L}$	$\hat{E}^c$	$\hat{Q}$	$\hat{U}^c$	$\hat{D}^c$	$\hat{H}^u$	$\hat{H}^d$
$SU(2)_L$	2	1	2	1	1	2	2
$A_4$	3	3	3	3	3	3	3

TABLE I: Basic multiplet assignments of the model

The most general renormalizable Yukawa Lagrangian for the charged fermions in the model is [14]

$$L_{\text{Yuk}} = y_{ijk}^l \hat{L}_i \hat{H}_j^d \hat{E}_k^c + y_{ijk}^d \hat{Q}_i \hat{H}_j^d \hat{D}_k^c + y_{ijk}^u \hat{Q}_i \hat{H}_j^u \hat{U}_k^c, \quad (4)$$

where  $y_{ijk}^{u,d,l}$  are  $A_4$ -tensors, assumed real at this stage.

The Higgs scalar potential invariant under  $A_4$  is

$$\begin{aligned} V = & (|\mu|^2 + m_{H_u}^2)(|H_1^u|^2 + |H_2^u|^2 + |H_3^u|^2) \\ & + (|\mu|^2 + m_{H_d}^2)(|H_1^d|^2 + |H_2^d|^2 + |H_3^d|^2) \\ & - [b(H_1^u H_1^d + H_2^u H_2^d + H_3^u H_3^d) + \text{c.c.}] \\ & + \frac{1}{8}(g^2 + g'^2)(|H_1^u|^2 + |H_2^u|^2 + |H_3^u|^2 + \\ & \quad - |H_1^d|^2 - |H_2^d|^2 - |H_3^d|^2)^2. \end{aligned} \quad (5)$$

Assuming that the Higgs doublet scalars take real vevs  $\langle H_i^{u,d} \rangle = v_i^{u,d}$  one can show that the minimization of the potential  $V$  gives as possible local minima the alignments  $\langle H^{0,u,d} \rangle \sim (1, 0, 0)$  and  $(1, 1, 1)$ . Only the first is viable and we verify that minimization leads to this solution within a wide region of parameters. By adding  $A_4$  soft breaking terms to the  $A_4$ -invariant scalar potential in eq. (5)

$$V_{\text{soft}} = \sum_{ij} (\mu_{ij}^u H_i^{u*} H_j^u + \mu_{ij}^d H_i^{d*} H_j^d) + \sum_{ij} b_{ij} H_i^d H_j^u,$$

one finds that

$$\langle H^u \rangle = (v^u, \varepsilon_1^u, \varepsilon_2^u), \quad \langle H^d \rangle = (v^d, \varepsilon_1^d, \varepsilon_2^d), \quad (6)$$

where  $\varepsilon_{1,2}^u \ll v^u$  and  $\varepsilon_{1,2}^d \ll v^d$ .

### Charged fermions

By using  $A_4$  product rules it is straightforward to show that the charged fermion mass matrix takes the following universal structure [14]

$$M_f = \begin{pmatrix} 0 & y_1^f \langle H_3^f \rangle & y_2^f \langle H_2^f \rangle \\ y_2^f \langle H_3^f \rangle & 0 & y_1^f \langle H_1^f \rangle \\ y_1^f \langle H_2^f \rangle & y_2^f \langle H_1^f \rangle & 0 \end{pmatrix}, \quad (7)$$

where  $f$  denotes any charged lepton, up- or down-type quarks. Note that, in addition to the ‘‘texture’’ zeros in the diagonal, one has additional relations among the

parameters. This may be seen explicitly by rewriting eq. (7) as

$$M_f = \begin{pmatrix} 0 & a^f \alpha^f & b^f \\ b^f \alpha^f & 0 & a^f r^f \\ a^f & b^f r^f & 0 \end{pmatrix}, \quad (8)$$

where  $a^f = y_1^f \varepsilon_1^f$ ,  $b^f = y_2^f \varepsilon_1^f$ , with  $y_{1,2}^f$  denoting the only two couplings arising from the  $A_4$ -tensor in eq. (4),  $r^f = v^f / \varepsilon_1^f$  and  $\alpha^f = \varepsilon_2^f / \varepsilon_1^f$ . Thanks to the fact that the same Higgs doublet  $H^d$  couples to the lepton and to the down-type quarks one has, in addition, the following relations

$$r^l = r^d, \quad \alpha^l = \alpha^d, \quad (9)$$

involving down-type quarks and charged leptons.

It is straightforward to obtain analytical expressions for  $a^f$ ,  $b^f$  and  $r^f$  from eq. (8) in terms of the charged fermion masses and  $\alpha^f$ ,

$$\frac{r^f}{\sqrt{\alpha^f}} \approx \frac{m_3^f}{\sqrt{m_1^f m_2^f}}, \quad a^f \approx \frac{m_2^f}{m_3^f} \frac{\sqrt{m_1^f m_2^f}}{\sqrt{\alpha^f}}, \quad b^f \approx \frac{\sqrt{m_1^f m_2^f}}{\sqrt{\alpha^f}}. \quad (10)$$

From eq. (9) and eq. (10) it follows that

$$\frac{m_\tau}{\sqrt{m_e m_\mu}} \approx \frac{m_b}{\sqrt{m_d m_s}},$$

a formula relating quark and lepton mass ratios (to a very good approximation this formula also holds for complex Yukawa couplings). This relation is a strict prediction of our model, and appears in a way similar to the celebrated  $SU(5)$  mass relation, despite the fact that we have not assumed any unified group, but just the  $SU(3) \times SU(2) \times U(1)$  gauge structure. It allows us to compute the down quark mass in terms of the charged fermion masses and the  $s$  and  $b$  quarks, as

$$m_d \approx m_e \frac{m_\mu}{m_s} \left( \frac{m_b}{m_\tau} \right)^2. \quad (11)$$

This mass formula predicts the down quark mass at the scale of the  $Z$  boson mass, to lie in the region

$$\begin{aligned} 1.71 \text{ MeV} &< m_d^{th} < 3.35 \text{ MeV} \\ 1.71 \text{ MeV} &< m_d < 4.14 \text{ MeV}, \end{aligned} \quad (12)$$

at  $1\sigma$  [15]. This is illustrated in Fig. 1 where, to guide the eye, we have also included the  $1\sigma$  experimental ranges from reference [15], as well as the best fit point and the GJ prediction.

Note also that, thanks to supersymmetry, we obtain a relation only among the charged lepton and down-type quark mass ratios, avoiding the unwanted relation found by Wilczek and Zee in Ref. [16].




FIG. 1: The shaded band gives our prediction for the down-strange quark masses at the  $M_z$  scale, eq. (11), vertical and horizontal lines are the  $1\sigma$  experimental ranges from ref. [15].

### Neutrinos

To the renormalizable model we have so far we now add an effective dimension-five  $A_4$ -preserving lepton-number violating operator

$$\mathcal{L}_{5d} = \frac{f_{ijklm}}{\Lambda} \hat{L}_i \hat{L}_j \hat{H}_l^u \hat{H}_m^u, \quad (13)$$

where the  $A_4$ -tensor  $f_{ijklm}$  takes into account all the possible contractions of the product of four  $A_4$  triplets<sup>1</sup>.

Neutrino masses are induced after electroweak symmetry breaking from the operator in eq. (13). In order to determine the flavor structure of the resulting mass matrix we take the limit where the vev hierarchy  $\langle H_1^u \rangle \gg \langle H_2^u \rangle, \langle H_3^u \rangle$  holds, leading to [14]

$$M_\nu = \begin{pmatrix} xr^{u2} & \kappa r^u & \kappa r^u \alpha^u \\ \kappa r^u & yr^{u2} & 0 \\ \kappa r^u \alpha^u & 0 & zr^{u2} \end{pmatrix}, \quad (14)$$

where  $x, y, z$  and  $\kappa$  are coupling constants, while  $r^u$  and  $\alpha^u$  already been introduced above in the up quark sector.

The best fit of neutrino oscillation data [2] yields maximally mixed  $\mu$  and  $\tau$  neutrinos. This is possible, in the basis where charged lepton is diagonal, if and only if the light-neutrino mass matrix is approximately  $\mu - \tau$  invariant. In turn this holds true if  $y \approx z$  and  $\alpha^u \approx 1$  [14]<sup>2</sup>. When  $\alpha^u < 1$  the ‘‘atmospheric angle’’ deviates from the maximality. We have verified that for  $\alpha^u \gtrsim 0.5$  the atmospheric angle is within its  $3\sigma$  allowed range.

### RELATING THE CABIBBO ANGLE TO $\theta_{13}$

In the CP conserving limit we have taken so far we have in total 14 free parameters to describe the fermion sector: six  $a^f$  and  $b^f$  parameters (three for each charged fermion-type), plus four  $r^f$  and  $\alpha^f$  (here only down-type are counted, in view of eq. (9)), plus four parameters describing the neutrino mass induced by the dimension-5 operator:  $x, y, z, \kappa$ . These parameters describe 18 observables, which may be taken as the 9 charged fermion masses, the two neutrino squared mass differences describing neutrino oscillations, the three neutrino mixing angles, the neutrinoless double beta decay effective mass parameter, the Cabibbo angle, in addition to  $V_{ub}$  and  $V_{cb}$ . Hence we have four relations.

The first of these we have already seen, namely the mass relation in eq. (1) and Fig. 1. The second is a quark-lepton mixing angle relation concerning the Cabibbo angle  $\lambda_C$  and the ‘‘reactor angle’’  $\theta_{13}$  describing neutrino oscillations. To derive it note first that the matrix in eq. (8) is diagonalized on the left by a rotation in the 12 plane, namely

$$\sin \theta_{12}^f \approx \sqrt{\frac{m_1^f}{m_2^f}} \frac{1}{\sqrt{\alpha^f}}. \quad (15)$$

In order to give an analytical expression for the relation between Cabibbo and reactor angles, we neglect mixing of the third family of quarks and go in the limit where our neutrino mass matrix, eq. (15) is  $\mu - \tau$  invariant, that is  $\alpha^u = 1$  and  $y = z$ . In this approximation, the reactor mixing angle is given by

$$\sin \theta_{13} = \frac{1}{\sqrt{2}} \sin \theta_{12}^t = \frac{1}{\sqrt{2}} \sqrt{\frac{m_e}{m_\mu}} \frac{1}{\sqrt{\alpha^t}}, \quad (16)$$

using our mass relation in eq. (1) one finds that the Cabibbo angle may be written as

$$\lambda_C = \frac{m_b}{m_s} \frac{\sqrt{m_e m_\mu}}{m_\tau} \frac{1}{\sqrt{\alpha^d}} - \sqrt{\frac{m_u}{m_c}}. \quad (17)$$

Comparing eq. (16) with eq. (17) leads immediately to equation (3). In order to display this prediction graphically we take the quark masses at  $1\sigma$ , obtaining the curved band shown in fig. 2. The narrow horizontal band indicates current determination of the Cabibbo angle, while the two vertical dashed lines represent the expected sensitivities of the Double-Chooz [18] and Daya-Bay [19] experiments on the ‘‘reactor mixing angle’’  $\theta_{13}$ . The curved line corresponds to the analytical approximation for the best fit value of the quark masses in eq. (3). Clearly the width of the curved band characterizing our prediction is dominated by quark mass determination uncertainties.

<sup>1</sup> Specific realizations of  $\mathcal{L}_{5d}$  within various seesaw schemes [17] can, of course, be envisaged.

<sup>2</sup> The charged lepton mass matrix is mainly diagonalized by a rotation in the 12 plane.




FIG. 2: The shaded band gives our predicted  $1\sigma$  correlation between the Cabibbo angle and the reactor angle, as above. Vertical lines give the expected sensitivities on  $\theta_{13}$  [18, 19].

Finally note that mixing parameters of the third family of quarks  $U_{13}^q \approx \frac{m_s^q}{m_3^q} \frac{\sqrt{m_1^q m_2^q}}{m_3^q} \frac{1}{\sqrt{\alpha^q}}$  and  $U_{23}^q \approx \frac{m_1^q (m_2^q)^2}{(m_3^q)^3} \frac{1}{\alpha^q}$  ( $q = u, d$ ) are negligible, and can not account for the measured values of  $V_{ub}$  and  $V_{cb}$ . The predicted values obtained for these are too small so that in its simplest presentation described above our model can not describe the CP violation found in the decays of neutral kaons. However there is a simple solution which maintains the good predictions described above, namely, adding colored vector-like  $SU(2)_L$  singlet states. In this case acceptable values for  $V_{ub}$  and  $V_{cb}$ , leading to adequate CP violation can arise solely from non-unitarity effects of the quark mixing matrix.

## OUTLOOK

We proposed a supersymmetric extension of the standard model with an  $A_4$  flavor symmetry, where all matter fields in the model transform as triplets of the flavor group. Charged fermion masses arise from renormalizable Yukawa couplings while neutrino masses are treated in an effective way. The scheme illustrates how, in combination with supersymmetry, flavor symmetry may relate quarks with leptons, even in the absence of a grand-unification group. Two good predictions emerge: (i) a relation between down-type quarks and charged lepton masses, and (ii) a correlation between the Cabibbo angle in the quark sector, and the reactor angle  $\theta_{13}$  characterizing CP violation in neutrino oscillations, which lies within the sensitivities of upcoming experiments.

Although the predicted values for the other mixing parameters  $V_{uc}$  and  $V_{cb}$  of the Cabibbo-Kobayashi-Maskawa matrix are too small, we mentioned a simple way to circumvent this, making the scheme fully realistic.

Finally note that, with few exceptions such as those in Refs. [20, 21], grand-unified flavor models are not more predictive than the novel idea proposed here and illustrate through this simple scheme. As it stands the model fits well with the idea that gauge coupling unification

may be an effect of the presence of extra dimensions rather than of grand-unified interactions [13]. Notwithstanding, we wish to stress that our model is manifestly embeddable into a standard Grand-Unified scenario, which would result in further predictive power. A detailed study of this particular model lies outside the scope of this letter and will be taken up elsewhere.

We thank M. Hirsch and M. Tanimoto for useful discussions. This work was supported by the Spanish MICINN under grants FPA2008-00319/FPA and MULTIDARK CSD2009-00064, by Prometeo/2009/091 (Generalitat valenciana), by the EU Network grant UNILHC PITN-GA-2009-237920. S. M. is supported by a Juan de la Cierva contract, E. P. is supported by CONACyT (Mexico) and Y.S. by the Japan Society of Promotion of Science, Grand-in-Aid for Scientific Research, No.22.3014.

---

\* Electronic address: morisi@ific.uv.es

† Electronic address: epeinado@ific.uv.es

‡ Electronic address: shimizu@muse.sc.niigata-u.ac.jp

§ Electronic address: valle@ific.uv.es

- [1] K. Nakamura *et al.*, Journal of Physics G: Nuclear and Particle Physics **37**, 075021 (2010).
- [2] T. Schwetz, M. Tortola and J. W. F. Valle, New J. Phys. **10**, 113011 (2008); New J. Phys. **13** (2011) 063004
- [3] H. Ishimori *et al.*, Prog. Theor. Phys. Suppl. **183**, 1 (2010), [1003.3552].
- [4] E. Ma and G. Rajasekaran, Phys. Rev. **D64**, 113012 (2001), [hep-ph/0106291].
- [5] K. S. Babu, E. Ma and J. W. F. Valle, Phys. Lett. **B552**, 207 (2003), [hep-ph/0206292].
- [6] A. Zee, Phys. Lett. **B630**, 58 (2005), [hep-ph/0508278].
- [7] G. Altarelli and F. Feruglio, Nucl. Phys. **B720**, 64 (2005), [hep-ph/0504165].
- [8] K. S. Babu and J. Kubo, Phys. Rev. **D71**, 056006 (2005).
- [9] H. Georgi and C. Jarlskog, Phys. Lett. **B86**, 297 (1979).
- [10] H. Nunokawa, S. J. Parke and J. W. F. Valle, Prog. Part. Nucl. Phys. **60**, 338 (2008).
- [11] ISS Physics Working Group, A. Bandyopadhyay *et al.*, Rep. Prog. Phys. **72**, 106201 (2009).
- [12] R. Gatto, G. Sartori and M. Tonin, Phys. Lett. **B28**, 128 (1968).
- [13] C. Munoz, JHEP **12**, 015 (2001); Mod. Phys. Lett. **A22**, 989 (2007).
- [14] S. Morisi and E. Peinado, Phys. Rev. **D80**, 113011 (2009), [0910.4389].
- [15] Z.-z. Xing, H. Zhang and S. Zhou, Phys. Rev. **D77**, 113016 (2008), [0712.1419].
- [16] F. Wilczek and A. Zee, Phys. Rev. Lett. **42**, 421 (1979).
- [17] For a recent seesaw review see, e.g. J. W. F. Valle, J. Phys. Conf. Ser. **53**, 473 (2006) [arXiv:hep-ph/0608101].
- [18] Double Chooz collaboration, F. Ardellier *et al.*, hep-ex/0606025.
- [19] Daya-Bay collaboration, X. Guo *et al.*, hep-ex/0701029.
- [20] S. F. King and G. G. Ross, Phys. Lett. B **574** (2003) 239
- [21] R. Dermisek and S. Raby, Phys. Lett. B **622** (2005) 327