



Visible-light photoredox catalyzed synthesis of pyrroloisoquinolines via organocatalytic oxidation/[3 + 2] cycloaddition/oxidative aromatization reaction cascade with Rose Bengal

Carlos Vila, Jonathan Lau and Magnus Rueping*

Letter

Open Access

Address:
Institute of Organic Chemistry, RWTH Aachen University,
Landoltweg 1, D-52074 Aachen, Germany

Email:
Magnus Rueping* - magnus.rueping@rwth-aachen.de

* Corresponding author

Keywords:
alkaloids; [3 + 2] cycloaddition; organocatalysis; oxidation;
photochemistry; photoredox catalysis; Rose Bengal; visible-light

Beilstein J. Org. Chem. **2014**, *10*, 1233–1238.
doi:10.3762/bjoc.10.122

Received: 20 February 2014
Accepted: 30 April 2014
Published: 27 May 2014

This article is part of the Thematic Series "Organic synthesis using photoredox catalysis".

Guest Editor: A. G. Griesbeck

© 2014 Vila et al; licensee Beilstein-Institut.
License and terms: see end of document.

Abstract

Pyrrolo[2,1-*a*]isoquinoline alkaloids have been prepared via a visible light photoredox catalyzed oxidation/[3 + 2] cycloaddition/oxidative aromatization cascade using Rose Bengal as an organo-photocatalyst. A variety of pyrroloisoquinolines have been obtained in good yields under mild and metal-free reaction conditions.

Introduction

Pyrrolo[2,1-*a*]isoquinolines constitute the core structure of the natural products family lamellarin alkaloids (Figure 1) [1-4]. These alkaloids display numerous biological activities such as inhibitor of human topoisomerase I by lamellarin D [5] or inhibition of HIV integrase by lamellarin α -20-sulfate [6,7]. Moreover lamellarin I and lamellarin K also showed potential anti-tumor activities [8,9]. Due to their potential biological activities, the synthesis of pyrrolo[2,1-*a*]isoquinolines has become a very interesting, important and attractive goal in organic synthesis [10-20]. For example, dipolar [3 + 2] cycloaddition using azomethine ylides [21] is a powerful class of reactions that

permits the synthesis of structural complex molecules in a straightforward way and has been used for the efficient synthesis of this type of compounds [22-26]. Recently, several metal mediated syntheses using a [3 + 2] cycloaddition reaction have been described in the literature. Porco Jr. et al. [27] described a silver-catalyzed cycloisomerization/dipolar cycloaddition for the synthesis of the pyrrolo[2,1-*a*]isoquinolines. Wang and co-workers described a copper catalyzed oxidation/[3 + 2] cycloaddition/aromatization cascade [28]. Also, Xiao disclosed a very elegant oxidation/[3 + 2] cycloaddition/aromatization cascade catalyzed by $[\text{Ru}(\text{bpy})_3]^{3+}$ under

irradiation with visible light [29]. In this context, very recently Zhao reported the same reaction using C₆₀-Bodipy hybrids [30] and porous material immobilized iodo-Bodipy [31] as photocatalysts, obtaining in both cases good yields for different pyrrolo[2,1-*a*]isoquinolines. Finally, Lu presented in 2013 a dirhodium complex for the synthesis of these compounds [32]. Despite these elegant and important syntheses of pyrrolo[2,1-*a*]isoquinolines through dipolar [3 + 2] cycloaddition, the development of metal-free syntheses using visible light photoredox catalysis with simple organic dyes remained unexplored. Visible-light photoredox catalysis has emerged as an important field and has attracted increasing attention in recent years [33–42]. Thus, in the last years spectacular advances in visible-light photoredox catalysis have been made and this kind of catalysis has become a powerful tool in organic synthesis. In this context, the use of organic dyes as photoredox catalysts [40–42] has been demonstrated by several groups [43–61] and became a useful alternative to the inorganic photoredox catalysts that are expensive and sometimes toxic. The organic dyes have very important qualities such as being inexpensive, environmentally friendly and easy to handle. As a part of our ongoing research on photoredox catalysis [62–72], we herein present a synthesis of pyrrolo[2,1-*a*]isoquinolines through an oxidation/[3+2] cycloaddition/aromatization cascade catalyzed by Rose Bengal under irradiation with green LEDs.

Results and Discussion

Initially, we focused on the reaction between methyl dihydroisoquinoline ester **1a** and *N*-methylmaleimide (**2a**) catalyzed by Rose Bengal. Although the [3 + 2] cycloaddition occurs smoothly in the presence of Rose Bengal (5 mol %) in acetonitrile under irradiation with visible light, the reaction was not selective affording the dihydropyrrolo[2,1-*a*]isoquinoline **3aa** in 35% yield and the hexahydropyrrolo[2,1-*a*]isoquinoline **4aa** in 26% yield, after column chromatography (Scheme 1).

In order to improve the selectivity of the reaction to the aromatized product **3aa**, *N*-bromosuccinimide was added to the reaction mixture when the starting materials were completely consumed [29–31,73]. In this case the desired product **3aa** was obtained in 72% yield (Table 1, entry 1). Other organic dyes such as Rhodamine B or Eosin Y were less efficient compared to Rose Bengal (Table 1, entries 2 and 3, respectively). Several solvents were tested without an improvement in the yield of the product (Table 1, entries 4–9). Finally, after tuning the relative amounts of the reagents, the product **3aa** was isolated in 76% yield (Table 1, entry 12).

With the optimal conditions in hand, we examined the substrate scope for the photoreaction catalyzed by Rose Bengal (Scheme 2). Various tetrahydroisoquinolines with different






Figure 1: Representative examples of lamellarin alkaloids.





Scheme 1: Photocatalytic metal free construction of pyrrolo[2,1-*a*]isoquinolines.

Table 1: Optimization of the reaction conditions.^a

Entry	Catalyst	Solvent	Yield (%) ^b
1	Rose Bengal	CH ₃ CN	72
2	Rhodamine B	CH ₃ CN	11
3	Eosin Y	CH ₃ CN	40
4	Rose Bengal	THF	29
5	Rose Bengal	CH ₂ Cl ₂	26
6	Rose Bengal	toluene	14
7	Rose Bengal	DMF	65
8	Rose Bengal	MeOH	52
9	Rose Bengal	EtOAc	16
10 ^c	Rose Bengal	CH ₃ CN	64
11 ^d	Rose Bengal	CH ₃ CN	60
12 ^e	Rose Bengal	CH ₃ CN	76

^aReaction conditions: **1a** (0.2 mmol), **2a** (0.2 mmol), organic dye (5 mol %), solvent (1 mL), green LEDs irradiation for 24 hours. NBS (1.1 equiv) was added to the reaction mixture and stirring was continued for 1 hour. ^bYields of the isolated products after column chromatography. ^c1.25 equiv of **1a** was used. ^d1.25 equiv of **2a** was used. ^e1.1 equiv of **1a** was used.

**Scheme 2:** Evaluation of the substrate scope.



Scheme 3: Evaluation of the substrate scope with activated alkynes.

electron-withdrawing groups (R²) such as methyl ester (**1a**), ethyl ester (**1b**), *tert*-butyl ester (**1c**), cyano (**1d**) or aromatic ketone (**1e**) were reacted with *N*-methylmaleimide (**2a**) and gave the corresponding products **3** in moderate to good yields. In addition, different *N*-substituted maleimides were tested under the optimized reaction conditions to give the corresponding products with good yields. Incorporation of methoxy groups at C-6 and C-7 in the dihydroisoquinoline core was well tolerated, affording the corresponding product **3fa** in 68% yield.

To demonstrate the synthetic utility of the oxidation/[3 + 2] cycloaddition/aromatization cascade we examined other dipolarophiles such as activated alkynes **5**. In this case, the addition of NBS was not necessary, and the corresponding products **6** were isolated in moderate yields (Scheme 3).

Conclusion

In conclusion, we have developed a metal-free photoredox oxidation/[3 + 2] dipolar cycloaddition/oxidative aromatization cascade catalyzed by Rose Bengal using visible-light. This protocol offers a “green” and straightforward synthesis of pyrrolo[2,1-*a*]isoquinolines starting from readily available maleimides and tetrahydroisoquinolines. Further investigations to expand the scope and potential of this methodology are underway in our laboratory.

Supporting Information

Supporting Information File 1

Experimental details and characterization of the synthesized compounds.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-10-122-S1.pdf>]

References

- Bailly, C. *Curr. Med. Chem.: Anti-Cancer Agents* **2004**, *4*, 363–378. doi:10.2174/1568011043352939
- Handy, S. T.; Zhang, Y. *Org. Prep. Proced. Int.* **2005**, *37*, 411–445. doi:10.1080/00304940509354977
- Fan, H.; Peng, J.; Hamann, M. T.; Hu, J.-F. *Chem. Rev.* **2008**, *108*, 264–287. doi:10.1021/cr078199m
- Pla, D.; Albericio, F.; Alvarez, M. *Anti-Cancer Agents Med. Chem.* **2008**, *8*, 746–760. doi:10.2174/187152008785914789
- Marco, E.; Laine, W.; Tardy, C.; Lansiaux, A.; Iwao, M.; Ishibashi, F.; Bailly, C.; Gago, F. *J. Med. Chem.* **2005**, *48*, 3796–3807. doi:10.1021/jm049060w
- Reddy, M. V. R.; Rao, M. R.; Rhodes, D.; Hansen, M. S. T.; Rubins, K.; Bushman, F. D.; Venkateswarlu, Y.; Faulkner, D. *J. Med. Chem.* **1999**, *42*, 1901–1907. doi:10.1021/jm9806650
- Aubry, A.; Pan, X.-S.; Fisher, L. M.; Jarlier, V.; Cambau, E. *Antimicrob. Agents Chemother.* **2004**, *48*, 1281–1288. doi:10.1128/AAC.48.4.1281-1288.2004
- Reddy, S. M.; Srinivasulu, M.; Satnarayana, N.; Kondapi, A. K.; Venkateswarlu, Y. *Tetrahedron* **2005**, *61*, 9242–9247. doi:10.1016/j.tet.2005.07.067
- Quesada, A. R.; Garcia Grávalos, M. D.; Fernández Puentes, J. L. *Br. J. Cancer* **1996**, *74*, 677–682. doi:10.1038/bjc.1996.421
- Heim, A.; Terpin, A.; Steglich, W. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 155–156. doi:10.1002/anie.199701551
- Ploypradith, P.; Mahidol, C.; Sahakitpichan, P.; Wongbundit, S.; Ruchirawat, S. *Angew. Chem., Int. Ed.* **2004**, *43*, 866–868. doi:10.1002/anie.200352043
- Boger, D. L.; Boyce, C. W.; Labroli, M. A.; Sehon, C. A.; Jin, Q. *J. Am. Chem. Soc.* **1999**, *121*, 54–62. doi:10.1021/ja982078+
- Banwell, M. G.; Flynn, B. L.; Stewart, S. G. *J. Org. Chem.* **1998**, *63*, 9139–9144. doi:10.1021/jo9808526
- Handy, S. T.; Zhang, Y.; Bregman, H. *J. Org. Chem.* **2004**, *69*, 2362–2366. doi:10.1021/jo0352833
- Ploypradith, P.; Kagan, R. K.; Ruchirawat, S. *J. Org. Chem.* **2005**, *70*, 5119–5125. doi:10.1021/jo050388m
- Ohta, T.; Fukuda, T.; Ishibashi, F.; Iwao, M. *J. Org. Chem.* **2009**, *74*, 8143–8153. doi:10.1021/jo901589e
- Gupton, J. T.; Clough, S. C.; Miller, R. B.; Lukens, J. R.; Henry, C. A.; Kanters, R. P. F.; Sikorski, J. A. *Tetrahedron* **2003**, *59*, 207–215. doi:10.1016/S0040-4020(02)01475-8
- Fujikawa, N.; Ohta, T.; Yamaguchi, T.; Fukuda, T.; Ishibashi, F.; Iwao, M. *Tetrahedron* **2006**, *62*, 594–604. doi:10.1016/j.tet.2005.10.014
- Chen, L.; Xu, M.-H. *Adv. Synth. Catal.* **2009**, *351*, 2005–2012. doi:10.1002/adsc.200900287
- Yadav, J. S.; Gayathri, K. U.; Reddy, B. V. S.; Prasad, A. R. *Synlett* **2009**, 43–46. doi:10.1055/s-0028-1087387

21. Najera, C.; Sansano, J. M. *Curr. Org. Chem.* **1998**, *7*, 1105–1150. doi:10.2174/1385272033486594
22. Ishibashi, F.; Miyazaki, Y.; Iwao, M. *Tetrahedron* **1997**, *53*, 5951–5962. doi:10.1016/S0040-4020(97)00287-1
23. Banwell, M. G.; Flynn, B. L.; Hockless, D. C. R. *Chem. Commun.* **1997**, 2259–2260. doi:10.1039/a705874h
24. Cironi, P.; Manzanares, I.; Albericio, F.; Álvarez, M. *Org. Lett.* **2003**, *5*, 2959–2962. doi:10.1021/ol0351192
25. Ploypradith, P.; Petchmanee, T.; Sahakitpichan, P.; Litvinas, N. D.; Ruchirawat, S. *J. Org. Chem.* **2006**, *71*, 9440–9448. doi:10.1021/jo061810h
26. Grigg, R.; Heaney, F. J. *Chem. Soc., Perkin Trans. 1* **1989**, 198–200. doi:10.1039/p19890000198
27. Su, S.; Porco, J. A., Jr. *J. Am. Chem. Soc.* **2007**, *129*, 7744–7745. doi:10.1021/ja072737v
28. Yu, C.; Zhang, Y.; Zhang, S.; Li, H.; Wang, W. *Chem. Commun.* **2011**, *47*, 1036–1038. doi:10.1039/c0cc03186k
29. Zou, Y.-Q.; Lu, L.-Q.; Fu, L.; Chang, N.-J.; Rong, J.; Chen, J.-R.; Xiao, W.-J. *Angew. Chem., Int. Ed.* **2011**, *50*, 7171–7175. doi:10.1002/anie.201102306
30. Huang, L.; Zhao, J. *Chem. Commun.* **2013**, *49*, 3751–3753. doi:10.1039/c3cc41494a
31. Guo, S.; Zhang, H.; Huang, L.; Guo, Z.; Xiong, G.; Zhao, J. *Chem. Commun.* **2013**, *49*, 8689–8691. doi:10.1039/c3cc44486d
32. Wang, H.-T.; Lu, C.-D. *Tetrahedron Lett.* **2013**, *54*, 3015–3018. doi:10.1016/j.tetlet.2013.04.004
33. Zeitler, K. *Angew. Chem., Int. Ed.* **2009**, *48*, 9785–9789. doi:10.1002/anie.200904056
34. Yoon, T. P.; Ischay, M. A.; Du, J. *Nat. Chem.* **2010**, *2*, 527–532. doi:10.1038/nchem.687
35. Narayanam, J. M. R.; Stephenson, C. R. J. *Chem. Soc. Rev.* **2011**, *40*, 102–113. doi:10.1039/b913880n
36. Xuan, J.; Xiao, W.-J. *Angew. Chem., Int. Ed.* **2012**, *51*, 6828–6838. doi:10.1002/anie.201200223
37. Shi, L.; Xia, W. *Chem. Soc. Rev.* **2012**, *41*, 7687–7697. doi:10.1039/c2cs35203f
38. Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, *113*, 5322–5363. doi:10.1021/cr300503r
39. Hu, J.; Wang, J.; Nguyen, T. H.; Zheng, N. *Beilstein J. Org. Chem.* **2013**, *9*, 1977–2001. doi:10.3762/bjoc.9.234
40. Ravelli, D.; Fagnoni, M. *ChemCatChem* **2012**, *4*, 169–171. doi:10.1002/cctc.201100363
41. Ravelli, D.; Fagnoni, M.; Albini, A. *Chem. Soc. Rev.* **2013**, *42*, 97–113. doi:10.1039/c2cs35250h
42. Nicewicz, D. C.; Nguyen, T. M. *ACS Catal.* **2014**, *4*, 355–360. doi:10.1021/cs400956a
43. Liu, H.; Feng, W.; Kee, C. W.; Zhao, Y.; Leow, D.; Pan, Y.; Tan, C.-H. *Green Chem.* **2010**, *12*, 953–956. doi:10.1039/b924609f
44. Pan, Y.; Kee, C. W.; Chen, L.; Tan, C.-H. *Green Chem.* **2011**, *13*, 2682–2685. doi:10.1039/c1gc15489c
45. Pan, Y.; Wang, S.; Kee, C. W.; Dubuisson, E.; Yang, Y.; Loh, K. P.; Tan, C.-H. *Green Chem.* **2011**, *13*, 3341–3344. doi:10.1039/c1gc15865a
46. Neumann, M.; Földner, S.; König, B.; Zeitler, K. *Angew. Chem., Int. Ed.* **2011**, *50*, 951–954. doi:10.1002/anie.201002992
47. Hari, D. P.; König, B. *Org. Lett.* **2011**, *13*, 3852–3855. doi:10.1021/ol201376v
48. Liu, Q.; Li, Y.-N.; Zhang, H.-H.; Chen, B.; Tung, C.-H.; Wu, L.-Z. *Chem.–Eur. J.* **2012**, *18*, 620–627. doi:10.1002/chem.201102299
49. Fidaly, K.; Ceballos, C.; Falguières, A.; Veitia, M. S.-I.; Guy, A.; Ferroud, C. *Green Chem.* **2012**, *14*, 1293–1297. doi:10.1039/c2gc35118h
50. Fu, W.; Guo, W.; Zou, G.; Xu, C. *J. Fluorine Chem.* **2012**, *140*, 88–94. doi:10.1016/j.jfluchem.2012.05.009
51. Hari, D. P.; Schroll, P.; König, B. *J. Am. Chem. Soc.* **2012**, *134*, 2958–2961. doi:10.1021/ja212099r
52. Hari, D. P.; Hering, T.; König, B. *Org. Lett.* **2012**, *14*, 5334–5337. doi:10.1021/ol302517n
53. Neumann, M.; Zeitler, K. *Org. Lett.* **2012**, *14*, 2658–2661. doi:10.1021/ol3005529
54. Hamilton, D. S.; Nicewicz, D. A. *J. Am. Chem. Soc.* **2012**, *134*, 18577–18580. doi:10.1021/ja309635w
55. Rueping, M.; Vila, C.; Bootwicha, T. *ACS Catal.* **2013**, *3*, 1676–1680. doi:10.1021/cs400350j
56. Grandjean, J.; Nicewicz, D. A. *Angew. Chem., Int. Ed.* **2013**, *52*, 3967–3971. doi:10.1002/anie.201210111
57. Riener, M.; Nicewicz, D. A. *Chem. Sci.* **2013**, *4*, 2625–2629. doi:10.1039/c3sc50643f
58. Wilger, D. J.; Gesmundo, N. J.; Nicewicz, D. A. *Chem. Sci.* **2013**, *4*, 3160–3165. doi:10.1039/c3sc51209f
59. Nguyen, T. M.; Nicewicz, D. A. *J. Am. Chem. Soc.* **2013**, *135*, 9588–9591. doi:10.1021/ja4031616
60. Perkowski, A. J.; Nicewicz, D. A. *J. Am. Chem. Soc.* **2013**, *135*, 10334–10337. doi:10.1021/ja4057294
61. Pitre, S. P.; McTiernan, C. D.; Ismaili, H.; Scaiano, J. C. *J. Am. Chem. Soc.* **2013**, *135*, 13286–13289. doi:10.1021/ja406311g
62. Rueping, M.; Vila, C.; Koenings, R. M.; Poschary, K.; Fabry, D. C. *Chem. Commun.* **2011**, *47*, 2360–2362. doi:10.1039/c0cc04539j
63. Rueping, M.; Zhu, S.; Koenings, R. M. *Chem. Commun.* **2011**, *47*, 8679–8681. doi:10.1039/c1cc12907d
64. Rueping, M.; Leonori, D.; Poisson, T. *Chem. Commun.* **2011**, *47*, 9615–9617. doi:10.1039/c1cc13660g
65. Rueping, M.; Zhu, S.; Koenings, R. M. *Chem. Commun.* **2011**, *47*, 12709–12711. doi:10.1039/c1cc15643h
66. Rueping, M.; Zoller, J.; Fabry, D. C.; Poschary, K.; Koenings, R. M.; Weirich, T. E.; Mayer, J. *Chem.–Eur. J.* **2012**, *18*, 3478–3481. doi:10.1002/chem.201103242
67. Rueping, M.; Koenings, R. M.; Poschary, K.; Fabry, D. C.; Leonori, D.; Vila, C. *Chem.–Eur. J.* **2012**, *18*, 5170–5174. doi:10.1002/chem.201200050
68. Rueping, M.; Vila, C.; Szadkowska, A.; Koenigs, R. M.; Fronert, J. *ACS Catal.* **2012**, *2*, 2810–2815. doi:10.1021/cs300604k
69. Zhu, S.; Rueping, M. *Chem. Commun.* **2012**, *48*, 11960–11962. doi:10.1039/c2cc36995h
70. Zhu, S.; Das, A.; Bui, L.; Zhou, H.; Curran, D. P.; Rueping, M. *J. Am. Chem. Soc.* **2013**, *135*, 1823–1829. doi:10.1021/ja309580a
71. Rueping, M.; Vila, C. *Org. Lett.* **2013**, *15*, 2092–2095. doi:10.1021/ol400317v
72. Vila, C.; Rueping, M. *Green Chem.* **2013**, *15*, 2056–2059. doi:10.1039/c3gc40587g
73. Tóth, J.; Váradi, L.; Dancsó, A.; Blaskó, G.; Töke, L.; Nyerges, M. *Synlett* **2007**, 1259–1263. doi:10.1055/s-2007-977461

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
[doi:10.3762/bjoc.10.122](https://doi.org/10.3762/bjoc.10.122)