



COACHING & SPORT SCIENCE REVIEW

The Official Tennis Coaching and Sport Science
Publication of the International Tennis Federation
24th Year, Issue 68, April 2016

www.itftennis.com/coaching/sportsscience
ISSN: 2225-4757

Pg.	Title / Author
2	Editorial
3	Coach-player communication: What, why, when and how? Janet Young (AUS)
5	Social media, learning, networking and promotion opportunities for tennis coaches Mauricio Cordova (ECU)
8	Specific nature of resistance training in tennis Josep Campos Rius and David Suarez (ESP)
11	Psychological factors related to choking under pressure Takehiro Iwatsuki (JAP) , Judy Van Raalte (USA) , Britton Brewer (USA) , Albert Petitpas (USA) and Masanori Takahashi (JAP)
13	Muscular imbalance and their effect on the health of the tennis player: Assessments of explosive strength in the lower limbs by means of jumps Javier Villaplana Velasco and Cristina Blasco Lafarga (ESP)
15	Is there a relationship between junior and senior tennis success? Michael Bane, Bruce Elliott and Machar Reid (AUS)
18	Helping junior tennis players to cope with their emotions Camilla Knight (GBR) , Francesca Lewis (GBR) and Stephen Mellalieu (GBR)
21	Enjoyment: Fresh clues for coaches Janet Young (AUS)
24	Tennis player development: From junior to professional transition stage Pancho Alvaríño, José Altur and Miguel Crespo (ESP)
27	The evolution of online education for sports coaches: A tennis perspective Richard Sackey-Addo (GBR) and Javier Perez Camarero (ESP)
29	Recommended e-books Editors
30	Recommended web links Editors
31	Guidelines for submitting articles to ITF Coaching & Sport Science Review Editors

CONTENTS ISSUE 68

Muscular imbalances and their impact on the health of the tennis player: Assessments of explosive strength in the lower limbs by means of jumps

Javier Villaplana Velasco and Cristina Blasco Lafarga (ESP)

ITF Coaching and Sport Science Review 2015; 68 (24): 13 - 14

ABSTRACT

This paper discusses the existence of asymmetry in tennis players lower limbs, if these asymmetries impact on determining capabilities as Explosive Strength and Elastic Explosive Strength, and if so, how they are affected by fatigue. This paper holds the hypothesis that there exist differences in the capabilities of useful strength in the lower limb muscles, and that these muscular differences or asymmetries are the source of injury for developing tennis players.

Key words: tennis, asymmetry, jumps, injury prevention, health

Corresponding author: javilla@gmail.com

Article received: 08 January 2016

Article accepted: 23 January 2016

INTRODUCTION

In order to continue being successful in competition, players must accelerate, decelerate, change direction, move quickly, and keep balance while repeatedly producing optimal strokes until the end (Girard, Millet, 2008). As a consequence of these characteristics, literature shows the presence of injuries. Even asymmetry, another essential characteristic of tennis, is admitted to be a source of injury. Pluim, Staal, Windler, Jayanthi (2006) express in their revision that tennis is characterized by varied injuries, most of them in the lower limbs. Confirming this, Hjelm, Werner, Renstrom (2012) literature states that lower limb injuries are the most frequent, reaching 51% of the total. As to asymmetry, we have found papers dealing with torso and upper limb asymmetry, but not lower limb related. Even so, we can infer the existence of lower limb asymmetry, Carpes, Mota, Faria (2010) explain the bilateral asymmetry even in cyclic activities such as continuous sprinting and pedalling, and its risk of injuries in healthy athletes. This makes us think that if in cyclic sports there is asymmetry in lower limbs, in acyclic and asymmetric sports such as tennis, this situation has to be valid too.

Within this context, the objective of the following research is a) to prove the existence of lower limb imbalance in developing tennis players, and, b) if so, to reflect on the consequences of muscular imbalance of the lower limbs and their possible impact on the health of the tennis player.

METHODOLOGY

Sample

There were 5 participants in the study, all of them were males, with media values in age between 14,2 (0,84), height 170,2 (6,06) and weight 59 (8,89). Their weekly training volume was 26 hours, from Monday to Friday, divided into three sessions, two in the morning and one in the afternoon. The morning sessions included 2h30' of on-court training, and 1h30' of physical training. The afternoon session included on-court training for 1h30'. There was no afternoon session on Wednesday.

Procedure

The subjects were analyzed in December 2012. All of them warmed-up for 10 minutes, 5 minutes of continuous running and dynamic movement exercises (Meylan, Nosaka, Green, Cronin, 2010). No static stretching due to the negative impact demonstrated in previous studies in the different variables of the jumps (Meylan et al., 2010).

The exercises were SJ (squat jump) to evaluate explosive strength and CMJ (counter-movement jump) to evaluate the elastic explosive strength.

Each exercise consisted of 9 attempts (18 jumps total), which were distributed as follows: 3 attempts with the non-dominant leg (ND), 3 attempts with the dominant leg (D) and 3 with both legs

simultaneously (2PR). The recovery time between each jump of the same type was 30s. The recovery time between 3 jump blocks was 120s. (Meylan et al., 2010).

The test was made just before beginning the morning tennis training and was repeated after finishing it. This second time there was no warming-up.

Each participant received a questionnaire so as to know age, sex, type and degree of the injury they could have suffered during the previous year.

Data treatment and analysis

The tests were evaluated by means of photoelectric cells (Optojump next, Micrograte Srl, Bolzano, Italy) and the data gathered were entered into SPSS.20 for statistics, setting (pre test and post test) fatigue, kinetic chain (non-dominant leg jump, dominant leg jump and both leg jumps), and the type of jump (SJ and CMJ) as independent intra subject variables. On the other hand, the athlete's health was taken as an independent inter subject variable. Height (cm) was taken as a dependant variable.

The statistic treatment that was used was an analysis of the variance (anova) for inter group comparisons. The greatest jumps per exercise, and per athlete were analysed. The significance level was set at $p < 0.05$.

RESULTS

Figure 1 shows the differences of the effect on fatigue, considering explosive strength and elastic explosive strength.




Figure 1. Graphic representation of the impact of fatigue on the type of jump. 1= SJ; 2= CMJ

After making the normality test (Kolgomorov-Smirnov) and sphericity test (Mauchly) the results of the anova intrasubject contrast test with repeated measurements, show significance for all factors analysed and their interactions.

Table 1 demonstrates that this significance is in favour of the pre-test, by comparing the differences between the total jumps prior to training with the same number of jumps when at the end of training. It also compares the difference between both types of jumps. Finally, it permits the analysis of the difference depending on the muscles involved.

		Media differences (I-I)	Error t.	P	IC (95 %)	
					Lower lim.	Upper lim.
Jump						
Total pre SJ	Total Post CMJ	1.677 * -0.890*	0.175 0.3	0.001 0.041	1.192 -1.1724	2.161 -0.056
Chain						
ND	D	-1.495*	0.174	0.003	-2.185	-0.805
	2PR	-12.200*	1.044	0.001	-16.335	-8.065
D	2PR	-10.705*	1.094	0.002	-15.036	-6.374

Table 1. Results of the comparison per pairs (Bonferroni adjustment) for the fatigue factors, jump and chain, considered in isolation.

When analysing the effect of fatigue on jumps, we observe that both exercises show training very significantly. And when analysing the chain, we note that both, for the non-dominant leg and for the two feet exercise, there were significant differences, while in the case of the dominant leg, the loss, though important, is not significant (table 2).

	Pre	Post	Media Differences (I-I)	Error t.	P	IC (95 %)	
						Lower lim.	Upper lim.
SJ	1	2	2.147*	0.241	0.001	1.478	2.816
CMJ	1	2	1.207*	0.207	0.004	0.632	1.781
ND	1	2	1.040*	0.315	0.03	0.165	1.915
D	1	2	0.85	0.378	0.088	-0.199	1.899
2PR	1	2	3.140*	0.213	0.000	2.549	3.731

Table 2. Results of the comparison per pairs (Bonferroni adjustment) for the fatigue*jump; fatigue*chain interactions.

Table 3 relates the interactions between fatigue, chain and jump, observing that for the SJ there is significance depending on fatigue for all kinetic chains proposed, however, for CMJ fatigue is only significant when jumping on two feet.

	Chain	(0)fatigue	(1)fatigue	Media differences (I-I)	Error tip.	p	IC (95%)	
							Lower lim.	Upper lim.
SJ	ND	pre	post	1,380*	0,292	0,009	0,569	2,191
	D	pre	post	1,860*	0,445	0,014	0,626	3,094
	2PR	pre	post	3,200*	0,305	0,000	2,353	4,047
CMJ	ND	pre	post	0,7	0,454	0,198	-0,56	1,96
	D	pre	post	-0,16	0,518	0,773	-1,599	1,279
	2PR	pre	post	3,080*	0,146	0,000	2,674	3,486

Table 3. Results of the comparison per pairs (Bonferroni adjustment) for the fatigue*chain* jump interactions.

DISCUSSION

In spite of the limitations of the sample, this paper confirms the hypothesis of the existence of asymmetry in the tennis players lower limbs. From our bibliographic search, it seems that this pilot study is the first one to show that asymmetry, since previous studies

centred on tennis players have studied the assessment of the trunk (Ellenbecker, Roetert, 2004) (Sanchis-Moysi, Idoate, Dorado, Alayon, Calbet, 2010). Some of these papers, Sanchis-Moysi et al. (2010) already say that the asymmetry of the trunk may be due to differences in the kinetic chain of the lower limbs, but none has evaluated those muscles directly. Later Sanchis-Moysi, Idoate, Izquierdo, Calbet, Dorado (2011) verified the existence of asymmetry in lower limbs for the muscles of the gluteus and iliopsoas but using magnetic resonance.

This functional muscular asymmetry has been related to the health of the athletes and their future risk of injuries (Menzel et al., 2012); the level of fatigue has been confirmed to be directly related to this risk (Goodall, Pope, Coyle, Neumayer, 2012). The results of our study confirm the loss of the explosive strength (SJ) and the elastic explosive strength (CMJ), after finishing training, either considering the jumps in isolation or together. This confirms that fatigue reduces the response capability of young players, increasing the risk of injury in the final part of training, or greater duration and intensity efforts.

What interested us most was what happened when comparing the dominant with the non-dominant leg. In this sense, we have found that when considering the total number of jumps with each modality, the loss of height has been significant, both, for the jump with the non-dominant leg, as well as for the jump with both legs, while there have been no significant differences for the dominant leg, when measuring pre- and post training. This makes us think that the fatigue of the non-dominant leg may be determining and may affect directly the jump with two legs. This disturbance in the application of the strength could be related to the injuries that occur as a consequence of the explosive actions in tennis.

REFERENCES

- Carpes, F. P., Mota, C. B., Faria, I. E. (2010). On the bilateral asymmetry during running and cycling - a review considering leg preference. *Phys Ther Sport*, 11(4), 136-142. doi: 10.1016/j.ptsp.2010.06.005
- Ellenbecker, T. S., Roetert, E. P. (2004). An isokinetic profile of trunk rotation strength in elite tennis players. *Med Sci Sports Exerc*, 36(11), 1959-1963.
- Girard, O., Millet, G. P. (2008). Neuromuscular fatigue in racquet sports. *Neurol Clin*, 26(1), 181-194; x. doi: 10.1016/j.ncl.2007.11.011
- Goodall, R. L., Pope, R. P., Coyle, J. A., Neumayer, R. (2012). Balance and agility training does not always decrease lower limb injury risks: a cluster-randomised controlled trial. *Int J Inj Contr Saf Promot*. doi: 10.1080/17457300.2012.717085
- Hjelm, N., Werner, S., Renstrom, P. (2012). Injury risk factors in junior tennis players: a prospective 2-year study. *Scand J Med Sci Sports*, 22(1), 40-48. doi: 10.1111/j.1600-0838.2010.01129.x
- Menzel, H. J., Chagas, M. H., Szmuchrowski, L. A., Araujo, S. R., de Andrade, A. G., de Jesus, F. R. (2012). Analysis of Lower Limb Asymmetries by Isokinetic and Vertical Jump Tests in Soccer Players. *J Strength Cond Res*. doi: 10.1519/JSC.obo13e318265a3c8
- Meylan, C. M., Nosaka, K., Green, J., Cronin, J. B. (2010). Temporal and kinetic analysis of unilateral jumping in the vertical, horizontal, and lateral directions. *J Sports Sci*, 28(5), 545-554. doi: 10.1080/02640411003628048
- Pluim, B. M., Staal, J. B., Windler, G. E., Jayanthi, N. (2006). Tennis injuries: occurrence, aetiology, and prevention. *Br J Sports Med*, 40(5), 415-423. doi: 10.1136/bjism.2005.023184
- Sanchis-Moysi, J., Idoate, F., Dorado, C., Alayon, S., Calbet, J. A. (2010). Large asymmetric hypertrophy of rectus abdominis muscle in professional tennis players. *PLoS One*, 5(12), e15858. doi: 10.1371/journal.pone.0015858
- Sanchis-Moysi, J., Idoate, F., Izquierdo, M., Calbet, J. A., Dorado, C. (2011). Iliopsoas and gluteal muscles are asymmetric in tennis players but not in soccer players. *PLoS One*, 6(7), e22858. doi: 10.1371/journal.pone.0022858

RECOMMENDED ITF TENNIS COACH CONTENT (CLICK BELOW)

Tennis*i*Coach