

Journal section: *Oral Medicine and Pathology*

doi:10.4317/jced.3.e166

Publication Types: *Case Report*

Oral Squamous Cell Carcinoma Associated with a Dental Implant – a case report and literature review

Mohammad Moshref¹, Abdolreza Jamilian², Ali Lotfi³, Rahman Showkatbakhsh⁴

¹ DDS; MS. Associate professor, Department of oral and maxillofacial pathology, Shahid Beheshti university of medical sciences.

² DDS; MS. Associate professor, Islamic Azad university. Fellow of Orthognathic surgery, Shahid Beheshti university of medical sciences.

³ DDS; MS. Assistant professor, Department of oral and maxillofacial pathology, Shahid Beheshti university of medical sciences.

⁴ DDS; MS. Associate professor, Department of Orthodontics, Director of Orthognathic surgery fellowship, Shahid Beheshti university of medical sciences.

Correspondence:

No 2713, jam tower next to jame-jam

Vali Asr St. Tehran 1966843133,

IRAN

E-mail: info@jamilian.net

Received: 08/07/2010

Accepted: 26/08/2010

Moshref M, Jamilian A, Lotfi A, Showkatbakhsh R. Oral Squamous Cell Carcinoma Associated with a Dental Implant – a case report and literature review. *J Clin Exp Dent*. 2011;3(2):e166-8.

<http://www.medicinaoral.com/odo/volumenes/v3i2/jcedv3i2p166.pdf>

Article Number: 50350 <http://www.medicinaoral.com/odo/indice.htm>
© Medicina Oral S. L. C.I.F. B 96689336 - eISSN: 1989-5488
eMail: jced@jced.es

Abstract

Objectives: Prosthodontic rehabilitation using dental implants has become a common practice in dentistry at the present time. Although severe complications related to dental osseointegrated implants are uncommon, in recent years several cases of oral squamous cell carcinoma adjacent to dental implants have been published.

Study Design: A 67-year-old edentulous woman developed an oral squamous cell carcinoma around right mandibular implant about 12 months after receiving dental implants. The treatments involve chemotherapy, explantation of the implants and radiotherapy.

Result and Conclusions: The use of endosseous implants has been associated with a low risk for the development of cancer. However, this case report serves as evidence to the importance of careful screening at routine dental examinations especially if the patient suffers from peri-implantitis, any mucosal erosion or discoloration, leukoplakia around implant, pain, implant loosening, bone loss around implant, mucosal erythema, and soft tissue inflammation.

Key words: *Dental Implant, Squamous Cell Carcinoma, Neoplasm, Cancer, Malignancy.*

Introduction

Endosseous dental implants have become a predictable and common practice for the replacement of missing teeth. The overall success rate for dental implants is more than 90 percent (1). A wide variety of metallic, ceramic, and polymeric synthetic biomaterials have been used for prosthetic implants in the human body. Both the safety and effectiveness of this technique have been extensively demonstrated in the literature (2). Nevertheless, the number of complications associated to dental implants has increased proportionally to the popularization of its employment (3). Complications, which are uncommon, include postoperative infection, peri-implantitis, implant fracture, bone loss, and failure of osseointegration with loosening of the implant. Recently, some reports have also been published about primary squamous cell carcinoma of oral mucosa (4-7), primary osteosarcoma in jaws (8), and metastatic carcinoma (9) in the vicinity of the implants.

The case report we present here describes a patient with squamous cell carcinoma around her dental implants.

Case Report

A diabetic 67-year-old woman sought treatment for pain and swelling of four months duration involving the right mandible in the distal area of the canines. 15 years previously the patient started wearing full denture. 16 months ago she referred to a dentist in order to fix her loose denture and she also complained from denture induced alveolar mucosal inflammation in the right mandible. Her dentist surgically placed two 6.0 × 13 millimeter dental implant (Nobel Biocare, Göteborg, Sweden) in the edentulous canines' area of the mandible; afterwards, the dentist placed the overdenture abutments onto the implants. (Figure 1) 12 months after the implants insertion she started having pain in the right mandibular canine area; which was so aggravating that prevented her from wearing her overdenture. Her dentist prescribed antibiotics; however, when the lesion did not respond to initial antibiotic therapy, the patient's dentist performed an incisional biopsy to establish a definitive diagnosis. Thus, she was referred to the oral and maxillofacial pathology department of a dental school for a more definitive diagnosis and treatment. She had had the lesions for two months. A fellow of orthognathic surgery recorded her medical and dental history, which included no smoking or alcohol consumption; however, she suffered from diabetes and high blood pressure both of which were under medical control. An exophytic mucosal mass overlying alveolar mucosa and extending from anterior floor of mouth to ventral surface of the tongue was apparent. (Fig. 1) Peri-implantitis, implant loosening, mucosal erythema, inflammation of the soft tissue, and bone loss around right mandibular implant were evident. The right submandibular lymph nodes were easily palpable



Fig. 1. Intra oral photo of the patient

by finger touch. An excisional biopsy was performed and the tissue was submitted for pathological evaluation to the department of oral and maxillofacial pathology.

Pathological findings: Microscopic examination of the submitted biopsy specimen revealed a moderately differentiated squamous cell carcinoma showing hyperchromatism of the nuclei, keratin pearl formation, pleomorphism, and numerous mitotic figures (Figures 2, 3).




Fig. 2. A, Section from the biopsy taken of the oral lesion (magnification ×4)




Fig. 3. A, Section from the biopsy taken of the oral lesion (magnification ×40)

Therapy: The patient received three initial courses of chemotherapy. Afterwards, the implants were explanted immediately. The tumor along with some local blood vessels and muscles were removed by pull-through technique. 34 lymph nodes were also removed, 3 of which were metastatic. The patient was then scheduled for radiotherapy.

Discussion

The development of neoplasia associated with alloplastic implant materials is a rare complication that a number of clinicians and researchers have reported, primarily in the orthopedic literature (10-11).

Although the materials composing these implants are relatively biologically inert and nontoxic they may induce a variety of tissue responses (12). By far the most serious complication of orthopaedic hardware and prosthetic material is malignancy. Animal studies analyzing the biological effects of the components of prosthetic joints revealed carcinogenic properties of beryllium, cadmium, chromium, cobalt, iron, lead, nickel, selenium, zinc, and titanium (13). In humans, prostheses containing cobalt, chromium, stainless steel, nickel, iron, manganese and silicon have been implicated as being carcinogenic. Other authors state that the placement of an implant may induce osteonecrosis, which in itself is a risk factor for the development of a sarcoma (8, 14).

Dental implants have also been of no exception. An electronic review of the literature suggests that an exhaustive differential diagnosis is required in implant patients who refer to dentist complaining from lesions, severe pain, halitosis, swelling and gum irritation, implant mobility, and numbness or paresthesia (1-9). Sometimes, the initial clinical appearance of lesions suggested peri-implantitis (7) with a red, hyperplastic or even ulcerated appearance while the patient was suffering from cancer. Detection of the oral lesions has great importance as they may be diagnosed first by the patient's dentist or by the maxillofacial surgeon. The lesion can indicate the presence of a primary squamous cell carcinoma, osteosarcoma and metastasis (9). Metastatic tumors of the jaws, from distant organs and tissues, may be diagnosed first by the oral surgeon, and most of the patients with oral metastasis may have seen another dentist recently but the condition has not been diagnosed (15).

Chronic inflammation, smoking and alcohol consumption as well as premalignant lesions and conditions such as leukoplakia, and oral lichen planus (OLP), have been recognized to be risk factors for the development of malignancy. It has also been reported that the majority of all presented cases had at least one recognized risk factor for oral cancer (4).

McGuff et al. have reported that there appears to be relation between implant-related sarcomas and sex of the patients. They also reported that the interval from im-

plant placement to the development of osteosarcoma has ranged from six months to 30 years, with a mean of nine years. (8)

In this case report we have presented an unusual patient who developed a SCC adjacent to a mandibular implant. The lesion described in this report initially was confused with peri-implantitis. In implant patients or implant candidates who suffer from peri-implantitis, any mucosal erosion or discoloration, leukoplakia around implant, pain, implant loosening, bone loss around implant, mucosal erythema, and soft tissue inflammation regular follow-up and careful exploration of the area is essential

References

1. Lekholm U, Gröndahl K, Jemt T. Outcome of oral implant treatment in partially edentulous jaws followed 20 years in clinical function. *Clin Implant Dent Relat Res* 2006;8:178-86.
2. Lemmerman KJ, Lemmerman NE. Osseointegrated dental implants in private practice: a long-term case series study. *J Periodontol* 2005;76:310-9.
3. McDermott NE, Chuang SK, Woo VV, Dodson TB. Complications of dental implants: identification, frequency, and associated risk factors. *Int J Oral Maxillofac Implants* 2003;18:848-55.
4. Czerninski R, Kaplan I, Almozino G, Maly A, Regev E. Oral squamous cell carcinoma around dental implants. *Quintessence Int* 2006;37:707-11.
5. Gallego L, Junquera L, Baladron J, Villarreal P. Oral squamous cell carcinoma associated with symphyseal dental implants: an unusual case report. *J Am Dent Assoc* 2008;139:1061-5.
6. Schache A, Thavaraj S, Kalavrezos N. Osseointegrated implants: a potential route of entry for squamous cell carcinoma of the mandible. *Br J Oral Maxillofac Surg* 2008;46:397-9.
7. Eguia del Valle A, Martinez-Conde Llamas R, Lopez Vicente J, Uribarri Etxebarria A, Aguirre Urizar JM. Primary oral squamous cell carcinoma arising around dental osseointegrated implants mimicking peri-implantitis. *Med Oral Patol Oral Cir Bucal* 2008;13:E489-91.
8. McGuff HS, Heim-Hall J, Holsinger FC, Jones AA, O'Dell DS, Hafemeister AC. Maxillary osteosarcoma associated with a dental implant: report of a case and review of the literature regarding implant-related sarcomas. *J Am Dent Assoc* 2008;139:1052-9.
9. Dib LL, Soares AL, Sandoval RL, Nannmark U. Breast metastasis around dental implants: a case report. *Clin Implant Dent Relat Res* 2007;9:112-5.
10. Visuri T, Pulkkinen P, Paavolainen P. Malignant tumors at the site of total hip prosthesis. Analytic review of 46 cases. *J Arthroplasty* 2006;21:311-23.
11. Kirkpatrick CJ, Alves A, Kohler H, Kriegsmann J, Bittinger F, Otto M, et al. Biomaterial-induced sarcoma: A novel model to study pre-neoplastic change. *Am J Pathol* 2000;156:1455-67.
12. Jacobs JJ, Skipor AK, Black J, Urban R, Galante JO. Release and excretion of metal in patients who have a total hip-replacement component made of titanium-base alloy. *J Bone Joint Surg Am* 1991;73:1475-86.
13. Heath JC, Freeman MA, Swanson SA. Carcinogenic properties of wear particles from prostheses made in cobalt-chromium alloy. *Lancet* 1971;1:564-6.
14. Adams JE, Jaffe KA, Lemons JE, Siegal GP. Prosthetic implant associated sarcomas: a case report emphasizing surface evaluation and spectroscopic trace metal analysis. *Ann Diagn Pathol* 2003;7:35-46.
15. Epker BN, Merrill RG, Henny FA. Breast adenocarcinoma metastatic to the mandible. Report of seven cases. *Oral Surg Oral Med Oral Pathol* 1969;28:471-9.