

ENTHALPY OF SOLVATION CORRELATIONS FOR GASEOUS SOLUTES
DISSOLVED IN DIMETHYL SULFOXIDE AND PROPYLENE CARBONATE
BASED ON THE ABRAHAM MODEL

Christina Mintz^a, Katherine Burton^a, William E. Acree, Jr.^{a*} and Michael H. Abraham^b

^a Department of Chemistry, P. O. Box 305070, University of North Texas,
Denton, TX 76203-5070 (U.S.A.)

^b Department of Chemistry, University College London, 20 Gordon Street,
London, WC1H 0AJ (U.K.)

Abstract

Introduction

The thermodynamic properties of molecules in various chemical and biological processes are greatly influenced by molecular interactions between the molecule and its solubilizing media. Such interactions may be either non-specific or specific in nature. Nonspecific interactions are described by a random distribution of molecules throughout the entire solution. Specific interactions, on the other hand, are generally much stronger and often result in a specific geometric orientation of one molecule with respect to an adjacent molecule. Even in systems known to contain specific interactions, the need to properly account for nonspecific interactions has been long recognized.

In previous studies (1-6) we have used the Abraham solvation parameter model to describe mathematically the various solute-solvent interactions found in a wide-range of different chemical and biological systems. We have reported partition coefficient correlations (both water-to-organic solvent and gas-to-organic solvent) for more than fifty common organic solvents. Our published correlations have for the most part pertained to 298.15 K. Manufacturing and biological processes are not restricted to 298.15 K, and there is a need to estimate partitioning properties in water and in organic solvents at other temperatures as well. From a thermodynamic standpoint, the gas-to-condensed phase partition coefficient, K , and water-to-organic solvent partition coefficient, P , can be estimated

$$\log K(at T) - \log K(at 298.15 K) = \frac{-\Delta H_{Solv}}{R} (1/T - 1/298.15) \quad (1)$$

and

$$\log P(at T) - \log P(at 298.15 K) = \frac{-\Delta H_{trans}}{R} (1/T - 1/298.15) \quad (2)$$

at other temperatures from measured partition coefficient data at 298.15 K and the solute's enthalpy of solvation, ΔH_{Solv} , or enthalpy of transfer, ΔH_{trans} , between the two condensed phases. The enthalpy of transfer needed in Eqn. 2 is defined as

$$\Delta H_{\text{trans}} = \Delta H_{\text{Solv,Org}} - \Delta H_{\text{Solv,W}} \quad (3)$$

the difference in the enthalpy of solvation of the solute in the specified organic solvent minus its enthalpy of solvation in water. The above equations assume zero heat capacity changes.

Mintz *et al.* (11) recently developed the correlation equations for enthalpies of solvation in water, $\Delta H_{\text{Solv,W}}$,

$$\begin{aligned} \Delta H_{\text{Solv,W}} \text{ (kJ/mole)} = & -13.310(0.457) + 9.910(0.814)\mathbf{E} + 2.836(0.807)\mathbf{S} \\ & - 32.010(1.102)\mathbf{A} - 41.816(0.781)\mathbf{B} - 6.354(0.200)\mathbf{L} \end{aligned} \quad (4)$$

(with N = 368, SD = 3.68, $R^2 = 0.964$, $R^2_{\text{adj}} = 0.964$, F = 1950.5)

$$\begin{aligned} \Delta H_{\text{Solv,W}} \text{ (kJ/mole)} = & -6.952(0.651) + 1.415(0.770)\mathbf{E} - 2.859(0.855)\mathbf{S} \\ & - 34.086(1.225)\mathbf{A} - 42.686(0.850)\mathbf{B} - 22.720(0.800)\mathbf{V} \end{aligned} \quad (5)$$

(with N = 369, SD = 4.04, $R^2 = 0.959$, $R^2_{\text{adj}} = 0.958$, F = 1688.2)

based on the Abraham model. Here and elsewhere, N corresponds to the number of solutes, R denotes the correlation coefficient, SD is the standard deviation and F corresponds to the Fisher F-statistic.

The independent variables in Eqns. 4 and 5 are as follows: **E** and **S** refer to the excess molar refraction and dipolarity/polarizability descriptors of the solute,

respectively, **A** and **B** are the solute overall or summation hydrogen bond acidity and basicity, **V** is the McGowan volume of the solute in units of $(\text{dm}^3 \text{ mol}^{-1})/100$, and **L** is the logarithm of the solute's gas phase dimensionless Ostwald partition coefficient into hexadecane at 298.15 K. The first four descriptors can be regarded as measures of the tendency of the given to undergo various solute-solvent interactions. The latter two descriptors, **V** and **L**, are both measures of solute size, and so will be measures of the solvent cavity term that will accommodate the dissolved solute. General dispersion interactions are also related to solute size, hence, both **V** and **L** will also describe the general solute-solvent interactions. The regression coefficients and constants (c , e , s , a , b , v and l) are obtained by regression analysis of experimental data for a given process (*i.e.*, a given partitioning processes, a given stationary phase and mobile phase combination, *etc.*).

As part of the study, Mintz *et al.* (11) also reported correlation equations for estimating enthalpies of solvation of organic solutes in 1-octanol. In follow-up studies the authors (12, 13) published the corresponding enthalpic equations for solutes dissolved in heptane, hexadecane, cyclohexane, benzene, toluene and carbon tetrachloride. Each of these latter correlations was based on experimental data for a minimum of 100 organic and inorganic solutes. Eventually, we hope to derive mathematical expressions for predicting enthalpies of solvation in most (if not all) of the organic solvents for which we have log K and log P correlations. The present study concerns developing ΔH_{Solv} equations for dimethyl sulfoxide and propylene carbonate. We have elected for the present time to correlate only enthalpies of solvation because there is considerably more experimental data available for ΔH_{Solv} than for ΔH_{trans} . To correlate enthalpies of transfer

one would need experimental enthalpy of solution data (or ΔH_{Solv} data) for the solute in both water and the organic solvent under consideration. One is more likely to find ΔH_{Solv} in water or the specified organic solvent, than to find both experimental values. Also, enthalpies of solvation allow one to estimate the temperature dependence of partition coefficients of solutes distributed between two immiscible organic solvents. Poole and coworkers [14-16] have recently reported partition coefficient data for neutral compounds between heptane and methanol, N,N-dimethylformamide, 2,2,2-trifluoroethanol, or 1,1,1,3,3,3-hexafluoro-2-propanol, and between hexane and acetonitrile. The correlations derived in the present study for the enthalpies of solvation in dimethyl sulfoxide and propylene carbonate can be used in conjunction with Eqn. (1) and our existing gas-to-organic solvent correlations for dimethyl sulfoxide and propylene carbonate to calculate log K for the solubility of gases and vapors in both solvents to be carried out for temperatures other than 298K.

Experimental Methods

Our search of the chemical literature found a large number of papers (17-82) that reported experimental partial molar enthalpies of solution of liquid and crystalline organic compounds in the two solvents of interest. The latter data were determined by either direct calorimetric methods or calculated based on the temperature dependence of measured infinite dilution activity coefficient data, and the published values were converted to gas-to-organic solvent enthalpies of transfer by

$$\text{Liquid solutes:} \quad \Delta H_{\text{Solv}} = \Delta H_{\text{Soln}} - \Delta H_{\text{Vap},298\text{K}} \quad (6)$$

$$\text{Crystalline solutes:} \quad \Delta H_{\text{Solv}} = \Delta H_{\text{Soln}} - \Delta H_{\text{Sub},298} \quad (7)$$

subtracting the solute's standard molar enthalpy of vaporization (83), $\Delta H_{\text{vap},298\text{K}}$, or standard molar enthalpy of sublimation (84), $\Delta H_{\text{sub},298\text{K}}$, at 298.15 K.

Based on an initial assessment of the available experimental data, we eliminated from consideration all experimental data that pertained to temperatures outside of the temperature range of 283 K – 318 K. Enthalpies of solvation are temperature dependent, and we did not want to introduce large errors in the database by including experimental data far removed from 298 K. Also excluded were values based on solubility measurements where the equilibrium solid phase might be either a solvated form of the solid solute. For several solutes there were multiple, independently determined values. In such cases, we selected direct calorimetric data over indirect values based on the temperature dependence of measured solubilities or infinite dilution activity coefficients. Using the fore-mentioned criteria, 150 molar enthalpies of solvation in dimethyl sulfoxide and 106 molar enthalpies of solvation in propylene carbonate were selected for regression analysis. The experimental $\Delta H_{\text{Solv,DMSO}}$ and $\Delta H_{\text{Solv,PC}}$ values are listed in Tables 1 and 2, respectively.

Molecular descriptors for all of the compounds considered in the present study are also tabulated in Tables 1 and 2. The tabulated values came from our solute descriptor database, which now contains values for more than 3,500 different organic and organometallic compounds. The descriptors were obtained exactly as described before, using various types of experimental data, including water to solvent partitions, gas to solvent partitions, solubility and chromatographic data (8). Solute descriptors used in the present study are all based on experimental data. There is also commercial software (85) and several published estimation schemes (86-89) available for calculating the numerical

values of solute descriptors from molecular structural information if one is unable to find the necessary partition, solubility and/or chromatographic data.

Results and Discussion

We have assembled in Table 1 values of $\Delta H_{\text{Solv,DMSO}}$ for 150 gaseous solutes dissolved in dimethyl sulfoxide covering a reasonably wide range of compound type and descriptor values. Analysis of the experimental data yielded the following correlation equations

$$\begin{aligned} \Delta H_{\text{Solv,DMSO}} \text{ (kJ/mole)} = & - 2.546(0.703) - 0.329(0.0.952) \mathbf{E} - 18.448(1.139) \mathbf{S} - \\ & 47.419(1.653) \mathbf{A} - 5.861(1.004) \mathbf{B} - 6.380(0.197) \mathbf{L} \\ \text{(with } N = 150, SD = 2.80, R^2 = 0.967, R^2_{\text{adj}} = 0.966, F = 850.6) \end{aligned} \quad (8)$$

$$\begin{aligned} \Delta H_{\text{Solv,DMSO}} \text{ (kJ/mole)} = & 2.184(0.845) - 7.233(0.951) \mathbf{E} - 24.071(1.175) \mathbf{S} - \\ & 50.992(1.744) \mathbf{A} - 5.182(1.051) \mathbf{B} - 22.301(0.723) \mathbf{V} \\ \text{(with } N = 150, SD = 2.92, R^2 = 0.965, R^2_{\text{adj}} = 0.963, F = 779.8) \end{aligned} \quad (9)$$

All regression analyses were performed using SPSS statistical software. Both Eqn. 8 and 9 are statistically very good with standard deviations of 2.80 and 2.92 kJ/mole for a data set that covers a range of 91.76 kJ/mole. See Figure 1 for a plot of the calculated values of $\Delta H_{\text{Solv,DMSO}}$ based on Eqn. 8 against the observed values. Equation 8 is slightly the better equation statistically, and from a thermodynamic standpoint Eqn. 8 is the enthalpic temperature derivative of the Abraham model's gas-to-condensed phase transfer

equation. Equation 9 might be more useful in some predictive applications in instances where the L -descriptor is not known. Equation 9 uses the McGowan volume, V -descriptor, which is easily calculable from the individual atomic sizes and numbers of bonds in the molecule (90). To our knowledge, equations 8 and 9 are the first expressions that allow one to predict the enthalpy of solvation of gaseous solutes in dimethyl sulfoxide.

Each of the equation coefficients in the Abraham model encodes chemical information. For example the large a -coefficient in Eqn. 8 indicates that dimethyl sulfoxide exhibits strong hydrogen-bonding basicity character, which is consistent with the molecule's molecular structure, $\text{CH}_3\text{S}(=\text{O})\text{CH}_3$. The two lone electron pairs on the oxygen atom serve as proton acceptor sites for hydrogen-bond formation. The small, nonzero b -coefficient suggests a very weak hydrogen-bond acidity character. While dimethyl sulfoxide is not normally considered to possess any acidic hydrogen(s), several published studies have mentioned the possibility that one of the C-H hydrogens may engage in hydrogen bonding. Fujimoto et al. [91] rationalized the proton NMR spectra of solutions of dimethyl sulfoxide, 2-propanol and/or acetone with tetrasulfonated derivatives of calix[4]resorcarene dissolved in D_2O in terms of well-defined complexes resulting from $\text{CH} - \pi$ interactions between the electron rich benzene rings on the calix[4]resorcarene derivatives and the polarized C-H bonds on the three guest molecules. The authors presented a very compelling argument for why $\text{CH} - \pi$ interactions should be regarded as $\text{C-H} \cdots \pi$ hydrogen bonding. Leggett [92] had earlier estimated acidity parameters for solvents like dimethyl sulfoxide, propylene carbonate, N,N-dimethylformamide and butyrolactone that were believed to not possess hydrogen donor

ability. At this time we do not place too much significance on the nonzero b-coefficient in Eqn. 8 as it possible to still obtain a very good correlation

$$\Delta H_{\text{Solv,DMSO}} \text{ (kJ/mole)} = - 2.767(0.778) + 2.477(0.911) \mathbf{E} - 22.053(1.062) \mathbf{S} - 50.701(1.724) \mathbf{A} - 0.000(0.000) \mathbf{B} - 6.563(0.216) \mathbf{L}$$

(with N = 150, SD = 3.11, R² = 0.969, R²_{adj} = 0.967, F = 539.1) (10)

by setting the b-coefficient equal to zero. The standard deviation increased slightly from SD = 2.80 (Eqn. 8) to SD = 3.11 (Eqn. 10). Given the likely experimental uncertainty in the measured enthalpy of solvation data there is virtually no difference in the two correlations.

In order to assess the predictive ability of Eqn. 8 we divided the 150 data points into a training set and a test set by allowing the SPSS software to randomly select half of the experimental data points. The selected data points became the training set and the compounds that were left served as the test set. Analysis of the experimental data in the training set gave

$$\Delta H_{\text{Solv,DMSO}} \text{ (kJ/mole)} = - 4.505(0.928) + 1.003(1.406) \mathbf{E} - 21.371(1.565) \mathbf{S} - 47.478(2.190) \mathbf{A} - 1.793(1.492) \mathbf{B} - 5.816(0.272) \mathbf{L}$$

with N = 75, SD = 2.49, R² = 0.969, R²_{adj} = 0.967 and F = 434.3 (11)

There is very little difference in the equation coefficients for the full dataset and training dataset correlations. The training set was then used to predict $\Delta H_{\text{Solv,DMSO}}$ values for the 75 compounds in the test set. For the predicted and experimental values, we find that SD

= 3.57, AAE (average absolute error) = 2.288, and AE (average error) = -0.797. There is therefore very little bias in the predictions using eq. 11 with AE equal to -0.878 kJ/mole.

In Table 2 are collected values of the enthalpies of solvation of 106 gaseous solutes in propylene carbonate. Regression analyses of the experimental $\Delta H_{\text{Solv,PC}}$ data in accordance with the Abraham model yielded

$$\Delta H_{\text{Solv,PC}} = -4.499(0.800) + 0.638(1.518) \mathbf{E} - 13.577(1.539) \mathbf{S} - 18.247(2.212) \mathbf{A} - 12.128(1.438) \mathbf{B} - 6.669(0.299) \mathbf{L}$$

(with N = 106, SD = 2.61, $R^2 = 0.962$, $R^2_{\text{adj}} = 0.960$, F = 509.9) (12)

$$\Delta H_{\text{Solv,PC}} = 1.409(0.987) - 7.886(1.336) \mathbf{E} - 18.776(1.535) \mathbf{S} - 20.632(2.205) \mathbf{A} - 11.636(1.413) \mathbf{B} - 24.199(1.056) \mathbf{V}$$

(with N = 106, SD = 2.55, $R^2 = 0.964$, $R^2_{\text{adj}} = 0.962$, F = 534.9) (13)

Both Eqn. (12) and (13) are statistically very good with standard deviations of 2.61 and 2.55 kJ/mole for a data set that covers a range of 103.13 kJ/mole. Figure 2 compares the calculated values of $\Delta H_{\text{Solv,PC}}$ based on Eqn. (12) against the observed values. To our knowledge there has been no previous attempt to correlate $\Delta H_{\text{Solv,PC}}$ data.

We assessed the predictive ability of eqn. (12) by dividing the 106 data points into a training set and a test set as before. Analysis of the experimental data in the training set gave

$$\Delta H_{\text{Solv,PC}} (\text{kJ/mole}) = -1.618(0.947) + 2.630(1.979) \mathbf{E} - 17.422(1.903) \mathbf{S} - 21.097(2.521) \mathbf{A} - 10.425(1.703) \mathbf{B} - 7.420(0.378) \mathbf{L}$$

(14)

with $N = 53$, $SD = 2.05$, $R^2 = 0.981$, $R^2_{adj} = 0.979$ and $F = 494.0$. There is very little difference in the equation coefficients for the full dataset and training dataset correlations. The training set was then used to predict $\Delta H_{Solv,PC}$ values for the 53 compounds in the test set. For the predicted and experimental values, we find that $SD = 3.50$, AAE (average absolute error) = 2.35, and AE (average error) = 0.793. There is therefore very little bias in the predictions using eq. 14 with AE equal to 0.793 kJ/mole. An uncertainty/error of ± 2 kJ/mole in the enthalpy of solvation results in an error of slightly less than 0.04 log units in extrapolating a log L value measured at 298.15 K to 313.15 K. This level of error will be sufficient for most practical chemical and engineering applications.

The correlations presented in this study further document the applicability of the Abraham solvation parameter model to different solute transfer properties. Past studies have shown that the basic model describes a wide range of equilibrium properties that are governed by the Gibbs energy for solute transfer between two condensed phases or between a gas and condensed phase. In the present study we find that the model also provides an accurate mathematical description of the enthalpic contributions to the Gibbs energy, and by inference, we assume that the model describes the entropic contributions as well.

References

1. Stovall, D. M., Acree, W. E. Jr., Abraham, M. H.: Solubility of 9-fluorenone, thianthrene and xanthene in organic solvents. *Fluid Phase Equilib.* **232**, 113-121 (2005).
2. Abraham, M. H., Acree, W. E. Jr.: Characterisation of the water-isopropyl myristate system. *Int. J. Pharm.* **294**, 121-128 (2005).
3. Abraham, M. H., Zissimos, A. M., Acree, W. E. Jr.: Partition of solutes into wet and dry ethers; an LFER analysis. *New J. Chem.* **27**, 1041-1044 (2003).
4. Abraham, M. H., Zissimos, A. M., Huddleston, J. G., Willauer, H. D., Rogers, R. D., Acree, W. E. Jr.: Some novel liquid partitioning systems: water-ionic liquids and aqueous biphasic systems. *Ind. Eng. Chem. Res.* **42**, 413-418 (2003).
5. Abraham, M. H., Acree, W. E. Jr.: Correlation and prediction of partition coefficients between the gas phase and water, and the solvents dodecane and undecane. *New J. Chem.* **28**, 1538-1543 (2004).
6. Abraham, M. H., Acree, W. E. Jr.: The correlation and prediction of butane/water and gas/butane partition coefficients. *Can J. Chem. Eng.* **83**, 362-365 (2005).
7. Abraham, M. H.: Scales of solute hydrogen-bonding: their construction and application to physicochemical and biochemical processes. *Chem. Soc. Rev.* **22**, 73-83 (1993).
8. Abraham, M. H., Ibrahim, A., Zissimos, A. M.: Determination of sets of solute descriptors from chromatographic measurements. *J. Chromatogr. A* **1037**, 29-47 (2004).

9. Abraham, M. H., Chadha, H. S., Whitting, G. S., Mitchell, R. C.: Hydrogen bonding. 32. An analysis of water-octanol and water-alkane partitioning and the $\Delta \log P$ parameter of Seiler. *J. Pharm. Sci.* **83**, 1085-1100 (1994).
10. Abraham, M. H., Le, J.: The correlation and prediction of the solubility of compounds in water using an amended solvation energy relationship. *J. Pharm. Sci.* **88**, 868-880 (1999).
11. Mintz, C., Clark, M., Acree, W. E. Jr., Abraham, M. H.: Enthalpy of solvation correlations for gaseous solutes dissolved in water and in 1-octanol based on the Abraham model. *J. Chem. Inf. Model.*, **47**, 115-121 (2007).
12. Mintz, C., Clark, M., Burton, K., Acree, W. E. Jr., Abraham, M. H.: Enthalpy of solvation correlations for gaseous solutes dissolved in benzene and in alkane solvents based on the Abraham model. *QSAR & Comb. Sci.*, accepted for publication.
13. Mintz, C., Clark, M., Burton, K., Acree, W. E. Jr., Abraham, M. H.: Enthalpy of solvation correlations for gaseous solutes dissolved in toluene and carbon tetrachloride based on the Abraham model. *J. Solution Chem.*, accepted for publication.
14. Ahmed, H.; Poole, C. F. Distribution of neutral organic compounds between n-heptane and methanol or N,N-dimethylformamide. *J. Sep. Sci.* (2006), 29, 2158-2165.
15. Qian, J.; Poole, C. F. Distribution of neutral organic compounds between n-heptane and fluorine-containing alcohols. *J. Chromatogr. A* (2007), 1143, 276-283.

16. Ahmed, H.; Poole, C. F. Model for the distribution of neutral organic compounds between n-hexane and acetonitrile. *J. Chromatogr., A* (2006), 1104, 82-90.
17. Stephenson, W. K.; Fuchs, R. Enthalpies of interaction of ketones with organic solvents. *Can. J. Chem.* (1985), 63, 336-41.
18. Stephenson, W. K.; Fuchs, R. Enthalpies of hydrogen bond formation of 1-octanol with aprotic organic solvents. A comparison of the solvation enthalpy, pure base, and non-hydrogen-bonding baseline methods. *Can. J. Chem.* (1985), 63, 342-348.
19. Krishnan, C. V.; Friedman, H. L. Solvation enthalpies of hydrocarbons and normal alcohols in highly polar solvents. *J. Phys. Chem.* (1971), 75, 3598-3606.
20. Kimura, T.; Tsuda, T.; Takagi, S. Excess enthalpies of some halogenized aromatic compounds + methyl methylthiomethyl sulfoxide or + dimethyl sulfoxide at 298.15 K XI. Thermodynamics of liquid mixtures containing methyl methylthiomethyl sulfoxide. *Thermochim. Acta* (1995) 267, 333-342.
21. Kimura, T.; Suzuki, Kenji; Takagi, Sadao. Excess enthalpies of o- and m-dihalogenated benzene + methyl methylthiomethyl sulfoxide, + dimethyl sulfoxide at 298.15 K. *Fluid Phase Equilib.* (1997), 136, 269-278.
22. Francesconi, R.; Comelli, F.; Bigi, A.; Rubini, K. Thermophysical properties of dimethyl sulfoxide + cyclic and linear ethers at 308.15 K. Application of an extended cell model. *Thermochim. Acta* (2006), 447, 154-160.
23. Hu, X.; Zhang, Y.; Huang, Z. Solvation of n-butane, n-but-1-ene, buta-1,3-diene in dipolar aprotic solvents and protic solvent. *Wuli Huaxue Xuebao* (1985), 1, 130-40.

24. Comelli, F.; Francesconi, R.; Bigi, A.; Rubini, K. Excess molar enthalpies, molar heat capacities, densities, viscosities, and refractive indices of dimethyl sulfoxide + esters of carbonic acid at 308.15 K and atmospheric pressure. *J. Chem. Eng. Data* (2006), 51, 665-670.
25. Comelli, F.; Ottani, S.; Francesconi, R.; Castellari, C. Excess molar enthalpies of binary mixtures containing glycols or polyglycols + dimethyl sulfoxide at 308.15 K. *J. Chem. Eng. Data* (2003), 48, 995-998.
26. Francesconi, R.; Bigi, A.; Rubini, K.; Comelli, F. Excess enthalpies, heat capacities, densities, viscosities and refractive indices of dimethyl sulfoxide + three aryl alcohols at 308.15 K and atmospheric pressure. *J. Chem. Eng. Data* (2005), 50, 1932-1937.
27. Topphoff, M.; Gruber, D.; Gmehling, J. Measurement of activity coefficients at infinite dilution using gas-liquid chromatography. 10. Results for various solutes with the stationary phases dimethyl sulfoxide, propylene carbonate, and N-ethylformamide. *J. Chem. Eng. Data* (1999), 44, 1355-1359.
28. Catalan, J.; Gomez, J.; Couto, A.; Laynez, J. Toward a solvent basicity scale: the calorimetry of the pyrrole probe. *J. Amer. Chem. Soc.* (1990), 112, 1678-1681.
29. Costa, F. S.; Eusebio, M. E.; Redinha, J. S.; Leitao, M. L. P. On the interpretation of the enthalpy of solvation of hydroxyl cyclohexane derivatives in different solvents. *J. Chem. Thermodyn.* (2000), 32, 311-317.
30. Nissema, A.; Kaivamo, T.; Karvo, M. Thermodynamic properties of binary and ternary systems. XII. Excess enthalpies of cyclohexane + p-xylene, p-xylene +

- dimethyl sulfoxide, and dimethyl sulfoxide + cyclohexane + p-xylene. *Journal of Chem. Thermodyn.* (1983), 15, 1083-1086.
31. Philippe, R. Excess enthalpy. Dimethyl sulfoxide-dichloromethane (methylene chloride) system. *Int. DATA Ser., Sel. Data Mix., Ser. A* (1975), 122.
 32. Philippe, R. Excess enthalpy. Dimethyl sulfoxide-1,1,1-trichloroethane (methylchloroform) system. *Int. DATA Ser., Sel. Data Mix., Ser. A* (1975), 126.
 33. Arnett, E. M.; McKelvey, D. R. Enthalpies of transfer from water to dimethyl sulfoxide for some ions and molecules. *J. Amer. Chem. Soc.* (1966), 88, 2598-2599.
 34. Abraham, M. H.; Nasehzadeh, A. Thermodynamic functions for solution of triethylamine in 30 solvents, and calculation of the solute-solvent interaction terms. *J. Chem. Thermodyn.* (1981), 13, 549-555.
 35. Benoit, R. L.; Frechette, M. Heats of solution of some pyrimidines, purines and related compounds in dimethylsulfoxide. *Thermochim. Acta* (1988), 126 155-164.
 36. Hallman, J. H.; Stephenson, W. K.; Fuchs, R. Hydrogen bonding and other polar interactions of 1-, 2-, and 4-octyne with organic solvents. *Can. J. Chem.* (1983), 61, 2044-2047.
 37. Takeki Matsui, Loren G. Hepler, and David V. Fenby, Thermodynamic investigation of complex formation by hydrogen bonding in binary liquid systems. Chloroform with triethylamine, dimethyl sulfoxide, and acetone *J. Phys. Chem.*, 77, 2397-2400 (1973).

38. Józwiak, M. The effect of properties of water–organic solvent mixtures on the solvation enthalpy of 12-crown-4, 15-crown-5, 8-crown-6 and benzo-15-crown-5 ethers at 298.15 K. *Thermochim. Acta*, 417, 31-41(2004).
39. Mikhail D. Borisover, Andrey A. Stolov, Felix D. Baitalov, Alexander I. Morozov and Boris N. Solomonov, Calorimetric and infrared study of methyl iodide and some monosubstituted butyl halides. Does methyl iodide form hydrogen bonds as a proton donor? *Thermochim. Acta*, 285, 199-209 (1996).
40. Felisbela S. Costa, M. Ermelinda Eusébio, J. S. Redinha and M. Luísa P. Leitão, Enthalpies of solvation of hydroxyl cyclohexane derivatives in different solvents. *J. Chem. Thermodyn.*, 31, 895-903 (1999).
41. Nunes, N.; Moreira, L.; Leitão, R. E.; Martins, F. Determination of solvation and specific interaction enthalpies of adamantane derivatives in aprotic solvents. *J. Chem. Thermodyn.*, in press (2007).
42. Bebahani, G. R.; Dunnion, D.; Falvey, P.; Hickey, K.; Meade, M.; McCarthy, Y.; Symons, M. C. R.; Waghorne, W. E. Nonelectrolyte solvation in aqueous dimethyl sulfoxide-a calorimetric and infrared spectroscopic study. *J. Solution Chem.* (2000), 29, 521-539.
43. Krishnan, C. V.; Friedman, H. L. Solvation enthalpies of various nonelectrolytes in water, propylene carbonate, and dimethyl sulfoxide. *J. Phys. Chem.* (1969), 73, 1572-1580.
44. Pathak, G.; Tripathi, A. D.; Pradhan, S. The excess enthalpies of mixing of 1,1,2,2-tetrachloroethane with acetone, n-dibutyl ether, acetonitrile and dimethyl sulfoxide. *Thermochim. Acta* (1992), 197, 329-333.

45. Figeys, D.; Koschmidder, M.; Benoit, R. L. Enthalpies of solution of naphthalene, N,N-dimethyl-1-naphthylamine, and 1,8-bis(dimethylamino)-naphthalene in 16 organic solvents. *Can. J. Chem.* (1992), 70, 1586-1589.
46. Manin, N. G.; Belichenko, S. Yu.; Korolev, V. P. Thermochemistry of benzene solution in alcohols, aprotic solvents, and mixtures of aprotic solvents with methanol. *Russ. J. Gen. Chem.* (2003), 73, 9-16.
47. Benoit, R. L.; Mackinnon, M. J.; Bergeron, L. Basicity of N-substituted anilines and pyridine in dimethyl sulfoxide. *Can. J. Chem.* (1981), 59, 1501-1504.
48. Kimura, T.; Takagi, S. Excess enthalpies of diethylether, ethylpropylether, dipropylether + methyl methylthiomethyl sulfoxide or + dimethyl sulfoxide at 298.15 K. *Thermochim. Acta* (1995), 253, 59-67.
49. Kimura, T.; Takagi, S. Excess enthalpies of binary mixtures with oxepane, oxetane, 3,4-dihydro-2H-pyran, and furan + methyl methylthiomethyl sulfoxide and dimethyl sulfoxide at 298.15 K. *Netsu Sokutei* (1996), 23, 53-59.
50. Ohtsu, K.; Ozutsumi, K. Thermodynamics of solvation of 18-crown-6 and its alkali-metal complexes in various solvents. *J. Incl. Phenom. Macrocyclic Chem.* (2003), 45, 217-224.
51. Pais, A. A. C. C.; Sousa, A.; Eusebio, M. E.; Redinha, J. S. Solvation of alkane and alcohol molecules. Energy contributions. *Phys. Chem. Chem. Phys.* (2001), 3, 4001-4009.
52. Cox, Brian G. Free energies, enthalpies, and entropies of transfer of nonelectrolytes from water to mixtures of water and dimethyl sulfoxide, water and

- acetonitrile, and water and dioxane. *J. Chem. Soc., Perkin Trans. 2* (1973), 607-610.
53. Batov, D. V.; Antonova, O. A.; Korolev, V. P. Thermochemistry of aniline solution in binary mixtures of water-aprotic solvent. *Russ. J. Gen. Chem.* (2001), 71, 689-695.
54. Benoit, R. L.; Frechette, M.; Lefebvre, D. 2,6-Di-tert-butylpyridine: an unusually weak base in dimethyl sulfoxide. *Can. J. Chem.* (1988), 66, 1159-1162.
55. G. A. Krestov, V. P. Korolyov and D. V. Batov, Thermochemistry of solvation of non-electrolytes in mono- and polyatomic alcohols and their mixtures with water. *Thermochim. Acta*, 169, 69-80 (1990).
56. Solomonov, B. N.; Khafizov, F. T.; Gorbachuk, V. V. Multiparameter equations for the description of enthalpy of nonspecific solvation of organic nonelectrolytes. *Zhur. Obsch. Khim.* 60, 1446-1452 (1990).
57. Stephenson, W. K.; Fuchs, R. Enthalpies of interaction of aromatic solutes with organic solvents. *Can. J. Chem.* (1985), 63, 2529-2534.
58. Stephenson, W. K.; Fuchs, R. Enthalpies of interaction of hydroxylic solutes with organic solvents. *Can. J. Chem.* (1985), 63, 2535-2539.
59. Philippe, R. Excess enthalpy. Dimethyl sulfoxide-dibromomethane (methylene bromide) system. *Int. DATA Series, Sel. Data Mix., Ser. A* (1975), 124.
60. Smirnov, V. I.; Kastorina, E. V.; Kiestov, G. A.; Fridman, A. Ya. Solvation of naphthalene and its hydroxy derivatives in nonaqueous media. *Zhur. Fiz. Khim.* (1993), 67, 1123-1126.

61. Lahti, M.; Kankaanpera, A.; Sapyska, H. Kinetic acidity of carbon acids. Initial and transition state enthalpies of transfer from water to aqueous dimethyl sulfoxide in the hydroxide ion catalyzed ionization of acetophenone. *J. Chem. Soc., Perkin Trans, 2* (1990), 1081-1083.
62. Abraham, M. H. Free energies, enthalpies, and entropies of solution of gaseous nonpolar nonelectrolytes in water and nonaqueous solvents. The hydrophobic effect. *J. Amer. Chem. Soc.* (1982), 104, 2085-2094.
63. Comelli, F.; Francesconi, R.; Ottani, S. Excess molar enthalpies of binary mixtures containing propylene carbonate + 23 alkanooates at 298.15 K. *J. Chem. Eng. Data* (1998), 43, 333-336.
64. Comelli, F.; Francesconi, R. Excess molar enthalpies for propylene carbonate + linear and cyclic ketones at 298.15 K. *J. Chem. Eng. Data* (1995), 40, 805-807.
65. Francesconi, R.; Comelli, F. Excess molar enthalpies, densities, and excess molar volumes of binary mixtures containing esters of carbonic acid at 298.15 and 313.15 K. *J. Chem. Eng. Data* (1995), 40, 811-14.
66. Comelli, F.; Francesconi, R.; Castellari, C. Excess molar enthalpies and excess molar volumes of propylene carbonate + seven pentanol isomers at 298.15 K. *J. Chem. Eng. Data* (1999), 44, 144-147.
67. Castellari, C.; Comelli, F.; Francesconi, R. Excess molar enthalpies of binary mixtures containing glycols or (polyethylene) glycols + propylene carbonate at 308.15 K. *Thermochim. Acta* (2004), 413, 249-253.

68. Ottani, S.; Comelli, F.; Castellari, C. Densities, viscosities, and excess molar enthalpies of propylene carbonate + anisole or + phenetole at (293.15, 303.15, and 313.15) K. *J. Chem. Eng. Data* (2001), 46, 125-129.
69. Comelli, F.; Francesconi, R.; Bigi, A.; Rubini, K. Excess molar enthalpies, molar heat capacities, densities, viscosities, and refractive indices of dimethyl sulfoxide + esters of carbonic acid at 308.15 K and atmospheric pressure. *J. Chem. Eng. Data* (2006), 51, 665-670.
70. Comelli, F.; Francesconi, R. Excess properties of binary mixtures of esters of carbonic acid + three aryl alcohols at 308.15 K. *J. Chem. Eng. Data* (2005), 50, 191-195.
71. Francesconi, R.; Comelli, F. Excess molar enthalpies and excess molar volumes of propylene carbonate + cyclic ethers. *J. Chem. Eng. Data* (1995), 40, 31-33.
72. Ohtsu, K.; Kawashima, T.; Ozutsumi, K. Thermodynamics of complexation of sodium, potassium, rubidium, cesium and ammonium ions with 18-crown-6 in acetonitrile and propylene carbonate. *J. Chem. Soc., Faraday Trans.* (1995), 91, 4375-4380.
73. Francesconi, R.; Comelli, F. Excess molar enthalpies of binary mixtures containing dimethylcarbonate, diethylcarbonate or propylene carbonate + three chloroalkenes at 298.15 K. *Thermochim. Acta* (1995), 260, 95-103.
74. de Namor, A. F. D.; Ng, J. C. Y.; Tanco, M. A. L.; Salomon, M. Thermodynamics of lithium-crown ether (12-crown-4 and 1-benzyl-1-aza-12-crown-4) interactions in acetonitrile and propylene carbonate. The anion effect on the coordination process. *J. Phys. Chem.* (1996), 100, 14485-14491.

75. Francesconi, R.; Comelli, F. Excess enthalpies and excess volumes of the liquid binary mixtures of propylene carbonate + six alkanols at 298.15 K. *J. Chem. Eng. Data* (1996), 41, 1397-1400.
76. Comelli, F.; Francesconi, R.; Castellari, C. Excess molar enthalpies of binary mixtures containing propylene carbonate + some n-alkoxy- and n-alkoxyethoxy-ethanols at 288.15, 298.15, and 313.15 K. *Thermochim. Acta* (2000), 354, 89-97.
77. Blanchard, F.; Carre, B.; Bonhomme, F.; Biensan, P.; Lemordant, D. Solubility of carbon dioxide in alkylcarbonates and lactones. *Can. J. Chem.* (2003), 81, 385-391.
78. Benoit, R. L.; Milanova, E. Vapor pressure and calorimetric data for the solution of sulfur dioxide in aprotic solvents. *Can. J. Chem.* (1979), 57, 1319-1323.
79. Fuchs, R.; Stephenson, W. K. Enthalpies of transfer of alkane solutes from the vapor state to organic solvents. *Can. J. Chem.* (1985), 63, 349-52.
80. Kimura, T.; Matsushita, T.; Kamiyama, T. Enthalpies of Solution of Aliphatic Amines, Aliphatic Benzene, and Alkane in Dimethyl Sulfoxide at 298.15 K. *J. Solution Chem.* (2004), 33, 875-886.
81. Kimura, T.; Matsushita, T.; Kamiyama, T. Enthalpies of solution of aliphatic compounds in dimethyl sulfoxide. *Thermochim. Acta* (2004), 416, 129-134.
82. Gennaro, A.; Isse, A. A.; Vianello, E. Solubility and electrochemical determination of carbon dioxide in some dipolar aprotic solvents. *J. Electroanal. Chem. Interfacial Electrochem.* (1990), 289, 203-215.

83. Chickos, J. S., Acree, W. E. Jr.: Enthalpies of vaporization of organic and organometallic compounds, 1880-2002. *J. Phys. Chem. Ref. Data* **32**, 519-878 (2003).
84. Chickos, J. S., Acree, W. E. Jr.: Enthalpies of sublimation of organic and organometallic compounds, 1910-2001. *J. Phys. Chem. Ref. Data* **31**, 537-698 (2002).
85. PharmaAlgorithms, ADME Boxes, Version 3.0, **2006**, PharmaAlgorithms Inc., 591 Indian Road, Toronto, ON M6P 2C4, Canada.
86. Platts, J. A., Butina, D., Abraham M. H., Hersey, A.: Estimation of molecular linear free energy relation descriptors using a group contribution approach. *J. Chem. Inf. Comp. Sci.* **39**, 835-845 (1999).
87. Jover, J.; Bosque, R.; Sales, J., Determination of Abraham solute properties from molecular structure. *J. Chem. Inf. Comp. Sci.* **2004**, *44*, 1098-1106 (2004).
88. Arey J. S., Green, W. H. Jr., Gschwend P. M.: The electrostatic origin of Abraham's solute polarity parameter. *J. Phys. Chem. B* **109**, 7564-7573 (2005).
89. Metelet, F., Rogalski, M.: Experimental determination and prediction of the gas-liquid n-hexadecane partition coefficients. *J. Chromatogr., A* **923**, 153-163 (2001).
90. Abraham, M. H., McGowan, J. C.: The use of characteristic volumes to measure cavity terms in reversed phase liquid chromatography. *Chromatographia* **23**, 245-246 (1987).
91. Fujimoto, T.; Yanagihara, R.; Kobayashi, K.; Aoyama, Y. C-H $\cdots\pi$ hydrogen bonding between electron-rich benzene rings and polarized C-H bonds: selectivity

in the complexation of highly hydrophobic guest molecules with calyx[4]-resorcarene hosts in water. *Bull Chem. Soc. Jpn.*, 68, 2113-2124 (1995).

92. Leggett, D. C. H-bond acidity of solvents: II. Transformation of the Snyder chi proton donor scale to the Taft-Kamlet alpha scale. *J. Solution Chem.* (1994), 23, 697-701.

TABLE 1. Values of the gas to dimethyl sulfoxide solvation enthalpy in kJ/mole at 298

K for 150 solutes, together with the solute descriptors

Solute	E	S	A	B	L	V	Exp	Ref
Butane	0.000	0.000	0.000	0.000	1.615	0.6722	-12.10	23
Pentane	0.000	0.000	0.000	0.000	2.162	0.8131	-15.23	19
Hexane	0.000	0.000	0.000	0.000	2.668	0.9540	-17.61	79
Heptane	0.000	0.000	0.000	0.000	3.173	1.0949	-21.13	79
Octane	0.000	0.000	0.000	0.000	3.677	1.2358	-24.10	79
Nonane	0.000	0.000	0.000	0.000	4.182	1.3767	-27.66	79
Decane	0.000	0.000	0.000	0.000	4.686	1.5176	-30.79	79
Cyclohexane	0.305	0.100	0.000	0.000	2.964	0.8454	-21.60	29
1-Butene	0.100	0.080	0.000	0.070	1.491	0.6292	-15.50	23
1-Pentene	0.093	0.080	0.000	0.070	2.047	0.7701	-20.31	27
1-Hexene	0.078	0.080	0.000	0.070	2.572	0.9110	-21.51	27
1-Octene	0.094	0.080	0.000	0.070	3.568	1.1928	-31.78	27
1,3-Butadiene	0.320	0.230	0.000	0.100	1.543	0.5862	-17.00	23
1-Octyne	0.155	0.220	0.090	0.100	3.521	1.1498	-37.15	36
2-Octyne	0.225	0.300	0.000	0.100	3.850	1.1498	-37.36	36
Benzene	0.610	0.520	0.000	0.140	2.786	0.7176	-30.71	43, 46
Toluene	0.601	0.520	0.000	0.140	3.325	0.8573	-34.40	80
Ethylbenzene	0.613	0.510	0.000	0.150	3.778	0.9982	-37.32	80
1,3-Dimethylbenzene	0.623	0.520	0.000	0.160	3.839	0.9982	-41.75	43
Propylbenzene	0.604	0.500	0.000	0.150	4.230	1.1391	-39.86	80
Isopropylbenzene	0.602	0.490	0.000	0.160	4.084	1.1391	-39.66	33
Mesitylene	0.649	0.520	0.000	0.190	4.344	1.1391	-39.37	57
Butylbenzene	0.600	0.510	0.000	0.150	4.730	1.2800	-42.66	80
Pentylbenzene	0.594	0.510	0.000	0.150	5.230	1.4210	-45.44	80
Hexylbenzene	0.591	0.500	0.000	0.150	5.720	1.5618	-48.72	80
Heptylbenzene	0.577	0.480	0.000	0.150	6.219	1.7027	-52.16	80
Octylbenzene	0.579	0.480	0.000	0.150	6.714	1.8436	-53.12	80
Naphthalene	1.340	0.920	0.000	0.200	5.161	1.0854	-53.19	45
Biphenyl	1.360	0.990	0.000	0.260	6.014	1.3240	-60.96	43
Anthracene	2.290	1.340	0.000	0.280	7.568	1.4540	-76.60	56
Methylamine	0.250	0.350	0.160	0.580	1.300	0.3493	-26.80	47
n-Propylamine	0.225	0.350	0.160	0.610	2.141	0.6311	-30.23	80
n-Butylamine	0.224	0.350	0.160	0.610	2.618	0.7720	-33.59	80
n-Pentylamine	0.211	0.350	0.160	0.610	3.139	0.9129	-36.48	80
n-Hexylamine	0.197	0.350	0.160	0.610	3.655	1.0538	-40.40	80
n-Heptylamine	0.197	0.350	0.160	0.610	4.153	1.1947	-43.15	80
Acetone	0.179	0.700	0.040	0.490	1.696	0.5470	-29.87	17
2-Butanone	0.166	0.700	0.000	0.510	2.287	0.6879	-32.47	17
2-Pentanone	0.143	0.680	0.000	0.510	2.755	0.8288	-35.49	27
2-Hexanone	0.136	0.680	0.000	0.510	3.286	0.9697	-39.80	56
2-Heptanone	0.123	0.680	0.000	0.510	3.760	1.1106	-41.00	17

4-Heptanone	0.110	0.660	0.000	0.510	3.705	1.1106	-39.83	17
2-Octanone	0.108	0.680	0.000	0.510	4.257	1.2515	-43.90	56
2-Nonanone	0.119	0.680	0.000	0.510	4.735	1.3924	-47.20	17
5-Nonanone	0.103	0.660	0.000	0.510	4.698	1.3924	-44.81	17
2,2,4,4-Tetramethyl-3-pentanone	0.099	0.560	0.000	0.520	4.370	1.3924	-35.15	17
Cyclohexanone	0.403	0.860	0.000	0.560	3.792	1.0020	-41.84	17
Diethyl ether	0.041	0.250	0.000	0.450	2.015	0.7309	-21.59	58
Dipropyl ether	0.008	0.250	0.000	0.450	2.954	1.0127	-29.59	48
Dibutyl ether	0.000	0.250	0.000	0.450	3.924	1.2945	-32.01	18
Ethyl propyl ether	0.000	0.250	0.000	0.450	2.493	0.8718	-25.25	48
n-Butyl methyl ether	0.045	0.250	0.000	0.440	2.658	0.8718	-25.36	57
Methyl heptyl ether	0.048	0.250	0.000	0.450	4.088	1.2945	-36.38	18
Methyl tert-butyl ether	0.024	0.210	0.000	0.590	2.380	0.8718	-24.23	58
Tetrahydrofuran	0.289	0.520	0.000	0.480	2.636	0.6223	-27.34	22
Tetrahydropyran	0.275	0.470	0.000	0.550	3.057	0.7672	-30.16	22
1,4-Dioxane	0.329	0.750	0.000	0.640	2.892	0.6810	-36.63	22
1,2-Dimethoxyethane	0.116	0.670	0.000	0.680	2.654	0.7896	-34.73	22
1,2-Diethoxyethane	0.008	0.730	0.000	0.790	3.310	1.0714	-39.15	22
Furan	0.369	0.510	0.000	0.130	1.913	0.5363	-29.99	49
12-Crown-4	0.420	0.990	0.000	1.390	5.135	1.3620	-65.98	38
15-Crown-5	0.410	1.200	0.000	1.750	6.779	1.7025	-78.85	38
18-Crown-6	0.400	1.340	0.000	2.130	7.919	2.0430	-103.86	50
1-Fluorooctane	-0.020	0.350	0.000	0.100	3.850	1.2538	-37.49	18
Dichloromethane	0.387	0.570	0.100	0.050	2.019	0.4943	-34.27	31
Chloroform	0.425	0.490	0.150	0.020	2.480	0.6167	-36.78	37
Carbon tetrachloride	0.458	0.380	0.000	0.000	2.823	0.7391	-32.40	33
1-Chlorobutane	0.210	0.400	0.000	0.100	2.722	0.7946	-27.68	39
1-Chlorooctane	0.191	0.400	0.000	0.090	4.708	1.3580	-39.79	18
2-Chloro-2-methylpropane	0.142	0.300	0.000	0.030	2.273	0.7946	-22.72	39
1,1,1-Trichloroethane	0.369	0.410	0.000	0.090	2.733	0.7576	-30.03	32
1,1,1,2-Tetrachloroethane	0.595	0.760	0.160	0.120	3.803	0.8800	-56.82	44
Dibromomethane	0.714	0.690	0.110	0.070	2.886	0.5995	-39.22	59
Methyl iodide	0.676	0.430	0.000	0.120	2.106	0.5077	-26.39	39
1-Iodobutane	0.628	0.400	0.000	0.150	3.628	0.9304	-33.49	39
Fluorobenzene	0.477	0.570	0.000	0.100	2.788	0.7341	-34.45	20
1,2-Difluorobenzene	0.390	0.630	0.000	0.060	2.843	0.7518	-34.93	21
1,3-Difluorobenzene	0.374	0.580	0.000	0.060	2.776	0.7518	-34.41	21
4-Fluorotoluene	0.488	0.550	0.000	0.140	3.366	0.6750	-36.99	20
Chlorobenzene	0.718	0.650	0.000	0.070	3.657	0.8388	-39.23	20
1,2-Dichlorobenzene	0.872	0.780	0.000	0.040	4.518	0.9612	-48.31	21
1,3-Dichlorobenzene	0.847	0.730	0.000	0.020	4.410	0.9612	-51.20	21
1,4-Dichlorobenzene	0.825	0.750	0.000	0.020	4.435	0.9612	-43.90	56
4-Chlorotoluene	0.705	0.740	0.000	0.050	4.205	0.9797	-43.13	20
Bromobenzene	0.882	0.730	0.000	0.090	4.041	0.8914	-42.32	20
Iodobenzene	1.188	0.820	0.000	0.120	4.502	0.9746	-48.30	19
Acetonitrile	0.237	0.900	0.070	0.320	1.739	0.4042	-33.24	81

Propanenitrile	0.162	0.900	0.020	0.360	2.082	0.5451	-34.57	81
1-Butanenitrile	0.188	0.900	0.000	0.360	2.548	0.6860	-36.87	81
1-Pentanenitrile	0.177	0.900	0.000	0.360	3.108	0.8269	-41.04	81
1-Hexanenitrile	0.166	0.900	0.000	0.360	3.608	0.9678	-43.54	81
1-Heptanenitrile	0.159	0.900	0.000	0.360	4.089	1.1087	-46.30	81
1-Octanenitrile	0.162	0.900	0.000	0.360	4.585	1.2496	-49.87	81
1-Nonanenitrile	0.159	0.900	0.000	0.360	4.970	1.3905	-53.36	81
1-Decanenitrile	0.156	0.900	0.000	0.360	5.460	1.5314	-56.98	81
1-Undecanenitrile	0.154	0.900	0.000	0.360	5.940	1.6723	-59.66	81
1-Dodecanenitrile	0.152	0.900	0.000	0.360	6.460	1.8132	-63.30	81
1-Tridecanenitrile	0.150	0.900	0.000	0.360	6.924	1.9541	-66.13	81
1-Tetradecanitrile	0.149	0.900	0.000	0.360	7.403	2.0950	-68.94	81
Methanol	0.278	0.440	0.430	0.470	0.970	0.3082	-39.17	81
Ethanol	0.246	0.420	0.370	0.480	1.485	0.4491	-41.13	81
1-Propanol	0.236	0.420	0.370	0.480	2.031	0.5900	-43.76	81
2-Propanol	0.212	0.360	0.330	0.560	1.764	0.5900	-41.56	43
1-Butanol	0.224	0.420	0.370	0.480	2.601	0.7309	-48.16	81
1-Pentanol	0.219	0.420	0.370	0.480	3.106	0.8718	-51.45	81
1-Hexanol	0.210	0.420	0.370	0.480	3.610	1.0170	-53.97	81
1-Heptanol	0.211	0.420	0.370	0.480	4.115	1.1536	-58.29	81
1-Octanol	0.199	0.420	0.370	0.480	4.619	1.2950	-60.25	18
1-Nonanol	0.193	0.420	0.370	0.480	5.124	1.4354	-66.92	81
1-Decanol	0.191	0.420	0.370	0.480	5.628	1.5763	-68.42	81
1-Undecanol	0.181	0.420	0.370	0.480	6.130	1.7173	-70.85	81
2-Methyl-2-propanol	0.180	0.300	0.310	0.600	1.963	0.7309	-40.34	33
2-methyl-2-butanol	0.194	0.300	0.310	0.600	2.630	0.8718	-44.52	58
Cyclohexanol	0.460	0.540	0.320	0.570	3.758	0.9040	-54.14	51
cis 1,2-Cyclohexanediol	0.600	0.860	0.500	0.860	4.203	0.9628	-69.70	40
Ethylene glycol	0.404	0.900	0.580	0.780	2.661	0.5078	-69.66	25
Propylene glycol	0.373	0.900	0.580	0.800	2.918	0.6487	-66.70	25
3-Methylphenol	0.822	0.880	0.570	0.340	4.310	0.9160	-74.56	58
Aniline	0.955	0.960	0.260	0.410	3.934	0.8162	-66.48	53
N-Methylaniline	0.948	0.940	0.170	0.470	4.494	0.9571	-57.70	47
N,N-Dimethylaniline	0.957	0.810	0.000	0.410	4.701	1.0980	-49.29	35
Dimethyl carbonate	0.142	0.540	0.000	0.570	2.328	0.6644	-34.65	24
Diethyl carbonate	0.060	0.580	0.000	0.530	3.412	0.9462	-38.49	24
Benzyl alcohol	0.803	0.870	0.330	0.560	4.221	0.9160	-70.30	26
2-Phenylethanol	0.811	0.910	0.300	0.640	4.628	1.0569	-70.47	26
Methyl acetate	0.142	0.640	0.000	0.450	1.911	0.6057	-30.38	42
Ethyl acetate	0.106	0.620	0.000	0.450	2.314	0.7466	-32.80	52
tert-Butyl acetate	0.025	0.540	0.000	0.470	2.802	1.0284	-33.46	52
N,N-Dimethyl formamide	0.367	1.310	0.000	0.740	3.173	0.6468	-45.64	42
N,N-Dimethyl acetamide	0.363	1.330	0.000	0.780	3.717	0.7880	-52.29	42
Acetophenone	0.818	1.010	0.000	0.480	4.501	1.0140	-51.35	61
Pyridine	0.631	0.840	0.000	0.520	3.022	0.6753	-39.00	35
2,6-Dimethylpyridine	0.607	0.700	0.000	0.630	3.760	0.9571	-42.60	54
Nitrobenzene	0.871	1.110	0.000	0.280	4.557	0.8906	-52.88	43

Nitromethane	0.313	0.950	0.060	0.310	1.892	0.4237	-38.96	55
(Trifluoromethyl)benzene	0.225	0.480	0.000	0.100	2.894	0.9104	-32.68	57
Anisole	0.710	0.750	0.000	0.290	3.890	0.9160	-44.27	57
Triethylamine	0.101	0.150	0.000	0.790	3.040	1.0538	-26.30	34
Pyrrrole	0.613	0.730	0.410	0.290	2.865	0.5770	-54.68	28
N-Methylpyrrrole	0.559	0.790	0.000	0.310	2.923	0.7180	-40.17	28
1,4-Dimethylbenzene	0.613	0.520	0.000	0.160	3.839	0.9982	-37.67	30
Diisopropyl ether	-0.060	0.160	0.000	0.580	2.530	1.0127	-26.53	27
Ethyl tert-butyl ether	-0.020	0.160	0.000	0.600	2.720	1.0127	-27.50	27
Methyl tert-amyl ether	0.050	0.210	0.000	0.600	2.916	1.0127	-29.07	27
Acetaldehyde	0.208	0.670	0.000	0.450	1.230	0.4060	-26.80	27
Vinyl acetate	0.223	0.640	0.000	0.430	2.152	0.7040	-30.38	27
1-Naphthol	1.520	1.050	0.600	0.370	6.130	1.1441	-90.60	60
1-Bromoadamantane	1.070	0.900	0.000	0.200	6.130	1.3668	-50.90	41
1-Adamantanol	0.940	0.900	0.310	0.660	5.634	1.2505	-66.00	41
Carbon dioxide	0.000	0.280	0.050	0.100	0.058	0.2809	-12.43	82

TABLE 2. Values of the gas to propylene carbonate solvation enthalpy in kJ/mole at 298

K for 106 solutes, together with the solute descriptors

Solute	E	S	A	B	L	V	Exp	Ref
Ethane	0.000	0.000	0.000	0.000	0.492	0.3904	-5.73	62
Propane	0.000	0.000	0.000	0.000	1.050	0.5313	-6.44	62
2-Methylpropane	0.000	0.000	0.000	0.000	1.409	0.6722	-9.71	62
Butane	0.000	0.000	0.000	0.000	1.615	0.6722	-10.83	62
Pentane	0.000	0.000	0.000	0.000	2.162	0.8131	-17.57	62
Hexane	0.000	0.000	0.000	0.000	2.668	0.9540	-21.00	62
Heptane	0.000	0.000	0.000	0.000	3.173	1.0949	-24.66	62
Octane	0.000	0.000	0.000	0.000	3.677	1.2358	-31.41	27
Decane	0.000	0.000	0.000	0.000	4.686	1.5176	-37.97	27
Cyclohexane	0.305	0.100	0.000	0.000	2.964	0.8454	-24.18	43
Methylcyclopentane	0.225	0.100	0.000	0.000	2.816	0.8454	-23.14	27
Methylcyclohexane	0.244	0.060	0.000	0.000	3.319	0.9863	-26.22	27
1-Butene	0.100	0.080	0.000	0.070	1.491	0.6292	-17.60	23
1-Hexene	0.078	0.080	0.000	0.070	2.572	0.9110	-24.81	27
1-Octene	0.094	0.080	0.000	0.070	3.568	1.1928	-30.56	27
Cyclohexene	0.395	0.200	0.000	0.100	3.021	0.8025	-27.01	27
1,3-Butadiene	0.320	0.230	0.000	0.100	1.543	0.5862	-19.30	23
Methyl acetate	0.142	0.640	0.000	0.450	1.911	0.6057	-32.50	63
Ethyl acetate	0.106	0.620	0.000	0.450	2.314	0.7466	-34.51	63
Propyl acetate	0.092	0.600	0.000	0.450	2.819	0.8875	-38.46	63
Vinyl acetate	0.223	0.640	0.000	0.430	2.152	0.7040	-34.28	63
Butyl acetate	0.071	0.600	0.000	0.450	3.353	1.0284	-41.22	63
tert-butyl acetate	0.025	0.540	0.000	0.470	2.802	1.0284	-35.42	63
Isobutyl acetate	0.052	0.570	0.000	0.470	3.161	1.0284	-36.13	63
Pentyl acetate	0.067	0.600	0.000	0.450	3.844	1.1693	-45.45	63
Hexyl acetate	0.056	0.600	0.000	0.450	4.351	1.3102	-47.36	63
Methyl propanoate	0.128	0.600	0.000	0.450	2.431	0.7466	-35.21	63
Ethyl propanoate	0.087	0.580	0.000	0.450	2.807	0.8875	-37.79	63
Propyl propanoate	0.070	0.560	0.000	0.450	3.338	1.0284	-40.40	63
Butyl propanoate	0.058	0.560	0.000	0.470	3.833	1.1693	-46.95	63
Methyl butanoate	0.106	0.600	0.000	0.450	2.893	0.8875	-38.08	63
Ethyl butanoate	0.068	0.580	0.000	0.450	3.271	1.0284	-39.85	63
Propyl butanoate	0.050	0.560	0.000	0.450	3.783	1.1693	-44.43	63
Methyl pentanoate	0.108	0.600	0.000	0.450	3.392	1.0284	-40.89	63
Ethyl pentanoate	0.049	0.580	0.000	0.450	3.769	1.1693	-43.63	63
Ethyl hexanoate	0.043	0.580	0.000	0.450	4.251	1.3102	-48.30	63
Ethyl heptanoate	0.027	0.580	0.000	0.450	4.733	1.4511	-47.61	63
Ethyl isobutyrate	0.034	0.550	0.000	0.470	3.072	1.0284	-37.63	63
Dimethyl carbonate	0.142	0.540	0.000	0.570	2.328	0.6644	-37.49	65
Diethyl carbonate	0.060	0.580	0.000	0.530	3.412	0.9462	-42.47	65
Acetone	0.179	0.700	0.040	0.490	1.696	0.5470	-30.79	64

2-Butanone	0.166	0.700	0.000	0.510	2.287	0.6879	-34.27	64
2-Pentanone	0.143	0.680	0.000	0.510	2.755	0.8288	-37.60	27
2-Hexanone	0.136	0.680	0.000	0.510	3.286	0.9697	-40.78	64
2-Octanone	0.108	0.680	0.000	0.510	4.257	1.2515	-47.43	64
2-Undecanone	0.101	0.680	0.000	0.510	5.732	1.6740	-59.14	64
Cyclohexanone	0.403	0.860	0.000	0.560	3.792	0.8610	-43.77	64
2-Methylcyclohexanone	0.372	0.830	0.000	0.560	4.050	1.0020	-45.28	64
Anisole	0.710	0.750	0.000	0.290	3.890	0.9160	-45.45	68
Phenetole	0.681	0.700	0.000	0.320	4.242	1.0570	-47.61	68
Dimethyl sulfoxide	0.522	1.720	0.000	0.970	3.459	0.6126	-51.44	69
Benzyl alcohol	0.803	0.870	0.330	0.560	4.221	0.9160	-57.40	43
2-Phenylethanol	0.811	0.860	0.310	0.650	4.628	1.0569	-62.37	70
3-Phenyl-1-propanol	0.821	0.940	0.310	0.650	5.312	1.1980	-65.00	70
Diethyl ether	0.041	0.250	0.000	0.450	2.015	0.7309	-25.50	27
Diisopropyl ether	-0.060	0.160	0.000	0.580	2.530	1.0127	-28.95	27
Methyl tert-butyl ether	0.024	0.210	0.000	0.590	2.380	0.8718	-29.09	27
Ethyl tert-butyl ether	-0.020	0.160	0.000	0.600	2.720	1.0127	-30.73	27
Methyl tert-amyl ether	0.050	0.210	0.000	0.600	2.916	1.0127	-33.78	27
Tetrahydropyran	0.275	0.470	0.000	0.550	3.057	0.7672	-34.33	71
Tetrahydrofuran	0.289	0.520	0.000	0.480	2.636	0.6223	-31.72	71
1,4-Dioxane	0.329	0.750	0.000	0.640	2.892	0.6810	-39.01	71
1,3-Dioxolane	0.298	0.510	0.000	0.620	1.830	0.5401	-36.10	71
12-Crown-4	0.420	0.990	0.000	1.390	5.135	1.3620	-68.73	74
18-Crown-6	0.400	1.340	0.000	2.130	7.919	2.0430	-108.86	72
Methanol	0.278	0.440	0.430	0.470	0.970	0.3082	-31.52	43
Ethanol	0.246	0.420	0.370	0.480	1.485	0.4491	-33.85	43
Propan-1-ol	0.236	0.420	0.370	0.480	2.031	0.5900	-36.80	43
Propan-2-ol	0.212	0.360	0.330	0.560	1.764	0.5900	-34.83	75
Butan-1-ol	0.224	0.420	0.370	0.480	2.601	0.7309	-41.31	43
2-Butanol	0.217	0.360	0.330	0.560	2.338	0.7309	-38.36	75
tert-Butanol	0.180	0.300	0.310	0.600	1.963	0.7309	-36.10	43
Pentan-1-ol	0.219	0.420	0.370	0.480	3.106	0.8718	-45.31	43
2-Pentanol	0.195	0.360	0.330	0.560	2.840	0.8718	-39.35	66
3-Pentanol	0.195	0.360	0.330	0.560	2.860	0.8718	-39.68	66
2-Methyl-1-butanol	0.219	0.390	0.370	0.480	3.011	0.8718	-39.90	43
2-Methyl-2-butanol	0.194	0.300	0.310	0.600	2.630	0.8718	-38.87	66
3-Methyl-1-butanol	0.192	0.390	0.370	0.480	3.011	0.8718	-39.95	66
3-Methyl-2-butanol	0.194	0.330	0.330	0.560	2.793	0.8718	-38.84	66
Ethan-1,2-diol	0.404	0.900	0.580	0.780	2.661	0.5078	-54.35	67
2-Methoxyethanol	0.269	0.500	0.300	0.840	2.490	0.6487	-40.20	76
2-Ethoxyethanol	0.237	0.520	0.310	0.810	2.792	0.7896	-42.42	76
Benzene	0.610	0.520	0.000	0.140	2.786	0.7176	-32.30	43
Toluene	0.601	0.520	0.000	0.140	3.325	0.8573	-35.65	43
Ethylbenzene	0.613	0.510	0.000	0.150	3.778	0.9982	-39.19	43
Isopropylbenzene	0.602	0.490	0.000	0.160	4.084	1.1391	-41.38	43
1,2-Dimethylbenzene	0.663	0.560	0.000	0.160	3.939	0.9982	-39.90	27
1,3-Dimethylbenzene	0.623	0.520	0.000	0.160	3.839	0.9982	-39.31	43

1,4-Dimethylbenzene	0.613	0.520	0.000	0.160	3.839	0.9982	-38.88	27
Biphenyl	1.360	0.990	0.000	0.260	6.014	1.3240	-62.65	43
Fluorobenzene	0.477	0.570	0.000	0.100	2.788	0.7341	-34.98	43
Chlorobenzene	0.718	0.650	0.000	0.070	3.657	0.8388	-39.24	43
Bromobenzene	0.882	0.730	0.000	0.090	4.041	0.8914	-41.55	43
Iodobenzene	1.188	0.820	0.000	0.120	4.502	0.9746	-46.17	43
Nitrobenzene	0.871	1.110	0.000	0.280	4.557	0.8906	-53.89	43
Dichloromethane	0.387	0.570	0.100	0.050	2.019	0.4943	-27.62	27
Chloroform	0.425	0.490	0.150	0.020	2.480	0.6167	-32.19	27
1,2-Dichloroethane	0.416	0.640	0.100	0.110	2.573	0.6352	-32.47	27
trans-1,2-Dichloroethylene	0.425	0.410	0.090	0.050	2.278	0.5922	-29.00	73
Trichloroethylene	0.524	0.370	0.080	0.030	2.997	0.7146	-33.78	73
Tetrachloroethylene	0.639	0.440	0.000	0.000	3.584	0.8370	-36.51	73
Carbon Dioxide	0.000	0.280	0.050	0.100	0.058	0.2809	-15.70	77
Nitromethane	0.313	0.950	0.060	0.310	1.892	0.4237	-38.64	55
Sulfur dioxide	0.370	0.660	0.280	0.100	0.778	0.3465	-31.38	78
1-Bromoadamantane	1.070	0.900	0.000	0.200	6.130	1.3668	-52.30	41
1-Adamantanol	0.940	0.900	0.310	0.660	5.634	1.2505	-61.10	41




Figure 1. A plot of the calculated values of $\Delta H_{\text{Solv,DMSO}}$ on eqn (8) against the observed values




Figure 2. A plot of the calculated values of $\Delta H_{\text{Solv,PC}}$ on eqn (12) against the observed values.