

ENTHALPY OF SOLVATION CORRELATIONS FOR ORGANIC SOLUTES AND GASES
DISSOLVED IN 1-PROPANOL AND TETRAHYDROFURAN

Timothy W. Stephens^a, Vicky Chou^a, Amanda N. Quay^a, William E. Acree, Jr.^{a*} and Michael H. Abraham^b

^a Department of Chemistry, 1155 Union Circle # 305070, University of North Texas,
Denton, TX 76203-5017 (U.S.A.)

^b Department of Chemistry, University College London, 20 Gordon Street,
London, WC1H 0AJ (U.K.)

Abstract

Data have been assembled from the published literature on the enthalpies of solvation for 103 organic vapors and gaseous solutes in 1-propanol and for 86 gaseous compounds in tetrahydrofuran. It is shown that an Abraham solvation equation with five descriptors can be used to correlate the experimental solvation enthalpies to within standard deviations of 2.35 kJ/mole and 2.10 kJ/mole for 1-propanol and tetrahydrofuran, respectively. The derived correlations provide very accurate mathematical descriptions of the measured enthalpy of solvation data at 298 K, which in the case of 1-propanol span a range of 119 kJ/mole. Division of the experimental values into a training set and a test set shows that there is no bias in predictions, and that the predictive capability of the correlations is better than 3.5 kJ/mole.

Key Words: Enthalpy of solvation, enthalpy of transfer, mathematical correlation, partition coefficient, solvation parameter model

* To whom correspondence should be addressed. e-mail: acree@unt.edu

1. Introduction

Linear free energy relationship (LFER) models represent a consolidated computational method to mathematically correlate thermodynamic transfer properties and permeabilities of chemical compounds to molecular features (called molecular descriptors) that range in type from topological and structural indices to electronic and quantum-chemical properties. Representative properties that have been correlated with LFER models include sorption of organic compounds on polydimethylsiloxane coated fibers [1-3], chromatographic retention times of organic compounds and drug molecules on immobilized artificial membrane chromatographic columns [4-6], solute partitioning between water and an immiscible (or partly miscible) organic solvent [7-10], solute partitioning between two partly miscible organic solvents [11-14], drug distribution from blood to the various body organs [15-20], permeability of organic compounds through human skin from aqueous solution [21], brain permeation of neutral molecules and ionic species [22], and enthalpies of solvation of organic vapors and gases in both water [23] and in organic solvent media [23-32]. The molecular descriptors may be of either experimental origin or calculated based solely on molecular structure considerations.

Previously we have applied the Abraham solvation parameter model to correlate experimental enthalpies of solvation of organic vapors and gases in water, $\Delta H_{\text{solv,w}}$, [23]:

$$\begin{aligned} \Delta H_{\text{solv,w}} \text{ (kJ/mol)} = & -13.310(0.457) + 9.910(0.814) \mathbf{E} + 2.836(0.807) \mathbf{S} - 32.010(1.102) \mathbf{A} \\ & - 41.816(0.781) \mathbf{B} - 6.354(0.200) \mathbf{L} \end{aligned} \quad (1)$$

(N = 368, SD = 3.68, R² = 0.964, F = 1950.5)

$$\begin{aligned} \Delta H_{\text{solv,w}} \text{ (kJ/mol)} = & -6.952(0.651) + 1.415(0.770) \mathbf{E} - 2.859(0.855) \mathbf{S} - 34.086(1.225) \mathbf{A} \\ & - 42.868(0.850) \mathbf{B} - 22.720(0.800) \mathbf{V} \end{aligned} \quad (2)$$

($N = 369$, $SD = 4.04$, $R^2 = 0.959$, $F = 1688.2$)

and in four alkanes (hexane [27], heptane [29], hexadecane [29] and cyclohexane [29]), in two aromatic hydrocarbons (benzene [29] and toluene [31]), in three chloroalkanes (trichloromethane [30], tetrachloromethane [31] and 1,2-dichloroethane [30]), in five alcohols (methanol [26], ethanol [26], 1-butanol [26], 1-octanol [23] and *tert*-butanol [25]), and in seven other organic solvents (acetone [24], propylene carbonate [32], dimethyl sulfoxide [32], dibutyl ether [28], ethyl acetate [28], acetonitrile [24] and *N,N*-dimethylformamide [25]). In total Abraham model, ΔH_{solv} correlations have been reported for total of 22 different organic solvents. Expressions have also been developed for predicting the enthalpies of solvation of organic vapors and gases into ionic liquid solvents based on the ionic-specific equation coefficient [33] and group contribution [34] versions of the Abraham model.

Each term in Eqns. 1 and 2 represents a different type of solute-solvent interaction contributing to the solute transfer process. The independent variables are solute descriptors, which are defined as follows: **E** denotes the solute excess molar refraction that reflects the solute's ability to interact with the surrounding solvent molecules through π - and lone-electron pairs, **S** is the solute dipolarity/polarizability parameter, **A** and **B** are measures of the solute's hydrogen-bond acidity and basicity, **V** is the McGowan volume of the solute in units of ($\text{dm}^3 \text{mol}^{-1}$)/100, and **L** is the logarithm of the solute's gas phase dimensionless Ostwald partition coefficient into hexadecane at 298 K. The first four descriptors can be regarded as measures of the tendency of the given compounds to undergo various solute-solvent interactions. The latter two descriptors, **V** and **L**, are both measures of solute size, and so will be measures of the solvent cavity term that will accommodate the dissolved solute. General dispersions are also related to solute size; hence both **V** and **L** will also describe the general solute-solvent interactions.

Numerical values of the six equation coefficients in Eqns. 1 and 2 are determined by multiple linear regression analyses of experimental ΔH_{solv} data for a series of organic solutes and gases in the solvent under consideration. The statistics of the derived correlation are the number of experimental values in the regressed database (N), the standard deviation (SD), the squared correlation coefficient (R^2) and the Fisher F-statistic. Standard uncertainties in the calculated equation coefficients are given in parenthesis immediately after the respective coefficient.

During the past two years we have been updating several of our existing gas-to-organic solvent partition coefficient, K, and water-to-organic solvent partition coefficient, P, correlations [35-38], and introducing additional terms into the equation ($j^+ \cdot \mathbf{J}^+$ and $j^- \cdot \mathbf{J}^-$) that enable the prediction of P values of neutral molecules, ions and ionic species at 298 K [39-41]. Previously $\log P$ the correlations had been limited in applicability to just neutral molecules. As part of these studies we have determined solute descriptors for several simple ions and several classes of organic ionic species (carboxylate anions, substituted phenoxide anions, substituted pyridinium cations and protonated amines). The descriptors for anions and cations are **E**, **S**, **A**, **B** and **V**, and are exactly on the same scale as for neutral molecules, together with an additional descriptor, \mathbf{J}^- , for anions and an additional descriptor, \mathbf{J}^+ , for cations.

Our published Abraham model partition coefficient correlations have for the most part pertained to 298 K. Manufacturing and biological processes are not restricted to 298 K, and there is a growing need to estimate gas-to-organic solvent and water-to-organic solvent partition coefficients at other temperatures as well. From a thermodynamic standpoint, ΔH_{Solv} data can be used to estimate the gas-to-condensed phase partition coefficient, K,

$$\log K(at T) - \log K(at 298.15 K) = \frac{-\Delta H_{\text{Solv}}}{2.303R} (1/T - 1/298.15) \quad (3)$$

and the water-to-organic solvent partition coefficient, P,

$$\log P(at T) - \log P(at 298.15 K) = \frac{-\Delta H_{trans}}{2.303R} (1/T - 1/298.15) \quad (4)$$

at other temperatures from measured partition coefficient data at 298.15 K and the solute's enthalpy of solvation, ΔH_{Solv} , or enthalpy of transfer, ΔH_{trans} , between the two condensed phases.

The enthalpy of transfer needed in Eqn. 4 is defined as

$$\Delta H_{trans} = \Delta H_{Solv,Org} - \Delta H_{Solv,W} \quad (5)$$

the difference in the enthalpy of solvation of the solute in the specified organic solvent minus its enthalpy of solvation in water. The above equations assume zero heat capacity changes. The 22 ΔH_{Solv} correlations that we have developed thus far allow us to extrapolate log P and log K values measured at 298 K to other temperatures. Eventually we hope to develop mathematical expressions for predicting enthalpies of solvation in most (if not all) of the more than 70 organic solvents for which we have log K and log P correlations. The present study concerns developing a ΔH_{Solv} correlation for solutes dissolved in 1-propanol and in tetrahydrofuran. The log K and log P correlations for both organic solvents were recently updated [35, 38].

2. Data Sets and Solute Descriptors

A search of the published chemical literature found a large number of papers [42-140] that reported experimental partial molar enthalpies of solution, ΔH_{soln} , of liquid solutes in 1-propanol and tetrahydrofuran, or excess molar enthalpies of binary mixtures containing either 1-propanol or tetrahydrofuran. The ΔH_{soln} data for the liquid solutes were determined by either direct calorimetric methods or calculated based on the temperature dependence of the measured

infinite dilution activity coefficient data. Calorimetric data was also found for the dissolution of a few gases in tetrahydrofuran [141,142], and for five crystalline organic solutes, ethylene carbonate [143], benzamide [144], benzoic acid [145], 1-bromoadamantane [146] and 1-adamantanol [147], in 1-propanol. Enthalpies of solution of several gas molecules [148-155], anthracene [156], acenaphthene [157], adipic acid [158], succinic acid [159] and sebacic acid [160] were calculated from the variation of mole fraction solubility with temperature. The ΔH_{soln} values were converted to gas-to-organic solvent enthalpies of transfer by

$$\text{Liquid solutes:} \quad \Delta H_{\text{Solv}} = \Delta H_{\text{Soln}} - \Delta H_{\text{Vap},298\text{K}} \quad (6)$$

$$\text{Crystalline solutes:} \quad \Delta H_{\text{Solv}} = \Delta H_{\text{Soln}} - \Delta H_{\text{Sub},298\text{K}} \quad (7)$$

subtracting the solute's standard molar enthalpy of vaporization [161, 162], $\Delta H_{\text{Vap},298\text{K}}$, or standard molar enthalpy of sublimation [162, 163], $\Delta H_{\text{Sub},298\text{K}}$, at 298.15 K.

We eliminated from consideration all experimental data that pertained to temperatures outside the temperature range of 283 K – 318 K. Enthalpies of solvation are temperature-dependent, and we did not want to introduce large errors in the database by including experimental data far removed from 298 K. For several solutes there were multiple, independently determined values. In such cases, we selected direct calorimetric data over indirect values based on the temperature dependence of measured solubilities or infinite dilution activity coefficients. Using the aforementioned criteria, 103 molar enthalpies of solvation in 1-propanol and 86 molar enthalpies of solvation in tetrahydrofuran were selected for regression analysis. The experimental $\Delta H_{\text{Solv,PrOH}}$ and $\Delta H_{\text{Solv,THF}}$ values are listed in Tables 1 and 2 along with the values of the solute descriptors of the compounds considered in the present study. The tabulated solute descriptors are of experimental origin and came from our solute descriptor

database, which now contains values for more than 5000 different organic and organometallic compounds. Large tabulations of solute descriptors are available in several earlier publications [23, 165, 166].

The characteristic McGowan volume, **V**, is calculated from the individual atomic sizes and numbers of bonds in the molecule.[164] For liquid solutes, the excess molar refraction descriptor, **E**, is obtained from the liquid refractive index. [167] In the case of solid solutes, one either estimates a hypothetical liquid refractive index using any of several available methods, or one can calculate **E** directly through addition of fragments or substructures. Numerical values of the three remaining descriptors, **S**, **A** and **B**, are determined through regression analysis using available organic solvent/water partition coefficients, chromatographic retention data, solubilities and infinite dilution activity coefficients as described elsewhere. [168-172]. If one is unable to locate sufficient experimental data for performing the fore-mentioned regression analysis, commercial software [173] is available for estimating the molecular solute descriptors from the structure of the compound. Several correlations [174, 175] have been reported for calculating the Abraham solute descriptors from the more structure-based, topological-based and/or quantum-based descriptors used in other QSAR and LFER treatments. Moreover, van Noort and coworkers [176] published an Abraham model correlation for estimating the **L** solute descriptor

$$L = - 0.882 + 1.183 \mathbf{E} + 0.839 \mathbf{S} + 0.454 \mathbf{A} + 0.157 \mathbf{B} + 3.505 \mathbf{V} \quad (8)$$

$$(N = 4785, SD = 0.31, R^2 = 0.992, F = 115279)$$

From known values of **E**, **S**, **A**, **B** and **V**. The authors cited a personnel communication from Dr. Abraham as the source of Eqn. 8.

3. Results and Discussion

We have assembled in Table 1 values of $\Delta H_{\text{Solv,PrOH}}$ for 103 organic vapors and gases dissolved in 1-propanol covering a reasonably wide range of compound type and descriptor value. Analysis of the experimental data yielded the following two Abraham model correlation equations:

$$\begin{aligned} \Delta H_{\text{Solv,PrOH}} \text{ (kJ/mole)} = & -8.713(0.462) - 2.593(1.086) \mathbf{E} + 5.190(1.113) \mathbf{S} - 53.042(1.210) \mathbf{A} \\ & - 7.852(1.094) \mathbf{B} - 8.108(0.193) \mathbf{L} \end{aligned} \quad (9)$$

(with N = 101, SD = 2.41, $R^2 = 0.989$, F = 1678)

$$\begin{aligned} \Delta H_{\text{Solv,PrOH}} \text{ (kJ/mole)} = & -0.114(0.801) - 12.428(1.272) \mathbf{E} - 2.052(1.495) \mathbf{S} - 55.258(1.571) \mathbf{A} \\ & - 7.964(1.426) \mathbf{B} - 30.202(0.955) \mathbf{V} \end{aligned} \quad (10)$$

(with N = 103, SD = 3.13, $R^2 = 0.981$, F = 986.2)

All regression analyses were performed using Version 17 of the SPSS statistical software. There is little intercorrelation between the descriptors in Eqs. 9 and 10. Both correlations provide a good statistical fit of the observed data with standard deviations of 2.41 and 3.13 kJ/mole for a data set that covers a range of about 119.0 kJ/mole. See Figure 1 for a plot of the calculated values $\Delta H_{\text{Solv,PrOH}}$ based on Eqn. 9 against the observed values. Equation 9 is the better equation statistically, and from a thermodynamic standpoint Eqn. 9 is the enthalpic derivative of the Abraham model's gas-to-condensed phase transfer equation. Equation 10 might be more useful in some predictive applications in instances where the **L**-descriptor is not known. Equation 10 uses the McGowan volume, **V**-descriptor, which is easily calculable from the individual atomic sizes and numbers of bonds in the molecule [164]. To our knowledge, Eqns. 9 and 10 are the

first expressions that allow one to predict the enthalpy of solvation of gaseous solutes in 1-propanol.

In order to assess the predictive ability of Eqn. 9 we divided the 103 data points into a training set and a test set by allowing the SPSS software to randomly select half of the experimental points. The selected data points became the training set and the compounds that were left served as the test set. Analysis of the experimental data in the training set gave

$$\begin{aligned} \Delta H_{\text{Solv,PrOH}} \text{ (kJ/mole)} = & -8.712(0.723) - 7.321(1.665) \mathbf{E} + 7.590(1.889) \mathbf{S} - 53.653(1.871) \mathbf{A} \\ & - 8.950(1.690) \mathbf{B} - 8.159(0.269) \mathbf{L} \end{aligned} \quad (11)$$

(with $N = 52$, $SD = 2.70$, $R^2 = 0.988$, $F = 767.4$)

There is very little difference in the equation coefficients for the full dataset and the training dataset correlations, thus showing that the training set of compounds is a representative sample of the total data set. The training set equation was then used to predict $\Delta H_{\text{Solv,PrOH}}$ values for the 51 compounds in the test set. For the predicted and experimental values, we find $SD = 2.26$, AAE (average absolute error) = 1.81 and AE (average error) = -0.10 kJ/mole. There is therefore very little bias in using Eqn. 11 with AE equal to -0.10 kJ/mole. The training set and test set analyses were performed two more times with similar results. Training and test validations were also performed for Eqn. 10. To conserve journal space, we give only the test set results. The derived training set correlation for Eqn. 10 predicted the 51 experimental $\Delta H_{\text{Solv,PrOH}}$ values in the test set to within a $SD = 3.14$, $AAE = 2.44$ and $AE = 0.01$ kJ/mole. Again, there is very little bias in the predictions using Eqn. 10 with AE equal to 0.01 kJ/mole. An error/uncertainty of ± 2 kJ/mole in the enthalpy of solvation results in an error of slightly less than 0.04 log units in

extrapolating a log K value measured at 298.15 K to a temperature of 313.15. This level of predictive error will be sufficient for most practical chemical and engineering applications.

In Table 2 are collected values of the enthalpies of solvation of 81 gaseous solutes in tetrahydrofuran. Preliminary analysis of the experimental $\Delta H_{\text{Solv,THF}}$ data yielded correlation equations having very small b coefficients, would be expected from the molecular structure considerations. Tetrahydrofuran does not have an acidic hydrogen. The b-coefficients were set equal to zero, and the final regression analyses performed to give:

$$\begin{aligned} \Delta H_{\text{Solv,THF}} \text{ (kJ/mole)} &= -6.040(0.437) + 3.640(1.223) \mathbf{E} - 14.478(1.252) \mathbf{S} - 40.652(1.455) \mathbf{A} \\ &\quad - 8.537(0.150) \mathbf{L} \end{aligned} \quad (12)$$

(with N = 86, SD = 2.10, $R^2 = 0.990$, F = 1997)

$$\begin{aligned} \Delta H_{\text{Solv,THF}} \text{ (kJ/mole)} &= 4.777(0.894) - 6.642(1.146) \mathbf{E} - 23.110(1.502) \mathbf{S} - 43.222(1.764) \mathbf{A} \\ &\quad - 33.683(0.724) \mathbf{V} \end{aligned} \quad (13)$$

(with N = 86, SD = 2.55, $R^2 = 0.985$, F = 1346)

Both Eqns. 12 and 13 are statistically very good with standard deviations of 2.10 and 2.55 kJ/mole for a dataset that covers an approximate range of 91.0 kJ/mole. Both equations were validated through training and test set analyses. Figure 2 compares the calculated values of $\Delta H_{\text{Solv,THF}}$ based on Eqn. 12 against the observed data. To our knowledge there has been no previous attempt to correlate enthalpies of solvation for gaseous solutes in tetrahydrofuran.

The predictive ability of Eqn. 13 was assessed as before by allowing the SPSS software to randomly divide the 86 experimental data points into training and test sets. Analyses of the experimental data in the training set gave

$$\Delta H_{\text{Solv,THF}} \text{ (kJ/mole)} = -6.549(0.596) + 5.800(1.797) \mathbf{E} - 15.407(1.876) \mathbf{S} - 42.258(2.357) \mathbf{A} \\ - 8.448(0.186) \mathbf{L} \quad (14)$$

(with N = 43, SD = 1.94, R² = 0.990, F = 942.6)

There is very little difference in the equation coefficients for the full dataset and the training dataset correlations, thus showing that the training set of compounds is a representative sample of the total data set. The training set equation was then used to predict $\Delta H_{\text{Solv,THF}}$ values for the 43 compounds in the test set. For the predicted and experimental values, we find SD = 2.44, AAE = 1.97 and AE = 0.35 kJ/mole. There is therefore very little bias in using Eqn. 14 with AE equal to 0.35 kJ/mole. The training set and test set analyses were performed two more times with similar results. Training and test validations were also performed for Eqn. 13. To conserve journal space, we give only the test set results. The derived training set correlation for Eqn. 13 predicted the 43 experimental $\Delta H_{\text{Solv,THF}}$ values in the test set to within a SD = 3.03, AAE = 2.26 and AE = 0.54 kJ/mole. Again, there is very little bias in the predictions using Eqn. 13 with AE equal to 0.54 kJ/mole.

The correlations presented in this study further document the applicability of the Abraham solvation parameter model to describe enthalpies of solvation for organic vapors and gaseous solutes dissolved in organic solvents. The derived ΔH_{Solv} correlations for 1-propanol and tetrahydrofuran will allow one to extrapolate gas-to-PrOH(or THF) and water-to-PrOH(or THF) measured at 298.15 K to other temperatures. Not all manufacturing applications occur at

298.15 K, and there is a growing need in the chemical industry to predict solute transfer and partition properties at other temperatures as well.

References

- [1] E.L. Di Filippo, R.P. Eganhouse, Assessment of PDMS-water partition coefficients: implications for passive environmental sampling of hydrophobic organic compounds, *Environ. Sci. Technol.* 44 (2010) 6917-6925.
- [2] P. Prikryl, J.G.K. Sevcik, Characterization of sorption mechanisms of solid-phase microextraction with volatile organic compounds in air samples using a linear solvation energy relationship approach, *J. Chromatogr. A* 1179 (2008) 24-32.
- [3] H.P.H. Arp, S. Endo, K.-U. Goss, Comment on "assessment of PDMS-water partition coefficients: implications for passive environmental sampling of hydrophobic compounds", *Environ. Sci. Technol.* 44 (2010) 8787-8788.
- [4] L. Sprunger, B.H. Blake-Taylor, A. Wairegi, W.E. Acree, Jr., M.H. Abraham, Characterization of the retention behavior of organic and pharmaceutical drug molecules on an immobilized artificial membrane column with the Abraham model, *J. Chromatogr. A* 1160 (2007) 235-245.
- [5] K. Valko, M. Plass, C. Bevan, D. Reynolds, M.H. Abraham, Relationships between the chromatographic hydrophobicity indices and solute descriptors obtained by using several reversed-phase, diol, nitrile, cyclodextrin and immobilized artificial membrane-bonded high-performance liquid chromatography columns, *J. Chromatogr. A* 797 (1998) 41-55.
- [6] H. Du, J. Watzl, J. Wang, X. Zhang, X. Yao, Z. Hu, Prediction of retention indices of drugs based on immobilized artificial membrane chromatography using projection pursuit regression and local lazy regression. *J. Sep. Sci.* 31 (2008) 2325-2333.

- [7] M.H. Abraham, W.E. Acree, Jr., A.J. Leo, D. Hoekman, Partition of compounds from water and from air into the wet and dry monohalobenzenes, *New J. Chem.* 33 (2009) 1685-1692.
- [8] M.H. Abraham, W.E. Acree, Jr., A.J. Leo, D. Hoekman, The partition of compounds from water and from air into wet and dry ketones, *New J. Chemistry* 33 (2009) 568-573.
- [9] L.M. Sprunger, A. Proctor, W.E. Acree, Jr., M.H. Abraham, N. Benjelloun-Dakhama, Correlation and prediction of partition coefficient between the gas phase and water, and the solvents dry methyl acetate, dry and wet ethyl acetate, and dry and wet butyl acetate. *Fluid Phase Equilibr.* 270 (2008) 30-44.
- [10] M.H. Abraham, A. Nasezadeh, W.E. Acree, Jr., Correlation and prediction of partition coefficients from the gas phase and from water to alkan-1-ols. *Ind. Eng. Chem. Res.* 47 (2008) 3990-3995.
- [11] T. Karunasekara, C.F. Poole, Model for the partition of neutral compounds between n-heptane and formamide. *J. Sep. Sci.* 33 (2010) 1167-1173.
- [12] J. Qian, C.F. Poole, Distribution of neutral organic compounds between n-heptane and fluorine-containing alcohols. *J. Chromatogr. A* 1143 (2007) 276-283.
- [13] H. Ahmed, C.F. Poole, Distribution of neutral organic compounds between n-heptane and methanol or N,N-dimethylformamide, *J. Sep. Sci.* 29 (2006) 2158-2165.
- [14] H. Ahmed, C.F. Poole, Model for the distribution of neutral organic compounds between n-hexane and acetonitrile, *J. Chromatogr. A* 1104 (2006), 82-90.
- [15] M.H. Abraham, A. Ibrahim, W.E. Acree, Jr., Air to lung partition coefficients for volatile organic compounds and blood to lung partition coefficients for volatile organic compounds and drugs. *Eur. J. Med. Chem.* (2008), 43, 478-485.

- [16] M.H. Abraham, A. Ibrahim, W.E. Acree, Jr., Air to liver partition coefficients for volatile organic compounds and blood to liver partition coefficients for volatile organic compounds and drugs, *Eur. J. Med. Chem.* 42 (2007) 743-751.
- [17] M.H. Abraham, A. Ibrahim, Air to fat and blood to fat distribution of volatile organic compounds and drugs: linear free energy analyses, *Eur. J. Med. Chem.* 41 (2006) 1430-1438.
- [18] M.H. Abraham, A. Ibrahim, Y. Zhao, W.E. Acree, Jr., A data base for partition of volatile organic compounds and drugs from blood/plasma/serum to brain, and an LFER analysis of the data, *J. Pharm. Sci.* 95 (2006) 2091-2100.
- [19] M.H. Abraham, A. Ibrahim, W. E. Acree, Jr., Air to brain, blood to brain and plasma to brain distribution of volatile organic compounds: linear free energy analyses, *Eur. J. Med. Chem.* 41 (2006) 494-502.
- [20] M. H. Abraham, A. Ibrahim, W.E. Acree, Jr., Air to muscle and blood/plasma to muscle distribution of volatile organic compounds and drugs: linear free energy analyses, *Chem. Res. Toxicol.* 19 (2006) 801-808.
- [21] M.H. Abraham, F. Martins, Human skin permeation and partition: general linear free-energy relationship analyses, *J. Pharm. Sci.* 93 (2004) 1508-1523.
- [22] M. H. Abraham, The permeation of neutral molecules, ions, and ionic species through membranes: brain permeation as an example, *J. Pharm. Sci.* (2011) in press.
- [23] C. Mintz, M. Clark, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in water and in 1-octanol based on the Abraham model, *J. Chem. Inf. Model.* 47 (2007) 115-121.

- [24] C. Mintz, J. Gibbs, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for organic solutes and gases dissolved in acetonitrile and acetone, *Thermochim. Acta* 484 (2009) 65-69.
- [25] C. Mintz, K. Burton, T. Ladlie, M. Clark, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for organic solutes and gases dissolved in N,N-dimethylformamide and tert-butanol, *J. Mol. Liq.* 144 (2009) 23-31.
- [26] C. Mintz, T. Ladlie, K. Burton, M. Clark, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in alcohol solvents based on the Abraham model, *QSAR Comb. Sci.* 27 (2008) 627-635.
- [27] C. Mintz, K. Burton, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in linear alkanes (C₅-C₁₆) based on the Abraham model, *QSAR Comb. Sci.* 27 (2008) 179-186.
- [28] C. Mintz, K. Burton, T. Ladlie, M. Clark, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in dibutyl ether and ethyl acetate, *Thermochim. Acta* 470 (2008) 67-76.
- [29] C. Mintz, M. Clark, K. Burton, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in benzene and in alkane solvents based on the Abraham model, *QSAR Comb. Sci.* 26 (2007) 881-888.
- [30] C. Mintz, K. Burton, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in chloroform and 1,2-dichloroethane based on the Abraham model, *Fluid Phase Equilibr.* 258 (2007) 191-198.

- [31] C. Mintz, M. Clark, K. Burton, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in toluene and carbon tetrachloride based on the Abraham model, *J. Solution Chem.* 36 (2007) 947-966.
- [32] C. Mintz, K. Burton, W.E. Acree, Jr., M.H. Abraham, Enthalpy of solvation correlations for gaseous solutes dissolved in dimethyl sulfoxide and propylene carbonate based on the Abraham model, *Thermochim. Acta* 459 (2007) 17-25.
- [33] L.M. Sprunger, S.S. Achi, W.E. Acree, Jr., M.H. Abraham, Linear free energy relationship correlations for enthalpies of solvation of organic solutes into room-temperature ionic liquids based on the Abraham model with ion-specific equation coefficients, *Ind. Eng. Chem. Res.* 48 (2009) 8704-8709.
- [34] L.M. Grubbs, W.E. Acree, Jr., M.H. Abraham, Correlation of enthalpies of solvation of organic vapors and gases in ionic liquid solvents using a group contribution version of the Abraham solvation parameter model, *Thermochim. Acta* 511 (2010) 96-101.
- [35] L.M. Sprunger, S.S. Achi, R. Pointer, B.H. Blake-Taylor, W.E. Acree, Jr., M.H. Abraham, Development of Abraham model correlations for solvation characteristics of linear alcohols, *Fluid Phase Equilibr.* 286 (2009) 170-174.
- [36] L.M. Sprunger, S.S. Achi, R. Pointer, W.E. Acree, Jr., M.H. Abraham, Development of Abraham model correlations for solvation characteristics of secondary and branched alcohols, *Fluid Phase Equilibria* 288 (2010) 121-127.
- [37] L.M. Grubbs, M. Saifullah, N.E. De La Rosa, S. Ye, S.S. Achi, W.E. Acree, Jr., M.H. Abraham, Mathematical correlations for describing solute transfer into functionalized alkane solvents containing hydroxyl, ether, ester or ketone solvents, *Fluid Phase Equilibr.* 298 (2010) 48-53.

- [38] M. Saifullah, S. Ye, L.M. Grubbs, N.E. De La Rosa, W.E. Acree, Jr., M.H. Abraham, Abraham model correlations for transfer of neutral molecules to tetrahydrofuran and to 1,4-dioxane, and for transfer of ions to tetrahydrofuran, *J. Solution Chem.* submitted for publication.
- [39] M.H. Abraham, W.E. Acree, Jr., Equations for the transfer of neutral molecules and ionic species from water to organic phases, *J. Org. Chem.* 75 (2010) 1006-1015.
- [40] M.H. Abraham, W.E. Acree, Jr., Solute descriptors for phenoxide anions and their use to establish correlations of rates of reaction of anions with iodomethane, *J. Org. Chem.* 75 (2010), 3021-3026.
- [41] M.H. Abraham, W.E. Acree, Jr., The transfer of neutral molecules, ions and ionic species from water to ethylene glycol and to propylene carbonate; descriptors for pyridinium cations, *New J. Chem.* 34 (2010) 2298-2305.
- [42] T.M. Letcher, J. Mercer-Chalmers, U.P. Govender, S. Radloff, Excess molar enthalpies and excess molar volumes of binary mixtures of 1-alkenes with 1-propanol and 2-propanol, *Thermochim. Acta* 224 (1993) 33-38.
- [43] T.M. Letcher, P.U. Govender, Excess molar enthalpies of an alkanol + a branched chain ether at the temperature 298.15 K, *J. Chem. Eng. Data* 40 (1995) 997-1000.
- [44] G.A. Krestov, V.P. Korolev, D.V. Batov, Thermochemistry of solvation of nonelectrolytes in mono- and polyatomic alcohols and their mixtures with water, *Thermochim. Acta* 169 (1990) 69-80.
- [45] J.C. Cobos, I. Garcia, J.A. Gonzalez, C. Casanova, Excess enthalpies of (3,6,9-trioxaundecane + an n-alkan-1-ol) at 298.15 K, *J. Chem. Thermodyn.* 22 (1990) 383-366.

- [46] L. Wang, G.C. Benson, B.C.Y. Lu, Excess enthalpies of {(ethanol or propan-1-ol or methyl 1,1-dimethylethyl ether) + (2,3-dimethylbutane or 2,2,4-trimethyl-pentane)}, J. Chem. Thermodyn. 24 (1992) 1305-1310.
- [47] L. Wang, G.C. Benson, B.C.Y. Lu, Excess enthalpies of 1-propanol + n-hexane + n-decane or n-dodecane at 298.15 K, J. Chem. Eng. Data 37 (1992) 403-406.
- [48] M.H. Abraham, A. Nasehzadeh, Thermodynamic functions for solution of triethylamine in 30 solvents, and calculation of the solute-solvent interaction terms, J. Chem. Thermodyn. 13 (1981) 549-555.
- [49] M.M. Mato, S.M. Cebreiro, J.L. Legido, M.I. Paz Andrade, Excess molar enthalpies of ternary and binary mixtures containing 2-methoxy-2-methylpropane, 1-propanol, and nonane, J. Chem. Eng. Data 54 (2009) 1692-1697.
- [50] M.M. Mato, S.M. Cebreiro, J.L. Legido, M.I. Paz Andrade, Study of the effect of increasing the chain length of the alkane in excess molar enthalpies of mixtures containing methyl *tert*-butyl ether, 1-propanol, alkane, Fluid Phase Equilibr. 271 (2008) 6-12.
- [51] M.A. Varfolomeev, K.V. Zaitseva, I.T. Rakipov, B.N. Solomonov, Enthalpy of cooperative hydrogen bonding in the complexes of triethyl- and tri-n-butylamines with alcohols: effect of the alkyl chain length, Russ. J. Gen. Chem. 80 (2010) 402-407.
- [52] E. Bich, M. Dzida, T. Vasiltsova, A. Heintz, Experimental and theoretical study of thermodynamic excess properties of associating liquid alkanol + 4-picoline mixtures based on quantum mechanical ab initio calculations, J. Mol. Liq. 136 (2007) 94-103.
- [53] D.M. Trampe, C.A. Eckert, Calorimetric measurement of partial molar excess enthalpies at infinite dilution, J. Chem. Eng. Data 36 (1991) 112-118.

- [54] A.V. Kustov, O.A. Antonova, V.P. Korolev, Enthalpies of 1,4-dioxane, 1,2-dimethoxyethane, and ethylacetate solvation in the water-1-propanol and water-Glycerol binary mixtures, *J. Solution Chem.* 31 (2001) 671-680.
- [55] R.M.C. Goncalves, L.M.P.C. Albuquerque, A.M.N. Simoes, J.J.M. Ramos, Calorimetric study of the solution process of tetramethylsilane in alcohols, *Thermochim. Acta* 209 (1992) 63-75.
- [56] L.M.P.C. Albuquerque, M.L.C.J. Moita, A.M.N. Simoes, R.M.C. Goncalves, The enthalpies of solution of 2-chloro-2-methylpropane, 2-chloro-2-methylbutane and 3-chloro-3-methylpentane in mono- and dialcohols at 298.15 K, *Thermochim. Acta* 275 (1996) 67-73.
- [57] H.C. Van Ness, M.M. Abbott, Excess enthalpy. 1-propanol (propyl alcohol)-toluene system, *Int. DATA Series, Selected Data on Mix., Ser. A* (1976) 33-34.
- [58] H.C. Van Ness, M.M. Abbott, Excess enthalpy. 1-propanol (propyl alcohol)-ethylbenzene system, *Int. DATA Ser., Sel. Data on Mix., Ser. A:* (1976) 35-36.
- [59] G.C. Benson, Excess enthalpy. 1-octanol (n-octyl alcohol)-1-propanol (propyl alcohol) system, *Int. DATA Ser., Sel. Data Mix., Ser. A* (1974) 97.
- [60] G.C. Benson, Excess enthalpy. 1-decanol (n-decyl alcohol)-1-propanol (propyl alcohol) system, *Int. DATA Ser., Sel. Data Mix. Ser. A* (1974) 98.
- [61] G.C. Benson, Excess enthalpy. 1-hexanol (n-hexyl alcohol)-1-propanol (propyl alcohol) system, *Int. DATA Ser., Sel. Data Mix., Ser. A* (1974) 96.
- [62] M.A. Villamanan, A.H. Roux, J.P.E. Grolier, J. P. E. Excess enthalpy. 1-propanol - dibutyl ether system, *Int. DATA Ser., Selected Data Mix., Ser. A* (1984) 120.

- [63] M.A. Villamanan, A.H. Roux, J.P.E. Grolier, Excess enthalpy. 1-propanol - butyl methyl ether system, *Int. DATA Ser., Sel. Data Mix., Ser. A* (1984) 124.
- [64] Y.X. Wang, J.P. Chao, M. Dai, Studies on the thermodynamic properties of binary mixtures containing an alcohol. XIII. Excess molar enthalpies of some mixtures of n-alcohols with aniline or chlorobenzene at 298.15 K, *Thermochim. Acta* 169 (1990) 161-169.
- [65] M.A. Villamanan, C. Casanova, A.H. Roux, J.P.E. Grolier, Calorimetric investigation of the interactions between oxygen and hydroxyl groups in (alcohol + ether) at 298.15 K, *J. Chem. Thermodyn.* 14 (1982) 251-258.
- [66] J.J. Christensen, R.L. Rowley, R.M. Izatt, *Handbook of Heats of Mixing: Supplementary Volume*, John Wiley and Sons, Inc., New York, NY, 524 (1988).
- [67] Yu. A. Sivolozhskaya, N.L. Potkina, V.P. Korolev, Thermochemistry of CCl₄ in systems alkanane - 1-alkanol, *Russ. J. Gen. Chem.* 70 (2000) 1399-1403.
- [68] D.V. Batov, O.A. Antonova, V.P. Slyusar, V. P. Korolev, Solvation and state of acetonitrile, propylene carbonate and nitromethane molecules in propanol and butanol isomers and in their mixtures with water at 298.15 K, *Russ. J. Gen. Chem.* 69 (1999) 1212-1218.
- [69] J. Mokrzan, Enthalpy of solution of benzoic aldehyde in aliphatic alcohols at 298.15 K, *Acta Univ. Lodzianis, Fol. Chim.* 4 (1985) 71-78.
- [70] E.D. Totchasov, M.Yu. Nikiforov, G.A. Al'per, Calculations of the enthalpy of solution of nonelectrolytes in mixed solvents within the framework of molecular association theory, *Russ. J. Phys. Chem. A* 83 (2009) 45-49.

- [71] I. McStravick, E. Duffy, W.E. Waghorne, Enthalpies of transfer of $-CH_2-$ between water and nonaqueous or mixed aqueous solvents, comparison of values from n-alcohols and tetraalkylammonium halides, *J. Mol. Liq.* 103-104 (2003) 121-131.
- [72] H. Iloukhani, Excess molar enthalpies for binary mixtures with trichloroethylene + alkan-1-ols (C_3-C_8) at 298.15 K, *J. Chem. Eng. Data* 42 (1997) 802-804.
- [73] R.M.C. Goncalves, L.M.P.C. Albuquerque, F.E.L. Martins, A.M.N. Simoes, J.J.M. Ramos, Enthalpies of solution and intermolecular forces. *tert*-butyl halides in hydroxylic solvents, *J. Phys. Org. Chem.* 5 (1992) 93-100.
- [74] T.M. Letcher, U.P. Govender, Excess molar enthalpies of an alkanol + a cyclic ether at 298.15 K, *J. Chem. Eng. Data* 40 (1995) 1097-1100.
- [75] T.M. Letcher, J.A. Nevines, R.P. Vijayan, S.E. Radloff, Excess enthalpies of (a chlorinated benzene + an alkanol) at the temperature 298.15 K, *J. Chem. Thermodyn.* 25 (1993) 379-383.
- [76] S.G. Collins, J.J. Christensen, R.M. Izatt, R.W. Hanks, The excess enthalpies of 10 n-pentane + n-alkanol mixtures at 298.15 K, *J. Chem. Thermodyn.* 12 (1980) 609-614.
- [77] T.M. Letcher, J. Mercer-Chalmers, A.K. Prasad, Excess molar enthalpies of mixtures of a cycloalkane and an alkanol, *Thermochim. Acta* 188 (1991) 157-162.
- [78] K. Tamura, I. Nagata, Excess molar enthalpies of (propan-1-ol + butan-2-one) and of (propan-1-ol or propan-2-ol + butan-2-one + benzene) at the temperature 298.15 K, *J. Chem. Thermodyn.* 23 (1991) 359-364.
- [79] I. Nagata, Excess molar enthalpies of binary and ternary mixtures containing aniline and 1-propanol, *Thermochim. Acta* 208 (1992) 73-82.

- [80] S.M. Cebreiro, M. Illobre, M.M. Mato, V.V. Verdes, J.L. Legido, M.I. Paz Andrade, Excess molar enthalpies at 298.15 K of the binary mixtures. *tert*-butyl methyl ether + alcohol ($n = 1, 2, 3, 5$), *J. Therm. Anal. Calorim.* 70 (2000) 251-254.
- [81] H.C. Van Ness, M.M. Abbott, Excess enthalpy. 1-propanol (propyl alcohol)-benzene system, *Int. DATA Ser., Sel. Data Mix., Ser. A* (1976) 22-23.
- [82] I. Nagata, K. Takeuchi, Excess molar enthalpies for 1-propanol + acetone and 1-propanol + acetone + benzene mixtures at 298.15 K, *Thermochim. Acta* 242 (1994) 103-110.
- [83] R. Garriga, F. Sanchez, P. Perez, M. Gracia, Vapor pressures at several temperatures T and excess functions at $T = 298.15$ K of (butanenitrile + propan-1-ol or propan-2-ol), *J. Chem. Thermodyn.* 27 (1995) 887-895.
- [84] K. Tamura, M.M.H. Bhuiyan, Excess molar enthalpies of (methanol + 1-propanol) + oxane or 1,4-dioxane mixtures at the temperature 298.15 K, *J. Chem. Thermodyn.* 35 (2003) 1657-1669.
- [85] K.C. Singh, K.C. Kalra, S. Maken, V. Gupta, Excess heat of mixing of 1-propanol or 2-propanol with benzene, toluene, *o*-, *m*- and *p*-xylenes at 298.15 K, *Thermochim. Acta* 276, (1996) 271-285.
- [86] E. Jimenez, L. Segade, C. Franjo, M.M. Pineiro, M.I. Paz Andrade, Experimental and predicted excess molar volumes and excess molar enthalpies for di-*n*-butyl ether + 1-propanol + *n*-octane at 298.15 and 308.15 K, *Fluid Phase Equilibr.* 179 (2001) 363-38.
- [87] A. Pineiro, A. Olvera, G.. Garcia-Miaja, M. Costas, Excess molar enthalpies of tetrahydrofuran or diisopropyl ether + 1-alkanols at 298.15 K, using a newly designed flow mixing cell for an isothermal microcalorimeter, *J. Chem. Eng. Data* 46 (2001) 1274-1279.

- [88] L.M.P.C. Albuquerque, M.L.C.J. Moita, A.M.N. Simoes, R.M.C. Goncalves, The enthalpies of solution of 2-bromo-2-methylpropane and 2-bromo-2-methylbutane in mono- and dialcohols at 298.15 K, *Thermochim. Acta* 322 (1998) 57-61.
- [89] J. Ortega, C. Gonzalez, J. Pena, S. Galvan, Thermodynamic study on binary mixtures of propyl ethanoate and an alkan-1-ol (C₂-C₄). Isobaric vapor-liquid equilibria and excess properties, *Fluid Phase Equilib.* 170 (2000) 87-111.
- [90] M. Jozwiak, The effect of properties of water-organic solvent mixtures on the solvation enthalpy of 12-crown-4, 15-crown-5, 18-crown-6 and benzo-15-crown-5 ethers at 298.15 K, *Thermochim. Acta* , 417 (2004) 31-41.
- [91] T. Kimura, T. Kitai, T. Kamiyama, M. Fujisawa, Excess enthalpies of binary mixtures of some propylamines + some propanols at 298.15 K, *Thermochim. Acta* 450(2006) 91-95.
- [92] S.-J. Liu, H.-L. Liu, Q.-Y. Chen, Z.-L. Xiao, D.-W. Yang, Molar excess enthalpies of binary mixtures of [tributyl phosphate + {CH₃(CH₂)_nOH (n =0 to 3)}] at 298.15K, *J. Chem. Thermodyn.* 39 (2007) 412-416.
- [93] K. Liu, J. Xu, S. Liu, Q. Chen, Z.-L. Xiao, D.-W. Yang, Excess molar enthalpies for binary mixtures of trimethyl phosphate with alkanols {CH₃(CH₂)_nOH, n = 0-3} at 298.15 K, *Thermochim. Acta* 471 (2008) 100-102.
- [94] T. Minamihonoki, H. Ogawa, H. Nomura, S. Murakami, Thermodynamic properties of binary mixtures of 2,2,2-trifluoroethanol with water or alkanols at T = 298.15K, *Thermochimica Acta* 459 (2007) 80-86.
- [95] J. Polak, S. Murakami, G.C. Benson, H.D. Pflug, Excess enthalpy, volume, and Gibbs free energy of n-propanol-isopropanol mixtures at 25.deg, *Can. J. Chem.* 48 (1970) 3782-3785.

- [96] G. Carthy, D. Feakins, C. O'Duinn, E.W. Waghorne, Enthalpies of transfer of formamide, N-methylformamide, N,N-dimethylformamide and urea from water to some aqueous alcohol solvent systems at 298 K, *J. Chem. Soc., Faraday Trans.* 87 (1991) 2447-2452.
- [97] G. C. Benson, Excess enthalpy. 1-propanol (propyl alcohol)-methanol (methyl alcohol) system, *Int. DATA Ser., Sel. Data Mix., Ser. A:* (1973) 77.
- [98] W.C. Duer, G.L. Bertrand, Thermochemical isotope effects. II. Methanol-methanol-d, ethanol-ethanol-d, and water-water-d₂ in selected solvents, *J. Am. Chem. Soc.* (1975)97 (1975) 3894-3897.
- [99] W.C. Duer, G.L. Bertrand, Thermochemical isotope effects. chloroform/chloroform-d and acetone/acetone-d₆ in selected solvents, *J. Am. Chem. Soc.* 96 (1974) 1300-1304.
- [100] B. Ruiz, S. Otin, C. Gutierrez Losa, Excess molar enthalpies at 298.15 K of chlorobenzene +, (chloromethyl)benzene +, and (2-chloroethyl)benzene + an ether, *J. Chem. Thermodyn.* 16 (1984) 25-32.
- [101] A. Inglese, W. Wilhelm, J.P.E. Grolier, H.V. Kehiaian, Thermodynamics of binary mixtures containing cyclic ethers. 1. excess enthalpies of oxolane, 1,3-dioxolane, oxane, 1,3-dioxane, and 1,4-dioxane with n-alkanes, *J. Chem. Thermodyn.* 12 (1980) 217-222.
- [102] J. Catalan, J. Gomez, A. Couto, J. Laynez, Toward a solvent basicity scale: the calorimetry of the pyrrole probe, *J. Am. Chem. Soc.* 112 (1990) 1678-1681.
- [103] H.C. Van Ness, M.M. Abbott, Excess enthalpy. trichloromethane (chloroform)-oxolane (tetrahydrofuran) system, *Int. DATA Series, Sel. Data Mix., Ser. A* (1974) 67.
- [104] H.C. Van Ness, M.M. Abbott, M. M. Excess enthalpy. dichloromethane (methylene chloride)-oxolane (tetrahydrofuran) system, *Int. DATA Ser., Sel. Data Mix., Ser. A* (1974) 65.

- [105] H.C. Van Ness, M.M. Abbott, Excess enthalpy. tetrachloromethane (carbon tetrachloride)-oxolane (tetrahydrofuran) system, *Int. DATA Ser., Sel. Data Mix., Ser. A* (1974) 63.
- [106] J. Catalan, A. Couto, J. Gomez, J.L. Saiz, J. Laynez, Towards a solvent acidity scale: the calorimetry of the N-methyl imidazole probe, *J. Chem. Soc., Perkin Trans. 2* (1992) 1181-5.
- [107] F. Comelli, R. Francesconi, Excess molar enthalpies and excess molar volumes of 1,2,4-trimethylbenzene + cyclic ethers at 298.15 K, *J. Chem. Eng. Data* 37 (1992) 319-322.
- [108] F. Comelli, R. Francesconi, Excess molar enthalpies of dimethyl carbonate or diethyl carbonate + cyclic ethers at 298.15 K, *J. Chem. Eng. Data* 40 (1995) 28-30.
- [109] F. Comelli, R. Francesconi, Calorimetric study of binary mixtures containing chlorobenzene + linear and cyclic ethers, *Thermochim. Acta* 196 (1992) 177-180.
- [110] F. Comelli, R. Francesconi, Liquid-phase excess enthalpies for the binary mixtures 1-chloronaphthalene + cyclic ethers, *J. Chem. Eng. Data* 36 (1991) 382-383.
- [111] R. Francesconi, F. Comelli, Excess molar enthalpies for the binary systems 1-methylnaphthalene + linear and cyclic ethers, *J. Chem. Eng. Data* 38 (1993) 221-223.
- [112] C. Lafuente, P. Cea, M. Dominguez, F.M. Royo, J.S. Urieta, Excess molar enthalpies of cyclic ethers with cyclohexane, methylcyclohexane, or chlorocyclohexane, *J. Solution Chem.* 30 (2001) 795-805.
- [113] S. Cabani, N. Ceccanti, Thermodynamic properties of binary mixtures of cyclohexane with cyclic amines or cyclic ethers at 298.15.deg.K, *J. Chem. Thermodyn.* 5 (1973) 9-20.

- [114] R. Francesconi, B. Lunelli, F. Comelli, Excess molar enthalpies and excess molar volumes of binary mixtures containing p-xylene + cyclic ethers, *Thermochim. Acta* 207 (1992) 45-52.
- [115] T.M. Letcher, P.U. Govender, U. Domanska, Excess molar enthalpies and volumes of diethylamine or dipropylamine + an ether at 298.15 K, *J. Chem. Eng. Data* 44(1999) 274-285.
- [116] A.M. Fanni, B. Marongiu, B. Pittau, S. Porcedda, DISQUAC predictions of excess enthalpies of the ternary mixture: tetrahydrofuran + cyclohexane + butanenitrile, *Fluid Phase Equilib.* 126 (1996) 163-175.
- [117] E.M. Arnett, L. Joris, E. Mitchell, T.S.S.R. Murty, T.M. Gorrie, P.v.R. Schleyer, Hydrogen-bonded complex formation. III. Thermodynamics of complexing by infrared spectroscopy and calorimetry, *J. Am. Chem. Soc.* 92 (1970) 2365-77.
- [118] S.M. Nazmul Hasan, D. Y. Peng, Excess molar enthalpies of dipropyl ether + dibutyl ether + (1-hexene or tetrahydrofuran) at 298.15 K, *J. Chem. Eng. Data*, ACS ASAP.
- [119] D. Kodama, Z. Wang, B.C.-Y. Lu, Excess molar enthalpies of the ternary mixtures: {tetrahydrofuran or 2-methyltetrahydrofuran + methyl *tert*-butyl ether + n-dodecane} at the temperature 298.15 K, *J. Chem. Thermodyn.* 37 (2005) 854-859.
- [120] I. Castro, M. Pintos, A. Amigo, R. Bravo, M.I. Paz Andrade, Excess enthalpies of (tetrahydrofuran or tetrahydropyran + an n-alkane) at the temperature 298.15 K, *J. Chemical Thermodyn.* 26 (1994) 29-33.
- [121] T.M. Letcher, U. Domanska, The excess enthalpies of (acetonitrile + an ether) at the temperature 298.15 K, *J. Chem. Thermodyn.* 26 (1994) 75-84.

- [122] R. Francesconi, F. Comelli, V. Malta, Isothermal vapor-liquid equilibria, densities, refractive indexes, excess enthalpies, and excess volumes of 1,3-dioxolane or oxolane + isooctane at 298.15 K, *J. Chem. Eng. Data* 38 (1993) 424-427.
- [123] Z. Wang, G.C. Benson, B.C.-Y. Lu, Excess molar enthalpies of the ternary mixtures: (diisopropyl ether or tetrahydrofuran) + 3-methylpentane + n-dodecane at 298.15 K, *Thermochim. Acta* 410 (2004) 57-64.
- [124] Z. Wang, G.C. Benson, B.C.-Y. Lu, Excess enthalpies of the ternary mixtures: {tetrahydrofuran + 3-methylpentane + (octane or decane)} at the temperature 298.15 K, *J. Chem. Thermodyn.* 35 (2003) 1635-1644.
- [125] Z. Wang, D. Kodama, G.C. Benson, B.C.-Y. Lu, Excess enthalpies of the ternary mixtures: tetrahydrofuran + (diisopropyl ether or 2-methyltetrahydrofuran) + n-heptane at 298.15 K, *Fluid Phase Equilibr.* 216 (2004) 293-299.
- [126] B. Giner, S. Martin, H. Artigas, M.C. Lopez, C. Lafuente, Study of weak molecular interactions through thermodynamic mixing properties, *J. Phys. Chem. B* 110 (2006) 17683-17690.
- [127] Z. Wang, G.C. Benson, B.C.-Y. Lu, Excess enthalpies of the ternary mixtures: {tetrahydrofuran + (2,2,4-trimethylpentane or heptane) + methylcyclohexane} at the temperature 298.15 K, *J. Chem. Thermodyn.* 34 (2002) 2073-2082.
- [128] Z. Wang, G.C. Benson, B.C.-Y. Lu, Excess enthalpies of the ternary mixtures: tetrahydrofuran + (hexane or cyclohexane) + decane at 298.15 K, *J. Chem. Eng. Data* 48 (2003) 190-194.

- [129] S.M. Nazmul Hasan, D.-Y. Peng, Excess molar enthalpies of ternary mixtures (1-hexene + tetrahydrofuran + 2,2-dimethylbutane or 2,3-dimethylbutane) at the temperature 298.15 K, *J. Chem. Thermodyn.* 43 (2011) 323-328.
- [130] C.Q. Lan, D. Kodama, Z. Wang, B.C.-Y. Lu, Excess molar enthalpies of the ternary mixtures (1-hexene + tetrahydrofuran or 2-methyltetrahydrofuran + methyl tert-butyl ether) at the temperature 298.15 K, *J. Chem. Thermodyn.* 38 (2006) 1606-1611.
- [131] H. Flores-Segura, L. A. Torres, Thermochemistry of amineborane adduct formation of pyrene and picoline isomers, *Struct. Chem.* 7 (1996) 363-366.
- [132] R. Chadha, A.D. Tripathi, Excess molar enthalpies of 1,1,2,2-tetrachloroethane + 2-methylfuran, + tetrahydrofuran, + 1,4-dioxane, and + cyclopentanone at 308.15 and 318.15 K, *J. Chem. Eng. Data* 40 (1995) 645-646.
- [133] G. Pathak, A.D. Tripathi, U.D. Phalgune, S.D. Pradhan, Enthalpies of mixing of tetrachloroethane (TCE) with furan, methylfuran, tetrahydrofuran, cyclopentanone and 1,4-dioxane, *Thermochim. Acta* 258 (1995) 109-115.
- [134] M.D. Borisover, A.A. Stolov, S.V. Izosimova, F.D. Baitalov, V.A. Breus, B.P. Solomonov, Heats of specific interactions of chloroethylenes in organic solvents, *Zhur. Fiz. Khim.* 65 (1991) 594-599.
- [135] J.N. Spencer, J.E. Mihalick, I.M. Paul, W.J. Nicholson, T.J. Nicholson, X. Ke, Q. He, F.J. Carter, S.E. Daniels, L.J. Fenton, J.L. Ealy, S. Puppala, C.H. Yonder, Solvent effects on host-guest complexation, *J. Solution Chem.* 23(1994) 721-734.
- [136] C.G. Osborne, K.W. Morcom, Thermodynamic behavior of (hexafluorobenzene + a cyclic ether). II. excess enthalpies, *J. Chem. Thermodyn.* 15(1983) 23-27.

- [137] Grolier, Jean Pierre E.; Inglese, Americo; Wilhelm, Emmerich Excess enthalpies of binary systems of cyclic ether + cyclohexene, *Journal of Chemical and Engineering Data* 27(1982) 333-335.
- [138] G. Conti, P. Gianni, L. Lepori, E. Matteoli, M.L. D'Amico, Excess molar enthalpies and excess molar heat capacities of (2-methoxyethanol + tetrahydrofuran + cyclohexane) at the temperature 298.15 K, *J. Chem. Thermodyn.* 30 (1998) 855-868.
- [139] R.M.C. Goncalves, A.M.N. Simoes, Enthalpies of solution of t-butyl chloride and bromide and carbon tetrachloride and tetrabromide in alcohols, *J. Solution Chem.* 16 (1987) 39-43.
- [140] T.M. Letcher, N. Deenadayalu, Excess molar enthalpies and excess molar volumes of binary mixtures of benzene and alkanols with quinoline, *J. Therm. Anal. Calorim.* 60 (2000) 857-860.
- [141] J. Cone, L.E.S. Smith, W. Van Hook, W. Alexander, Enthalpies of solution of gases in liquids. The calorimeter and measurements on lower hydrocarbons and carbon dioxide in benzene, tetrachlorocarbon, and tetrahydrofuran, *J. Chem. Thermodyn.* 11 (1979) 277-285.
- [142] R.L. Benoit, E. Milanova, Vapor pressure and calorimetric data for the solution of sulfur dioxide in aprotic solvents, *Can. J. Chem.* 57 (1979) 1319-1323.
- [143] M. Iwahashi, T. Nozaki, K. Kamaya, K. Taguchi, M. Fujita, Y. Kasahara, H. Minami, H. Matsuzawa, S. Nakamura, K. Harada, Y. Ozaki, T. Inoue, Phase behavior of binary mixtures composed of ethylene carbonate and various organic solvents, *J. Chem. Thermodyn.* 43 (2011) 80-87.

- [144] S. Taniewska-Osinska, L. Kaminska-Bartel, H. Piekarski, T.M. Krygowski, Application of the Hancock steric constants to interpretation of heat of solution of benzamide in aliphatic alcohols, *Can. J. Chem.* 59 (1981) 817-820.
- [145] T. Martins, N. Nunes, M.L. Moita, R.E. Leitao, Solution enthalpies of 1-bromoadamantane in monoalcohols at 298.15K, *Thermochim. Acta* 444 (2006) 83-85.
- [146] F. Martins, L. Moreira, N. Nunes, R.E. Leitao, Solvent effects on solution enthalpies of adamantyl derivatives. A multiparameter approach, *J. Therm. Anal. Calorim.* 100 (2010) 483-491.
- [147] Y. Miyano, T. Kobashi, H. Shinjo, S. Kumada, Y. Watanabe, W. Niya, Y. Tateishi, Henry's law constants and infinite dilution activity coefficients of *cis*-2-butene, dimethyl ether, chloroethane, and 1,1-difluoroethane in methanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, isobutanol, *tert*-butanol, 1-pentanol, 2-pentanol, 3-pentanol, 2-methyl-1-butanol, 3-methyl-1-butanol, and 2-methyl-2-butanol, *J. Chem. Thermodyn.* 38 (2006) 724-731.
- [148] Y. Miyano, Henry's constants and infinite dilution activity coefficients of propane, propene, butane, isobutene, 1-butene, isobutene, *trans*-2-butene and 1,3-butadiene in 1-propanol at T = (260 to 340) K, *J. Chem. Thermodyn.* 36 (2004) 101-106.
- [149] M.H. Abraham, Free energies, enthalpies, and entropies of solution of gaseous nonpolar nonelectrolytes in water and nonaqueous solvents. the hydrophobic effect, *J. Am. Chem. Soc.* 104 (1982) 2085-2094.
- [150] G.L. Pollack, J.F. Himm, J.J. Enyeart, Solubility of xenon in liquid n-alkanols: thermodynamic functions in simple polar liquids, *J. Chem. Phys.* 81 (1984) 3239-3246.

- [151] J.M.M.V. Sousa, A.G.M. Ferreira, H.C. Fachada, I.M.A. Fonseca, Solubility of CF₄ in lower alcohols, *Fluid Phase Equilibr.* 296 (2010) 95-98.
- [152] C.S.O. Silva, I.M.A. Fonseca, L.Q. Lobo, Solubility of methyl fluoride in some alcohols, *Fluid Phase Equilibr.* 135(1997) 137-144.
- [153] F. Gibanel, M.C. Lopez, F.M. Royo, J. Santafe, J.S. Urieta, Solubility of nonpolar gases in tetrahydrofuran at 0 to 30°C and 101.33 kPa partial pressure of gas, *J. Solution Chem.* 22 (1993) 211-217.
- [154] J. Tokunaga, Solubilities of oxygen, nitrogen and carbon dioxide in aqueous alcohol solutions, *J. Chem. Eng. Data* 20 (1975) 41-46.
- [155] J.M.M.V. Sousa, J.P.B. Almeida, A.G.M. Ferreira, H. C. Fachada and I.M.A. Fonseca, Solubility of HFCs in lower alcohols, *Fluid Phase Equilibr.* in press (2011).
- [156] A. Jouyban, J.L. Manzoori, J. Soleymani, V. Panahi-Azar, M.A.A. Fakhree, Mohammad S. Ahmadian, A. Shayanfar, W.E. Acree, Jr., Solubility of anthracene in C₁-C₃ alcohols from (298.2 to 333.2) K and their mixtures with 2-propanone at 298.2 K, *J. Chem. Eng. Data* 55 (2010), 5319-5322.
- [157] Z. Wang, P. Jiang, Solubility of Acenaphthene in Different Solvents from (283.00 to 323.00) K, *J. Chem. Eng. Data* 54 (2009) 150-151.
- [158] A.N. Gaivoronskii, V.A. Granzhan, Solubility of adipic acid in organic solvents and water, *Russ. J. Appl. Chem.* 78 (2005) 404-408.
- [159] Q. Yu, S. Black, H. Wei, Solubility of butanedioic acid in different solvents at temperatures between 283 K and 333 K, *J. Chem. Eng. Data* 54 (2009) 2123-2125.
- [160] S.-N. Chen, Q. Xia, L.-F. Lu, D. Li, F.-B. Zhang, G.-L. Zhang, Solubility of sebacic acid in binary mixtures containing alcohols, *Chem. Eng. Data* 54 (2009) 776-780.

- [161] J.S. Chickos, W.E. Acree, Jr., Enthalpies of vaporization of organic and organometallic compounds, 1880-2002, *J. Phys. Chem. Ref. Data* 32 (2003) 519-878.
- [162] W.E. Acree, Jr.; J.S. Chickos, Phase transition enthalpy measurements of organic and organometallic compounds. sublimation, vaporization and fusion enthalpies from 1880 to 2010, *J. Phys. Chem. Ref. Data* 39 (2010) 043101/1-043101/942.
- [163] J.S. Chickos, W.E. Acree, Jr., Enthalpies of sublimation of organic and organometallic compounds. 1910-2001, *J. Phys. Chem. Ref. Data* 31 (2002), 537-698.
- [164] M.H. Abraham, J.C. McGowan, The use of characteristic volumes to measure cavity terms in reversed phase liquid chromatography, *Chromatographia* 23 (1987) 243-246.
- [165] M. H. Abraham, Scales of solute hydrogen-bonding: their construction and application to physicochemical and biochemical processes, *Chem. Soc. Rev.* 22 (1993) 73-83.
- [166] M.H. Abraham, H.S. Chadha, G.S. Whiting, R.C. Mitchell, Hydrogen bonding. 32. An analysis of water-octanol and water-alkane partitioning and the $\Delta \log P$ parameter of Seiler, *J. Pharm. Sci.* 83 (1994) 1085-1100.
- [167] M.H. Abraham, G.S. Whiting, R.M. Doherty, W.J. Shuely, W.J. Hydrogen bonding. Part 14. The characterization of some N-substituted amides as solvents: Comparison with gas-liquid chromatographic stationary phases, *J. Chem. Soc., Perkin Trans. 2*, (1990) 1851-1857.
- [168] A.M. Zissimos, M.H. Abraham, C.M. Du, K. Valko, C. Bevan, D. Reynolds, J. Wood, K.Y. Tam, Calculation of Abraham descriptors from experimental data from seven HPLC systems: Evaluation of five different methods of calculation. *J. Chem. Soc., Perkin Trans. 2* (2002) 2001-2010.

- [169] C.E. Green, M.H. Abraham, W.E. Acree, Jr., K.M. De Fina, T.L. Sharp, Solvation descriptors for pesticides from the solubility of solids: diuron as an example *Pest Manag. Sci.* 56 (2000) 1043-1053.
- [170] M.H. Abraham, A. Ibrahim, A.M. Zissimos, Determination of sets of solute descriptors from chromatographic measurements. *J. Chromatogr. A* 1037 (2004) 29-47.
- [171] A.M. Zissimos, M.H. Abraham, M.C. Barker, K.J. Box, K.Y. Tam, K.Y. Calculation of the Abraham descriptors for solvent-water partition coefficients in four different systems: Evaluation of different methods of calculation. *J. Chem. Soc., Perkin Trans. 2* (2002) 470-477.
- [172] C.F. Poole, S.N. Atapaltu, S. K. Poole, A. K. Bell, Determination of solute descriptors by chromatographic methods, *Anal. Chem. Acta* 652 (2009) 32-53.
- [173] ADME Boxes, version*, Advanced Chemistry Development, 110 Yonge Street, 14th Floor, Toronto, Ontario, M5C 1T4, Canada.
- [174] A.M. Zissimos, M.H. Abraham, A. Klamt, F. Eckert, J.A. Wood, J.A. A comparison between two general sets of linear free energy descriptors of Abraham and Klamt. *J. Chem. Inf. Comput. Sci.* 42 (2002) 1320-1331.
- [175] J. Jover, R. Bosque, J. Sales, Determination of the Abraham solute descriptors from molecular structure. *J. Chem. Inf. Comput. Sci.* 44 (2004) 1098-1106.
- [176] P.C.M. van Noort, J.J.H. Haftka, J.R. Parsons, Updated Abraham solvation parameters for polychlorinated biphenyls. *Environ. Sci. Technol.* 44 (2010) 7037-7042.

Table 1. Values of the gas-to-1-propanol solvation enthalpy in kJ/mole at 298 K, $\Delta H_{\text{solv,PrOH}}$, for 103 solutes, together with the solute descriptors

Solute	E	S	A	B	L	V	ΔH_{solv}	Ref
Methane	0.000	0.000	0.000	0.000	-0.323	0.2495	-2.09	149
Ethane	0.000	0.000	0.000	0.000	0.492	0.3904	-10.33	149
Propane	0.000	0.000	0.000	0.000	1.050	0.5313	-18.60	148
2-Methylpropane	0.000	0.000	0.000	0.000	1.409	0.6722	-20.79	148
Butane	0.000	0.000	0.000	0.000	1.615	0.6722	-22.10	148
Pentane	0.000	0.000	0.000	0.000	2.162	0.8131	-25.39	76
Heptane	0.000	0.000	0.000	0.000	3.173	1.0949	-33.76	149
Octane	0.000	0.000	0.000	0.000	3.677	1.2358	-38.45	70,86
Nonane	0.000	0.000	0.000	0.000	4.182	1.3767	-43.75	49
Decane	0.000	0.000	0.000	0.000	4.686	1.5176	-47.35	47, 50
Dodecane	0.000	0.000	0.000	0.000	5.696	1.7994	-58.09	47
2,3-Dimethylbutane	0.000	0.000	0.000	0.000	2.495	0.9540	-27.86	46
2,2,4-Trimethylpentane	0.000	0.000	0.000	0.000	3.106	1.2358	-33.37	46
Cyclohexane	0.305	0.100	0.000	0.000	2.964	0.8454	-31.14	53
Cyclooctane	0.413	0.100	0.000	0.000	4.329	1.1272	-40.98	77
1-Propene	0.100	0.080	0.000	0.070	0.946	0.4883	-18.75	148
1-Butene	0.100	0.080	0.000	0.070	1.491	0.6292	-19.78	148
<i>cis</i> 2-Butene	0.140	0.080	0.000	0.050	1.737	0.6292	-21.55	147
<i>trans</i> -2-Butene	0.126	0.080	0.000	0.050	1.664	0.6292	-21.39	148
2-Methyl-1-propene	0.120	0.080	0.000	0.080	1.579	0.6292	-21.67	148
1-Hexene	0.078	0.080	0.000	0.070	2.572	0.9110	-29.10	42
1-Heptene	0.092	0.080	0.000	0.070	3.063	1.0519	-34.00	42
1,3-Butadiene	0.320	0.230	0.000	0.100	1.543	0.5862	-20.93	148
Methanol	0.278	0.440	0.430	0.470	0.970	0.3082	-42.20	97
Ethanol	0.246	0.420	0.370	0.480	1.485	0.4491	-42.20	71
Propan-1-ol	0.236	0.420	0.370	0.480	2.031	0.5900	-47.50	ΔH_{vap}
Butan-1-ol	0.224	0.420	0.370	0.480	2.601	0.7309	-52.19	71
Pentan-1-ol	0.219	0.420	0.370	0.480	3.106	0.8718	-56.81	71
Hexan-1-ol	0.210	0.420	0.370	0.480	3.610	1.0127	-60.48	61
Octan-1-ol	0.199	0.420	0.370	0.480	4.619	1.2950	-70.07	59, 71
Decan-1-ol	0.191	0.420	0.370	0.480	5.628	1.5763	-80.00	60
2-Propanol	0.212	0.360	0.330	0.560	1.764	0.5900	-45.52	95
Benzene	0.610	0.520	0.000	0.140	2.786	0.7176	-31.77	81
Toluene	0.601	0.520	0.000	0.140	3.325	0.8573	-35.98	57
1,4-Dimethylbenzene	0.613	0.520	0.000	0.160	3.839	0.9982	-40.50	85

Difluoromethane	-0.320	0.490	0.060	0.050	0.040	0.2489	-9.41	155
Trifluoromethane	-0.430	0.180	0.110	0.030	-0.274	0.3026	-12.50	155
Carbon tetrachloride	0.458	0.380	0.000	0.000	2.823	0.7391	-32.88	66,67,139
Chloroethane	0.227	0.400	0.000	0.100	1.678	0.5128	-23.50	147
2-Chloro-2-methylpropane	0.142	0.300	0.000	0.030	2.273	0.7946	-27.32	56
2-Chloro-2-methylbutane	0.171	0.270	0.000	0.150	2.858	0.9355	-32.93	56
2-Chloro-2-methylpentane	0.207	0.390	0.000	0.130	3.520	1.0764	-37.62	56
Trichloroethene	0.520	0.370	0.080	0.030	2.997	0.7146	-33.94	72
2-Methyl-2-bromopropane	0.305	0.290	0.000	0.070	2.609	0.8472	-29.81	73,88
2-Methyl-2-bromobutane	0.343	0.400	0.000	0.150	3.400	0.9881	-35.55	88
2-Methyl-2-iodopropane	0.589	0.350	0.000	0.190	3.439	0.9304	-33.37	73
1-Bromoadamantane	1.070	0.900	0.000	0.200	6.130	1.3668	-56.01	145
Acetone	0.179	0.700	0.040	0.490	1.696	0.5470	-25.00	82
2-Butanone	0.166	0.700	0.000	0.510	2.287	0.6879	-28.93	78
Dimethyl ether	0.000	0.270	0.000	0.410	1.285	0.4491	-18.30	147
Diethyl ether	0.041	0.250	0.000	0.450	2.015	0.7309	-26.10	66
Diisopropyl ether	-0.060	0.160	0.000	0.580	2.530	1.0127	-32.46	43,87
Dibutyl ether	0.000	0.250	0.000	0.450	3.924	1.2950	-43.13	62
Butyl methyl ether	0.045	0.250	0.000	0.440	2.658	0.8718	-31.09	63
Methyl <i>tert</i> -butyl ether	0.024	0.210	0.000	0.590	2.380	0.8718	-28.64	43,80,84
Methyl <i>tert</i> -amyl ether	0.050	0.210	0.000	0.600	2.916	1.0127	-34.40	43
Tetrahydrofuran	0.289	0.520	0.000	0.480	2.636	0.6223	-28.97	87
Tetrahydropyran	0.275	0.470	0.000	0.550	3.057	0.7672	-34.44	74,84
1,4-Dioxane	0.329	0.750	0.000	0.640	2.892	0.6810	-30.79	44, 54
15-Crown-5	0.410	1.200	0.000	1.750	6.779	1.7025	-70.22	90
Aniline	0.955	0.960	0.260	0.410	3.934	0.8162	-53.89	79
Argon	0.000	0.000	0.000	0.000	-0.688	0.1900	-1.67	149
Xenon	0.000	0.000	0.000	0.000	0.378	0.3290	-9.62	150
Nitrogen	0.000	0.000	0.000	0.000	-0.978	0.2222	1.17	149
Carbon monoxide	0.000	0.000	0.000	0.040	-0.836	0.2220	-1.72	149
Carbon dioxide	0.000	0.280	0.050	0.100	0.058	0.2809	-9.75	154
Oxygen	0.000	0.000	0.000	0.000	-0.723	0.1830	0.50	149
Ethyl acetate	0.106	0.620	0.000	0.450	2.314	0.7466	-29.88	54
Propyl acetate	0.092	0.600	0.000	0.450	2.819	0.8875	-33.81	89
Butyronitrile	0.180	0.900	0.000	0.360	2.548	0.6860	-31.80	83
Triethylamine	0.101	0.150	0.000	0.790	3.040	1.0538	-44.40	48, 51
Chlorobenzene	0.718	0.650	0.000	0.070	3.657	0.8388	-39.49	64
Nitromethane	0.313	0.950	0.060	0.310	1.892	0.4237	-29.67	44, 68
4-Methylpyridine	0.630	0.820	0.000	0.540	3.640	0.8162	-46.80	52
1,3-Dichlorobenzene	0.847	0.730	0.000	0.020	4.410	0.9612	-45.66	75

1,3-Dimethoxybenzene	0.816	1.010	0.000	0.450	5.022	1.1160	-52.05	66
Ethylbenzene	0.613	0.510	0.000	0.150	3.778	0.9982	-40.07	58
1,2-Dimethoxyethane	0.116	0.670	0.000	0.680	2.654	0.7896	-33.60	54
1,2-Diethoxyethane	0.008	0.730	0.000	0.790	3.310	1.0714	-40.00	66
Tetramethylsilicon	-0.057	0.080	0.000	0.000	1.812	0.9179	-23.00	55
Tetramethyltin	0.324	0.110	0.000	0.100	2.651	1.0431	-30.46	149
Benzo-15-crown-5	1.055	1.940	0.000	1.590	9.403	2.0285	-82.42	90
Tributylamine	0.051	0.150	0.000	0.790	5.983	1.8992	-66.20	51
Carbon tetrafluoride	-0.580	-0.260	0.000	0.000	-0.817	0.3203	-1.12	151
Succinic acid	0.370	1.320	1.030	0.710	3.951	0.8210	-95.70	159
Adipic acid	0.350	1.210	1.130	0.760	4.474	1.1028	-105.90	158
Sebacic acid	0.350	1.360	1.120	0.870	6.920	1.6664	-118.05	160
1,3-Diaminopropane	0.446	0.610	0.430	1.140	2.852	0.7309	-69.50	91
3,6,9-Trioxoundecane	0.040	0.870	0.000	1.200	4.815	1.4111	-51.40	45
Digylme	0.113	0.760	0.000	1.170	3.920	1.1301	-42.63	65
2,2,2-Trifluoroethanol	0.015	0.600	0.570	0.250	1.224	0.5022	-48.33	94
N-Methylformamide	0.405	1.360	0.400	0.550	2.863	0.5059	-51.21	96
Anthracene	2.290	1.340	0.000	0.280	7.568	1.4544	-72.34	156
Fluoromethane	0.070	0.350	0.000	0.090	0.057	0.2672	-8.12	152
Trimethylphosphate	0.113	1.360	0.000	0.930	3.850	0.9707	-40.50	93
Tributyl phosphate	-0.100	0.110	0.000	1.260	7.539	2.2390	-79.71	92
Ethylene carbonate	0.381	1.250	0.000	0.580	2.670	0.5558	-31.70	143
Benzaldehyde	0.820	1.000	0.000	0.390	4.008	0.8730	-44.16	69
Acenaphthene	1.604	1.050	0.000	0.220	6.469	1.2586	-57.97	157
1-Adamantanol	0.940	0.900	0.310	0.660	5.634	1.2505	-74.59	146
Quinoline	1.268	0.970	0.000	0.540	5.457	1.0443	-58.65	140
Benzamide	0.990	1.500	0.490	0.670	5.767	0.9728	-81.53	144
Benzoic acid	0.730	0.900	0.590	0.400	4.510	0.9317	--74.77	144

Table 2. Values of the gas-to-tetrahydrofuran solvation enthalpy in kJ/mole at 298 K, $\Delta H_{\text{solv,THF}}$, for 86 solutes, together with the solute descriptors

Solute	E	S	A	B	L	V	ΔH_{solv}	Ref
Methane	0.000	0.000	0.000	0.000	-0.323	0.2495	-3.02	153
Ethane	0.000	0.000	0.000	0.000	0.492	0.3904	-13.74	153
Propane	0.000	0.000	0.000	0.000	1.050	0.5313	-16.69	153
Hexane	0.000	0.000	0.000	0.000	2.668	0.9540	-28.39	128
Heptane	0.000	0.000	0.000	0.000	3.173	1.0949	-33.38	127
Octane	0.000	0.000	0.000	0.000	3.677	1.2358	-37.55	120,124
Nonane	0.000	0.000	0.000	0.000	4.182	1.3767	-42.14	120
Decane	0.000	0.000	0.000	0.000	4.686	1.5176	-46.13	101
Dodecane	0.000	0.000	0.000	0.000	5.696	1.7994	-56.53	123
Tetradecane	0.000	0.000	0.000	0.000	6.705	2.0812	-64.36	101
3-Methylpentane	0.000	0.000	0.000	0.000	2.581	0.9540	-27.28	124
2,2-Dimethylbutane	0.000	0.000	0.000	0.000	2.352	0.9540	-24.83	129
2,3-Dimethylbutane	0.000	0.000	0.000	0.000	2.495	0.9540	-26.21	129
Cyclohexane	0.305	0.100	0.000	0.000	2.964	0.8454	-29.39	112,113
Methylcyclohexane	0.244	0.060	0.000	0.000	3.319	0.9863	-32.87	112,127
2,2,4-Trimethylpentane	0.000	0.000	0.000	0.000	3.106	1.2358	-32.90	122,127
Ethene	0.107	0.100	0.000	0.070	0.289	0.3474	-11.42	153
1-Hexene	0.078	0.080	0.000	0.070	2.572	0.9110	-29.18	130
Cyclohexene	0.395	0.200	0.000	0.100	3.021	0.8025	-32.33	137
Dichloromethane	0.387	0.570	0.100	0.050	2.019	0.4943	-34.17	104
Trichloromethane	0.425	0.490	0.150	0.020	2.480	0.6167	-39.62	99, 103
Tetrachloromethane	0.458	0.380	0.000	0.000	2.823	0.7391	-35.72	105
1-Chlorobutane	0.210	0.400	0.000	0.100	2.722	0.7946	-34.46	117
1,1,2,2-Tetrachloroethane	0.595	0.760	0.160	0.120	3.803	0.8800	-57.03	132,133
Chlorocyclohexane	0.448	0.480	0.000	0.100	3.832	0.9678	-43.10	112
Trichloroethene	0.524	0.370	0.080	0.030	2.997	0.7146	-39.69	134
Tetrachloroethene	0.639	0.440	0.000	0.000	3.584	0.8370	-40.66	134
Dipropyl ether	0.008	0.250	0.000	0.450	2.954	1.0127	-34.60	118
Diisopropyl ether	-0.060	0.160	0.000	0.580	2.530	1.0127	-31.40	125
Dibutyl ether	0.000	0.250	0.000	0.450	3.924	1.2945	-44.14	118
Methyl tert-butyl ether	0.024	0.110	0.000	0.630	2.380	0.8718	-29.97	119
Tetrahydrofuran	0.289	0.520	0.000	0.480	2.636	0.6223	-32.00	ΔH_{vap}
Methanol	0.278	0.440	0.430	0.470	0.970	0.3082	-33.90	98
Ethanol	0.246	0.420	0.370	0.480	1.485	0.4491	-37.99	98
1-Butanol	0.224	0.420	0.370	0.480	2.601	0.7309	-47.86	117

2-Propanol	0.212	0.360	0.330	0.560	1.764	0.5900	-38.23	74
1-Hexanol	0.210	0.420	0.370	0.480	3.610	1.0170	-56.98	87
1-Heptanol	0.211	0.420	0.370	0.480	4.115	1.1536	-61.76	87
1-Octanol	0.199	0.420	0.370	0.480	4.619	1.2950	-65.59	87
1-Nonanol	0.193	0.420	0.370	0.480	5.120	1.4354	-71.16	87
1-Decanol	0.191	0.420	0.370	0.480	5.628	1.5763	-75.31	87
2-Methoxyethanol	0.269	0.500	0.300	0.840	2.490	0.6487	-42.80	138
Acetonitrile	0.237	0.900	0.070	0.320	1.739	0.4042	-33.70	121
Butyronitrile	0.180	0.900	0.000	0.360	2.548	0.6860	-39.66	116
Benzene	0.610	0.520	0.000	0.140	2.786	0.7176	-35.24	102
Toluene	0.601	0.520	0.000	0.140	3.325	0.8573	-38.77	102
1,4-Dimethylbenzene	0.613	0.520	0.000	0.160	3.839	0.9982	-43.23	114
1,2,4-Trimethylbenzene	0.677	0.560	0.000	0.190	4.441	1.1391	-48.70	107
Anisole	0.710	0.750	0.000	0.290	3.890	0.9160	-48.93	117
Chlorobenzene	0.718	0.650	0.000	0.070	3.657	0.8388	-43.77	109,100
Fluorobenzene	0.477	0.570	0.000	0.100	2.788	0.7341	-37.60	126
Nitrobenzene	0.871	1.110	0.000	0.280	4.557	0.8906	-57.72	66
Benzyl chloride	0.821	0.860	0.000	0.140	4.353	0.9797	-52.75	100
Pyrrrole	0.613	0.730	0.410	0.290	2.865	0.5770	-53.26	102
N-Methylpyrrrole	0.559	0.790	0.000	0.310	2.923	0.7180	-41.56	102,106
Aniline	0.955	0.960	0.260	0.410	3.934	0.8162	-62.86	66
Helium	0.000	0.000	0.000	0.000	-1.741	0.0680	8.12	153
Neon	0.000	0.000	0.000	0.000	-1.575	0.0850	7.01	153
Argon	0.000	0.000	0.000	0.000	-0.688	0.1900	-0.04	153
Krypton	0.000	0.000	0.000	0.000	-0.211	0.2460	-3.39	153
Xenon	0.000	0.000	0.000	0.000	0.378	0.3290	-8.85	153
Hydrogen	0.000	0.000	0.000	0.000	-1.200	0.1086	4.48	153
Deuterium	0.000	0.000	0.000	0.000	-1.200	0.1100	6.01	153
Nitrogen	0.000	0.000	0.000	0.000	-0.978	0.2222	1.32	153
Carbon dioxide	0.000	0.280	0.050	0.100	0.058	0.2809	-17.02	141
Carbon tetrafluoride	-0.580	-0.260	0.000	0.000	-0.817	0.3203	-0.02	153
Sulfur hexafluoride	-0.600	-0.200	0.000	0.000	-0.120	0.4643	4.83	153
Pyridine	0.631	0.840	0.000	0.520	3.022	0.6753	-39.60	131
2-Methylpyridine	0.598	0.750	0.000	0.580	3.422	0.8162	-42.00	131
3-Methylpyridine	0.631	0.810	0.000	0.540	3.631	0.8162	-44.00	131
4-Methylpyridine	0.630	0.820	0.000	0.540	3.640	0.8162	-44.30	131
Triethylamine	0.101	0.150	0.000	0.790	3.040	1.0538	-33.30	48
Hexafluorobenzene	0.088	0.560	0.000	0.010	2.345	0.8226	-35.50	136
Sulfur dioxide	0.370	0.660	0.280	0.100	0.778	0.3465	-39.30	142
15-Crown-5	0.410	1.200	0.000	1.750	6.779	1.7025	-78.89	135

Phenol	0.805	0.890	0.600	0.300	3.766	0.7751	-71.00	117
4-Fluorophenol	0.670	0.970	0.630	0.230	3.844	0.7928	-74.61	117
1-Chloronaphthalene	1.417	1.000	0.000	0.140	5.856	1.2078	-68.35	110
1-Methylnaphthalene	1.337	0.940	0.000	0.220	5.802	1.2260	-60.24	111
Dimethyl carbonate	0.142	0.540	0.000	0.570	2.328	0.6644	-36.90	108
Diethyl carbonate	0.060	0.580	0.000	0.530	3.412	0.9462	-42.28	108
Diethylamine	0.154	0.300	0.080	0.690	2.395	0.7720	-33.27	115
Dipropylamine	0.124	0.300	0.080	0.690	3.351	1.0538	-40.13	115
Hexanoic acid	0.174	0.630	0.620	0.440	3.697	1.0284	-71.78	66
Octanoic acid	0.150	0.640	0.620	0.450	4.680	1.3102	-82.88	66
4-Fluoroanisole	0.571	0.740	0.000	0.280	3.904	0.9337	-51.84	117

Figure 1




Figure 1. A plot of the calculated values of $\Delta H_{\text{Sol},\text{PrOH}}$ based on eqn 9 against the observed values.

Figure 2.




Figure 2. A plot of the calculated values of $\Delta H_{\text{Soln,THF}}$ based on eqn 12 against the observed values.