

SUPPORTING INFORMATION FOR
ABRAHAM MODEL CORRELATIONS FOR SOLUTE PARTITIONING INTO *o*-XYLENE,
m-XYLENE AND *p*-XYLENE FROM BOTH WATER AND THE GAS PHASE

Timothy W. Stephens^a, Nohelli E. De La Rosa^a, Mariam Saifullah^a, Shulin Ye^a, Vicky Chou^a,
Amanda N. Quay^a, William E. Acree, Jr.^{a*} and Michael H. Abraham^b

^a Department of Chemistry, 1155 Union Circle # 305070, University of North Texas,
Denton, TX 76203-5017 (U.S.A.)

^b Department of Chemistry, University College London, 20 Gordon Street,
London, WC1H 0AJ (U.K.)

Figure S1




Figure S1. Comparison of observed $\log P$ data for solutes dissolved in *m*-xylene and predicted values based on Eqn. 10.

FIGURE S2




Figure S2. Comparison of observed $\log K$ data for solutes dissolved in *m*-xylene and predicted values based on Eqn. 11.

FIGURE S3




Figure S3. Comparison of observed $\log P$ data for solutes dissolved in *p*-xylene and predicted values based on Eqn. 14.

FIGURE S4




Figure S4. Comparison of observed $\log P$ data for solutes dissolved in *p*-xylene and predicted values based on Eqn. 15.