

2F/1254

PNNL-11286
UC-000

Approved for public release; distribution is unlimited

Evaluation of Dredged Material Proposed for Ocean Disposal from Port Chester, New York

E. S. Barrows
H. L. Mayhew
J. Q. Word
J. J. S. Tokos

Battelle Marine Sciences Laboratory
Sequim, Washington

August 1996

Prepared for the
U.S. Army Corps of Engineers - New York District
under a Related Services Agreement
with the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest National Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute



DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401.

Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161



3 3679 00047 3910

**EVALUATION OF DREDGED MATERIAL
PROPOSED FOR OCEAN DISPOSAL FROM
PORT CHESTER, NEW YORK**

**E. S. Barrows
H. L. Mayhew
J. Q. Word
J. J. S. Tokos**

**Battelle Marine Sciences Laboratory
Sequim, Washington 98382**

August 1996

**Prepared for the
U.S. Army Corps of Engineers - New York District
under a Related Services Agreement
with the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830**

**Pacific Northwest National Laboratory
Richland, Washington 99352**

EVALUATION OF PROPOSED MATERIAL
PROCESSED FOR OCEAN DISPOSAL FROM
PORT CHESTER, NEW YORK

E. S. Barows
H. L. Mayhew
J. O. Ford
J. B. Tokos

British Marine Sciences Laboratory
Seaford, West Sussex BN27 3RQ

August 1988

Prepared for the
U.S. Army Corps of Engineers - New York District
under a Related Services Agreement
with the U.S. Department of Energy
under Contract DE-AC05-81OR21400

Pacific Northwest Regional Laboratory
Richland, Washington 98882

Summary

Port Chester (FP No. 1) was one of seven waterways that the U.S. Army Corps of Engineers-New York District (USACE-NYD) requested the Battelle Marine Sciences Laboratory (MSL) to sample and evaluate for dredging and disposal in March 1994. Sediment samples were collected from Port Chester, as well as from the Hudson River, Gravesend Bay Anchorage, South Brother Island, Buttermilk, Eastchester, and Brown's Creek, during a survey conducted from March 7 through 14, 1994. Combining sample collection and concurrent evaluation of multiple dredged material projects was more cost-effective for the USACE-NYD, because the expense of reference site testing and quality control analyses could be shared among project budgets.

Tests and analyses were conducted on Port Chester sediment core samples according to the manual developed by the USACE and the U.S. Environmental Protection Agency (EPA), *Evaluation of Dredged Material Proposed for Ocean Disposal (Testing Manual)*, commonly referred to as the "Green Book," and the regional manual developed by the USACE-NYD and EPA Region II, *Guidance for Performing Tests on Dredged Material to be Disposed of in Ocean Waters*. Because the Port Chester area is located on the border between New York and southeast Connecticut, its dredged material may also be considered for disposal at the Central Long Island Sound Disposal Site (CLIS). Therefore, Port Chester sediment was also tested for possible disposal at the CLIS according to the USACE-New England Division (NED) guidelines, *Guidance for Performing Tests on Dredged Material to be Disposed of in Open Waters (Draft)*.

The evaluation of proposed dredged material from Port Chester consisted of bulk sediment chemical analyses, chemical analyses of site water and dredged material elutriate preparations, water-column and benthic acute toxicity tests, and bioaccumulation studies. Individual sediment core samples collected from Port Chester were analyzed for grain size, moisture content, and total organic carbon (TOC). Additionally, a composite sediment sample, representing the entire area proposed for dredging, was analyzed for bulk density, specific gravity, metals, chlorinated pesticides, polychlorinated biphenyl (PCB) congeners, polynuclear aromatic hydrocarbons (PAHs), and 1,4-dichlorobenzene. Additional testing requirements specified by the USACE-NED consisted of an extended list of sediment chemical parameters of additional metals, chlorinated pesticides, and PAHs. Site water and elutriate water, prepared from the suspended-particulate phase (SPP) of Port Chester sediment, were analyzed for metals, pesticides, and PCBs. Water-column or SPP toxicity tests were performed with three species, the mysid *Mysidopsis bahia*, the juvenile silverside *Menidia beryllina*, and larvae of the mussel *Mytilus galloprovincialis*. Benthic acute toxicity tests were performed with two amphipods, *Ampelisca abdita* and *Rhepoxynius abronius*, as well as with the mysid *M. bahia*. The amphipod benthic toxicity test procedures followed EPA guidance for reduction of total ammonia concentrations in test systems prior to test

initiation. A similar procedure was followed for the mysid toxicity test. Bioaccumulation tests were conducted with the burrowing, polychaete worm *Nereis virens* and the surface-feeding, bent-nose clam, *Macoma nasuta*.

Port Chester sediment core samples were either a black, silt and clay material or a black, sand and gravel mixture. The Port Chester sediment composite sample contained elevated levels of metals, pesticides (particularly the DDD/DDE/DDT group of compounds), PCBs, PAHs, and 1,4-dichlorobenzene.

In water-column toxicity tests, 100% SPP treatments were acutely toxic to all three species tested. The LC₅₀ values ranged from 75.1% SPP for *M. beryllina* to >100% SPP for *M. bahia* and *M. galloprovincialis* survival. The EC₅₀ value for *M. galloprovincialis* normal development, a more sensitive measure than survival, was 53.7% SPP. In the static renewal test with *A. abdita* exposed to Port Chester sediment, test organism survival was statistically significant and >20% lower than the Mud Dump Reference and the CLIS Reference. In the static renewal test with *R. abronius*, survival was statistically significantly different when compared with both reference site sediments, but there was not a >20% reduction in survival. In the *M. bahia* static test, that was not manipulated to reduce porewater or overlying ammonia concentrations prior to test initiation, test organisms survival was 77% in the Port Chester sediment treatment, which was not statistically significant nor was there a 10% or greater difference in survival between the test and either of the two reference sediments.

Concentrations of some contaminants were elevated in tissues of *N. virens* and *M. nasuta* that were exposed to Port Chester sediment in 28-day bioaccumulation tests. Concentrations of metals, pesticides, PCBs, and PAHs were generally the same or slightly higher in *M. nasuta* than in *N. virens*. Tissues of both species exposed to Port Chester sediment had tissue body burdens that were lower than the U.S. Food and Drug Administration (FDA) action levels for poisonous or deleterious substances in fish and shellfish for human consumption for selected pesticides, and FDA levels of concern for chronic shellfish consumption for selected metals. Tissue burdens of organisms exposed to Port Chester sediment compared with those exposed to Mud Dump Reference Site and the Central Long Island Sound Reference Site sediment were significantly higher for selected metals, pesticides, PCBs, and PAHs. Therefore, Port Chester sediment requires further evaluation to determine limiting permissible concentration (LPC) and benthic effects compliance.

Contents

Summary	iii
1.0 Introduction	1.1
1.1 Project Objectives	1.1
1.2 Project Background	1.2
1.3 Organization of This Report	1.2
2.0 Materials and Methods	2.1
2.1 Sediment and Water Collection	2.1
2.1.1 Test Sediment and Site Water Sampling	2.1
2.1.2 Reference and Control Sediment Sampling	2.3
2.2 Test Organism Collection	2.3
2.3 Sediment Sample Preparation	2.4
2.3.1 Laboratory Preparation and Safety Considerations	2.5
2.3.2 Preparation of Sediment for Benthic Testing and Bulk Sediment Analyses	2.6
2.3.3 Preparation of Suspended-Particulate Phase and Elutriate	2.6
2.4 Physical and Chemical Analytical Procedures	2.7
2.4.1 Grain Size and Percentage of Moisture	2.8
2.4.2 Bulk Density and Specific Gravity	2.8
2.4.3 TOC	2.8
2.4.4 Metals	2.12
2.4.5 Chlorinated Pesticides and PCBs	2.13
2.4.6 PAHs and 1,4-Dichlorobenzene	2.14
2.4.7 Lipids	2.14
2.5 Biological Testing Procedures	2.15
2.5.1 Water-Column Toxicity Tests	2.15
2.5.1.1 Water-Column Toxicity Test with <i>Menidia beryllina</i>	2.15
2.5.1.2 Water-Column Toxicity Test with <i>Mysidopsis bahia</i>	2.16
2.5.1.3 Water-Column Toxicity Test with <i>Mytilus galloprovincialis</i> Larvae	2.17
2.5.2 Benthic Acute Toxicity Tests	2.19
2.5.2.1 Static Renewal Tests with <i>Ampelisca abdita</i> and <i>Rhepoxynius</i> <i>abronius</i>	2.19
2.5.2.2 Static Test With <i>Mysidopsis bahia</i>	2.21

Contents (contd)

2.5.3 Bioaccumulation Testing.....	2.23
2.6 Data Analysis and Interpretation Procedures	2.24
2.6.1 Randomization.....	2.24
2.6.2 Statistical Analysis of Water-Column Tests	2.24
2.6.3 Statistical Analysis of Benthic Toxicity Tests	2.25
2.6.4 Statistical Analysis of Bioaccumulation	2.25
2.7 Quality Assurance/Quality Control Procedures	2.26
3.0 Results	3.1
3.1 Sample Collection and Processing	3.1
3.2 Physical and Chemical Analyses	3.2
3.2.1 Sediment Core Sample Description	3.2
3.2.2 Grain Size, Percentage of Moisture, Bulk Density, and Specific Gravity	3.3
3.2.3 Total Organic Carbon	3.3
3.2.4 Metals	3.5
3.2.5 Chlorinated Pesticides	3.7
3.2.6 PCBs.....	3.7
3.2.7 PAHs and 1,4-Dichlorobenzene	3.8
3.3 Site Water and Elutriate Analyses.....	3.16
3.3.1 Metals	3.16
3.3.2 Chlorinated Pesticides and PCBs.....	3.16
3.4 Water-Column Toxicity Testing.....	3.17
3.4.1 <i>Menidia beryllina</i> Water-Column Toxicity Test	3.19
3.4.2 <i>Myxidopsis bahia</i> Water-Column Toxicity Test	3.19
3.4.3 <i>Mytilus galloprovincialis</i> Water-Column Toxicity Test	3.20
3.5 Benthic Acute Toxicity Testing	3.20
3.5.1 <i>Ampelisca abdita</i> Benthic Acute Toxicity Test	3.20
3.5.2 <i>Rhepoxynius abronius</i> Benthic Acute Toxicity Test.....	3.21
3.5.3 <i>Myxidopsis bahia</i> Benthic Acute Toxicity Test.....	3.22
3.6 Bioaccumulation Tests with <i>Macoma nasuta</i> and <i>Nereis virens</i>	3.23
3.6.1 Bioaccumulation of Metals in <i>Macoma nasuta</i>	3.23
3.6.2 Bioaccumulation of Pesticides in <i>Macoma nasuta</i>	3.24
3.6.3 Bioaccumulation of PCBs in <i>Macoma nasuta</i>	3.24

Contents (contd)

3.6.4	Bioaccumulation of PAHs and 1,4-Dichlorobenzene in <i>Macoma nasuta</i>	3.26
3.6.5	Bioaccumulation of Metals in <i>Nereis virens</i>	3.27
3.6.6	Bioaccumulation of Pesticides in <i>Nereis virens</i>	3.27
3.6.7	Bioaccumulation of PCBs in <i>Nereis virens</i>	3.28
3.6.8	Bioaccumulation of PAHs and 1,4-Dichlorobenzene in <i>Nereis virens</i>	3.28
3.6.9	Magnification Factors of Compounds in <i>Macoma nasuta</i> and <i>Nereis virens</i> Tissues	3.28
4.0	Discussion and Conclusions	4.1
4.1	Sediment Physical and Chemical Characterization	4.1
4.2	Site Water and Elutriate Chemical Characterization	4.2
4.3	Toxicity	4.2
4.4	Bioaccumulation	4.5
5.0	References	5.1
Appendix A.	Quality Assurance/Quality Control Data for Sediment Physical/ Chemical Analyses, Port Chester Project	A.1
Appendix B.	Quality Assurance/Quality Control Data for Water and Elutriate Chemical Analyses, Port Chester Project	B.1
Appendix C.	Water-Column Toxicity Test Data, Port Chester Project	C.1
Appendix D.	Benthic Acute Toxicity Test Data, Port Chester Project	D.1
Appendix E.	Bioaccumulation Test Data, Port Chester Project	E.1
Appendix F.	<i>Macoma nasuta</i> Tissue Chemical Analyses and Quality Assurance Quality Control Data; Port Chester Project	F.1
Appendix G.	<i>Nereis virens</i> Tissue Chemical Analyses and Quality Assurance Quality Control Data; Port Chester Project	G.1

Figures

FIGURE 1.1	Location of Port Chester Project Area and Sample Collection Stations	1.3
FIGURE 2.1.	Testing Containers for Amphipod Static Renewal Toxicity Tests	2.20
FIGURE 3.1.	GC/FID Trace Showing Total Petroleum Hydrocarbons in COMP PC	3.12

Contents (contd)

FIGURE 3.2	Distribution of Unsubstituted PAH Compounds and their Alkylated Homologs in COMP PC.....	3.14
FIGURE 3.3	Typical PAH Distributions for Oil Products	3.15
FIGURE 4.1.	Summary Matrix of Port Chester Sediment Toxicity and Bioaccumulation Potential Compared with the Mud Dump Reference Site	4.3
FIGURE 4.2.	Summary Matrix of Port Chester Sediment Toxicity and Bioaccumulation Potential Compared with the Central Long Island Sound Reference Site.....	4.4

Tables

TABLE 2.1.	List of Analytes, Methods, and Target Detection Limits	2.8
TABLE 3.1	Summary of Sediment Sample Data for Port Chester	3.2
TABLE 3.2	Sediment Compositing Scheme for USACE-NED Chemistry Composites	3.3
TABLE 3.3	Port Chester Sediment Core Descriptions	3.4
TABLE 3.4	Results of Analysis of Port Chester Sediment Samples for Grain Size and Percentage of Moisture	3.5
TABLE 3.5	Results of Analysis of Port Chester Sediment Samples for Metals	3.6
TABLE 3.6	Results of Analysis of Port Chester Sediment for Chlorinated Pesticides	3.8
TABLE 3.7	Results of Analysis of Port Chester Sediment for PCBs	3.9
TABLE 3.8	Results of Analysis of Port Chester Sediment for PAHs and 1,4-Dichlorobenzene	3.11
TABLE 3.9	Extended List of PAH Compounds Analyzed in COMP PC	3.13
TABLE 3.10	Results of Analysis of Port Chester Site Water and Elutriate for Metals.....	3.17
TABLE 3.11	Results of Analysis of Port Chester Site Water and Elutriate for Chlorinated Pesticides and PCBs	3.18
TABLE 3.12	Summary of Water-Column Toxicity Tests Performed with Port Chester Sediment	3.19

Contents (contd)

TABLE 3.13	Summary of Benthic Acute Toxicity Tests Performed with Port Chester Sediment	3.21
TABLE 3.14	Mean Concentrations of Metals in <i>M. nasuta</i> Tissues Exposed to Port Chester Sediment	3.24
TABLE 3.15.	Mean Concentrations of Pesticides and PCBs in <i>M. nasuta</i> Tissues Exposed to Port Chester Sediment	3.25
TABLE 3.16	Mean Concentrations of PAHs in <i>M. nasuta</i> Tissues Exposed to Port Chester Sediment	3.26
TABLE 3.17	Mean Concentrations of Metals in <i>N. virens</i> Tissues Exposed to Port Chester Sediment	3.27
TABLE 3.18	Mean Concentrations of Pesticides and PCBs in <i>N. virens</i> Tissues Exposed to Port Chester Sediment	3.29
TABLE 3.19	Mean Concentrations of PAHs in <i>N. virens</i> Tissues Exposed to Port Chester Sediment	3.30
TABLE 3.20	Magnification Factors of All Analyzed Compounds in <i>Macoma nasuta</i> and <i>Nereis virens</i> Tissues Exposed to the Port Chester Composite Relative to Tissues Exposed to the Mud Dump Reference Site and Central Long Island Sound Reference Site.....	3.31
TABLE 4.1	Comparison of Contaminant Concentrations in <i>N. virens</i> and <i>M. nasuta</i> Tissues Exposed to Proposed Dredged Material from Port Chester with FDA Action Levels and Levels of Concern	4.6

Contents (cont'd)

<p>TABLE 2-13 Summary of Benign Aortic Toxicity Tests Performed with Port Chester Sediment 2-21</p>	<p>TABLE 2-13 Summary of Benign Aortic Toxicity Tests Performed with Port Chester Sediment 2-21</p>
<p>TABLE 2-14 Mean Concentrations of Metals in M. galeata Tissues Exposed to Port Chester Sediment 2-24</p>	<p>TABLE 2-14 Mean Concentrations of Metals in M. galeata Tissues Exposed to Port Chester Sediment 2-24</p>
<p>TABLE 2-15 Mean Concentrations of Pesticides and PCBs in M. galeata Tissues Exposed to Port Chester Sediment 2-25</p>	<p>TABLE 2-15 Mean Concentrations of Pesticides and PCBs in M. galeata Tissues Exposed to Port Chester Sediment 2-25</p>
<p>TABLE 2-16 Mean Concentrations of PAHs in M. galeata Tissues Exposed to Port Chester Sediment 2-29</p>	<p>TABLE 2-16 Mean Concentrations of PAHs in M. galeata Tissues Exposed to Port Chester Sediment 2-29</p>
<p>TABLE 2-17 Mean Concentrations of Metals in W. vernalis Tissues Exposed to Port Chester Sediment 2-27</p>	<p>TABLE 2-17 Mean Concentrations of Metals in W. vernalis Tissues Exposed to Port Chester Sediment 2-27</p>
<p>TABLE 2-18 Mean Concentrations of Pesticides and PCBs in W. vernalis Tissues Exposed to Port Chester Sediment 2-23</p>	<p>TABLE 2-18 Mean Concentrations of Pesticides and PCBs in W. vernalis Tissues Exposed to Port Chester Sediment 2-23</p>
<p>TABLE 2-19 Mean Concentrations of PAHs in W. vernalis Tissues Exposed to Port Chester Sediment 2-30</p>	<p>TABLE 2-19 Mean Concentrations of PAHs in W. vernalis Tissues Exposed to Port Chester Sediment 2-30</p>
<p>TABLE 2-20 Mean Tissue Factors of All Analyzed Organisms in A. saccharinum and W. vernalis Tissues Exposed to the Port Chester Contaminated Sediment Tissues Exposed to the Port Chester Contaminated Sediment Tissues Exposed to the Port Chester Contaminated Sediment 2-31</p>	<p>TABLE 2-20 Mean Tissue Factors of All Analyzed Organisms in A. saccharinum and W. vernalis Tissues Exposed to the Port Chester Contaminated Sediment Tissues Exposed to the Port Chester Contaminated Sediment Tissues Exposed to the Port Chester Contaminated Sediment 2-31</p>
<p>TABLE 2-1 Comparison of Concentrations of Organisms in W. vernalis and M. galeata Tissues Exposed to Port Chester Contaminated Sediment with FDA Action Levels and Levels of Concern 2-40</p>	<p>TABLE 2-1 Comparison of Concentrations of Organisms in W. vernalis and M. galeata Tissues Exposed to Port Chester Contaminated Sediment with FDA Action Levels and Levels of Concern 2-40</p>

1.0 Introduction

1.1 Project Objectives

The objective of the Port Chester project (FP No. 1) was to evaluate proposed dredged material from Port Chester in the Byram River and Port Chester Harbor to determine its suitability for unconfined ocean disposal at the Mud Dump Site or at the Central Long Island Sound Disposal Site (CLIS). Tests and analyses for Mud Dump Site disposal were conducted on Port Chester sediment core samples according to the manual developed by the U.S. Army Corps of Engineers (USACE) and the U.S. Environmental Protection Agency (EPA), *Evaluation of Dredged Material Proposed for Ocean Disposal (Testing Manual)* (EPA/USACE 1991), commonly referred to as the "Green Book," and the regional manual developed by the USACE-New York District (NYD) and EPA Region II, *Guidance for Performing Tests on Dredged Material to be Disposed of in Ocean Waters* (USACE-NYD/EPA Region II 1992), hereinafter referred to as the "Regional Guidance Manual." The Regional Guidance Manual provides specifications for the use of local or appropriate test species in biological tests and identifies chemical contaminants of concern. Because the Port Chester area is located on the border between New York and southeast Connecticut, its dredged material may also be considered for disposal at the Central Long Island Sound Site. Therefore, Port Chester sediment was also tested for possible disposal at the Central Long Island Sound Site according to the USACE-New England Division (NED) guidelines (USACE-NED/EPA Region I 1989).

As required by the Regional Guidance Manual, the evaluation of proposed dredged material from Port Chester consisted of bulk sediment chemical analyses, chemical analyses of site water and dredged material elutriate preparations, water-column and benthic acute toxicity tests, and bioaccumulation studies. Individual sediment core samples collected from Port Chester were analyzed for grain size, moisture content, and total organic carbon (TOC). An additional composite sediment sample, representing the entire area proposed for dredging, was analyzed for bulk density, specific gravity, metals, chlorinated pesticides, polychlorinated biphenyl (PCB) congeners, polynuclear aromatic hydrocarbons (PAH), and 1,4-dichlorobenzene. Site water and elutriate water, prepared from the suspended-particulate phase (SPP) of Port Chester sediment, were analyzed for metals, pesticides, and PCBs. Water-column or SPP toxicity tests were performed with three species, the mysid *Mysidopsis bahia*, the juvenile silverside *Menidia beryllina*, and larvae of the mussel *Mytilus galloprovincialis*. Benthic acute toxicity tests were performed with two amphipods, *Ampelisca abdita*, and *Rhepoxynius abronius*, as well as with the mysid *M. bahia*. Bioaccumulation tests were conducted with the burrowing worm *Nereis virens* and the surface-feeding clam *Macoma nasuta*.

Additional testing requirements specified by the USACE-NED consisted of an extended list of sediment chemical parameters of additional metals, chlorinated pesticides, and PAHs.

1.2 Project Background

The proposed Port Chester project area is located on the border of southwest Connecticut and New York in the Byram River and Port Chester Harbor. The Byram River empties into Long Island Sound (Figure 1.1). The project requires dredging and disposal of an estimated 40,000 cu yd of sediment. Project depth of the channel is -10 ft mean low water (MLW) plus 2 ft of overdepth. Port Chester was one of seven waterways that the USACE-NYD requested the Battelle Marine Sciences Laboratory (MSL) to evaluate in a series of dredged material projects that became known as the New York/New Jersey Federal Projects 2 program. The projects evaluated under the Federal Projects 2 program were Port Chester, the Hudson River, South Brothers Island, Gravesend Bay Anchorage, Brown's Creek and Buttermilk Channel survey. Sediment samples from 12 reaches in these waterways were collected during a survey that took place from March 7 through March 14, 1994. Combining sample collection and concurrent evaluation of multiple dredged material projects was more cost-effective for the USACE-NYD, because the expense of reference site testing and quality control analyses could be shared among project budgets. The results of each project were reported separately.

1.3 Organization of This Report

Following this introduction, Section 2 presents the methods and materials used for sample collection, sample processing, sediment sample analysis of physical and chemical parameters, and quality assurance. Results of all physical/chemical analyses and bioassays are presented in Section 3. A discussion of the results and conclusions are provided in Section 4. Section 5 lists the literature cited in this report. Appendix A contains tabulated quality control data for all physical and chemical sediment analyses. Appendix B contains results of replicate sample analyses and quality control data for site water and elutriate chemical parameters. Appendix C contains raw data associated with water-column toxicity tests, such as water quality measurements, test animal survival data, and results of reference toxicant tests. Similar data for benthic acute toxicity tests are provided in Appendix D. Appendix E contains water quality measurements, test animal survival data, and results of reference toxicant tests for the bioaccumulation tests. Appendix F contains replicate sample results and quality control data for chemical analyses of *M. nasuta* tissue samples generated by the bioaccumulation tests, and Appendix G contains replicate sample results and quality control data for chemical analyses of *N. virens* tissue samples.




FIGURE 1.1. Location of Port Chester Project Area and Sample Collection Stations




FIGURE 1. Location of Fort Chesapeake and its various buildings.

2.0 Materials and Methods

2.1 Sediment and Water Collection

The sampling plan required collection of sediment from 13 stations, but samples from stations 12 and 13 were not collected because the depth (MLW) of those stations was greater than project depth. Numerous unsuccessful attempts were made to locate shoaled areas in the vicinity of stations 12 and 13 using the ship's fathometer. Samples were collected from 11 stations within Port Chester. Sampling locations were selected by the USACE-NYD based on recent bathymetric surveys. The locations, their coordinates, and water and core sampling depths are presented with the sampling results in Section 3.0. Water samples were collected at a representative location in Port Chester and the Mud Dump Site. Reference sediments were collected from the Mud Dump Reference Site and the Central Long Island Sound Reference Site. All samples were collected aboard the *M/V Gelberman*, a vessel owned and operated by USACE-NYD at Caven Point, New Jersey.

2.1.1 Test Sediment and Site Water Sampling

Test sediment core samples were collected using a vibracore sampler deployed from the *Gelberman*. The approximate sampling locations were first determined with the aid of reference to landmarks, such as shoreline features or buoys, as well as by water depth. Then, a hand-held Magellan Global Positioning System (GPS) was used to identify and record (within 30 m) each sampling station. The vessel's LORAN was available as a backup system. Water depth at the time of sampling was measured by a fathometer on the ship. The actual water depth was corrected to MLW depth by correcting to the tide height at the time the depth was recorded. The difference between the MLW depth and the project depth, plus 2 ft overdepth, yields the amount of core required. Sediment from stations representative of the upper west shore sites was difficult to collect with the vibracore sampler and caused damage to the core cutter head. To evaluate the problem, van Veen grab samples were taken at those sites. The sediment was characterized as gravel and stone, which is similar to material present on the shore at various commercial facilities. At the stations where this problem occurred, sediment was collected using a van Veen grab sampler.

Core samples were collected aboard the *Gelberman* using the MSL's vibracore sampler. The vibracore sampler consisted of a 4-in. outer diameter, steel core barrel attached to an electric vibratory hammer. The vibratory hammer could be fitted to steel core barrels of various lengths, depending on the length of core needed. To collect a core sample, the core barrel was fitted with a 3.125-in. interior diameter, steam-cleaned, Lexan polycarbonate tube. The vibracore was then

suspended by the ship's crane. Once the coring apparatus was directly above the sampling station, the core was lowered through the water to the sediment surface. At this point, the station coordinates were recorded from the Magellan GPS, and water depth was recorded from the ship's fathometer. The vibratory hammer was switched on until the corer penetrated through the sediment to the desired project depth. Adequate penetration was determined relative to marks on the outside of the core barrel and on the cable suspending the vibracore from the crane. The vibracore apparatus was then pulled out of the sediment and lowered onto the ship's deck. A cutter-head and core-catcher assembly prevented loss of the sediment through the bottom of the core liner. After each core was brought on board, the liner was pulled from the barrel and the length of cored sediment was measured from the mudline to determine whether the appropriate depth had been reached. If not, the liner was replaced and a second core sample was attempted. If the sediment core length achieved project depth plus 2 ft overdepth, it was capped, sealed with tape, and labeled. While on board the sampling vessel, cores were kept cool (~4°C) in a freezer on the deck of the ship. If necessary, cores were cut into shorter sections to fit in the freezer.

A surface-water sample for site water chemical analysis was collected at one station in Port Chester (PC-5). Site water was also collected from the Mud Dump Site for chemical analysis and used as dilution water in water-column toxicity tests. Water samples were collected using a clean, epoxy-coated bucket below the surface of the water. Water was then transferred to precleaned, 20-L polypropylene carboys. (Prior to the sampling survey, carboys were washed with hot water and detergent, acid-rinsed with dilute hydrochloric acid, then rinsed with distilled water, followed by acetone and methylene chloride). The carboys were rinsed with site water three times before filling. Water samples were labeled and stored at ambient temperature (in the shade) while on board the ship.

A log book was maintained containing records of each sample collected, including station designation, coordinates, replicate number, date, sampling time, water depth, core length, and number of core sections per core. At the end of each sampling day, all sediment cores and water samples were loaded into a refrigerated van, thermostatically controlled to maintain approximately 4°C. Sample identification numbers were logged on chain-of-custody forms daily.

At the conclusion of the sample collection survey, sediment cores and water samples were shipped by refrigerated van from Caven Point, New Jersey, to the MSL in Sequim, Washington. The shipment departed from Caven Point on March 14, 1994, and arrived at the MSL on March 18, 1994.

2.1.2 Reference and Control Sediment Sampling

Reference sediment for toxicity and bioaccumulation tests was collected from the Mud Dump and the Central Long Island Sound Reference Sites. Four 5-gal containers of surficial sediment were collected from each site using a van Veen grab sampler. After recovery, the sediments were transferred to epoxy-coated steel buckets. The buckets were covered, labeled, and stored in a freezer at 4°C on the deck of ship, and then were transferred to the refrigerated van at the end of the sampling day.

Records of the collected reference sediment included latitude/longitude, replicate number, date, sampling time, and water depth. Sample identification numbers were logged on chain-of-custody forms.

Control sediments were used in each toxicity and bioaccumulation test to validate test procedures. Control sediment used in *M. nasuta* and *M. bahia* tests was collected from Sequim Bay, Washington, using a van Veen sampler deployed from an MSL research vessel. *R. abronius* control sediment was collected from West Beach, at Whidbey Island, Washington, using a small anchor-dredge sampler specially designed for collecting the amphipods and their sediment. Locations of these control sites were determined by reference to known shoreline features. While in transit from the sampling site, all control sediments were placed in coolers maintained at ambient temperature which were then stored in a walk-in cold room at 4°C±2°C upon arrival at the MSL. Native sediment for *A. abdita* and *N. virens* were supplied with the test organisms by their respective suppliers.

2.2 Test Organism Collection

Seven species of test organisms were used to evaluate sediment samples from the Port Chester project area:

- *Ampelisca abdita*, a tube-dwelling, surface detrital-feeding amphipod
- *Rhepoxynius abronius*, a free-burrowing, subsurface detrital-feeding amphipod
- *Mysidopsis bahia*, a juvenile (1-to 5-day old) mysid shrimp
- *Menidia beryllina*, a juvenile (9-to 14-day old) silverside fish
- *Mytilus galloprovincialis*, the larval zooplankton stage of the mussel
- *Macoma nasuta*, the bent-nose clam, a burrowing, detrital-surface feeder
- *Nereis virens*, a burrowing, deposit-feeding polychaete.

All test organisms except mysids, silversides, and mussels were wild-captured animals, collected either by a commercial supplier or by MSL personnel. The amphipod *A. abdita* was supplied by East Coast Amphipod, Kingston, Rhode Island. *A. abdita* and its native sediment were collected from Narragansett Bay, Rhode Island, by dragging a large dipnet along the sediment surface. Test organisms were carefully removed from their tubes for counting, and then placed in clean, native sediment for overnight transport to the MSL. The amphipod *R. abronius*

was collected by MSL personnel from West Beach, at Whidbey Island, using the same anchor-dredge sampler that was used for collecting the amphipod's native sediment. The amphipods were transported to the MSL in clean coolers containing approximately 10 cm of sediment and 5 gal of clean seawater at a temperature approximating natural conditions. Mysids were purchased from Aquatic Biosystems, Fort Collins, Colorado. Mysids that were less than 24 h old were shipped via overnight delivery in plastic bags containing oxygen-supersaturated seawater maintained at approximately 15°C with "blue ice." Silversides were supplied by Aquatic Research Organisms in Hampton, New Hampshire, and were shipped via overnight delivery in plastic bags containing oxygen-supersaturated seawater maintained at approximately 22°C with blue ice. Mussels used for obtaining *M. galloprovincialis* larvae were purchased from the commercial supplier Johnson and Gunstone, Quilcene, Washington. Mussels were wrapped in moist paper towels and transported in a Styrofoam cooler packed with blue ice to maintain an ambient temperature of approximately 15°C. Clams (*M. nasuta*) were collected from intertidal zones in Discovery Bay, Washington, by Johnson and Gunstone. The clams were kept in large containers filled with sediment and seawater obtained from the collection site and transported to the MSL. Worms (*N. virens*) were purchased through Enviro Systems, Inc., Hampton, New Hampshire, and were collected from an intertidal region in Newcastle, Maine. The worms were packed in insulated boxes with mats of moist seaweed and shipped at ambient temperature to the MSL via overnight delivery.

All organisms were shipped or transported in native sediment or under conditions designed to ensure their viability. After arrival at the MSL, the test organisms were gradually acclimated to test conditions. Animals with abnormal behavior or appearance were not used in toxicological tests. All acclimation and animal care records are part of the raw data files for these projects.

2.3 Sediment Sample Preparation

Sediment sample preparation consists of all steps performed in the laboratory between receipt of the samples at the MSL and the preparation of samples for biological testing and physical/chemical analyses. Sediment samples for physical, chemical, and biological analysis were prepared from individual core samples, composites of a number of core samples, reference sediment, and control sediment. All sediment samples were assigned random, unique code numbers to ensure that samples are handled without bias by staff in the biology or chemistry laboratories.

Sediment for biological testing was used within the 6-week holding period specified in the Green Book. During this holding time, the sediment samples were received at the MSL;

inventoried against chain-of-custody forms; processed and used for benthic and water-column toxicity tests, elutriate analysis, and bioaccumulation tests; and subsampled for sediment physical/chemical analyses. This section describes procedures for equipment preparation, compositing strategy, and preparation of sediments for biological testing and chemical analyses.

2.3.1 Laboratory Preparation and Safety Considerations

All glassware, stainless-steel or titanium utensils, Nalgene, Teflon, and other laboratory containers and equipment underwent stringent cleaning procedures to avoid contamination of samples. Glassware (e.g., test containers, aquaria, sediment transfer dishes) was washed with hot water and detergent, rinsed with deionized water, then soaked in a 10% solution of reagent grade nitric acid for a minimum of 4 h and rinsed again with deionized water before it was allowed to air dry. Glassware was then rinsed with methylene chloride and allowed to dry under a fume hood. Polyvinyl chloride (PVC), Nalgene, and Teflon tools were treated in the same manner as glassware. Stainless-steel bowls, spoons, spatulas, and other utensils were washed with hot water and detergent, rinsed with deionized water, and allowed to air dry. They were then solvent-rinsed with methylene chloride and allowed to dry under a fume hood.

Neoprene stoppers and polyethylene sheets or other porous materials were washed with hot water and detergent and rinsed with deionized water. These items were then "seasoned" by continuous soaking in 0.45- μ m filtered seawater for at least 2 days prior to use. Large pieces of laboratory equipment, such as the epoxy-coated sediment mixer, were washed with a dilute solution of detergent, and thoroughly rinsed with tap water followed by deionized water.

Equipment used for determining water quality, including the meters for pH, dissolved oxygen (DO), temperature, and salinity, were calibrated according to the manufacturers' specifications and internal MSL standard operating procedures (SOPs). Calibration records are part of the raw data files for this project.

Because the potential toxicity of the Port Chester sediment was unknown, sediment processing and testing were segregated from other laboratory activities. Specific areas at the MSL were established for sample storage and for core-cutting, sediment mixing, and sediment sieving. Work areas were covered with plastic sheeting to contain any waste sediment. Wastewater generated during all operations was retained in 55-gal barrels and periodically pumped through activated charcoal filters and into the MSL's wastewater treatment system. These procedures minimized any potential for cross-contamination of sediment samples.

Laboratory staff members were protected by personal safety equipment such as Tyvek suits, plastic aprons, and rubber gloves. Those who were likely to have the most exposure to the potential volatile compounds in the bulk sediment (i.e., those responsible for opening, homogenizing, and compositing core samples) were also provided with half-mask respirators.

2.3.2 Preparation of Sediment for Benthic Testing and Bulk Sediment Analyses

Each core was opened by scoring the Lexan core liner longitudinally with a circular saw and splitting the liner with a clean linoleum knife to expose the sediment. As each sediment core sample was opened, it was examined for physical characteristics (e.g., sediment type and consistency, color, odor). In particular, the presence of any strata in the cores was noted. All core observations were recorded in the sediment preparation log book. The sediment between the mudline and project depth was then transferred from the core liner to a clean, stainless-steel bowl by scooping the sediment from the core liner with a spoon or spatula. The sediment was mixed by hand with stainless-steel utensils until the color and consistency appeared homogenous, creating a sample representative of the individual sampling station. Sieving was not necessary because organisms that might interfere with the benthic toxicity tests were not present in the sediment samples.

Aliquots of the homogenized sediment were then transferred to the appropriate sample jar(s) for physical or chemical analyses required on individual core samples. A portion of each homogenized core sample was also retained as an archive sample. The remainder of the homogenized sediment from the individual core stations was combined to create a composite sample representing the entire Port Chester project area, designated COMP PC. The composite sediment was homogenized in an epoxy-coated mixer. Aliquots of homogenized composite sediment were transferred to the appropriate sample jar(s) for physical or chemical analyses required on the composite sample. Additional composites were created for chemical analysis as required for USACE-NED. The compositing scheme for these samples is provided in Section 3. A portion of the homogenized composited sediment was also retained as an archive sample. The remainder was stored in labeled epoxy-coated pails, tightly covered, at $4^{\circ}\text{C}\pm 2^{\circ}\text{C}$ until used for SPP/elutriate preparation or benthic toxicity and bioaccumulation tests.

The Mud Dump Reference Site sediment, Central Long Island Reference Site sediment, *M. nasuta* native control sediment, and *N. virens* native control sediment were homogenized in the large, epoxy-coated mixer, but prior to mixing, these sediments were pressed through a 1-mm mesh to remove live organisms that might affect the outcome of toxicity tests. After mixing, aliquots for physical and chemical analyses were removed. Control sediments for *A. abdita*, *R. abronius*, and *M. bahia* were sieved through a 0.5-mm mesh to remove live organisms and mixed in stainless-steel bowls after sieving. All reference and control sediments were stored at $4^{\circ}\text{C}\pm 2^{\circ}\text{C}$ until use in benthic toxicity and bioaccumulation tests.

2.3.3 Preparation of Suspended-Particulate Phase and Elutriate

Toxicological effects of dredged sediment contaminants dissolved and suspended in the water-column at an open-water disposal site were simulated in the laboratory by preparation of the SPP. To prepare the SPP, a sediment-water slurry was created and centrifuged at low speed. Low speed centrifugation provided a timely SPP preparation and maintained consistency between projects. The supernatant was decanted and reserved for testing with water-column organisms. The elutriate phase was prepared by centrifuging the SPP at a higher speed and collecting the supernatant. This liquid was analyzed for chemical constituents to identify water-soluble contaminants that could remain in the water-column after dredge and disposal operations.

The SPP was prepared by creating a 4:1 (volume:volume) water-to-sediment slurry in 1-L glass jars with Teflon-lined lids. The jars were marked at 200 mL and 400 mL and filled to the 200-mL mark with 0.45- μ m-filtered Sequim Bay seawater. Sequim Bay seawater was substituted for dredging site water to maintain consistency in salinity among the dredging projects tested. Homogenized COMP PC sediment was added until the water was displaced to the 400-mL mark. Each jar was then filled to 1 L with filtered seawater, placed on a shaker table, and agitated for 30 min at 120 to 150 cycles/min. The slurry was then transferred to 500-mL Teflon jars, tightly sealed, and centrifuged at approximately 1750 rpm for 10 min, at a relative centrifugal force of approximately 1000 g. Following centrifugation, the supernatant was poured into 4-L glass jars. The Teflon jars were rinsed after each use and the above process continued until an adequate amount of SPP was produced from each composite. Between SPP preparations, all glass and Teflon containers were cleaned according to procedures described in Section 2.3.1. When all SPP for a treatment was prepared, portions were taken for elutriate preparation. The remaining SPP was either used immediately for biological tests or stored at $4^{\circ}\text{C}\pm 2^{\circ}\text{C}$ and used within 24 h for testing. The 100% COMP PC SPP was mixed with Mud Dump Site water to yield three dilutions: 0%, 10%, and 50% SPP, for a total of four concentrations.

To prepare elutriate for chemistry analyses, a 1-L aliquot of the SPP was collected in an acid-washed Teflon bottle for trace metals analysis, and three 1-L aliquots were collected in EPA-certified amber glass bottles for analysis of organic compounds. The SPP for metals analysis was transferred to acid-washed polycarbonate centrifuge jars, and the SPP for analysis of organic compounds was transferred to Teflon centrifuge jars. Both were centrifuged at 2000 rpm for 30 min at a relative centrifugal force of approximately 1200 g. The decanted supernatant liquid was the elutriate phase. One liter of elutriate was submitted for triplicate trace metals analysis and three 1-L portions were submitted for analysis of organic compounds.

2.4 Physical and Chemical Analytical Procedures

Sediment cores, composited bulk sediment, water, elutriate, and tissue samples were analyzed for physical and chemical parameters. Table 2.1 lists the parameters measured in each sample type, the method used for each analysis, and the target analytical detection limits. The following sections briefly describe the procedures used for physical and chemical analyses. Procedures followed those required by the Regional Guidance Manual unless otherwise noted.

2.4.1 Grain Size and Percentage of Moisture

Grain size was measured following two methods described by Plumb (1981). The wet sieve method was used to determine the size distribution of sand or coarser-grained particles larger than a U.S. No. 230 standard sieve (62.5- μm mesh). The size distribution of particles smaller than a U.S. No. 230 sieve was determined using the pipet method. Grain size was reported as percentages within four general size classes:

gravel	>2000- μm diameter
sand	$\geq 62.5\text{-}\mu\text{m}$ diameter and <2000- μm diameter
silt	$\geq 3.9\text{-}\mu\text{m}$ diameter and < 62.5- μm diameter
clay	< 3.9- μm diameter.

Percentage of moisture was obtained using the Plumb (1981) method for determining total solids. The procedure involves drying a sediment sample at 100°C until a constant weight is obtained. Percentage of moisture was calculated by subtracting the percentage of total solids from 100%.

2.4.2 Bulk Density and Specific Gravity

Bulk density, or unit weight, was determined according to EM 111-2-1906 (USACE 1970). Specific gravity, the ratio of the mass of a given volume of material to an equal volume of water at the same temperature, was measured according to ASTM D-854.

2.4.3 TOC

Samples were analyzed for TOC according to the EPA Edison, New Jersey, Laboratory Procedure (EPA 1986). Inorganic carbon was removed from the sediment sample by acidification. The sample was combusted and the evolved carbon dioxide was quantitated using a carbon-hydrogen-nitrogen (CHN) analyzer. TOC was reported as a percentage of the dry weight of the unacidified sample.

TABLE 2.1. List of Analytes, Methods, and Target Detection Limits

<u>Analyte</u>	<u>Methods</u>	<u>Sediment Detection Limit (a)</u>	<u>Tissue Detection Limit (b)</u>	<u>Water Detection Limit</u>
PHYSICAL PARAMETERS				
Grain Size	Plumb (1981)	1.0%		
Specific Gravity	ASTM D-854			
Bulk Density	EM 1110-2-1906 (USACE 1970)			
Percent Moisture	Sediment: Plumb (1981)	1.0 %		
	Tissue: Freeze-dry		1.0 %	
METALS				
Arsenic	EPA 200.2, -.3, -.8 (c)	0.1 mg/kg	1.0 mg/kg	---
Cadmium	EPA 200.2, -.3, -.8 (c)	0.01 mg/kg	0.1 mg/kg	0.025 µg/L
Chromium	EPA 200.2, -.3, -.8 (c)	0.02 mg/kg	0.2 mg/kg	1.0 µg/L
Copper	EPA 200.2, -.3, -.8 (c)	0.1 mg/kg	1.0 mg/kg	0.35 µg/L
Lead	EPA 200.2, -.3, -.8 (c)	0.1 mg/kg	0.1 mg/kg	0.35 µg/L
Mercury	EPA 245.5 (sed.); 245.6 (tiss.) (c)	0.02 mg/kg	0.02 mg/kg	
	Bloom and Crecelius (1983) (water)			0.002 µg/L
Nickel	EPA 200.2, -.3, -.8 (c)	0.1 mg/kg	0.1 mg/kg	0.30 µg/L
Silver	EPA 200.2, -.3, -.9 (c)	0.1 mg/kg	0.1 mg/kg	0.25 µg/L
Zinc	EPA 200.2, -.3, -.8 (c)	0.1 mg/kg	1.0 mg/kg	0.15 µg/L
METALS (Required for Central Long Island Sound Disposal Site Testing)				
Antimony	EPA 200.2, -.3, -.8, -.9 (c)	0.1 µg/kg		
Beryllium	EPA 200.2, -.3, -.8, -.9 (c)	0.1 µg/kg		
Selenium	EPA 200.2, -.3, -.8, -.9 (c)	0.1 µg/kg		
Thallium	EPA 200.2, -.3, -.8, -.9 (c)	0.1 µg/kg		
ORGANIC COMPOUNDS				
<u>TOC</u>	EPA (1986)	0.1%		
<u>Pesticides</u>				
Aldrin	EPA 8080 (sediment, tissue)	1.0 ng/g	0.4 ng/g	
	EPA 608 (water) (c)			0.004 µg/L
α-Chlordane	EPA 8080 (sediment, tissue)	1.0 ng/g	0.4 ng/g	
	EPA 608 (water) (c)			0.014 µg/L
<i>trans</i> -Nonachlor	EPA 8080 (sediment, tissue)	1.0 ng/g	0.4 ng/g	
	EPA 608 (water) (c)			0.014 µg/L
Dieldrin	EPA 8080 (sediment, tissue)	1.0 ng/g	0.4 ng/g	
	EPA 608 (water) (c)			0.002 µg/L

TABLE 2.1. (contd)

Analyte	Method(s)	Sediment Detection Limit	Tissue Detection Limit	Water Detection Limit
4,4'-DDT	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.012 µg/L
2,4'-DDT	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.020 µg/L
4,4'-DDD	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.011 µg/L
2,4'-DDD	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.020 µg/L
4,4'-DDE	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.004 µg/L
2,4'-DDE	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.020 µg/L
Endosulfan I	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.014 µg/L
Endosulfan II	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.004 µg/L
Endosulfan sulfate	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.010 µg/L
Heptachlor	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.003 µg/L
Heptachlor epoxide	EPA 8080 (sediment, tissue) EPA 608 (water) (c)	1.0 ng/g	0.4 ng/g	0.100 µg/L

PESTICIDES (Required For Central Long Island Sound Disposal Site Testing)

Endrin	EPA 8080	0.02 mg/kg
Endrin aldehyde	EPA 8080	0.02 mg/kg
α-Hexachlorocyclohexane	EPA 8080	0.02 mg/kg
β-Hexachlorocyclohexane	EPA 8080	0.02 mg/kg
δ-Hexachlorocyclohexane	EPA 8080	0.02 mg/kg
γ-Hexachlorocyclohexane	EPA 8080	0.02 mg/kg
Methoxychlor	EPA 8080	0.02 mg/kg
Toxaphene	EPA 8080	0.02 mg/kg

TABLE 2.1. (contd)

<u>Analyte</u>	<u>Method(s)</u>	<u>Sediment Detection Limit</u>	<u>Tissue Detection Limit</u>	<u>Water Detection Limit</u>
PCBs				
PCB 8	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 18	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 28	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 44	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 49	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 52	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 66	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 87	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 101	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 105	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 118	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 128	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 138	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 153	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 170	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 180	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 183	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 184	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 187	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 195	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 206	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PCB 209	NYSDEC (1992) ^(d)	1.0 ng/g	0.4 ng/g	0.0005 µg/L
PAHs				
Acenaphthene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Acenaphthylene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Anthracene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Fluorene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Naphthalene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Phenanthrene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Benzo[a]anthracene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Benzo[a]pyrene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Benzo[b]fluoranthene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Benzo[g,h,i]perylene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Benzo[k]fluoranthene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Chrysene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Dibenzo[a,h]anthracene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Fluoranthene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Indeno[1,2,3-cd]pyrene	EPA 8270(C)	10 ng/g	4.0 ng/g	
Pyrene	EPA 8270(C)	10 ng/g	4.0 ng/g	

TABLE 2.1. (contd)

Analyte	Method(s)	Sediment Detection Limit	Tissue Detection Limit	Water Detection Limit
PAHS (REQUIRED FOR Central Long Island Sound Disposal Site Testing)				
Biphenyl	EPA 8270(C)	0.02 µg/g		
2,6 dimethylnaphthalene	EPA 8270(C)	0.02 µg/g		
1-methylphenanthrene	EPA 8270(c)	0.02 µg/g		
1-methylnaphthalene	EPA 8270(C)	0.02 µg/g		
2-methylnaphthalene	EPA 8270(C)	0.02 µg/g		
Industrial Chemicals				
1,4-Dichlorobenzene	EPA 8270(C)	1 ng/g	0.4 ng/g ©	
Lipids				
	Bligh & Dyer (1959) Randall (1988)		0.1%	

(a) Detection limits are in dry weight for all sediment parameters except Hg and Lipids

(b) Detection limits are in wet weight for all organic and inorganic tissue parameters.

(c) Equivalent MSL standard operating procedures were substituted for the methods cited.

2.4.4 Metals

Preparation and analysis of water samples for Cd, Cr, Cu, Pb, Ni, Ag, and Zn were conducted according to MSL SOPs equivalent to EPA Methods 200.2 and 200.9 (EPA 1991). Samples were chelated with 2% ammonium pyrrolidinedithiocarbamate (APDC), precipitated out of solution, and filtered. The filter was digested in concentrated nitric acid and the digestate was analyzed by graphite furnace atomic absorption (GFAA) spectroscopy for Cr and Zn, or by inductively coupled plasma/mass spectrometry (ICP/MS) for Cd, Cu, Pb, Ni, and Ag. Water samples were analyzed for Hg directly by cold vapor atomic fluorescence (CVAF) according to the method of Bloom and Crecelius (1983). This CVAF technique is based on emission of 254-nm radiation by excited elemental Hg atoms in an inert gas stream. Mercuric ions in an oxidized sample were reduced to elemental Hg with tin chloride (SnCl₂), then purged onto gold-coated sand traps to preconcentrate the Hg and remove interferences. Mercury vapor was thermally desorbed to a second "analytical" gold trap, and from that into the fluorescence cell. The amount of fluorescence (indicated by peak area) is proportional to the quantity of Hg collected, and was quantified using a standard curve as a function of the quantity of the sample purged.

Sediment samples for analysis of USACE-NYD metals, As, Cd, Cr, Cu, Pb, Ni, and Zn and the USACE-NED metals, Tl, Be, Se, and Sb, were prepared according to an MSL SOP equivalent to EPA Method 200.2 (EPA 1991). Solid samples were first freeze-dried and blended

in a Spex mixer mill. A 0.2- to 0.5-g aliquot of dried homogeneous sample was then digested using peroxide and nitric acid. Samples were heated in sealed Teflon bombs overnight at approximately 130°C. Sediment samples were analyzed for As, Cd, Cr, Cu, Pb, Ni, Zn, Ti, Be, Se, and Sb using ICP/MS, following an MSL SOP based on EPA Method 200.8 (EPA 1991). Sediment samples were analyzed for Ag by GFAA according to an MSL SOP based on EPA Method 200.9 (EPA 1991). Sediments were analyzed for Hg by CVAA according to an MSL procedure for total Hg determination equivalent to EPA Method 245.5 (EPA 1991).

Sediment samples initially showed poor matrix spike recovery for Ag. (Refer to Appendix A, QA/QC Summary for analysis of metals in sediment.) EPA Method 200.2 was modified by the addition of aqua regia to the digestion procedure and all samples were reanalyzed for Ag. Matrix spike recoveries improved and concentrations of Ag in the dredging site sediments increased slightly. The low recovery of Ag appears to occur in analysis of marine sediment samples having high (in excess of approximately 5 µg/g) Ag concentrations. During the EPA Method 200.2 digestion procedure, a precipitate of AgCl can form with the Ag in the sediment and the Cl in the seawater. The sample reanalyses showed little change between the EPA Method 200.2 digestion and the aqua regia-modified digestion because the dredging site sediments tested had fairly low levels of Ag. (Most samples were approximately 0.1 µg/g to 3 µg/g, with a few as high as 9 µg/g.) However, the aqua regia modification resulted in improved recovery of Ag in the matrix spike samples that were spiked with higher concentrations of Ag (20 µg/g). The additional metals required by USACE-NED (Sb, Be, Se, and Ti) were also analyzed in the sample extracts obtained from the aqua regia-modified digestion procedure.

Tissue samples were prepared according to an MSL SOP based on EPA Method 200.3 (EPA 1991). Solid samples were freeze-dried and blended, and a 0.2- to 0.5-g aliquot of dried homogeneous sample was then digested in a microwave using nitric acid, hydrogen peroxide, and hydrochloric acid. Tissue samples were analyzed for As, Cd, Cr, Cu, Pb, Ni, Ag, and Zn using the ICP/MS method (EPA Method 200.8 [EPA 1991]). Tissue samples were analyzed for Hg by CVAA following an MSL procedure equivalent to EPA Method 245.6 (EPA 1991).

2.4.5 Chlorinated Pesticides and PCBs

Water samples were prepared and analyzed for chlorinated pesticides and PCBs according to an MSL procedure equivalent to EPA Method 8080 (EPA 1990), and incorporating techniques developed by the National Oceanic and Atmospheric Administration (NOAA) National Status and Trends "Mussel Watch" Program (NOAA 1993). Samples were extracted with methylene chloride. Extract volumes were reduced and solvent exchanged to hexane. The sample extracts underwent cleanup by alumina and silica column chromatography; further interferences were removed by an additional cleanup treatment using high-performance liquid

chromatography (HPLC). Sample extracts were concentrated and analyzed using gas chromatography with electron capture detection (GC-ECD) using the internal standard technique.

Sediment and tissue samples for analysis of pesticides and PCBs required by both the USACE-NYD and USACE-NED guidance manuals were extracted and analyzed according to an MSL procedure similar to EPA Method 8080 for pesticides and the New York State Department of Environmental Conservation (NYSDEC) Congener-Specific Method 91-11 (NYSDEC 1992). The method also uses techniques from the NOAA Mussel Watch procedure. A 20- to 50-g sample of homogenized sediment or macerated tissue was first combined with sodium sulfate in a sample jar to remove water. Samples were extracted by adding successive portions of methylene chloride and agitating sample jars at ambient temperature using a roller technique. Extract volumes were reduced and solvent-exchanged to hexane, followed by Florisil column chromatography cleanup. Interferences were removed using HPLC cleanup; tissue sample extracts underwent an additional cleanup by gel permeation chromatography (GPC). Sample extracts were concentrated and analyzed using GC-ECD by the internal standard technique.

The concentration of total PCB in each matrix was estimated by taking the sum of the 22 congeners (x) and multiplying by two. The procedure for calculation of total PCBs was established in 1996 (Mario Del Vicario, Chief of the Marine and Wetlands Protection Branch, U.S. Environmental Protection Agency Region 2, Feb 14, 1996, letter to John F. Tavolaro, Chief Operations Support Branch, U.S. Army Corps of Engineers, New York District). One-half of the detection limit was used in summation when an analyte was undetected.

2.4.6 PAHs and 1,4-Dichlorobenzene

Sediment samples were prepared for the analysis of 16 PAHs and 1,4-dichlorobenzene, and an additional seven PAHs required by the USACE-NED guidance manual according to an MSL method based on the NOAA Mussel Watch procedure (NOAA 1993). A 20- to 50-g sample of homogenized sediment or macerated tissue was first combined with sodium sulfate in a sample jar to remove water. Samples were extracted by adding successive portions of methylene chloride and agitating sample jars at ambient temperature using an ambient shaker technique. Extract volumes were reduced and solvent-exchanged to hexane, followed by column chromatography cleanup. Interferences were removed using HPLC cleanup; tissue sample extracts underwent an additional cleanup by GPC. Sample extracts were concentrated and analyzed using gas chromatography with mass spectrometry (GC/MS) in the selective ion monitoring (SIM) mode.

2.4.7 Lipids

The lipid content of *M. nasuta* and *N. virens* was determined by the analysis of unexposed background tissue samples of each species. The lipid analysis procedure is a

modification of the Bligh and Dyer (1959) methods, which involves a chloroform extraction followed by gravimetric measurement of lipids. Randall (1988) modified the original Bligh and Dyer method to accommodate a smaller tissue sample size. Lipid analysis was performed in triplicate, once for each species. Lipid concentration was reported as a percentage on both a wet and dry weight basis.

2.5 Biological Testing Procedures

2.5.1 Water-Column Toxicity Tests

Water-column effects of open-water dredged-material disposal were evaluated by exposing three species of water-column organisms to the SPP of the Port Chester sediment composite. The three test species were juvenile *M. beryllina* (silverside) and *M. bahia* (mysid), and larval *M. galloprovincialis* (mussel). Total ammonia monitoring was not performed during water-column toxicity tests, but prior to test initiation total ammonia concentrations were measured for the 100% SPP concentration and is presented in Section 3.4.

2.5.1.1 Water-Column Toxicity Test with *Menidia beryllina*

Upon receipt, the *M. beryllina* were placed in a 10-gal glass aquarium and gradually acclimated from 27.5‰ seawater to 30.0‰ Sequim Bay seawater over a 24-h period. *M. beryllina* were received and held at $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$ prior to testing and were fed concentrated brine shrimp nauplii daily. During acclimation and holding, 2% to 3% mortality of the silversides was observed.

Test containers for the water-column toxicity test with silversides were 500-mL glass jars, labeled with sediment treatment code, concentration, position number, and replicate number. Five replicates of each concentration (0%, 10%, 50%, and 100%) were tested. The 300-mL test volume of SPP was placed in each of the five replicate test chambers. Each test chamber was then placed in a randomly assigned position on a water table at $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$ and allowed to equilibrate to test temperature for several hours. After the concentrations were prepared and placed on the water table, water quality parameters were measured and recorded for all replicates of all concentrations for each sediment treatment.

To initiate the test, *M. beryllina* were transferred from the holding tank to test chambers using a wide-bore pipet and small transfer cups. Ten individuals were introduced to each test chamber, creating a test population of 50 silversides per concentration for each treatment. Ten animals per test chamber were used, rather than the 20 animals per chamber as described in the Regional Guidance Manual, because it is not possible to make accurate daily observations of *M. beryllina* behavior when using 20 animals. Test initiation time and date were recorded.

Following test initiation, water quality parameters were recorded in one replicate of each concentration daily. Because several treatments had DO levels lower than 40% saturation prior to test initiation, all test chambers were aerated to maintain consistency in handling DO concentration among test containers. Acceptable parameters for this test were as follows:

Temperature	20°C±2°C
DO	>40% saturation (>3.04 mg/L at 20°C, 30‰)
pH	7.8±0.5
Salinity	30.0‰±2.0‰.

The test was run under a 16-h light/8-h dark photoperiod, and silversides were fed brine shrimp nauplii daily during the test. Observations of the animals were performed at 2 h, 24 h, 48 h, and 72 h, and the number of live, dead, and missing organisms was recorded. At the end of the 96-h test period, water quality parameters were measured for all test chambers, and the number of live, dead, and missing silversides was recorded on termination forms. As a quality control check, a second observer confirmed surviving test organisms on at least 10% of the termination counts.

A 96-h, water-only, reference toxicant test was performed concurrently with the toxicity test with each population of *M. beryllina* to establish the health and expected response of the test organisms. The reference toxicant test was conducted in the same manner as the water-column toxicity test. *M. beryllina* were exposed to a seawater control plus four concentrations of copper sulfate: 16, 64, 160, and 400 µg/L copper, using three replicates of each concentration.

2.5.1.2 Water-Column Toxicity Test with *Mysidopsis bahia*

Upon receipt, the *M. bahia* were placed in a 10-gal aquarium and gradually acclimated from 28.0‰ seawater to 30‰ Sequim Bay seawater over a 24-h period. Mysids were received and held at 20°C±2°C until testing and were fed concentrated brine shrimp nauplii twice daily prior to testing. Mortality of the *M. bahia* during holding was less than 1%.

The water-column toxicity test with the mysid was performed in 200 mL of test solution in 400-mL jars, labeled with sediment treatment code, concentration, position number, and replicate number. Five replicates of each concentration were tested. Each of the test chambers received 200 mL of test solution, then was placed randomly in a recirculating water bath and allowed to equilibrate to test temperature for several hours. Prior to test initiation, water quality parameters were measured in each replicate of each sediment treatment concentration. Acceptable water quality parameters for this test were as follows:

Temperature	20°C±2°C
DO	>40% saturation (>3.04 mg/L at 20°C, 30‰)
pH	7.8±0.5
Salinity	30.0‰±2.0‰.

To initiate the test, *M. bahia* were transferred from the holding tank to test chambers using a wide-bore pipet via small transfer cups. Ten individuals were introduced to each test chamber, creating a test population of 50 mysids per concentration (200 mysids per treatment). Ten animals per test chamber were used, rather than the 20 animals per chamber as described in the Regional Guidance Manual, because it is not possible to make accurate daily observations of *M. bahia* behavior when using 20 animals. Test initiation time and date were documented on data forms. Observations of test organisms were performed at 4 h, 24 h, 48 h, and 72 h, using a fluorescent light table to enhance visibility of the *M. bahia*. After test initiation, water quality parameters were measured daily in one replicate concentration of all concentrations for each sediment treatment. During the 96-h exposure, *M. bahia* were fed <24-h-old brine shrimp twice daily. Excess food was removed daily with a small pipet, taking care not to disturb test animals. Molted exoskeletons and any particles from the SPP solutions were also removed.

Prior to test termination, water quality parameters were measured in all replicates. At 96 h, the number of live versus dead animals was recorded for each test container. An animal was considered dead if it did not respond to gentle probing. As a quality control check, a second observer confirmed surviving test organisms on at least 10% of the termination counts.

A 96-h, water-only, reference toxicant test was performed concurrently with the SPP toxicity test, to establish the health and expected response of the test organisms. Water quality conditions were the same as for the SPP test, and animals were fed daily over the 96-h exposure. *M. bahia* were exposed to a seawater control plus four concentrations of copper sulfate: 50, 100, 150, and 200 µg/L copper, using three replicates of each concentration.

2.5.1.3 Water-Column Toxicity Test with *Mytilus galloprovincialis* Larvae

Prior to testing, adult *M. galloprovincialis* were held in flowing, unfiltered Sequim Bay seawater at ambient temperatures for approximately 5 days.

Chambers for the bivalve larvae test were 500-mL glass jars labeled with sediment treatment code, concentration, position number, and replicate number. Dilutions of SPP from the sediment composites (0%, 10%, 50%, and 100%) were prepared with Mud Dump Site water in a 2000-mL graduated cylinder, then 300 mL of test solution was transferred into each test chamber. Test chambers were placed in random positions on a water table and allowed to equilibrate to test temperature for several hours. Initial water quality parameters were measured in all replicates once test chambers reached testing temperatures (16°C±2°C).

Spawning was induced by placing *M. galloprovincialis* into 15°C, filtered Sequim Bay seawater and rapidly raising the holding water temperature to 20°C. Spawning generally occurs within 1 h of temperature elevation; however, on the first day of spawning, gametes were shed after 3 h to 4 h. For this group of mussels, the water bath was changed when DO levels fell

below 3.0 mg/L. When spawning began, males and females were identified and isolated in individual jars containing filtered Sequim Bay seawater and allowed to shed gametes for approximately 45 min. Eggs from each female were filtered through a 75- μ m Nytex screen into separate jars to remove feces, detritus, and byssal fibers. Sperm from at least three males was pooled and 10 mL of sperm solution was then added to each of the egg stocks. Egg-sperm solutions were gently mixed every 10 min with a perforated plunger. Fertilization proceeded for 1 h, then fertilization rate (percentage of fertilized eggs) was determined by removing a subsample and observing the number of multicell-stage embryos. Fertilization was considered successful if greater than 90% of the embryos were in the multicell stage. Egg stocks with greater than 90% fertilization were combined and rinsed on a 20- μ m Nytex screen to remove excess sperm. Stock embryo solution density was estimated by removing a 0.1-mL subsample and counting all multicell embryos, then multiplying by 10 to yield embryo density (embryos/mL). Stock solution was diluted or concentrated to yield 7500 to 9000 embryos/mL. The test was initiated by introducing 1 mL of stock solution into each test chamber, to produce embryo densities of 25 to 30 embryos/mL. Test initiation date and time were recorded on data sheets. Following initiation, 10-mL stocking-density subsamples were removed from each container and preserved in 5% formaldehyde to determine actual stocking density later.

Water quality parameters were measured in one replicate of each concentration per treatment daily throughout the test. Acceptable ranges for water quality parameters were as follows:

Temperature	16°C \pm 2°C
DO	>60% saturation (>4.93 mg/L at 16°C, 30‰)
pH	7.8 \pm 0.
Salinity	30.0‰ \pm 2.0‰.

Each chamber was provided with gentle aeration to maintain consistency in handling DO concentration among test containers. The bivalve test was terminated after 72 h when greater than 80% of the larvae in the controls had reached the D-cell stage. Final water quality parameters were recorded for all replicates. The contents of each chamber were then homogenized with a perforated plunger, and a 10-mL subsample was removed and placed into a 20-mL scintillation vial. The subsample was then fixed with 1 mL of 50% solution of formaldehyde in seawater. Samples were scored for the appearance of normal and abnormal D-shaped larvae, blastula larvae, and total number of larvae. At least 10% of the counts were confirmed by a second observer.

A 72-h reference toxicant test was conducted to establish the health and expected response of the test organisms. The reference toxicant test was set up and conducted in the same manner as the liquid-phase tests. *M. galloprovincialis* larvae were exposed to a filtered

Sequim Bay seawater control plus copper sulfate concentrations of 1, 4, 16, and 64 µg/L copper, with three replicates per concentration.

2.5.2 Benthic Acute Toxicity Tests

Deposited sediment effects of open-water dredged material disposal were evaluated by benthic acute toxicity tests with three marine amphipod species, *A. abdita* and *R. abronius*, and the mysid *M. bahia*.

2.5.2.1 Static Renewal Tests with *Ampelisca abdita* and *Rhepoxynius abronius*

Upon receipt, the *A. abdita* were placed in a tub of clean sand from their collection area and gradually acclimated with flowing Sequim Bay seawater from 28‰ to 30.5‰, over a period of 2 days. *A. abdita* were received at approximately 11°C and acclimated to 20°C±2°C over 4 days. They were held at 20°C±2°C for one day and were not fed prior to testing. The *R. abronius* were also placed in a tub of clean sand from their collection area and held under flowing seawater upon arrival at the laboratory. They were received and held at a salinity of 30‰±2‰ and a temperature of 15°C±2°C until testing. *R. abronius* were not fed during the 11-day holding period.

All amphipod static renewal tests were performed in 1-L glass jars modified for use as flow-through test chambers. The test chambers were fitted with funneled lids and screened outflow and overflow ports (Figure 2.1). The static-renewal system was turned on long enough to deliver the seawater at a rate of two chamber exchanges per day. Five replicates of COMP PC, Mud Dump Reference Site, Central Long Island Sound Reference Site, and native test animal control treatments were tested.

Concentrations of ammonia have been encountered in the porewater of sediment core samples from New York/New Jersey waterways at concentrations high enough to affect survival of amphipods in benthic toxicity tests (Barrows et al. 1996). Therefore, the amphipod tests were conducted according to the ammonia protocols issued by EPA and the USACE (EPA/USACE 1993). This guidance requires postponing test initiation (exposure of test animals) until porewater total ammonia concentrations are <30 mg/L for *A. abdita* and *R. abronius*. During this "purging" period, test chambers were set up and maintained under test conditions, and the overlying water was exchanged twice daily until the porewater ammonia concentrations reached the level appropriate for the particular amphipod. Porewater ammonia measurements were made on "dummy" containers that were set up and maintained in the same manner as the actual test containers but without animals added to them. The porewater was obtained by siphoning off the overlying water in the dummy jar and centrifuging the sediment in a Teflon jar for at least 20 min at approximately 3000 rpm. Salinity, temperature, and pH were also determined in the porewater samples.




FIGURE 2.1. Testing Containers for Amphipod Static Renewal Toxicity Tests

The amphipod benthic toxicity tests were initiated by the addition of 20 organisms to each test chamber for a test population of 100 amphipods per sediment treatment. Amphipods were gently sieved from their native sediment in holding tanks and transferred to shallow baking dishes. For each test chamber, five animals were counted and transferred by pipet into each of four small, plastic cups. The organisms in each transfer cup were recounted by a second analyst and then placed in the test chamber by dipping the cup below the water surface of release the amphipods.

Salinity, temperature, DO, and pH were measured in all replicates prior to test initiation, in at least one replicate per treatment daily, and in all replicates at test termination. Measurements of total ammonia levels in the overlying water and porewater also continued during testing. Overlying water ammonia was measured in all replicates prior to test initiation (Day 0), in at least one replicate per treatment daily, and in all replicates at test termination (Day 10). Porewater ammonia was measured on Day 0 and Day 10. Flow rates to each test chamber were calibrated once at the start on the renewal process. The water-system was turned on for 15 min twice a day. Test chambers were renewed for 9 days before testing and continued daily throughout the 10-day test. The following were the acceptable ranges for water quality parameters during the amphipod tests:

	<u><i>A. abdita</i></u>	<u><i>R. abronius</i></u>
Temperature	20°C±2°C	14°C±2°C
DO	>60% saturation	>60% saturation
pH	7.8±0.5	7.8±0.5
Salinity	30‰±2‰	30‰±2‰
Ammonia	<30 mg/L	<30 mg/L
Renewal Rate	2 exchanges/day	2 exchanges/day.

Gentle aeration was provided throughout the test, and the amphipods were not fed during testing. At the end of the 10-day period, the contents of each chamber were gently sieved through 0.5-mm mesh, and the number of live, dead, and missing amphipods was recorded on termination forms. An animal was considered dead if it did not respond to gentle probing. As a quality control check, a second observer confirmed surviving organisms on at least 10% of the termination counts.

Reference toxicant tests with cadmium chloride were performed concurrently with each species. The reference toxicant tests were 96-h, water-only exposures that were otherwise conducted following the same procedures as for the static tests with sediment. *A. abdita* were exposed to nominal concentrations of 0, 0.25, 0.5, 1, and 2 mg/L cadmium. *R. abronius* were exposed to nominal concentrations of 0, 0.5, 1, 2, and 4 mg/L cadmium.

2.5.2.2 Static Test with *Mysidopsis bahia*

Upon receipt at the laboratory, *M. bahia* were placed in 10-gal aquaria and acclimated from 28‰ seawater to 30‰ with Sequim Bay seawater over a 24-h period. Mysids were received

and held for 4 days at $20^{\circ}\text{C}\pm 2^{\circ}\text{C}$ until testing and were fed concentrated brine shrimp nauplii twice daily prior to testing. Mortality of the *M. bahia* during holding was less than 1%.

The 10-day static benthic acute toxicity test with *M. bahia* was performed in 1-L glass jars. To prepare each test container, 200 mL of clean seawater was placed in each jar. Sediment was added until water was displaced up to the 400-mL mark, then seawater was added up to the 750-mL mark. Five replicates of each Port Chester composite, Mud Dump Reference Site sediment, Central Long Island Sound Reference Site, and control sediment (Sequim Bay sediment) were tested. Static jars were renewed twice daily for 8 days. At the start of the test the overlying water ammonia concentrations were all less than 14.5 mg/L.

The mysid benthic toxicity test was initiated by the addition of 20 organisms to each test chamber for a test population of 100 mysids per sediment treatment. Mysids were transferred from holding tanks to shallow glass dishes. For each test chamber, five animals were counted and transferred by pipet into each of four small, plastic cups. The animals in each transfer cup were recounted by a second analyst. The animals were placed in the test chamber by dipping the cup below the surface of the water to release the mysids.

Salinity, temperature, DO, pH, and total ammonia in overlying water were measured in all replicates prior to test initiation, in at least one replicate per treatment daily, and in all replicates at test termination. The total ammonia concentrations in the overlying water were <20 mg/L in each test chamber. The following were the acceptable ranges for water quality parameters during the *M. bahia* benthic test:

Temperature	$20^{\circ}\text{C}\pm 2^{\circ}\text{C}$
DO	$>40\%$ saturation
pH	7.8 ± 0.5
Salinity	$30\text{‰}\pm 2\text{‰}$.

Gentle aeration was provided to all test chambers during the test to maintain consistency in handling DO concentration among test containers. Animals were fed 1-2 mL of brine shrimp nauplii (<24 -h old) in suspension daily. At the end of the 10-day period, the contents of each chamber were gently sieved through 0.5-mm mesh, and the number of live and dead or missing mysids was recorded on termination forms. An animal was considered dead if it did not respond to gentle prodding. As a quality control check, a second observer confirmed surviving test organisms on at least 10% of the termination counts.

A 96-h, water-only, reference toxicant test was performed concurrently with the benthic test to establish the health and expected response of the test organisms. Water quality conditions were the same as for the SPP test, and animals were fed daily over the 96-h exposure. *M. bahia* were exposed to a seawater control plus four concentrations of copper sulfate: 50, 100, 150, and 200 $\mu\text{g/L}$ copper, using three replicates of each concentration.

2.5.3 Bioaccumulation Testing

The bivalve *M. nasuta* and the polychaete *N. virens* were used to evaluate the potential bioaccumulation of contaminants from dredged material. The bioaccumulation tests were 28-day flow-through exposures to sediment followed by a 24-h depuration period that allowed the organisms to void their digestive tracts of sediment. *M. nasuta* and *N. virens* were tested in separate 10-gal flow-through aquaria. Animals were exposed to five replicates of COMP PC, Mud Dump Reference Site sediment, Central Long Island Reference Site sediment, and native control sediment. Each chamber contained 25 *M. nasuta* or 20 *N. virens*. Water quality parameters (temperature, DO, pH, and salinity) were measured in all replicates at test initiation, in at least one replicate per treatment daily, and in all replicates at test termination. Flow rates were measured daily in all chambers.

Upon receipt at the laboratory, *M. nasuta* were received damp and held in control sediment with flowing Sequim Bay seawater at $15^{\circ}\text{C}\pm 2^{\circ}\text{C}$ until testing and were not fed. *N. virens* were placed in holding trays of control sediment with heated Sequim Bay seawater flowing into the trays. *N. virens* were received at 17°C and gradually acclimated to $20^{\circ}\text{C}\pm 2^{\circ}\text{C}$. *N. virens* were not fed prior to testing. Mortality of *M. nasuta* and *N. virens* during holding was less than 1%.

The Regional Guidance Manual provides an acceptable temperature range of $13^{\circ}\text{C}\pm 1^{\circ}\text{C}$ for *M. nasuta*; however, laboratory logistics required that *M. nasuta* share a 15°C flow-through water supply with *R. abronius*. This alteration of test temperature was not expected to affect the outcome of the test; bioaccumulation tests with *M. nasuta* have been conducted at $15^{\circ}\text{C}\pm 2^{\circ}\text{C}$ successfully. After discussion with the USACE-NYD project manager, the following ranges for water quality parameters were established as acceptable for the *M. nasuta* and *N. virens* tests:

	<u><i>M. nasuta</i></u>	<u><i>N. virens</i></u>
Temperature	$14^{\circ}\text{C}\pm 2^{\circ}\text{C}$	$20^{\circ}\text{C}\pm 2^{\circ}\text{C}$
DO	> 60% saturation	> 60% saturation
pH	7.8 ± 0.5	7.8 ± 0.5
Salinity	$30\text{‰}\pm 2\text{‰}$	$30\text{‰}\pm 2\text{‰}$
Flow Rate	125 ± 10 mL/min	125 ± 10 mL/min.

Aeration was provided to all test chambers to maintain consistency in handling DO concentrations among test chambers. Water quality, organism behavior (e.g., burrowing activity, feeding) and organism mortality were recorded daily. Dead organisms were removed daily and at the end of the 28-day testing period, *M. nasuta* and *N. virens* were placed in clean, flowing seawater for 24 h, after which the tissues were transferred into the appropriate chemistry jars for metals, and organic compound analyses. All tissue samples were frozen immediately and stored at less than -20°C until analysis.

Water-only reference toxicant tests (96-h) were also performed using copper sulfate in six geometrically increasing concentrations. The exposures were conducted using a test volume of 5 L in static 9.5-L (2.5-gal) aquaria. Three replicates of each concentration were tested, each containing 10 organisms. Water quality parameters were monitored at the same frequency and maintained within the same limits as the 28-day test, except that there were no flow rates. The *M. nasuta* reference toxicant test was conducted with treatments of 0, 0.25, 0.50, 0.75, 1.0, 1.5 and 2.5 mg/L copper; the *N. virens* test was conducted with treatments of 0, 0.05, 0.075, 0.15, 0.20, 0.25, and 0.30 mg/L copper.

2.6 Data Analysis and Interpretation Procedures

Statistical analyses were conducted to determine the magnitude and significance of toxicity and bioaccumulation in test treatments relative to the reference treatment. Each statistical test was based on a completely random design that allowed unbiased comparison between treatments.

2.6.1 Randomization

All water-column and benthic toxicity tests were designed as completely random tests. Organisms were randomly allocated to treatments, and treatments were randomly positioned on water tables. To determine randomization, a random-number table was generated for each test using the discrete random-number generator in Microsoft *Excel* spreadsheet software.

2.6.2 Statistical Analysis of Water-Column Tests

Two statistical analyses are presented in the Green Book for the interpretation of SPP (water-column) tests. The first is a one-sided t-test between survival in control test replicates and survival in the 100% SPP test replicates. A significant difference indicates acute toxicity in the 100% SPP treatment. This analyses was performed only when survival in the 100% SPP is less than the control (0% SPP) survival, and when control survival is >90% for nonlarval tests and >70% for larval tests. Prior to conducting the t-test, angular transformation (arcsine of the square root) of the proportion surviving in test replicates was performed to reduce possible heterogeneity of variance between mean survival of test organisms in the control and in the 100% SPP. The second analysis required by the Green Book is estimation of the medium lethal concentration (LC_{50}) or median effective concentration (EC_{50}). The LC_{50} or EC_{50} values for these tests were estimated using the trimmed Spearman-Kärber method (Finney 1971) and are expressed in percentage of SPP. The Spearman-Kärber estimator is appropriate only if there was increasing mortality (or effect) with increasing concentration, and if $\geq 50\%$ mortality (or effect) was observed in at least one test concentration when normalized to control survival. If 50%

mortality (or effect) did not occur in the 100% SPP concentrations for any treatments, then LC₅₀ or EC₅₀ values were reported as >100% SPP.

2.6.3 Statistical Analysis of Benthic Toxicity Tests

Benthic toxicity of all sediment treatments was compared to a single reference treatment using Dunnett's test (Dunnett 1964). The arcsine square root of the proportion of organisms surviving the test was used to stabilize the within-class variances. As recommended by the Green Book an experiment-wise error $\alpha=0.05$ was used. Acute toxicity for the amphipod test indicates that the test treatment was statistically significant relative to the reference treatment and had a greater than 20% difference in survival from the reference treatment. Acute toxicity for the mysid test indicates that the test treatment was statistically significant relative to the reference treatment and had a greater than 10% difference in survival from the reference treatment.

2.6.4 Statistical Analysis of Bioaccumulation

The results of the chemical analyses of test organism tissues exposed to the dredged sediment treatments was statistically compared with those of tissues similarly exposed to the Mud Dump Reference Site treatment using Dunnett's test with an experiment-wise error of $\alpha=0.05$. The Dunnett's tests determined whether or not the concentrations of contaminants of concern in the organisms exposed to the dredged sediments statistically exceeded those of organisms exposed to the reference sediment.

Statistical analyses were performed on the dry weight concentrations. When a compound (metals, pesticides, PCBs, and PAHs) was undetected (indicated by a "Q" flag in the report tables and a "U" flag in the appendix tables), one-half the detection limit of a compound was used in numerical calculations. If the compound was undetected in all five replicates of a test treatment, or if the mean concentration of a compound was greater in tissue samples from the reference treatment than in tissue samples from the test treatments, no further analysis was necessary. If a compound was undetected in all five replicates of the reference treatment, a one-sided, one-sample t-test ($\alpha=0.05$) was used to determine if the tissue concentrations from organisms exposed to dredged sediment treatments were statistically greater than the mean detection limit for that compound from the reference tissue. Results of background and control tissues were not statistically compared with the reference.

Magnification factors were calculated for each compound as the dry weight ratio of the mean tissue concentration from organisms exposed to dredged sediment treatments to the mean tissue concentration from organisms exposed to the Mud Dump Reference Site sediment. Whole detection limits were used for non-detects in this calculation.

2.7 Quality Assurance/Quality Control Procedures

The quality assurance/quality control (QA/QC) procedures for the Port Chester project were consistent with the Regional Guidance Manual and the Green Book, and were documented in the Work/Quality Assurance Project Plan, *Evaluation of Dredged Material Proposed for Ocean Disposal from Federal Projects in New York*, prepared by the MSL and submitted to the USACE-NYD for this program. This document describes all QA/QC procedures that were followed for sample collection, sample tracking and storage, and physical/chemical analyses. A member of Pacific Northwest National Laboratory's quality engineering staff was present throughout all phases of this program to observe procedures, review and audit data, and ensure that accepted protocols were followed. Data accumulation notebooks were assigned to each portion of these studies and served as records of day-to-day project activities.

3.0 Results

This section presents results of sample collection and processing, and physical and chemical analyses conducted on sediment samples collected from the proposed Port Chester dredging area.

3.1 Sample Collection and Processing

Sediment core and grab samples were collected from the Port Chester project area on March 8, 1994. The western shore of the project area is in New York and the eastern shore is in Connecticut. The sediment collected on the eastern shore was generally muddy and contained leaves and twigs. The sediment collected on the western shore was difficult to collect with the vibracore sampler. Several core cutters were damaged trying to collect sediment from this area. A van Veen grab was used to collect sediment representative of the western shore and was approved by the USACE project manager. Sediment collected with the van Veen grab was characterized by gravel and stones. This material was also present on the shore at the commercial facilities; it could have fallen in the water during loading onto barges, or fallen off barges that may not have been properly secured in transit. During sampling, oil was noted in the sediment samples, and two samples taken from PC-4 and PC-6 smelled of pesticides. The field crew also noted the presence of an oil sheen on the water surface at the time that core or grab samples were collected. The presence of oil was reported to state agencies and the Coast Guard. A site water sample was collected from PC-10.

Table 3.1 lists each sampling station within the Port Chester project area, sampling coordinates, collection date, length of core required for testing, and length of core actually collected. All core and grab samples were collected aboard the *Gelberman* for the 11 Port Chester samples. Seven core samples were collected to a project depth of -10 ft MLW and overdepths ranging from 0.3 ft to 4 ft. Four stations were collected with a van Veen grab sampler. Two of these stations were collected to project depth and varying overdepth lengths whereas the two remaining stations were not.

Upon receipt of the sediment samples at the MSL on March 18, 1994, samples were prepared for the physical and chemical analyses according to the procedures described in Section 2. Individual sediment samples were analyzed for grain size, moisture content, and TOC. One composited sediment sample representing the entire Port Chester project area (COMP PC) was analyzed for bulk density, specific gravity, metals, chlorinated pesticides, PCBs, PAHs, and 1,4-dichlorobenzene. Prior to sample analysis, the USACE-NED and State

of Connecticut requested the USACE-NYD to analyze the Port Chester sediment samples according to a different compositing scheme. Table 3.2 lists the stations or samples included in each of the five USACE-NED chemistry composites (CT COMP). The CT COMP samples were also analyzed for an extended list of metals, chlorinated pesticides, PCBs, PAHs, and 1,4-dichlorobenzene.

3.2 Physical and Chemical Analyses

3.2.1 Sediment Core Sample Description

Table 3.3 lists physical characteristics of each sediment core sample. Port Chester sediment samples were generally black or gray-black, silty-clayey material.

TABLE 3.1. Summary of Sediment Sample Data for Port Chester

Station	Collection Date	Station Coordinates		Core Length Required (ft)	Core Length Collected (ft)	Mudline Depth (-MLW ft)
		Latitude N	Longitude W			
Core Samples						
PC-1	3/8/94	40° 59.93' N	73° 39.59' W	3.5	1.8	8.5
PC-2	3/8/94	40° 59.91' N	73° 39.58' W	3.3	1.5	8.7
PC-5	3/8/94	40° 59.84' N	73° 39.56' W	3.0	2.75	9.0
PC-8	3/8/94	40° 59.64' N	73° 39.59' W	1.0	1.0	11.0
PC-9	3/8/94	40° 59.62' N	73° 39.58' W	3.3	2.6	8.7
PC-10(a)	3/8/94	40° 59.65' N	73° 39.53' W	4.5	5.0	7.5
PC-11	3/8/94	40° 59.51' N	73° 39.45' W	4.0	6.0	8.0
Grab Samples						
PC-3	3/8/94	40° 59.08' N	73° 35.58' W	3.3	1.25	8.7
PC-4	3/8/94	40° 59.81' N	73° 39.58' W	2.9	1.7	9.1
PC-6	3/8/94	40° 59.79' N	73° 39.58' W	2.5	1.8	9.5
PC-7	3/8/94	40° 59.77' N	73° 39.59' W	5.0	1.75	7.0
MDRS(b)	3/13/94	40° 20.19' N	73° 52.20' W	---(c)	---	67
MDRS	3/13/94	40° 20.21' N	73° 52.19' W	---	---	65
MDRS	3/13/94	40° 20.22' N	73° 52.19' W	---	---	66
MDRS	3/13/94	40° 20.22' N	73° 52.19' W	---	---	66
MDRS	3/13/94	40° 20.21' N	73° 52.23' W	---	---	65
MDRS	3/13/94	40° 20.21' N	73° 52.23' W	---	---	64
MDRS	3/13/94	40° 20.22' N	73° 52.23' W	---	---	66
MDRS	3/13/94	40° 20.21' N	73° 52.24' W	---	---	66
MDRS	3/13/94	NR(d)	NR	---	---	66
MDRS	3/13/94	NR	NR	---	---	66
MDRS	3/13/94	NR	NR	---	---	NR
MDRS	3/13/94	NR	NR	---	---	NR
CLIS(e)	3/7/94	NR	NR	---	---	NR

(a) Site water samples collected at this station.

(b) MDRS Mud Dump Reference Site.

(c) --- Not applicable.

(d) NR Data not recorded during sample collection.

(e) CLIS Central Long Island Sound Reference Site.

TABLE 3.2. Sediment Compositing Scheme for USACE-NED Chemistry Composites

Station	CT Comp
PC-1 PC-2 PC-3	CT COMP PC-I
PC-4 PC-6	CT COMP PC-II
PC-5 PC-7	CT COMP PC-III
PC-8 PC-9 PC-10	CT COMP PC-IV
PC-11	CT COMP PC-V

3.2.2 Grain Size, Percentage of Moisture, Bulk Density, and Specific Gravity

Table 3.4 shows the results of the analysis of Port Chester sediment samples for grain size percentage of moisture, and TOC. A quality control sample summary and associated quality control data for grain size and TOC measurements are provided in Appendix A. The upriver Port Chester samples PC-1, PC-2, PC-3, PC-4, PC-6 and PC-7, sediments were predominantly coarse-grained. Port Chester samples PC-5, PC-8, PC-9, PC-10, and PC-11, were predominately fine-grained: percentages of sand ranged from 21% to 34%; silt ranged from 34% to 48%; and clay ranged from 20% to 33%. The Mud Dump Reference Site sediment was composed of 98% sand and the Central Long Island Sound Reference Site sediment was 60% silt and 34% clay. The total percentage of solids ranged from 30% to 96%. Bulk density was analyzed on COMP PC. The results were reported in wet weight and dry weight with values of 100 lb/cu ft and 58 lb/cu ft respectively. The specific gravity value for COMP PC was 2.49.

3.2.3 Total Organic Carbon

The upriver Port Chester sediments PC-1 through PC-4 had $\leq 1.51\%$ TOC (Table 3.4) in Stations PC-5 through PC-11 had percentages of TOC ranging from 5.34% to 6.99%, which are higher than both that of the Mud Dump Reference sediment (0.01%) and of the Central Long Island Sound Reference sediment (1.64%).

TABLE 3.3. Port Chester Sediment Core Descriptions

Station	Depth Below Mudline (-ft MLW)		Project Depth(a)	Description of Observations
	Core Top	Core Bottom		
PC-1	8.5	10.3	12.0	Black sand and gravel at top of core; remaining core black silty sand.
PC-2	8.7	10.2	12.0	Black, slippery, silty material at top of core; black sandy gravel in middle section of core; remaining core had a layer of gray silty clay followed by a layer of slippery, black silty/clayey material.
PC-3	8.7	9.95	12.0	All black pea-size gravel.
PC-4	9.1	10.8	12.0	All black gravel and stone.
PC-5	9.0	11.75	12.0	All uniform black silty material.
PC-6	9.5	11.3	12.0	All black gravel bound together with black sand. Some rocks, mussels, and shells interspersed through the core.
PC-7	7.0	8.75	12.0	All rocks, leaves, shells bound together with black silty material.
PC-8	11.0	12.0	12.0	All uniform black silty material; high water content, almost liquid.
PC-9	8.7	11.3	12.0	All uniform black silty material.
PC-10	7.5	12.0	12.0	All uniform slippery, black, silty/clayey material with a band at -8.5 ft to -10.0 ft MLW where rocks and shells were mixed in with black material.
PC-11	8.0	4.0	12.0	Few rocks and mussel shells on top surface of core. From mudline to -11 ft MLW all uniform slippery, black, silty/clayey material; remaining of core was a band of grayish silty clay followed by band of black slippery material.

(a) Project depth of 10 ft plus 2 ft overdepth.

TABLE 3.4. Results of Analysis of Port Chester Sediment Samples for Grain Size and Percentage of Moisture

Sediment Treatment	Total Percent (dry weight)				Percent Moisture	Percent Total Organic Carbon
	Gravel >2000 μm	Sand 62.4-2000 μm	Silt 3.9-62.4 μm	Clay <3.9 μm		
PC-1	14	82	2	2	18	0.23
PC-2	17	62	18	3	20	1.09
PC-3	93	1	5	1	26	0.24
PC-4	99	0	1	0	4	1.51
PC-5	1	34	42	23	63	5.34
PC-6	55	37	5	3	20	ND(a)
PC-7	80	11	5	4	33	6.78
PC-8	2	22	48	28	70	6.07
PC-9	0	21	47	32	69	6.35
PC-10	13	33	34	20	54	6.18
PC-11	1	22	44	33	67	6.99
Mud Dump Reference	1	98	0	1	16	0.01
Central Long Island Sound	0	6	60	34	52	1.64

(a) ND No data.

3.2.4 Metals

Table 3.5 shows the results of the analysis of COMP PC, the five CT COMPs, the Mud Dump Reference Site, and the Central Long Island Sound Reference Site sediment samples for metals. The initial analysis of metals in sediment resulted in low spike recoveries for Ag. It was determined that sodium chloride in marine sediments caused an interference during ICP-MS analysis. The sediment digestion was repeated on the Port Chester samples with the addition of aqua regia to the procedure. This step alleviated the sodium chloride interference problem and improved spike recovery. The results of the Ag reanalysis are presented in Table 3.5. A quality control summary and quality control data associated with the metals, including the reanalysis of sediment samples for Ag, are provided in Appendix A.

Levels of all nine metals in COMP PC exceeded those found in the Mud Dump Reference Site sediment. Concentrations of Ag, As, Cr, Cu, Ni, Pb, and Zn were an order of magnitude higher in COMP PC than in the reference sediment. Mercury levels were two orders of magnitude greater in COMP PC than in the reference site sediment; Cd levels were eight orders of magnitude greater in COMP PC than in the reference site sediment.

Concentrations of eight metals in CT COMPs PC-III, PC-IV, and PC-V exceeded those found in the Central Long Island Sound Reference Site sediment. Levels of seven metals in CT COMPs PC-I and PC-II were lower than those found in the Central Long Island Sound Reference Site sediment.

TABLE 3.5. Results of Analysis of Port Chester Sediment Samples for Metals

Sediment Treatment	Metals (mg/kg dry weight)												
	Ag ^(a)	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn	Be ^(b)	Sb ^(b)	Se ^(b)	Tl ^(b)
COMP PC	3.45	8.58	72.8	90.1	183	1.29	65.9	307	431	NA	NA	NA	NA
CT COMP PC-I	0.342	3.44	1.47	16.7	25.2	0.099	12.3	38.9	58.6	0.291 J	0.263	0.21	0.071
CT COMP PC-II	0.262	2.40	0.748	10.8	127	0.067	11.8	24.6	75.6	0.236 J	0.196	0.21	0.094
CT COMP PC-III	2.54	10.4	19.8	90.0	158	0.170	42.9	292	386	0.643	1.31	1.03	0.415
CT COMP PC-IV	5.25	12.3	45.8	148	250	2.20	88.5	404	633	0.913	2.33	0.91	0.572
CT COMP PC-V	9.06	16.3	227	188	334	4.79	189	413	967	1.14	2.84	0.90	0.582
Mud Dump Reference ^(c)	0.062	5.64	0.085	10.0	1.90	0.006	3.10	6.50	14.1	NA	NA	NA	NA
Central Long Island Sound ^(d)	0.689	7.01	0.523	58.3	46.0	0.202	27.2	43.0	116	NA	NA	NA	NA

(a) Ag analyzed by Aqua Regia.

(b) Selected metals were analyzed only for the USACE-NED Composites.

(c) MDRS - Mud Dump Reference Site.

(d) CLIS - Central Long Island Sound Reference Site.

3.2.5 Chlorinated Pesticides

Table 3.6 shows the results of the analysis of COMP PC, the five CT COMPs, the Mud Dump Reference Site, and the Central Long Island Sound Reference Site sediments for chlorinated pesticides. A quality control sample summary and associated quality control data are provided in Appendix A.

The COMP PC sediment contained concentrations of pesticides at concentrations elevated over those found in the Mud Dump Reference site sediment. The dominant pesticides found in COMP PC were the DDT family of compounds (167 $\mu\text{g}/\text{kg}$ total DDTs), followed by dieldrin, α -chlordane, and *trans*-nonachlor. Endosulfan I and 2,4-DDE coeluted in the primary GC analysis of these samples, but examination of the confirmatory analysis using a second GC column revealed that neither compound was detected. The value shown is the detection limit for 2,4-DDE. Pesticides were undetected or detected at concentrations near or below the target detection limit (1.0 $\mu\text{g}/\text{kg}$) in sediment from the Mud Dump Reference Site.

Concentrations of pesticides were generally higher in the sediments from CT COMPs PC-III, PC-IV, and PC-V relative to those found in the Central Long Island Sound Reference Site sediment. The dominant pesticides found in the five CT COMPs were generally the same as found in COMP PC sediment. Pesticides were either undetected or detected at concentrations near or below the target detection limit (1.0 $\mu\text{g}/\text{kg}$) in sediment from the Central Long Island Sound Reference Site.

3.2.6 PCBs

Table 3.7 shows the results of the analysis of COMP PC, the five CT COMPs, the Mud Dump Reference Site, and the Central Long Island Sound Reference Site sediment for PCBs. A quality control sample summary and associated quality control data are provided in Appendix A.

All of the 22 PCB congeners analyzed were detected in COMP PC sediment, with only two congeners (PCB 8 and PCB 18) found at a concentration below the detection limit. The total PCB concentration calculated for COMP PC was 1060 $\mu\text{g}/\text{kg}$, about two orders of magnitude higher than in reference site sediment. PCBs were either undetected or detected at concentrations near or below the target detection limit (1.0 $\mu\text{g}/\text{kg}$) in Mud Dump Reference Site sediment.

Concentrations of the 22 PCB congeners in CT COMPs PC-I and PC-II were generally undetected or detected at concentrations near or below the target detection limit. All of the 22 PCB congeners analyzed were detected in CT COMPs PC-III, PC-IV, and PC-V sediment and at concentrations higher than those in the Central Long Island Sound Reference

TABLE 3.6. Results of Analysis of Port Chester Sediment for Chlorinated Pesticides

Treatment	Concentration in µg/kg dry weight						Mud Dump Reference	Central Long Island Sound
	COMP PC	CT COMP PC-I	CT COMP PC-II	CT COMP PC-III	CT COMP PC-IV	CT COMP PC-V		
2,4'-DDD	21.2	1.41	0.91	9.60	30.5	24.9 D ^(a)	0.01 J ^(c)	1.14 U
2,4'-DDT	3.64	0.50 U	0.47 U	0.84 U	1.50 U	1.59 U	0.60 U	1.07 U
4,4'-DDD	118	3.92	1.26	35.1	122	2.72 U	0.06 J	0.57 J
4,4'-DDE	21.9	2.01	1.00 J	12.9	29.8	82.5 D	0.01 J	1.04 J
4,4'-DDT	1.21 J	0.37 J	0.31 J	3.60 J	3.30 J	9.08 U	3.45 U	0.53 J
Total DDT^(f)	167	8.62	4.34	62.7	188	116	2.91	4.65
Aldrin	0.95 U	0.48 U	0.45 U	0.80 U	1.44 U	1.53 U	0.58 U	1.02 U
α-Chlordane	18.7	1.65	0.59 J	11.9	33.4	7.94 D	0.01 J	1.49 U
Dieldrin	19.7	1.39	0.52 J	8.90	36.4	9.46 D	0.21 J	0.78 J
Endosulfan I /2,4-DDE ^(d)	2.59 U	1.31 U	1.23 U	2.21 U	3.94 U	4.18 U	1.59 U	2.80 U
Endosulfan II	1.93 U	0.98 U	0.84 J	2.81	2.94 U	21.4 D	0.05 J	2.09 U
Endosulfan Sulfate	1.83 U	0.48 J	0.87 U	3.45	16.1	25.4 D	1.12 U	1.98 U
Endrin ^(e)	NA	1.78 U	1.67 U	2.99 U	5.34 U	5.67 U	NA	3.80 U
Endrin Aldehyde ^(e)	NA	1.06 U	0.99 U	1.79 U	3.19 U	3.39 U	NA	2.27 U
Heptachlor	2.12 U	1.07 U	1.01 U	1.81 U	3.22 U	0.86 J,D	1.30 U	2.30 U
Heptachlor epoxide	1.18 U	0.23 J	0.06 J	1.00 U	1.97	1.90 U	0.72 U	1.27 U
α-BHC ^(e)	NA	0.66 U	0.62 U	1.11 U	1.99 U	0.22 J,D	NA	1.42 U
β-BHC ^(e)	NA	0.98 U	0.92 U	1.65 U	2.94 U	3.12 U	NA	2.09 U
δ-BHC ^(e)	NA	0.89 U	0.83 U	1.50 U	2.67 U	2.84 U	NA	1.90 U
Lindane ^(e)	NA	0.78 U	0.73 U	1.31 U	2.33 U	0.75 J,D	NA	0.89 J
Methoxychlor ^(e)	NA	1.11 U	1.04 U	1.87 U	3.34 U	3.55 U	NA	2.38 U
Toxaphene ^(e)	NA	33.7 U	31.6 U	56.7 U	101 U	107 U	NA	72.1 U
<i>trans</i> -Nonachlor	8.59	0.93 J	0.41 J	6.36	18.5	3.28 D	0.00 J	2.19 U

(a) D Dilution required; results in higher detection limit.

(b) U Undetected at or above given concentration.

(c) J Concentration estimated; analyte detected below method detection limit.

(d) Endosulfan I and 2,4'-DDE coelute.

(e) Analyte required only in samples designated for Central Long Island Disposal Testing Site.

(f) Sum of 2,4'-DDD, 2,4'-DDE, 2,4'-DDT, 4,4'-DDD, 4,4'-DDE, and 4,4'-DDT; one-half of the detection limit used in summation when analyte was undetected.

Site sediment. PCBs were either undetected or detected at concentrations near or below the target detection limit (1.0 µg/kg) in Central Long Island Sound Reference Site sediment.

3.2.7 PAHs and 1,4-Dichlorobenzene

Table 3.8 shows the results of the analysis of COMP PC, the five CT COMPs, the Mud Dump Reference Site, and the Central Long Island Sound Reference Site sediment for PAHs. A quality control sample summary and associated quality control data are provided in Appendix A.

TABLE 3.7. Results of Analysis of Port Chester Sediment for PCBs

Treatment	Concentration in $\mu\text{g}/\text{kg}$ dry weight						Mud Dump Reference	Central Long Island Sound
	COMP PC	CT COMP PC-I	CT COMP PC-II	CT COMP PC-III	CT COMP PC-IV	CT COMP PC-V		
PCB 8	1.15 J ^(a)	0.51 J	0.25 J	1.22 J	4.04 J	1.98 J,D ^(b)	2.91 U ^(c)	1.39 J
PCB 18	2.91 J	1.53 U	1.43 U	1.51 J	7.26	3.11 J,D	1.85 U	3.26 U
PCB 28	4.51	1.00 U	0.94 U	1.93	8.12	3.72 D	1.21 U	2.13 J
PCB 44	13.4	0.61 J	0.33 J	5.40	19.9	17.9 D	0.22 J	0.53 J
PCB 49	8.07	0.38 J	0.18 J	2.99	11.3	11.0 D	0.04 J	0.40 J
PCB 52	28.4	0.98	1.26	8.36	40.6	43.3 D	0.06 J	0.17 J
PCB 66	68.9	2.05	0.76	15.7	66.2	22.3 D	0.04 J	1.57 J
PCB 87	25.9	0.73	0.29 J	5.59	22.0	44.6 D	0.05 J	0.18 J
PCB 101	67.7	1.82	0.88	16.2	79.7	81.9 D	0.04 J	1.12
PCB 105	20.9	0.40	0.25	3.86	13.4	30.9 D	0.03 J	0.19 J
PCB 118	53.0	1.36	0.59 J	11.3	42.2	82.8 D	0.02 J	0.89 J
PCB 128	8.67	0.55 J	0.34 J	3.76	10.6	16.0 D	0.92 U	0.65 J
PCB 138	63.5	2.07	0.89	19.3	70.9	92.9	0.07 J	1.45
PCB 153	46.1	1.43 J	0.60 J	11.5	45.5	70.2 D	0.03 J	1.51 J
PCB 170	21.4	1.01	1.11	6.33	21.0	14.8 D	0.97 U	1.08 J
PCB 180	35.8	1.12	0.56	9.67	37.0	41.7 D	0.65 U	0.59 J
PCB 183	4.60	0.27 J	0.09 J	1.98	6.07	9.64 D	0.72 U	0.07 J
PCB 184	2.37	0.14 J	0.06 J	0.63 J	5.56	6.21	0.01 J	0.20 J
PCB 187	17.5	0.45 U	0.42 U	5.53	18.0	9.53 D	0.01 J	0.67 J
PCB 195	6.32	0.14 J	0.10 J	1.11 J	5.09	5.73 D	0.83 U	0.14 J
PCB 206	9.56	0.19 J	0.10 J	2.16	12.2	12.0 D	1.26 U	0.66 J
PCB 209	21.2	0.30 J	0.29 J	2.57	8.21	6.80 D	0.79 U	0.94 J
Total PCB^(d)	1060	35.1	20.7	277	1110	1260	13.4	36.3

(a) J Concentration estimated; analyte detected below method detection limit.

(b) D Dilution required; results in higher detection limit.

(c) U Undetected at or above given concentration.

(d) Total PCB = 2.0 (x), where x = sum of all PCB congeners detected; one-half of the detection limit used in summation when analyte was undetected.

All 16 PAHs analyzed were detected in COMP PC sediment. Low-molecular-weight PAH (LPAH) made up approximately 20% of the total PAH concentration, whereas high-molecular-weight PAH (HPAH) made up 80% of the total. The COMP PC PAH levels ranged from 1 to 3 orders of magnitude higher than those found in the reference site sediments. Concentrations of PAH compounds in Mud Dump Reference Site sediment were either undetected or detected at concentrations below the target detection limit (0.01 $\mu\text{g}/\text{kg}$). The COMP PC concentration of 1,4-dichlorobenzene was two orders of magnitude higher than that in the Mud Dump Reference Site sediment samples.

All 23 PAHs analyzed were detected in the five CT COMPs. CT COMPs PC-I, PC-II, PC-III, and PC-IV had concentrations of LPAH compounds that made up approximately 20% of the total PAH concentration and HPAH compounds that made up approximately 80% of the total PAH concentration. CT COMP PC-V had 43% LPAH compounds and 57% HPAH compounds. The concentrations of PAH compounds in the CT COMPs PC-I, PC-II, and PC-III were \leq six times higher than those found in the Central Long Island Sound Reference Site sediment. Concentrations of PAH compounds in Central Long Island Sound Reference Site sediment ranged from 214 $\mu\text{g}/\text{kg}$ (LPAH) to 1310 $\mu\text{g}/\text{kg}$ (HPAH). All five CT COMPs had detected concentrations of 1,4-dichlorobenzene that were higher than those found in the Central Long Island Sound Reference Site sediment.

In addition to the list of PAH analytes required for analysis in the Regional Guidance Manual, hydrocarbon fingerprint characterization of the Port Chester sediment composite was accomplished using a tiered analytical approach. This fingerprinting characterization was done because of the visible oil sheen present on the water and in the sediment at the time of collection. As the first tier analysis, COMP PC was evaluated by gas chromatography with flame ionization detection (FID) to determine total petroleum hydrocarbon levels and to obtain a qualitative chromatographic representation of potential product type(s) present in the samples. As a second tier analysis, individual petroleum-specific PAH were measured using GC/MS for more detailed information used for petroleum source identification.

High resolution GC/FID analysis is a useful means for measuring total petroleum hydrocarbon (TPH), for making preliminary determinations about product type, and for evaluating weathering state of products in environmental samples. Figure 3.1 is the GC chromatogram for COMP PC. Based on a comparison of this chromatogram with those of petroleum product standards, the GC trace suggests that the Port Chester sediment sample appears to be contaminated with a petroleum product in the lubricating oil/fuel oil range. The product in the sample appears to be heavily degraded.

GC/MS analysis of PAH can confirm product identification and source allocation, because the PAH compounds are substantially more resistant to weathering and degradation and therefore retain the characteristic chemical pattern of the product in the samples than the aliphatic compounds measured by GC/FID. In this analysis, samples are monitored for the parent (unsubstituted) PAH compounds as well as for their alkylated homologs. This extended list of PAH compounds and concentrations in $\mu\text{g}/\text{kg}$ dry weight is shown in Table 3.9. The distribution of parent PAH compounds and their alkyl homologs in COMP PC are presented graphically in Figure 3.2. This PAH distribution was matched against distribution graphs from Battelle's Standard Oil Library to identify the source of the PAH input to the

TABLE 3.8. Results of Analysis of Port Chester Sediment for PAHs and 1,4-Dichlorobenzene

Treatment	Concentration in ug/kg dry weight										Mud Dump Reference	Central Island	Long Sound
	COMP PC	CT PC-I	COMP PC-II	CT PC-III	COMP PC-IV	CT PC-V	COMP	CT	COMP	CT			
naphthalene	237	7.99	5.91 J ^(a)	55.8	320	326		1.13	J	21.6			
1-methylnaphthalene	NA ^(b)	11.9	3.72 J	28.0	94.3	264	NA			6.26 J			
2-methylnaphthalene	NA	18.8	5.15 J	63.3	189	530	NA			11.1 J			
biphenyl	60.9	3.85 J	1.55 J	15.9	68.6	104	6.94	U ^(c)		3.77 J			
2,6-dimethylnaphthalene	NA	15.1 J	2.32 J	89.0	237	472	NA			2.12 J			
acenaphthylene	144	10.6	5.62	57.3	252	123	6.61	U		29.0			
acenaphthene	816	5.67 J	13.5	52.3	1410	197	8.59	U		4.03 J			
fluorene	214	11.4	11.9	71.6	271	347	7.11	U		8.62 J			
phenanthrene	986	80.4	130.2	558	1330	1320	0.720	J		79.6			
anthracene	752	26.2	22.9	227	1280	514	6.96	U		29.2			
1-methylphenanthrene	NA	24.3	14.8	103	369	638	NA			18.8			
TOTAL LPAH^(d)	3,210	216	218	1,320	5,820	4,830	20.0			214			
fluoranthene	4260	181	256	1460	7340	1550	0.528	J		193			
pyrene	3220	169	215	1290	5400	1360	0.554	J		205			
benzo[a]anthracene	809	71.0	91.0	530	1320	422	0.621	J		82.8			
chrysene	996	97.3	120	648	1520	600	9.42	U		115			
benzo[b]fluoranthene	1170	113	139	839	1680	565	0.499	J		175			
benzo[k]fluoranthene	372	40.3	53.1	280	606	201	8.42	U		62.5			
benzo[e]pyrene	NA	56.5	68.6	439	846	302	NA			97.3			
benzo[a]pyrene	712	70.5	91.0	534	1040	327	6.58	U		120			
perylene	NA	20.9	22.4	147	297	186	NA			39.1			
indeno[1,2,3-cd]pyrene	143	53.8	66.8	478	231	106	5.68	U		100			
dibenzo[a,h]anthracene	180	13.0	16.5	119	293	138	5.77	U		21.1			
benzo[g,h,i]perylene	729	51.0	60.3	440	1170	565	4.77	U		94.2			
TOTAL HPAH^(d)	12,600	937	1,100	7,210	21,700	6,320	22.5			1,310			
TOTAL PAH^(d)	15,800	1,150	1,320	8,530	27,500	11,200	42.4			1,520			
1,4-Dichlorobenzene	76.0	3.92	2.48	19.1	127	44.7			0.79	U ^(b)	1.40	U	

- (a) J Concentration estimated; analyte detected below the method detection limit.
 (b) NA Not applicable; USACE-NYD sample not analyzed for extended list of PAHs.
 (c) U Undetected at or above given concentration.
 (d) One-half detection limit used in summation for undetected values.

sediment sample. The Standard Oil Library contains PAH distributions prepared from analyses of a number of control oil products; examples of typical PAH distributions for various oil products are shown in Figure 3.3.




Figure 3.1. GC/FID Trace Showing Total Petroleum Hydrocarbons in COMP PC

TABLE 3.9. Extended List of PAH Compounds Analyzed in COMP PC

PAH list	Concentration µg/kg dry weight
naphthalene	237
C1-naphthalenes	227
C2-naphthalenes	853
C3-naphthalenes	2410
C4-naphthalenes	3340
biphenyl	60.7
acenaphthylene	144
acenaphthene	816
fluorene	214
C1-fluorenes	544
C2-fluorenes	1460
C3-fluorenes	2490
anthracene	752
phenanthrene	986
C1-phenanthrenes/anthracenes	1670
C2-phenanthrenes/anthracenes	2870
C3-phenanthrenes/anthracenes	2780
C4-phenanthrenes/anthracenes	2030
dibenzothiophene	197
C1-dibenzothiophene	593
C2-dibenzothiophene	1230
C3-dibenzothiophene	1401
fluoranthene	4260
pyrene	3220
C1-fluoranthenes/pyrenes	2200
benzo[a]anthracene	810
chrysene	996
C1-chrysene	719
C2-chrysene	806
C3-chrysene	470
C4-chrysene	ND(a)
benzo[b]fluoranthene	1170
benzo[k]fluoranthene	372
benzo[a]pyrene	712
indeno[1,2,3-c,d]pyrene	143
dibenzo[a,h]anthracene	180
benzo[g,h,i]perylene	729

(a) ND Not detected above the method detection limit.




FIGURE 3.2. Distribution of Unsubstituted PAH Compounds and their Alkylated Homologs in COMP PC. (N: naphthalene; N1: C1-naphthalenes; N2: C2-naphthalenes; N3: C3-naphthalenes; N4: C4-naphthalenes; AY: acenaphthylene; AN: acenaphthene; F: fluorene; F1: C1-fluorenes; F2: C2-fluorenes; F3: C3-fluorenes; A: anthracene; P: phenanthrene; P1: C1-phenanthrenes/anthracenes; P2: C2-phenanthrenes/anthracenes; P3: C3-phenanthrenes/anthracenes; P4: C4-phenanthrenes/anthracenes; D: dibenzothiophene; D1: C1-dibenzothiophenes; D2: C2-dibenzothiophenes; D3: C3-dibenzothiophenes; FL: fluoranthene; PY: pyrene; FP1: C1-fluoranthenes/pyrenes; BA: benzo[a]anthracene; C: chrysene; C1: C1-chrysenes; C2: C2-chrysenes; C3: C3-chrysenes; C4: C4-chrysenes; BB: benzo[b]fluoranthene; BK: benzo[k]fluoranthene; BAP: benzo[a]pyrene; IP: indeno[1,2,3-cd]pyrene; DA: dibenzo[a,h,]anthracene; BP: benzo[g,h,i]perylene.)




FIGURE 3.3. Typical PAH Distributions for Oil Products

Qualitatively, the distribution of PAHs suggests that COMP PC contains a mixture of hydrocarbons. The PAH distribution between naphthalene and dibenzothiophene resembles that of marine diesel fuel or a lubricating oil where the alkylated homologs of parent PAHs increase in concentration with level of alkylation (C1 thru C4). The presence of 4- and 5-ring PAHs also suggest inputs of coal tar or creosote material to the Port Chester sediment.

3.3 Site Water and Elutriate Analyses

Metals, chlorinated pesticides, and PCBs were analyzed in dredging site water collected from Port Chester and in elutriate samples prepared from clean seawater (Sequim Bay) and the Port Chester sediment composite. Sequim Bay seawater was used in place of dredging site water to maintain consistency in salinity among the dredging projects. Mud Dump Site water and Sequim Bay control water were also analyzed. All water and elutriate samples were analyzed in triplicate. Mean results of the triplicate analyses are presented and discussed in the following sections. Complete results of site water and elutriate samples, and quality control summary and associated quality control data are provided in Appendix B.

3.3.1 Metals

Results of analysis of Sequim Bay control water, Mud Dump Site water, Port Chester site water, and Port Chester sediment elutriates are shown in Table 3.10. Concentrations of Cd, Cr, and Zn were similar between the control water and Mud Dump Site water, whereas concentrations of Ag, Cu, Hg, Ni, and Pb were at least twice as high in the Mud Dump Site water than in the control. In particular, Hg and Pb were about an order of magnitude higher in the Mud Dump Site than in the control water.

Port Chester Site water had elevated levels of all metals measured when compared with Mud Dump Site Water. Concentrations of Hg, Ni, and Zn were only slightly elevated, whereas concentrations of Ag, Cd, Cr, Cu, and Pb were at least twice as high in Port Chester site water than in Mud Dump Site water. Port Chester elutriate concentrations for metals were generally more similar to those found in the Mud Dump Site water than to those in the Port Chester site water. Concentrations of all metals, with the exception of Cd and Ni, were lower in the Port Chester elutriate than in the Port Chester site water.

3.3.2 Chlorinated Pesticides and PCBs

Results of analysis of Sequim Bay control water, Mud Dump Site water, Port Chester site water, and Port Chester elutriate are shown in Table 3.11. With few exceptions, pesticides and PCB congeners were not detected in any of the samples. Traces of 4,4'-DDD,

TABLE 3.10. Results of Analysis of Port Chester Site Water and Elutriate for Metals

Analyte	Concentration in $\mu\text{g/L}$ (a)			
	Control Water	Mud Dump Site Water	Port Chester Site Water	Port Chester Elutriate
Ag	0.0035 Q(b)	0.0223	0.0860	0.0200
Cd	0.0557	0.0603	0.340	0.530
Cr	0.180	0.27	1.79	0.727
Cu	0.471	2.06	8.28	1.63
Hg	0.0003	0.0096	0.0249	0.0227
Ni	0.469	1.27	2.39	3.54
Pb	0.0430	0.931	10.1	1.73
Zn	9.20	10.3	23.8	6.94

(a) Value shown is the mean of triplicate analyses; one-half the detection limit used when analyte was undetected.

(b) Q undetected at or above twice the given concentration.

4,4'-DDE, 4,4'-DDT, α -chlordane, and dieldrin were found in Port Chester site water samples above the detection limits for these compounds. Concentrations of pesticides in the Port Chester elutriate samples were elevated by a factor of 50 for 4,4'-DDD, and by a factor of 10 to 15 times for 4,4'-DDE, dieldrin and α -chlordane. Only PCB 28 was detected above the detection limit in Port Chester site water samples. Nineteen PCB congeners were detected in the Port Chester elutriate samples at concentrations ranging from 1.3 to 84.6 times above those found in the Port Chester site water and the control and Mud Dump site water.

3.4 Water-Column Toxicity Testing

Water-column tests were performed on four concentrations of an SPP preparation made from the Port Chester composite. SPP tests were conducted with the silverside, *M. beryllina*, the mysid, *M. bahia*, and larvae of the bivalve *M. galloprovincialis*. This section discusses the results of all water-column and reference toxicant testing. Complete test results, water quality measurements, and the results of the reference toxicant tests are presented in Appendix C. Throughout this section, the term "significant difference" is used to express *statistically* significant differences only. Tests for statistical significance between test and reference treatments were performed following methods outlined in Section 2.6.

TABLE 3.11. Results of Analysis of Port Chester Site Water and Elutriate for Chlorinated Pesticides and PCBs

Analyte	Concentration in ng/L ^(a)			
	Control Water	Mud Dump Site Water	Port Chester Site Water	Port Chester Elutriate
2,4'-DDD	0.39 Q ^(b)	0.38 Q	0.38 Q	0.40 Q
2,4'-DDT	0.40 Q	0.39 Q	0.39 Q	16.5
4,4'-DDD	0.57 Q	0.56 Q	1.85	8.09
4,4'-DDE	0.49 Q	0.47 Q	0.68	11.8
4,4'-DDT	0.49 Q	0.48 Q	1.03	55.3
Total DDT^(c)	2.76	2.69	4.74	91.2
Aldrin	0.36 Q	0.36 Q	0.36 Q	14.1
α-Chlordane	0.46 Q	0.45 Q	1.83	15.8
Dieldrin	0.48 Q	0.47 Q	1.64	0.54 Q
Endosulfan I/2,4-DDE	0.42 Q	0.41 Q	0.41 Q	5.03
Endosulfan II	5.51 Q	5.38 Q	4.11	0.46 Q
Endosulfan sulfate	4.03 Q	3.94 Q	3.94 Q	5.45
Heptachlor	1.02	0.32 Q	0.32 Q	14.3
Heptachlor epoxide	0.42 Q	0.41 Q	0.41 Q	0.35 Q
<i>trans</i> -Nonachlor	0.47 Q	0.46 Q	0.46 Q	0.46 Q
PCB 8	0.43 Q	0.42 Q	0.42 Q	0.47 Q
PCB 18	0.52 Q	0.51 Q	0.51 Q	0.58 Q
PCB 28	0.59 Q	0.57 Q	3.31	6.03
PCB 44	0.60 Q	0.59 Q	0.59 Q	15.5
PCB 49	0.51 Q	0.50 Q	0.50 Q	8.83
PCB 52	0.60 Q	0.59 Q	0.59 Q	31.1
PCB 66	0.47 Q	0.46 Q	0.46 Q	14.4
PCB 87	0.53 Q	0.51 Q	0.69	29.6
PCB 101	0.53 Q	0.52 Q	0.52 Q	88.0
PCB 105	0.63 Q	0.62 Q	0.62 Q	31.6
PCB 118	0.50 Q	0.49 Q	0.49 Q	59.5
PCB 128	0.56 Q	0.55 Q	0.55 Q	11.4
PCB 138	0.67 Q	0.66 Q	0.66	73.0
PCB 153	0.64 Q	0.63 Q	0.74	47.5
PCB 170	0.19	0.56 Q	0.56 Q	16.4
PCB 180	0.50 Q	0.49 Q	0.49 Q	33.5
PCB 183	0.52 Q	0.51 Q	0.51 Q	6.50
PCB 184	0.49	0.51 Q	0.51 Q	0.57 Q
PCB 187	0.49 Q	0.48 Q	0.48 Q	22.0
PCB 195	0.57 Q	0.55 Q	0.55 Q	3.93
PCB 206	0.55 Q	0.54 Q	0.54 Q	8.08
PCB 209	0.61 Q	0.60 Q	0.60 Q	1.54
Total PCB^(d)	23.4	23.7	29.8	1020

(a) Value shown is the mean of triplicate analyses; one-half the detection limit used when analyte was undetected.

(b) Q undetected at or above twice the given concentration.

(c) Total DDT is the sum of 4,4'-DDT, 4,4'-DDE, 4,4'-DDD, 2,4'-DDT, 2,4'-DDE, and 2,4'-DDD; one-half of the detection limit was used in summation when analyte was not detected.

(d) Total PCBs = 2(x), where x is the sum of all PCB congeners detected; one-half of the detection limit used in summation when an analyte was undetected.

3.4.1 *Menidia beryllina* Water-Column Toxicity Test

Results of the *M. beryllina* water-column toxicity test are summarized in Table 3.12. Complete test results as well as water quality data are presented in Appendix C, Tables C.1 through C.4. Control survival was 100%, validating this test. Survival in the 100% SPP preparation was 26% and was significantly lower than in the controls. The *M. beryllina* LC₅₀ of the Port Chester composite was 75.1% SPP.

Water quality parameters were within acceptable ranges throughout testing, except for one pH measurement above the acceptable range. Ammonia concentrations in the 100% SPP preparation reached 15.6 mg/L. The copper reference toxicant test produced an LC₅₀ of 98.1 µg/L Cu, which is within the control limits established at the MSL (71µg/L to 136 µg/L Cu).

3.4.2 *Mysidopsis bahia* Water-Column Toxicity Test

Results of the *M. bahia* water-column toxicity test are summarized in Table 3.12. Complete test results, as well as water quality data are presented in Appendix C, Tables C.5 through C.8. This test was validated by a control survival of 98%. Survival in the 100% SPP preparation was 58% and was significantly lower than the controls. The LC₅₀ was greater than 100% SPP.

All water quality parameters were within acceptable ranges throughout the test, with the exception of pH, which rose to 8.33 and 8.39 in the 50% and 100% treatments,

TABLE 3.12. Summary of Water-Column Toxicity Tests Performed with Port Chester Sediment

Test Organism	Percentage of Survival in 0% SPP	Percentage of Survival in 100% SPP	0% and 100% Significantly Different	LC ₅₀ (%SPP)
<i>Menidia beryllina</i>	100	26	Yes	75.1
<i>Mysidopsis bahia</i>	98	58	Yes	>100
<i>Mytilus galloprovincialis</i> (Survival)	99	84	Yes	>100
<i>Mytilus galloprovincialis</i> (Normal development)	99	0	Yes	53.7(a)

(a) Median effective concentration (EC₅₀) based on normal development to the D-cell, prodissoconch stage.

respectively. The ammonia concentration at test termination in the 100% SPP preparation was 15.6 mg/L. An LC₅₀ could not be calculated for the copper reference toxicant test, because there was not a greater than 50% reduction in mortality.

3.4.3 *Mytilus galloprovincialis* Water-Column Toxicity Test

Results of the *M. galloprovincialis* water-column toxicity test are summarized in Table 3.12. Complete test results and water quality data are presented in Appendix C, Tables C.9 through C.12. This test was validated by 99% survival and 99% normal development in the controls. Survival was 84% in the 100% SPP preparation and was significantly lower than the controls. The LC₅₀ was >100% SPP. Normal development, considered a more sensitive indicator of toxicity, was also significantly reduced in the 100% SPP, with 0% normal prodissoconch in this treatment. The EC₅₀ was 53.7% SPP.

All water quality parameters were within acceptable ranges throughout the test. The ammonia concentration in the 100% SPP preparation was 7.9 mg/L. The copper reference toxicant test revealed an LC₅₀ of 45.6 µg/L Cu and an EC₅₀ of 6.5 µg/L Cu, which were within the control limits (LC₅₀: 5.8 µg/L to 35 µg/L copper; EC₅₀: 5.7 µg/L to 21µg/L copper) established at the MSL.

3.5 Benthic Acute Toxicity Testing

Benthic acute toxicity tests were performed on the Port Chester composite, Mud Dump Reference Site, and Central Long Island Sound Reference Site sediment. Benthic toxicity tests were conducted with the amphipods *A. abdita* and *R. abronius*, and the mysid, *M. bahia*. This section discusses the results of all benthic and reference toxicant testing. Complete test results, water quality measurements, and the results of the reference toxicant tests are presented in Appendix D. Throughout this section the term "significant difference" is used to express *statistically* significant differences only. In addition to statistical significance, a sediment is considered acutely toxic if a greater than 20% difference in amphipod (10% for mysids) survival is demonstrated between a test and reference sediment. Tests for statistical significance between the treatment and reference treatment were performed following methods outlined in Section 2.6.

3.5.1 *Ampelisca abdita* Benthic Acute Toxicity Test

Results of the benthic acute toxicity test with *A. abdita* are summarized in Table 3.13. Complete test results and water quality data are presented in Appendix D, Tables D.1 through D.5. Prior to test setup, total ammonia concentrations measured in the Port Chester

TABLE 3.13. Summary of Benthic-Acute Toxicity Tests Performed with Port Chester Sediment

Test Organism	Mean % Survival	Significantly Different Than MDRS Reference	≥ 20% Difference (Amphipod) ≥ 10% Difference (Mysid)	Significantly Different Than CLIS Reference	≥ 20% Difference (Amphipod) ≥ 10% Difference (Mysid)
<i>A. abdita</i>	0%	Yes	Yes	Yes	Yes
<i>R. abronius</i>	78%	Yes	Yes	Yes	No
<i>M. bahia</i>	77%	No	No	No	No

bulk sediment composite was about 156 mg/L. Test chambers containing sediment and overlying water were set up (March 25, 1994) and maintained under test temperatures with aeration during the ammonia purging period. Overlying water was exchanged twice daily, delivered via a flow-through system (i.e., two times each day, the seawater flow into the test chambers was turned on long enough to displace the volume of the water in the test chamber once). Porewater ammonia was measured in "dummy" jars every few days until concentrations were 30 mg/L or less. The test was initiated after 10 days (April 4, 1994) when the porewater ammonia concentration was 10.7 mg/L.

Survival in the *A. abdita* control sediment was 97%, validating this test. Survival in the Port Chester composite was 0%, which was statistically significant and represented a > 20% reduction in survival compared with that in the Mud Dump (93%) and Central Long Island Sound (97%) reference sediments.

Temperature, pH, and DO were outside of acceptable water quality ranges in two test treatments during testing. However, test organism survival in the reference and control treatments were acceptable, thus these exceedences do not appear to have affected test interpretation. Ammonia concentrations were less than 1.0 mg/L in the overlying water during the 10-day test, and were 10.7 mg/L in the porewater at test termination. The cadmium reference toxicant test produced an LC₅₀ of 0.66 mg/L Cd, which was within the control limits established at the MSL (0.5 mg/L to 1.4 mg/L Cd).

3.5.2 *Rhepoxynius abronius* Benthic Acute Toxicity Test

Results of the benthic toxicity test with *R. abronius* are summarized in Table 3.13. Complete test results and water-quality data are presented in Appendix D, Tables D.6 through D.10. The same procedure that was followed to reduce the bulk sediment porewater

ammonia concentration from 156 mg/L to 30 mg/L or less in the *A. abdita* test was used in the *R. abronius* test. Test chambers containing sediment and overlying water were set up (March 25, 1994) and maintained under test temperatures with aeration during the ammonia purging period. Overlying water was exchanged twice daily. The test was initiated after 11 days (April 5, 1994) when the porewater ammonia concentration was 11.0 mg/L.

Survival in the West Beach control sediment was 97%, validating this test. Survival in the Port Chester composite was 78% which was statistically significant and represented a $\geq 20\%$ reduction in survival relative to that in both the Mud Dump (98% survival) and the Central Long Island Sound (91% survival) reference sediments.

All water quality parameters were within acceptable ranges throughout the test, with the exception of pH, which rose to 8.48 and 8.40 in COMP PC and C-WB sediments, respectively. Ammonia concentrations were less than 1.0 mg/L in the overlying water during the 10-day test, and were 11.0 mg/L in the porewater at test termination. The cadmium reference toxicant test produced an LC_{50} of 1.14 mg/L Cd, which was within the control limits established at the MSL (0.48 mg/L to 1.70 mg/L Cd).

3.5.3 *Mysidopsis bahia* Benthic Acute Toxicity Test

Results of the static benthic toxicity test with *M. bahia* are summarized in Table 3.13. Complete test results and water-quality data are presented in Appendix D, Tables D.11 through D.14. The mysid static test was not manipulated in any way to reduce porewater or overlying water ammonia concentrations prior to test initiation. Survival in the Port Chester composite was 77% and was not statistically significant, nor was the survival greater than 10% different from that in the Mud Dump Reference (76% survival) and the Central Long Island Sound Reference (74% survival) sediments. A control sediment treatment was not run concurrently with the Port Chester sediment treatments. However, the mysid benthic test was rerun 2 weeks later using other New York Federal Project-2 sediments. Mysid survival in the control treatment of that test was 93%.

All water quality parameters were within acceptable ranges throughout the test, with the exception of pH, which rose to 8.53 in COMP PC sediments. Ammonia concentrations were less than 1.0 mg/L in the overlying water during the 10-day static test; therefore, the *M. bahia* static renewal benthic acute toxicity test was not performed using Port Chester sediments. The copper reference toxicant test produced an LC_{50} of 151 $\mu\text{g/L}$ Cu, which is within the control limits of 116 $\mu\text{g/L}$ to 229 $\mu\text{g/L}$ Cu, established at the MSL.

3.6 Bioaccumulation Tests with *Macoma nasuta* and *Nereis virens*

Bioaccumulation tests with *M. nasuta* and *N. virens* were conducted using the Port Chester composite, Mud Dump Reference Site, Central Long Island Sound Reference Site, and control sediments. Both *M. nasuta* and *N. virens* were exposed for 28 days under flow-through conditions. Survival was greater than 90% in the *M. nasuta* control sediment, and 89% in the *N. virens* control sediment. Statistical analysis of the survival data was not conducted, since the purpose of the 28-day solid-phase test was to provide results as to the bioaccumulation potential in *M. nasuta* and *N. virens* tissues. Complete test results and water quality data are presented in Appendix E.

The tissues of the exposed organisms were analyzed for metals and selected organic contaminants (pesticides, PCBs, and PAHs), the results of which are summarized in this section. Complete test results and water quality data are tabulated in Appendix E for both species. Analytical results, including a quality control summary and associated quality control data, are presented in Appendix F for *M. nasuta* and in Appendix G for *N. virens*. The statistical analysis of tissue data was performed using sample dry weight concentrations to remove any variance associated with water content in each sample. Statistical difference between reference site and test sediment exposures is shown in the following tables with the results of sample analysis on a wet weight basis. Reporting data in this manner allows for comparison of wet weight concentrations obtained from this study with regulatory levels such as the FDA action levels reported in section 4.0 of this report. Lipids were analyzed on the background samples of the *M. nasuta* and *N. virens* tissues. These samples were triplicated and the average dry weight lipid content in wet weight of tissues for *M. nasuta* and *N. virens* were 0.59% and 2.10%, respectively. At the end of this section magnification tables are presented that show a comparison of Port Chester tissue concentrations with the reference tissue concentrations. Whole detection limit and dry weight of samples were used to create the magnification tables.

3.6.1 Bioaccumulation of Metals in *Macoma nasuta*

All nine metals were detected in tissues exposed to the Port Chester composite and to the Mud Dump and CLIS Reference sediments, except Pb, which was undetected in the tissues exposed to Mud Dump Reference sediment (Table 3.14). Of these, Cd, Cu, Ni, and Pd were statistically significant and elevated relative to tissues exposed to the Mud Dump Reference Site sediment, and Pb was at statistically significant and elevated relative to tissues exposed to the Central Long Island Sound Reference Site sediment. The magnification factor, the magnitude by which a contaminant concentration in the test composite

TABLE 3.14. Mean Concentrations of Metals in *M. nasuta* Tissues Exposed to Port Chester Sediment

Analyte	Concentration (mg/kg wet weight) (a)			SD(d)	SD
	MDRS(b)	CLIS(c)	COMP PC	MDRS	CLIS
Silver	0.0372	0.0294	0.0368	No	No
Arsenic	3.16	2.78	3.03	No	No
Cadmium	0.0355	0.0236	0.208	Yes	Yes
Chromium	0.408	0.451	0.436	No	No
Copper	1.78	2.31	2.49	Yes	No
Mercury	0.0180	0.0153	0.0148	No	No
Nickel	0.402	0.576	0.559	Yes	No
Lead	0.157 Q(e)	0.848	0.721	Yes	No
Zinc	13.1	11.2	14.0	No	No

(a) Value shown is a mean of five replicate s; one-half the detection limit used when analyte was undetected.

(b) MDRS- Mud Dump Reference Site.

(c) CLIS-Central Long Island Sound Reference Site.

(d) SD Significantly different.

(e) Undetected at or above twice the given concentration.

tissues exceeds that of the reference-exposed tissues, was below three for all metals, except Cd, which was 6.77 (MDRS) and 9.22 (CLIS).

3.6.2 Bioaccumulation of Pesticides in *Macoma nasuta*

Of the 15 pesticides analyzed, seven were detected in tissues exposed to the Port Chester composite (Table 3.15). With respect to both the Mud Dump Reference and Central Long Island Sound Reference Site tissues, aldrin, dieldrin, α -chlordane, *trans*-nonachlor, 2,4'-DDD, 4,4'-DDD, and 4,4'-DDE were statistically significant and elevated in the Port Chester composite tissues. Concentrations of α -chlordane, 2,4'-DDD, 4,4'-DDE, and 4,4'-DDD concentrations in Port Chester composite tissues exceeded those in tissues exposed to the Mud Dump Reference sediment by a factor of greater than 10. Concentrations of α -chlordane, 2,4'-DDD, and 4,4'-DDD concentrations exceeded those in the Central Long Island Reference tissues by a factor of greater than 10. The total concentration of DDT in the tissues exposed to the Port Chester composite was 21.9 μ g/kg.

3.6.3 Bioaccumulation of PCBs in *Macoma nasuta*

Eighteen of 22 PCBs analyzed were detected in *M. nasuta* tissues exposed to the Port Chester composite (Table 3.15). Twelve PCBs were observed at concentrations that were statistically significant and elevated relative to tissues exposed to the Mud Dump Reference composite. Nine PCBs were significantly elevated relative to the Central Long Island Sound Reference Site tissues. Port Chester tissue concentrations of six PCB congeners (87, 101, 105,

TABLE 3.15. Mean Concentrations of Pesticides and PCBs in *M. nasuta* Tissues Exposed to Port Chester Sediment

Analyte	Concentration ($\mu\text{g}/\text{kg}$ wet weight) ^(a)					SD ^(d)	SD
	MDRS ^(b)		CLIS ^(c)		COMP PC	MDRS	CLIS
2,4'-DDD	0.12	Q ^(e)	0.13	Q	5.05	Yes	Yes
2,4'-DDE	0.18		0.13	Q	0.13	No	No
2,4'-DDT	0.09	Q	0.09	Q	0.09	No	No
4,4'-DDD	0.13	Q	0.16		11.1	Yes	Yes
4,4'-DDE	0.34		1.27		5.41	Yes	Yes
4,4'-DDT	1.23		7.20		0.08	No	No
Total DDT ^(f)	2.09		8.98		21.9	Yes	Yes
α -Chlordane	0.05	Q	0.05	Q	3.71	Yes	Yes
Aldrin	0.35		0.06	Q	1.16	Yes	Yes
Dieldrin	0.26	Q	0.33		3.77	Yes	Yes
Endosulfan I	0.09	Q	0.09	Q	0.09	No	No
Endosulfan II	0.09	Q	0.09	Q	0.09	No	No
Endosulfan Sulfate	0.09	Q	0.09	Q	0.09	No	No
Heptachlor	0.09	O	0.09	O	0.30	No	No
Heptachlor Epoxide	0.06	Q	0.07	Q	0.07	No	No
<i>trans</i> -Nonachlor	0.07	Q	0.07	Q	0.67	Yes	Yes
PCB 8	0.87		0.20	Q	0.21	No	No
PCB 18	0.21	Q	0.21	Q	0.57	No	No
PCB 28	0.62		0.77		1.24	No	No
PCB 44	0.08	Q	0.15		0.70	Yes	Yes
PCB 49	0.17		0.62		1.68	Yes	No
PCB 52	0.81		0.74		5.08	Yes	Yes
PCB 66	0.18		0.96		0.05	No	No
PCB 87	0.16		0.19		3.09	Yes	Yes
PCB 101	0.45		0.97		6.92	Yes	Yes
PCB 105	0.09		0.10		1.79	Yes	Yes
PCB 118	0.17		0.41		4.26	Yes	Yes
PCB 128	0.07	Q	0.13		0.61	Yes	Yes
PCB 138	0.18		0.59		2.84	Yes	Yes
PCB 153	0.15		1.06		2.22	Yes	No
PCB 170	0.12		0.10		0.24	No	No
PCB 180	0.09	Q	0.26		0.66	Yes	Yes
PCB 183	0.12	Q	0.12	Q	0.22	No	No
PCB 184	0.12	Q	0.12	Q	0.12	No	No
PCB 187	0.06	Q	1.01		0.56	Yes	No
PCB 195	0.05	Q	0.05	Q	0.05	No	No
PCB 206	0.05	Q	0.06	Q	0.11	No	No
PCB 209	0.05	Q	0.05	Q	0.31	No	No
Total PCB ^(g)	9.74		17.1		67.1	Yes	Yes

(a) Value shown is a mean of five replicates; one-half the detection limit used when analyte was undetected.

(b) MDRS- Mud Dump Reference Site.

(c) CLIS-Central Long Island Sound Reference Site.

(d) SD Significantly different.

(e) Undetected at or above twice the given concentration.

(f) Total DDT is the sum of 4,4'-DDT, 4,4'-DDE, 4,4'-DDD, 2,4'-DDT, 2,4'-DDE, and 2,4'-DDD; one-half of the detection limit was used in summation when analyte was not detected.

(g) Total PCB = 2(x), where x=sum of all PCB congeners; one-half of the detection limit used in summation when analyte was undetected.

118, 138, and 153) exceeded those of the Mud Dump Reference tissues by a factor of greater than 10. Port Chester tissue concentrations of PCBs 87 and 105 exceeded those of the Central Long Island Sound Reference Site tissues by a factor of greater than 10. The concentration of total PCBs in the tissues exposed to the Port Chester composite was 67.1 µg/kg.

3.6.4 Bioaccumulation of PAHs and 1,4 Dichlorobenzene in *Macoma nasuta*

All PAHs analyzed were detected in *M. nasuta* tissues exposed to the Port Chester composites at statistically significant and elevated concentrations, relative to tissues exposed to both reference sediments (Table 3.16) except for acenaphthylene and benzo[k]fluoranthene. Nine PAHs were elevated above the Mud Dump Reference tissues by factors ranging from 12.7 times to 126 times. Acenaphthene, anthracene, fluoranthene, pyrene, benzo(a)anthracene, and chrysene were elevated above the Central Long Island Sound

TABLE 3.16. Mean Concentrations of PAHs in *M. nasuta* Tissues Exposed to Port Chester Sediment

Analyte	Concentration (µg/kg wet weight) (a)					SD	
	MDRS(b)	CLIS(c)	COMP PC	MDRS	CLIS	MDRS	CLIS
Naphthalene	1.12	0.93	Q(e)	3.71	Yes	Yes	
Acenaphthylene	0.36	Q	0.99	0.52	No	No	
Acenaphthene	0.64	Q	0.65	Q	17.7	Yes	
Fluorene	0.61	Q	0.62	Q	5.99	Yes	
Phenanthrene	1.26	Q	3.29	29.6	Yes	Yes	
Anthracene	1.10	Q	3.05	34.9	Yes	Yes	
Fluoranthene	2.64	Q	9.18	616	Yes	Yes	
Pyrene	2.25	Q	11.6	455	Yes	Yes	
Benzo[a]anthracene	2.36	Q	5.23	98.0	Yes	Yes	
Chrysene	1.12	Q	5.19	115	Yes	Yes	
Benzo[b]fluoranthene	3.37	Q	13.2	74.9	Yes	Yes	
Benzo[k]fluoranthene	1.83	Q	5.64	6.19	No	No	
Benzo[a]pyrene	1.21	Q	5.98	29.7	Yes	Yes	
Indeno[123-cd]pyrene	0.87	Q	4.38	10.7	Yes	Yes	
Dibenzo[a,h]anthracene	0.62	Q	0.76	3.31	Yes	Yes	
Benzo[g,h,i]perylene	0.99	Q	4.42	10.6	Yes	Yes	
1,4-Dichlorobenzene	0.92	Q	0.93	Q	0.96	No	

- (a) Value shown is a mean of five replicates; one-half the detection limit used when analyte was undetected.
- (b) MDRS- Mud Dump Reference Site.
- (c) CLIS-Central Long Island Sound Reference Site.
- (d) SD Significantly different.
- (e) Undetected at or above twice the given concentration.

Reference Site tissues by a factor of greater than 10. The compound 1,4-dichlorobenzene was undetected in all replicates of the Port Chester composite tissues.

3.6.5 Bioaccumulation of Metals in *Nereis virens*

All metals analyzed except silver and chromium were detected in *N. virens* tissues exposed to the Port Chester composite (Table 3.17). Of these, only cadmium was measured at concentrations that were statistically significant and higher than those measured in tissues exposed to the Mud Dump Reference Site sediments. Magnification factors were ≤ 3.0 for all metals.

3.6.6 Bioaccumulation of Pesticides in *Nereis virens*

Of the 15 pesticides analyzed, 8 were detected in the Port Chester tissues (Table 3.18). Aldrin, dieldrin, α -chlordane, *trans*-nonachlor, 2,4'-DDD, 4,4'-DDE, and 4,4'-DDD were detected at concentrations that were statistically significant and higher than in tissues exposed to both reference sediments. Several pesticides were detected in Port Chester

TABLE 3.17. Mean Concentrations of Metals in *N. virens* Tissues Exposed to Port Chester Sediment

Analyte	Concentration mg/kg wet weight) (a)				
	MDRS ^(b)	CLIS ^(c)	COMP PC	SD ^(d) MDRS	SD CLIS
Silver	0.0224	0.0122 Q ^(e)	0.0124 Q	No	No
Arsenic	2.07	2.08	1.85	No	No
Cadmium	0.0619	0.0548	0.102	Yes	Yes
Chromium	0.103 Q	0.107 Q	0.109 Q	No	No
Copper	3.30	1.52	1.61	No	No
Mercury	0.0121	0.0104	0.0072	No	No
Nickel	0.0928 Q	0.153	0.302	No	No
Lead	0.311	0.361	0.388	No	No
Zinc	11.2	26.2	31.4	No	No

(a) Value shown is a mean of five replicates; one-half the detection limit used when analyte was undetected.

(b) MDRS- Mud Dump Reference Site.

(c) CLIS-Central Long Island Sound Reference Site.

(d) SD Significantly different.

(e) Undetected at or above twice the given concentration.

tissues at concentrations greater than 10 times those of the Mud Dump Reference tissues (dieldrin, α -chlordane, 2,4'-DDD, 4,4'-DDE, and 4,4'-DDD) and Central Long Island Sound Reference Site tissues (α -chlordane and 4,4'-DDD).

3.6.7 Bioaccumulation of PCBs in *Nereis virens*

Nineteen of 22 PCBs analyzed were detected in *N. virens* tissues exposed to the Port Chester composite (Table 3.18). Of these, 16 were statistically significant and elevated relative to the Mud Dump Reference tissues, and 13 relative to the Central Long Island Sound Reference Site tissues. Six PCBs (52, 87, 101, 105, 118, and 138) were observed at concentrations greater than 10 times those of the tissues exposed to the Mud Dump Reference composite. The magnification factors, relative to the CLIS Reference were all below 10. The concentration of total estimated PCBs in *N. virens* tissues exposed to the Port Chester composite was 189 µg/kg wet weight.

3.6.8 Bioaccumulation of PAHs and 1,4-Dichlorobenzene in *Nereis virens*

All PAHs analyzed were detected in tissues exposed to the Port Chester composites (Table 3.19). Of these, eight were detected in statistically significant and elevated concentrations relative to tissues exposed to the Mud Dump Reference composite. Fluoranthene, pyrene, and chrysene were detected at concentrations over 10 times higher in *N. virens* exposed to Port Chester sediments than in the two reference composites. The compound 1,4-dichlorobenzene was undetected in the Port Chester and both reference composite tissues.

3.6.9 Magnification Factors of Compounds in *Macoma nasuta* and *Nereis virens* Tissues

Table 3.20 shows the calculated magnification factors of all compounds analyzed in tissues of *M. nasuta* and *N. virens*. Magnification factors were calculated with the dry weight concentrations of the compounds in the tissues of the test organism. These factors show the magnification of the Port Chester-exposed tissues over the Mud Dump Reference Site-exposed tissues and the Central Long Island Site-exposed tissues. When all replicate analysis of a compound showed that the compound was undetected, the magnification factor displays the magnification of the Port Chester-exposed tissues above the detection limit of either of the reference tissue values.

TABLE 3.18. Mean Concentrations of Pesticides and PCBs in *N. virens* Tissues Exposed to Port Chester Sediment

Analyte	Concentration (µg/kg wet weight) ^(a)						SD ^(d) MDRS	SD CLIS
	MDRS ^(b)		CLIS ^(c)		COMP PC			
2,4'-DDD	0.18		1.11		9.67		Yes	No
2,4'-DDE	0.14	Q ^(e)	0.13	Q	0.13	Q	No	No
2,4'-DDT	0.09	Q	0.09	Q	0.09	Q	No	No
4,4'-DDD	0.51		1.90		32.8		Yes	Yes
4,4'-DDE	0.15		0.62		6.12		Yes	Yes
4,4'-DDT	0.08	Q	0.08	Q	0.08	Q	No	No
Total DDT	1.15		3.93		48.9		Yes	Yes
Aldrin	0.07	Q	0.82		1.22		Yes	No
α-Chlordane	0.05	Q	0.12		5.15		Yes	Yes
Dieldrin	0.58		0.90		8.72		Yes	Yes
Endosulfan I	0.09	Q	0.09	Q	0.09	Q	No	No
Endosulfan II	0.09	Q	0.09	Q	0.09	Q	No	No
Endosulfan Sulfate	0.09	Q	0.12		1.76		No	No
Heptachlor	0.10	Q	0.09	Q	0.10	Q	No	No
Heptachlor Epoxide	0.07	Q	0.11		0.07	Q	No	No
<i>trans</i> - Nonachlor	0.54		0.61		3.10		Yes	Yes
PCB 8	0.21	Q	0.20	Q	0.21	Q	No	No
PCB 18	0.22	Q	0.21	Q	0.65		No	No
PCB 28	0.11	Q	0.27		1.06		Yes	No
PCB 44	0.09	Q	0.11		1.40		Yes	Yes
PCB 49	0.12	Q	0.53		2.04		Yes	Yes
PCB 52	0.32		1.81		9.41		Yes	Yes
PCB 66	0.05	Q	0.05	Q	0.05	Q	No	No
PCB 87	0.11		0.23		2.18		Yes	Yes
PCB 101	0.46		2.99		16.4		Yes	Yes
PCB 105	0.18		0.86		6.63		Yes	Yes
PCB 118	0.15	Q	1.95		9.51		Yes	Yes
PCB 128	0.25		0.55		1.99		No	No
PCB 138	1.18		2.87		13.7		Yes	Yes
PCB 153	2.01		3.79		12.9		Yes	No
PCB 170	0.28		0.61		2.73		Yes	Yes
PCB 180	0.58		1.17		5.76		Yes	Yes
PCB 183	0.17		0.44		2.41		Yes	Yes
PCB 184	0.12	Q	0.12	Q	0.12	Q	No	No
PCB 187	0.50		0.97		3.73		Yes	Yes
PCB 195	0.05	Q	0.05	Q	0.46		Yes	Yes
PCB 206	0.23		0.32		1.05		Yes	No
PCB 209	0.16		0.19		0.25		No	No
Total PCB	15.1		40.6		189		Yes	Yes

(a) Value shown is a mean of five replicates; one-half the detection limit used when analyte was undetected.

(b) MDRS- Mud Dump Reference Site.

(c) CLIS-Central Long Island Sound Reference Site.

(d) SD Significantly different.

(e) Undetected at or above twice the given concentration.

(f) Total DDT is the sum of 4,4'-DDT, 4,4'-DDE, 4,4'-DDD, 2,4'-DDT, 2,4'-DDE, and 2,4'-DDD; one-half of the detection limit was used in summation when analyte was not detected.

(g) Total PCB = 2(x), where x=sum of all PCB congeners; one-half of the detection limit used in summation when analyte was undetected.

TABLE 3.19. Mean Concentrations of PAHs in *N. virens* Tissues Exposed to Port Chester Sediment

Analyte	Concentration ($\mu\text{g}/\text{kg}$ wet weight) ^(a)			SD ^(d)	SD
	MDRS ^(b)	CLIS ^(c)	COMP PC	MDRS	CLIS
Naphthalene	4.49	1.85	4.14	No	No
Acenaphthylene	0.88	0.36 Q ^(e)	1.57	No	Yes
Acenaphthene	2.02	1.40	14.6	Yes	Yes
Fluorene	1.85	0.61 Q	3.11	No	Yes
Phenanthrene	3.01	1.55	5.02	No	No
Anthracene	1.17 Q	1.11 Q	4.75	Yes	Yes
Fluoranthene	2.80 Q	2.66 Q	100	Yes	Yes
Pyrene	3.86	3.74	69.8	Yes	Yes
Benzo[a]anthracene	3.43	1.73	7.42	No	Yes
Chrysene	1.18 Q	1.91	26.4	Yes	Yes
Benzo[b]fluoranthene	2.66	3.33	7.79	Yes	No
Benzo[k]fluoranthene	1.09	2.36	5.01	Yes	No
Benzo[a]pyrene	0.78 Q	1.05	4.60	Yes	Yes
Indeno[123-cd]pyrene	1.43	1.70	3.39	No	No
Dibenzo[a,h]anthracene	0.66 Q	0.63 Q	1.80	No	No
Benzo[g,h,i]perylene	1.27	1.55	3.86	No	No
1,4-Dichlorobenzene	0.97 Q	0.92 Q	0.95 Q	No	No

(a) Value shown is a mean of five replicates; one-half the detection limit used when analyte was undetected.

(b) MDRS- Mud Dump Reference Site.

(c) CLIS-Central Long Island Sound Reference Site.

(d) SD Significantly different.

(e) Undetected at or above twice the given concentration.

TABLE 3.20. Magnification Factors of All Analyzed Compounds in *Macoma nasuta* and *Nereis virens* Tissues Exposed to the Port Chester Composite Relative to Tissues Exposed to the Mud Dump Reference Site and the Central Long Island Sound Reference Site

Analyte	Magnification Factors(a)			
	<i>Macoma nasuta</i>		<i>Nereis virens</i>	
	MDRS	CLIS	MDRS	CLIS
Ag	0.99	1.30	0.94	1.00
As	1.10	1.14	0.84	0.87
Cd	<u>6.77</u>	<u>9.22</u>	1.55	1.82
Cr	1.21	1.01	1.00	1.00
Cu	1.63	1.13	0.47	1.03
Hg	0.95	1.01	0.55	0.66
Ni	1.61	1.02	1.55	1.41
Pb	2.60	0.89	1.06	1.14
Zn	1.20	1.30	2.74	1.22
2,4'-DDD	22.2	21.2	32.5	<u>8.27</u>
2,4'-DDE	1.02	1.08	0.92	1.00
2,4'-DDT	1.13	1.07	0.92	1.00
4,4'-DDD	47.1	43.9	54.7	16.4
4,4'-DDE	16.7	4.47	25.4	<u>9.35</u>
4,4'-DDT	0.14	0.02	0.91	1.01
α-Chlordane	40.9	39.7	46.6	36.7
Aldrin	3.35	<u>9.54</u>	<u>8.56</u>	1.47
Dieldrin	<u>7.99</u>	<u>7.43</u>	11.7	<u>8.97</u>
Endosulfan I	1.12	1.07	0.92	1.00
Endosulfan II	1.12	1.07	0.92	1.00
Endosulfan sulfate	1.12	1.07	<u>9.21</u>	<u>9.61</u>
Heptachlor	2.04	1.96	0.92	1.01
Heptachlor epoxide	1.15	1.09	0.92	0.82
trans-Nonachlor	4.96	4.75	<u>5.41</u>	4.93
PCB 8	0.45	1.08	0.92	1.01
PCB 18	1.74	1.67	1.56	1.71
PCB 28	2.18	1.67	4.70	3.37
PCB 44	4.61	3.38	<u>7.49</u>	<u>7.71</u>
PCB 49	<u>7.62</u>	2.83	<u>7.72</u>	3.54
PCB 52	<u>6.85</u>	<u>7.14</u>	20.6	4.87
PCB 66	0.43	0.10	0.90	1.00
PCB 87	16.2	13.5	11.3	<u>7.42</u>
PCB 101	17.1	<u>7.51</u>	32.8	<u>5.28</u>
PCB 105	16.6	15.6	30.8	<u>7.50</u>
PCB 118	16.0	<u>9.06</u>	29.2	4.62
PCB 128	4.44	<u>3.97</u>	<u>5.58</u>	3.57
PCB 138	10.6	<u>5.04</u>	10.9	4.63
PCB 153	14.5	2.19	<u>6.07</u>	3.30
PCB 170	1.76	1.76	<u>8.81</u>	4.40
PCB 180	4.02	2.62	<u>9.47</u>	4.77
PCB 183	1.32	1.26	<u>9.08</u>	<u>5.08</u>
PCB 184	1.13	1.08	0.92	1.01
PCB 187	4.75	0.59	<u>7.02</u>	3.73
PCB 195	1.14	1.09	4.32	4.69
PCB 206	1.37	1.30	4.15	3.24
PCB 209	4.45	4.29	1.52	1.29

TABLE 3.20. (contd)

Analyte	Magnification Factors(a)			
	<i>Macoma nasuta</i>		<i>Nereis virens</i>	
	MDRS	CLIS	MDRS	CLIS
Naphthalene	2.17	2.10	0.76	1.78
Acenaphthylene	1.16	0.81	1.24	2.14
Acenaphthene	15.1	14.4	<u>5.92</u>	<u>7.78</u>
Fluorene	<u>5.32</u>	<u>5.08</u>	1.33	2.55
Phenanthrene	12.7	<u>8.77</u>	1.21	1.89
Anthracene	17.2	12.0	1.90	2.09
Fluoranthene	126	70.4	16.8	18.5
Pyrene	110	41.1	12.5	13.3
Benzo[a]anthracene	44.9	19.7	2.02	3.67
Chrysene	55.8	23.2	10.5	11.0
Benzo[b]fluoranthene	24.2	<u>5.95</u>	2.43	2.26
Benzo[k]fluoranthene	3.73	1.27	2.61	2.06
Benzo[a]pyrene	21.2	<u>5.21</u>	2.76	2.72
Indeno[1,2,3-cd]pyrene	<u>6.68</u>	2.56	1.54	1.54
Dibenzo[a,h]anthracene	2.89	2.75	1.36	1.50
Benzo[g,h,i]perylene	<u>8.14</u>	2.51	1.94	1.88
1,4-Dichlorobenzene	1.14	1.08	0.92	1.01

(a) Magnification factors are the number of times the test treatment concentration is greater than the reference treatment concentration. When the analyte is undetected in one or more replicates, the achieved detection limit value is used in the calculation. Calculations are based on dry weight concentrations. Underlined values are between 5 and <10 times reference site values, values shown in bold are ≥ 10 times reference site values.

4.0 Discussion and Conclusions

In this section, physical and chemical analyses, and bioassays performed on the Port Chester sediment composite are evaluated relative to the Mud Dump Reference Site and the Central Long Island Sound Reference Site sediment by the guidelines of the Green Book Tier III. Tier III evaluations include water-column toxicity tests, benthic toxicity tests, and whole-sediment bioaccumulation studies. Tier III evaluations assess the impact of contaminants in the dredged material on marine organisms to determine whether there is potential for the material to have an unacceptable environmental effect during ocean disposal. The Green Book provides the following guidance for determining whether the proposed dredged material is unacceptable for ocean disposal based on the Tier III test:

- **Water-Column Toxicity.** The limiting permissible concentration (LPC) of dissolved plus suspended contaminants cannot exceed 0.01 of the acutely toxic concentration at the boundaries of the disposal site within the first 4 h after disposal, or at any point in the marine environment after the first 4 h. The acutely toxic concentration in this case is taken to the LC_{50} ; therefore, acute toxicity in SPP tests would require at least 50% mortality in an SPP treatment to be evaluated according to the Green Book. A numerical mixing model should be used to predict whether concentrations greater than 0.01 of the acutely toxic SPP concentrations are likely to occur beyond the boundaries of the disposal site within the first 4 h after disposal.
- **Benthic Acute Toxicity.** The proposed dredged material does not meet the LPC for benthic toxicity when organism survival in the test sediment and the reference site sediment is statistically significant, and the decrease in survival is at least 20% for *A. abdita* and *R. abronius* or at least 10% for *M. bahia*.
- **Bioaccumulation.** The proposed dredged material does not meet the LPC for bioaccumulation if tissue concentrations of one or more contaminants of concern are greater than the applicable FDA levels. Regional guidance (USACE 1981) for interpretation of bioaccumulation was also considered. When the bioaccumulation of contaminants in the dredged material exceeds that in the reference material exposures, further case-specific evaluation criteria listed in the Green Book should be consulted to determine LPC and benthic effects compliance.

Sections 4.1 through 4.4 discuss the proposed Port Chester dredged material in terms of sediment characterization and Tier III evaluations. The contribution of the Port Chester composite to water-column or benthic acute toxicity and potential for bioaccumulation relative to the Mud Dump Reference Site and the Central Long Island Sound Reference Site is also presented.

4.1 Sediment Physical and Chemical Characterization

Port Chester sediment core samples were either a black, silt and clay material or a black, sand and gravel mixture. The upriver Port Chester samples PC-1, PC-2, PC-3, PC-4, PC-6 and PC-7, sediments were predominantly coarse-grained. Port Chester samples PC-5, PC-8, PC-9,

PC-10, and PC-11, were predominately fine-grained, with percentages of sand ranging from 21% to 34%; silt ranging from 34% to 48%; and clay ranging from 20% to 33%. Sediment moisture contents varied from 4% to 70% in individual cores. Levels of metals (excluding As) in Port Chester sediment exceeded those found in the Mud Dump Reference Site sediment. The dominant pesticides found were those in the DDD/DDE/DDT group of compounds, α -chlordane, dieldrin, and *trans*-nonachlor. All of the 22 PCB congeners analyzed were detected in Port Chester sediment, with a total PCB concentration of 1060 $\mu\text{g}/\text{kg}$ dry weight. All 17 PAHs analyzed were detected in Port Chester sediment. Total PAH was 15,800 $\mu\text{g}/\text{kg}$ dry weight; 20% of the total was LPAH; 80% of the total was HPAH. The concentration of 1,4-dichlorobenzene was 76 $\mu\text{g}/\text{kg}$ dry weight.

4.2 Site Water and Elutriate Chemical Characterization

Metals concentrations were consistently higher in the Port Chester site water than in Mud Dump site water. Port Chester elutriate concentrations of metals were generally lower than Mud Dump Site water, except for Cr, Hg, Ni, and Pb, which were elevated by a factor of two in the Port Chester elutriate. The dominant pesticides found in the Port Chester site water and elutriate samples were those in the DDD/DDE/DDT group of compounds, and α -chlordane. Measurable amounts of the PCB congener PCB (28) were found in Port Chester site water, and all the PCB congeners except PCB(08), PCB(18), and PCB(184) was found in the Port Chester elutriate.

4.3 Toxicity

The contribution of the Port Chester composite to water-column toxicity relative to the Mud Dump Reference Site and the Central Long Island Sound Reference Site are presented in Figures 4.1 and 4. In water-column toxicity tests, 100% SPP treatments were acutely toxic to all three species tested. The LC_{50} values ranged from 75.1% SPP for *M. beryllina* to >100% SPP for *M. bahia* and *M. galloprovincialis* survival. The EC_{50} value for *M. galloprovincialis* normal development, a more sensitive measure than survival, was 53.7% SPP. The LPC for water-column effects outside of the disposal site boundaries after 4 h is 0.75% SPP for Port Chester sediment. A projection of SPP concentrations exceeding 0.75% SPP after 4 h at the Mud Dump Site boundary would be unacceptable.

The contribution of the Port Chester composite to benthic acute toxicity relative to the Mud Dump Reference Site and the Central Long Island Sound Reference Site are also presented in Figures 4.1 and 4.2.

		Sediment Treatment	Port Chester vs R-MUD	
Acute Toxicity	<i>A. abdita</i> Benthic Static-Renewal Test		AT ^(a)	
	<i>R. abronius</i> Benthic Static-Renewal Test		-	
	<i>M. bahia</i> Benthic Static Test		-	
	<i>M. beryllina</i> SPP Test		S ^(c)	
	<i>M. bahia</i> SPP Test		S ^(c)	
	<i>M. galloprovincialis</i> SPP Test		S ^(c)	
		Test Species ^(d)	<i>N. virens</i>	<i>M. nasuta</i>
Any Significant Bioaccumulation	# of Metals (9 total)		1	4
	# of Pesticide compounds (15 total)		7	7
	# of PCB congeners (22 total)		16	12
	# of PAH compounds (16 total)		8	14
	1,4-dichlorobenzene		-	-
Bioaccumulation ≤ 2 times Ref.	# of Metals (9 total)		1	2
	# of Pesticide compounds (15 total)		-	-
	# of PCB congeners (22 total)		-	-
	# of PAH compounds (16 total)		1	-
	1,4-dichlorobenzene		-	-
Bioaccumulation >2.5 times Ref.	# of Metals (9 total)		-	1
	# of Pesticide compounds (15 total)		-	2
	# of PCB congeners (22 total)		3	4
	# of PAH compounds (16 total)		3	2
	1,4-dichlorobenzene		-	-
Bioaccumulation >5 ≤ 10 times Ref.	# of Metals (9 total)		-	1
	# of Pesticide compounds (15 total)		2	1
	# of PCB congeners (22 total)		7	2
	# of PAH compounds (16 total)		1	3
	1,4-dichlorobenzene		-	-
Bioaccumulation >10 times Ref.	# of Metals (9 total)		-	-
	# of Pesticide compounds (15 total)		5	4
	# of PCB congeners (22 total)		6	6
	# of PAH compounds (16 total)		3	9
	1,4-dichlorobenzene		-	-

- (a) AT Significantly different from reference (R-MUD) and mortality 20% (10% for mysids) greater than reference.
 (b) - No significant difference/no significant bioaccumulation at this level.
 (c) Significantly different mortality between 0% and 100% SPP.
 (d) Number of compounds bioaccumulating in tissues of test species.

FIGURE 4.1. Summary Matrix of Port Chester Sediment Toxicity and Bioaccumulation Potential Compared with the Mud Dump Reference Site.

		Sediment Treatment	Port Chester vs R-CLIS
Acute Toxicity	<i>A. abdita</i> Benthic Static-Renewal Test		AT ^(a)
	<i>R. abronius</i> Benthic Static-Renewal Test		-(b)
	<i>M. bahia</i> Benthic Static Test		-
	<i>M. beryllina</i> SPP Test		S ^(c)
	<i>M. bahia</i> SPP Test		S ^(c)
	<i>M. galloprovincialis</i> SPP Test		S ^(c)

Any Significant Bioaccumulation	Test Species ^(d)	<i>N. virens</i>	<i>M. nasuta</i>
		# of Metals (9 total)	1
	# of Pesticide compounds (15 total)	5	7
	# of PCB congeners (22 total)	13	9
	# of PAH compounds (16 total)	9	14
	1,4-dichlorobenzene	-	-

Bioaccumulation ≤ 2 times Ref.	# of Metals (9 total)	1	-
	# of Pesticide compounds (15 total)	-	-
	# of PCB congeners (22 total)	-	-
	# of PAH compounds (16 total)	-	-
	1,4-dichlorobenzene	-	-

Bioaccumulation >2≤5 times Ref.	# of Metals (9 total)	-	-
	# of Pesticide compounds (15 total)	1	2
	# of PCB congeners (22 total)	8	3
	# of PAH compounds (16 total)	5	4
	1,4-dichlorobenzene	-	-

Bioaccumulation >5≤10 times Ref.	# of Metals (9 total)	-	1
	# of Pesticide compounds (15 total)	2	2
	# of PCB congeners (22 total)	5	4
	# of PAH compounds (16 total)	1	4
	1,4-dichlorobenzene	-	-

Bioaccumulation >10 times Ref.	# of Metals (9 total)	-	-
	# of Pesticide compounds (15 total)	2	3
	# of PCB congeners (22 total)	-	2
	# of PAH compounds (16 total)	3	6
	1,4-dichlorobenzene	-	-

- (a) AT Significantly different from reference (R-CLIS) and mortality 20% (10% for mysids) greater than reference.
- (b) - No significant difference/no significant bioaccumulation at this level.
- (c) Significantly different mortality between 0% and 100% SPP.
- (d) Number of compounds bioaccumulating in tissues of test species.

FIGURE 4.2. Summary Matrix of Port Chester Sediment Toxicity and Bioaccumulation Potential Compared with the Central Long Island Sound Reference Site

In the static renewal test with *A. abdita* exposed to Port Chester sediment, test organism survival was statistically significant and $\geq 20\%$ different from that in the Mud Dump Reference and the Central Long Island Sound Reference Sites.

In the static renewal test with *R. abronius*, survival was statistically significant when compared with both reference site sediments, but there was not a $\geq 20\%$ difference in survival relative to the Central Long Island Sound Reference Site. In the *M. bahia* static test, which was not manipulated to reduce porewater or overlying ammonia concentrations prior to test initiation, test organisms survival was 77% in the Port Chester sediment treatment. The survival percentage statistically significant, nor was this difference $\geq 10\%$ from that in either of the two references.

4.4 Bioaccumulation

Results of *N. virens* And *M. nasuta* tissue analyses from test sediment bioaccumulation studies were compared with action levels for poisonous or deleterious substances in fish and shellfish for human consumption published by the FDA and with USACE-NYD (1981) bioaccumulation matrix tables. Concentrations of As, Cd, Cr, Ni, and Pb were also compared with the FDA level of concern for chronic shellfish consumption (FDA 1993a, 1993b, 1993c, 1993d, 1993e) for each of these metals. Results of tissue analyses from test sediment bioaccumulation studies were also compared with contaminant concentrations in tissues of organisms similarly exposed to Mud Dump Reference Site and Central Long Island Sound Reference Site sediments.

When *N. virens* and *M. nasuta* were exposed to Port Chester sediment in 28-day bioaccumulation tests, concentrations of some contaminants were elevated in tissues of both species. Concentrations of metals were generally higher in *M. nasuta* than in *N. virens*. Pesticide and PCB concentrations were similar in the two species, with some analytes higher in the *N. virens*, and others higher in the *M. nasuta*. Concentrations of PAHs were higher in *M. nasuta*, many compounds by a factor of 4 to 10 times, than in *N. virens*.

When tissue burdens of organisms exposed to Port Chester sediment were compared with those exposed to Mud Dump Reference Site and Central Long Island Sound Reference Site sediment, the tissue burdens were statistically significant and elevated for metals, pesticides, PCBs, and PAHs in both *M. nasuta* and *N. virens* tissues. Figures 4.1 and 4.2 show bioaccumulation potential as the number of contaminants that were elevated in the tissues of

M. nasuta and *N. virens* at certain magnitudes (i.e. 2, 5, or 10 times) above tissues of each species exposed to each reference sediment. This format clearly indicates where and to what degree similar classes of contaminants were accumulated in both species.

Table 4.1 compares the FDA action levels for poisonous or deleterious substances in fish and shellfish for human consumption for selected pesticides, and FDA levels of concern for chronic shellfish consumption for selected metals with the mean concentration of these contaminants found in tissues of each test species. The *N. virens* and *M. nasuta* tissues exposed to Port Chester sediment had tissue body burdens that were lower than the FDA levels for each of these selected contaminants.

TABLE 4.1. Comparison of Contaminant Concentrations in *N. virens* and *M. nasuta* Tissues Exposed to Proposed Dredged Material from Port Chester with FDA Action Levels and Levels of Concern

Substance	FDA Level (mg/kg wet wt)	Concentrations ^(a) in <i>N. virens</i> Tissues (mg/kg wet wt)	Concentrations ^(b) in <i>M. nasuta</i> Tissues (mg/kg wet wt)
		Port Chester Composite	Port Chester Composite
Chlordane ^(b)	0.3 ^(c)	0.005	0.004
Total DDT ^(d)	5.0 ^(c)	0.0489	0.0219
Dieldrin + Aldrin	0.3 ^(c)	0.010	0.005
Heptachlor+			
Heptachlor epoxide	0.3 ^(c)	<0.0002	0.0004
Total PCBs ^(e)	2.0 ^(c)	0.189	0.067
Arsenic	86 ^(f)	1.85	3.03
Cadmium	3.7 ^(f)	0.102	0.208
Chromium	13 ^(f)	<0.218	0.436
Lead	1.7 ^(f)	0.388	0.721
Nickel	80 ^(f)	0.302	0.559
Methyl Mercury	1.0 ^(f)	0.007 ^(g)	0.015 ^(g)
Total DDT ^(d)	0.04 ^(h)	0.0489	0.0219
Total PCBs ^(e)	0.40 ^(h)	0.189	NA ⁽ⁱ⁾
Total PCBs ^(e)	0.10 ^(h)	NA	0.067
Total Mercury	0.20 ^(h)	0.007	0.015
Cadmium	0.30 ^(h)	0.102	0.208

- (a) Concentration shown is the mean of five replicate tissue analysis. If any constituents were undetected, one-half of the detection limit was used in calculation of the mean concentration.
- (b) Sum of *a*-chlordane and *trans*-nonachlor only, whereas FDA action level is a sum of nine chlordane analytes.
- (c) FDA action levels for poisonous and deleterious substances in fish and shellfish for human food.
- (d) Sum of mean values for 2,4'-DDT, 4,4'-DDT, 2,4'-DDE, and 4,4'-DDE, 2,4'-DDD, and 4,4'-DDD. One-half of the detection limit was used in the summation when mean values were undetected in a replicate.
- (e) Total PCBs= 2.0(x), where x equals the sum of the 22 congeners. One-half of the detection limit was used in summation when mean values were undetected in a replicate.
- (f) FDA level of concern for chronic shellfish consumption.
- (g) Value reported here is for total mercury.
- (h) NYD bioaccumulation matrix value designated in 1981 (USACE 1981).
- (i) NA Not applicable.

5.0 References

- Barrows, E.S., M.R. Pinza, W.W. Gardiner, N.P. Kohn, B.D. Gruendell, H.L. Mayhew, J.Q. Word, and L.B. Rosman. 1996. *Evaluation of Dredged Material Proposed for Ocean Disposal from Federal Projects in New York and New Jersey and the Military Ocean Terminal (MOTBY)*. Prepared for Department of the Army, U.S. Army Corps of Engineers, New York District, Water Quality Branch, under a Related Services Agreement with the U.S. Department of Energy by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory.
- Bligh, E.G., and W.J. Dyer. 1959. "A Rapid Method of Total Lipid Extraction and Purification." *Canadian Journal of Biochemistry and Physiology*. 37:8 911-917.
- Bloom, N.S., and E.A. Crecelius. 1983. "Determination of Mercury in Seawater at Sub-Nanogram per Liter Levels." *Mar. Chem.* 14:49-59.
- Dunnett, C.W. 1964. "New Tables for Multiple Comparisons with a Control." *Biometrics* 20:482-491.
- EPA (U.S. Environmental Protection Agency). 1986. *Determination of Total Organic Carbon in Sediment*. U.S. Environmental Protection Agency, Region II, Environmental Services Division, Monitoring Management Branch, Edison, New Jersey.
- EPA (U.S. Environmental Protection Agency). 1990. *Test Methods for Evaluating Solid Waste, Physical/Chemical Methods, SW-846*. 3rd Ed. U.S. Environmental Protection Agency, Washington, D.C.
- EPA (U.S. Environmental Protection Agency). 1991. *Methods for the Determination of Metals in Environmental Samples*. EPA/600/4-91-010. U.S. Environmental Protection Agency, Office of Research and Development, Washington, D.C.
- EPA/USACE (U.S. Environmental Protection Agency/U.S. Army Corps of Engineers). 1991. *Evaluation of Dredged Material Proposed for Ocean Disposal (Testing Manual)*. EPA-503/8-91/001. U.S. Environmental Protection Agency, Office of Marine and Estuarine Protection, Washington, D.C.
- EPA/USACE (U.S. Environmental Protection Agency/U.S. Army Corps of Engineers). 1993. Memorandum: Technical Panel Recommendations Concerning Use of Acute Amphipod Tests in Evaluation of Dredged Material. December 21, 1993. U.S. Environmental Protection Agency/U.S. Army Corps of Engineers, Washington, D.C.
- FDA (U.S. Food and Drug Administration). 1993a. Guidance Document for Arsenic in Shellfish. U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Washington, D.C.
- FDA (U.S. Food and Drug Administration). 1993b. Guidance Document for Cadmium in Shellfish. U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Washington, D.C.
- FDA (U.S. Food and Drug Administration). 1993c. Guidance Document for Chromium in Shellfish. U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Washington, D.C.
- FDA (U.S. Food and Drug Administration). 1993d. Guidance Document for Nickel in Shellfish. U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Washington, D.C.

FDA (U.S. Food and Drug Administration). 1993e. Guidance Document for Lead in Shellfish. U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Washington, D.C.

Finney, D. J. 1971. *Probit Analysis*. 3rd Ed. Cambridge University Press, Boston, Massachusetts.

NOAA (National Oceanic and Atmospheric Administration). 1989. *A Summary of Data on Tissue Contamination from the First Three Years (1986-1988) of the Mussel Watch Program*. NOAA Technical Memorandum NOS OMA 49. U.S. Department of Commerce, NOAA National Ocean Service, Rockville, Maryland.

NOAA (National Oceanic and Atmospheric Administration). 1993. *Sampling and Analytical Methods of the National Status and Trends Program National Benthic Surveillance and Mussel Watch Projects 1984-1992. Volume IV. Comprehensive Descriptions of Trace Organic Analytical Methods*. G. G. Lauenstein and A.Y. Cantillo, eds. NOAA Technical Memorandum NOS ORCA 71. National Oceanic and Atmospheric Administration, Coastal Monitoring and Bioeffects Assessment Division, Office of Ocean Resources Conservation and Assessment, Silver Spring, Maryland.

NYSDEC (New York State Department of Environmental Conservation). 1992. *Analytical Method for the Determination of PCB congeners by Fused Silica Capillary Column Gas Chromatography with Electron Capture Detector*. NYSDEC Method 91-11. New York State Department of Environmental Conservation, Albany, New York.

Plumb, R. H., Jr. 1981. *Procedure for Handling and Chemical Analysis of Sediment and Water Samples*. Tech. Rep. EPA/CE-81-1. Prepared by Great Lakes Laboratory, State University College at Buffalo, New York, for the U.S. Environmental Protection Agency/U.S. Army Corps of Engineers Technical Committee on Criteria for Dredged and Fill Material. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.

Randall, R. C. 1988. *Lipid Standard Operating Procedure, Modified Bligh/Dyer*. U.S. Environmental Protection Agency Office of Research and Development, Environmental Research, Washington D.C.

USACE (U.S. Army Corps of Engineers). 1970 (Revised 1980). *Engineer Manual, Engineering and Design, Laboratory Soils Testing*. EM 110-2-1906. Headquarters, Department of the Army, Office of the Chief of Engineers, Washington, D.C.

USACE (U.S. Army Corps of Engineers). 1981. Memorandum: Final Interpretative Guidance for Bioaccumulation of Petroleum Hydrocarbons, DDT, Cadmium, and Mercury in the New York Bight. April 7, 1981. USACE Office of The Chief Engineers, New York, New York.

USACE-NED/EPA Region I (U.S. Army Corps of Engineers, New England Division/U.S. Environmental Protection Agency, Region I). 1989. *Guidance for Performing Tests on Dredged Material to be Disposed of in Open Waters (Draft)*. U.S. Army Corps of Engineers, New England Division, Waltham, Massachusetts, and U.S. Environmental Protection Agency, Region I, Boston, Massachusetts.

USACE-NYD/EPA Region II (U.S. Army Corps of Engineers, New York District/U.S. Environmental Protection Agency, Region II). 1992. *Guidance for Performing Tests on Dredged Material Proposed for Ocean Disposal*. 18 December 1992 Draft. U.S. Army Corps of Engineers, New York District and U.S. Environmental Protection Agency, Region II. New York, New York.

Appendix A

**Quality Assurance/Quality Control Data for
Sediment Physical/Chemical Analyses,
Port Chester Project**

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Grain Size, Bulk Density, Specific Gravity and Total Solids
LABORATORY: Soil Technology, Bainbridge Island, Washington
MATRIX: Sediment

QA/QC DATA QUALITY OBJECTIVES

	<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (dry wt)</u>
Grain Size	ASTM D-2217 and D-422	N/A	N/A	≤20%	1.0%
Bulk Density	ASTM D-854	N/A	N/A	≤20%	N/A
Specific Gravity	EM 1110-2-1906	N/A	N/A	≤20%	N/A
Total Solids	Plumb 1981	N/A	N/A	N/A	1.0%

METHOD Grain size was measured for four fractions using a combination of sieve and pipet techniques, following ASTM method D-2217 and D-422 for wet sieving. Bulk density was measured in accordance with ASTM method D-854. Specific gravity was measured in accordance with USACE Method EM 1110-2-1906. Total solids were measured gravimetrically following Plumb (1981).

HOLDING TIMES Samples were analyzed within the 6 month holding time.

DETECTION LIMITS Target detection limits of 1.0% by weight for each fraction were met for all samples.

METHOD BLANKS Not applicable.

MATRIX SPIKES Not applicable.

REPLICATES Six samples were analyzed in triplicate for grain size for the entire set of NY/NJ Federal Projects-2 program. Precision was measured by calculating the relative standard deviation (RSD) among triplicate results. The RSD's ranged from 0% to 10%, indicating acceptable precision. Two samples were analyzed in duplicate for bulk density and specific gravity. Precision was measured by calculating the relative percent difference (RPD) between the replicate results. The RPDs for bulk density were 0% and 2% while the RPDs for specific gravity were both 1%, indicating acceptable precision of the methods.

For total solids, three samples were analyzed in duplicate and four samples were analyzed in triplicate. All RSDs and RPDs were 0%.

**QA/QC SUMMARY/GRAIN SIZE, BULK DENSITY, SPECIFIC GRAVITY and
TOTAL SOLIDS (continued)**

SRMs Not applicable.

REFERENCES

ASTM D-2217. Standard Method for Wet Preparation of Soil Samples for Particle-size Analysis and Determination of Soil Constants.

ASTM D-422. Standard Method for Particle-Size Analysis of Soils.

ASTM D-854. Standard Method for Specific Gravity

EM 1110-2-1906. USACE (U.S. Army Corps of Engineers). 1970. Engineering and Design Laboratory Soils Testing.

Plumb R.H. 1981. Procedure for Handling and Chemical Analysis of Sediment and Water Samples. Tech. Rep. EPA/CE-81-1. Prepared for Great Lakes Laboratory, State University College at Buffalo, Buffalo, New York, for the U.S. Environmental Protection Agency/U.S. Army Corps of Engineers Technical Committee on Criteria for Dredged and Fill Material. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Total Organic Carbon (TOC)
LABORATORY: Global Geochemistry, Canoga Park, California
MATRIX: Sediment

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (dry wt)</u>
EPA 1986	N/A	≤20%	≤10%	0.1%

METHOD TOC was analyzed in accordance with EPA (1986). Analysis was performed by combustion and quantitation of evolved carbon dioxide using a LECO analyzer.

HOLDING TIMES Samples were analyzed within the 6 month holding time.

DETECTION LIMITS Target detection limits of 0.1% was met for all samples.

METHOD BLANKS Thirty-four method blanks were analyzed with the entire set of NY/NJ Federal Projects-2 program sediment samples. TOC levels detected in blanks ranged from 0.001% to 0.008% which were less than the established detection limit.

MATRIX SPIKES Not applicable.

REPLICATES Four samples were analyzed in triplicate and three samples were analyzed in duplicate. Precision was measured by calculating the relative standard deviation (RSD) or relative percent difference (RPD) between the replicate results. All RSDs and RPDs were between 1% and 10% indicating acceptable precision of the method.

SRMs Standard reference material MESS-1, obtained from the National Research Council of Canada, was analyzed at least once per batch of sediment samples. Although MESS-1 is not certified for TOC, accuracy was measured by calculating the percent difference (PD) from the in-house consensus value. PD values reported ranged from 1% to 8%.

REFERENCES

EPA (U.S. Environmental Protection Agency) 1986. Determination of Total Organic Carbon in Sediment. Environmental Protection Agency, Region II, Environmental Services Division, Monitoring Management Branch, Edison, New Jersey.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Metals
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Sediment

QA/QC DATA QUALITY OBJECTIVES

	<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Achieved Detection Limit (mg/kg dry wt)</u>
Arsenic	ICP/MS	75-125%	≤20%	≤20%	0.572
Cadmium	ICP/MS	75-125%	≤20%	≤20%	0.020
Chromium	ICP/MS	75-125%	≤20%	≤20%	0.401
Copper	ICP/MS	75-125%	≤20%	≤20%	0.525
Lead	ICP/MS	75-125%	≤20%	≤20%	0.136
Mercury	CVAA	75-125%	≤20%	≤20%	0.001
Nickel	ICP/MS	75-125%	≤20%	≤20%	0.849
Silver	ICP/MS	75-125%	≤20%	≤20%	0.119
Zinc	ICP/MS	75-125%	≤20%	≤20%	2.55

METHOD

A total of nine metals was analyzed: silver (Ag), arsenic (As), cadmium (Cd), chromium (Cr), copper (Cu), mercury (Hg), nickel (Ni), lead (Pb) and zinc (Zn). Hg was analyzed using cold-vapor atomic absorption spectroscopy (CVAA) according to the method of Bloom and Creclius (1983). The remaining metals were analyzed by inductively coupled plasma mass spectrometry (ICP/MS) following EPA Method 200.8 (EPA 1991)

To prepare sediment samples for analysis, samples were freeze-dried and blended in a Spex mixer-mill. Approximately 5 g of mixed sample was ground in a ceramic ball mill. For ICP/MS and CVAA analyses, 0.2- to 0.5-g aliquots of dried homogenous sample were digested using nitric acid following modified EPA Method 200.2 (EPA 1991). Sediment samples initially showed poor matrix spike recovery for Ag. (Refer to Matrix Spike section of this QA/QC Summary.) EPA Method 200.2 was modified by the addition of aqua regia to the digestion procedure and all samples were reanalyzed for Ag.

HOLDING TIMES

Samples were received on 3/30/94 and were logged into Battelle's log-in system. Samples were frozen to -80°C and subsequently freeze dried. Samples were all analyzed within 180 days of collection. The following list summarizes all analysis dates:

<u>Task</u>	<u>Date Performed</u>
Sample Digestion	5/5/94
ICP-MS	5/20/94
CVAA-Hg	5/9/94

QA/QC SUMMARY/METALS (continued)

- DETECTION LIMITS** Target detection limits were exceeded for some metals; however, metals were detected above the MDLs in all samples with the exception of Ag in one sample. MDLs were determined by multiplying the standard deviation of the mean of four replicate low level sediment spikes by 3.5.
- METHOD BLANKS** Two method blanks were analyzed. No metals were detected above the MDL in either blank with the exception of Pb in Blank-2. The value was less than three times the MDL and all sample values were detected at levels greater than five times the blank concentration, so no data were flagged. All data were blank corrected.
- MATRIX SPIKES** Two samples were spiked with all nine metals. In the original set of matrix spikes, recoveries of all metals, with the exception of Ag, were within the QC limits of 75% to 125%. Recoveries of Ag in the original spikes were low (3% and 10%). After reanalysis of the matrix spikes with the addition of aqua regia to the digestion procedure (see Methods section of this QA/QC Summary), matrix spike recoveries improved (93%) and concentrations of Ag in the dredging site sediments increased slightly. The low recovery of Ag appears to occur in analysis of marine sediment samples having high (in excess of approximately 5 µg/g) Ag concentrations. During the EPA Method 200.2 digestion procedure, a precipitate of AgCl can form with the Ag in the sediment and the Cl in the seawater.
- REPLICATES** Two samples were digested and analyzed in triplicate. Precision of triplicate analyses is reported by calculating the relative standard deviation (RSD) between the replicate results. RSD values ranged from 1% to 5%, within the QC limits of ±20%, indicating acceptable precision.
- SRM** Standard Reference Material (SRM) 1646 (estuarine sediment from the National Institute of Standards and Technology [NIST]), was analyzed for all metals. Only results for Cd, Cu and Hg were within ±20% of the certified value (Ag is not certified). Results for As, Ni, and Pb were between 20 and 30% of the certified values. The poorest result was with Cr, where the mean was 46% of the certified value. Values for the remaining metals were low because the digestion method used is not as strong as the method (perchloric acid) used to certify the SRM; thus, the results of this analysis should not be expected to match the SRM certified values. Therefore, no corrective actions were taken.

REFERENCES

- Bloom, N. S., and E.A. Crecelius. 1983. "Determination of Mercury in Seawater at Sub-Nanogram per Liter Levels." *Mar. Chem.* 14:49-59.
- EPA (U.S. Environmental Protection Agency). 1991. *Methods for the Determination of Metals in Environmental Samples*. EPA-600/4-91-010. Environmental Services Division, Monitoring Management Branch., Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Additional Metals
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Sediment

QA/QC DATA QUALITY OBJECTIVES

	<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Achieved Detection Limit (mg/kg dry wt)</u>
Antimony	ICP/MS	75-125%	≤20%	≤20%	0.03
Beryllium	ICP/MS	75-125%	≤20%	≤20%	0.5
Selenium	GFAA	75-125%	≤20%	≤20%	0.13
Thallium	ICP/MS	75-125%	≤20%	≤20%	0.024

METHOD An additional four metals were analyzed for a subset of sediment samples: Antimony (Sb), Beryllium (Be), Selenium (Se) and Thallium (Tl).

To prepare sediment samples for analysis, samples were freeze-dried and blended in a Spex mixer-mill. Approximately 5 g of mixed sample was ground in a ceramic ball mill. For inductively coupled plasma mass spectrometry (ICP/MS) and graphite furnace atomic absorption (GFAA) analyses, 0.2- to 0.5-g aliquots of dried homogenous sample were digested according to EPA Method 200.2 (EPA 1991), modified by the addition of aqua regia to the digestion procedure. Se was analyzed using GFAA. The other three metals were analyzed by ICP/MS following EPA Method 200.8 (EPA 1991).

HOLDING TIMES Samples were received on 3/30/94 and was logged into Battelle's log-in system. Samples were frozen to -80°C and subsequently freeze-dried. According to instructions from the program manager, 21 samples were composited into 8 samples. A subset of 17 samples (the Port Chester and Eastchester sediment composites) were analyzed for an additional four metals as requested in a memo from the program manager dated 1/11/95. The following list summarizes all analysis dates:

<u>Task</u>	<u>Date Performed</u>
Aqua Regia	2/1/95
ICP/MS - Sb, Be, Tl	3/7/95
GFAA - Se	2/7/95

DETECTION LIMITS Target detection limits were met for Sb, Se, and Tl. The detection limit (DL) for Be exceeds the target detection limit. However, all but three values were greater than the estimated DL and these values were flagged with a J to indicate an estimation.

QA/QC SUMMARY/ADDITIONAL METALS (continued)

- METHOD BLANKS** Two method blanks were analyzed. Only Sb was detected in one of the blanks; however, the values were less than three times the MDL and all sample values were detected at levels greater than five times the blank concentration. Therefore, no data were flagged and all data were blank corrected.
- MATRIX SPIKES** One sample was spiked with all four metals. Recoveries of all metals except Sb (228%) were within the QC limits of 75% to 125%.
- REPLICATES** One sample was digested and analyzed in triplicate. Precision for triplicate analyses is reported by calculating the relative standard deviation (RSD) between replicate results. RSD values ranged from 2% to 12%, which is within the QC limits of $\pm 20\%$, indicating acceptable precision.
- SRM** SRM 1646 (estuarine sediment from the National Institute of Standards and Technology [NIST]), was analyzed for all metals. None of the four additional metals are certified. However, non-certified values are reported and all four metals, with the exception of one replicate for Sb, are within 39% of the non-certified values.

REFERENCES

EPA (U.S. Environmental Protection Agency). 1991. *Methods for the Determination of Metals in Environmental Samples*. EPA-600/4-91-010. Environmental Services Division, Monitoring Management Branch, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Chlorinated Pesticides, PCB Congeners, and 1,4-Dichlorobenzene
LABORATORY: Battelle Ocean Sciences, Duxbury, Massachusetts
MATRIX: Sediment

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>MS Recovery</u>	<u>Surrogate Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (dry wt)</u>
GC/ECD	50-120%	30-150%	≤30%	≤30%	1.0 - 20 ng/g

METHOD Sediment samples were extracted with methylene chloride according to a modified version of EPA Method 8080 and the National Oceanic and Atmospheric Administration (NOAA) Status and Trends cleanup procedure (Krahn et al. 1988). Extracts were analyzed using gas chromatography with electron capture detection (GC/ECD) following a modified version of EPA Method 8270. Pesticide detections were qualitatively confirmed on a secondary column.

HOLDING TIMES Samples were collected from 3/22/94 through 3/25/94, and after compositing, were held frozen at -20°C until shipment to the analytical laboratory. Sediment samples were received by Battelle Ocean Sciences on 4/22/94. Samples were held frozen at -20°C until extraction and analysis. Samples were extracted by 5/6/94 and analyzed from 6/2/94 to 6/29/94.

DETECTION LIMITS Target detection limits were exceeded for most of the analytes. Actual detection limits were determined by the Method Detection Limit (MDL) verification study. Four sediment samples with very low background concentrations of contaminants were spiked with target compounds. For each analyte, the standard deviation of the four spiked replicates was multiplied by 3.5.

METHOD BLANKS One method blank was extracted with batch of samples. No pesticides or PCB congeners were detected in the blank.

SURROGATES Two compounds, DBOFB and PCB congener 112, were added to all samples prior to extraction to assess the efficiency of the analysis. The mean recoveries of DBOFB and PCB 112 were 71% and 60%, respectively. Recoveries of these compounds were within the QC guidelines of 30% -150% for all samples analyzed.

MATRIX SPIKES One sample in each batch was spiked with pesticides and PCB congeners. Recoveries for PCB congener CL₂ (25% and 47%) fell below the acceptable criteria of 50% to 120%. The reason for this low recovery is probably that the PCB congener CL₂ coeluted with alpha-BHC. All other PCB congener recoveries ranged from 54% to 121%. Recoveries for all pesticides and 1,4-dichlorobenzene ranged from 57% to 115%. Since >80% of all analytes were between 50% and 120%, no corrective action was taken.

QA/QC SUMMARY/CHLORINATED PESTICIDES and PCB CONGENERS (continued)

REPLICATES

One sample from each batch was extracted in triplicate. Precision was measured by calculating the relative standard deviation (RSD) between the replicate results. RSDs were evaluated only when pesticides or PCB congeners were detected in all three replicates. RSDs ranged from 5% to 114%. Six of the RSDs were greater than 30% but of those six, only three were for analytes that were >10 times the MDL. These three were 31% for CL₃(18), 114% for CL₅(105) and 52% for CL₆(138).

SRMs

One SRM, 1941a, a marine sediment sample obtained from the National Institute of Science and Technology (NIST) was analyzed with each batch. Many of the values exceeded the acceptable criteria of ≤30%; however all were <10 times the MDL. Percent differences were calculated using SRM concentrations that were corrected for surrogate recovery.

REFERENCES

Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.

EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Polynuclear Aromatic Hydrocarbons (PAH)
LABORATORY: Battelle Ocean Sciences, Duxbury, Massachusetts
MATRIX: Sediment

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>MS/MSD Recovery</u>	<u>Surrogate Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (dry wt)</u>
GC/MS/SIM	50-120%	30-150%	≤30%	≤30%	10 ng/g

METHOD Sediment samples were extracted according to a modified version of EPA Method 8080 and the NOAA Status and Trends cleanup procedure (Krahn et al. 1988). Extracts were analyzed using gas chromatography/mass spectrometry (GC/MS) in the selected ion mode (SIM) following a modified version of EPA Method 8270.

HOLDING TIMES Samples were collected from 3/22/94 through 3/25/94, and after compositing, were held frozen at -20°C until shipment to the analytical laboratory. Sediment samples were received by Battelle Ocean Sciences, Duxbury, Massachusetts, on 4/22/94. Samples were held frozen at approximately -20°C until extraction and analysis. Samples were extracted by 5/6/94 and analyzed from 5/16/94 to 6/28/94.

DETECTION LIMITS Target detection limits of 10 ng/g dry weight were met for most of the PAH compounds. Actual detection limits were determined by the Method Detection Limit (MDL) verification study. Four sediment samples with very low background concentrations of contaminants were spiked with target compounds. For each analyte, the standard deviation of the four spiked replicates was multiplied by 3.5. Actual detection limits ranged from 7.18 to 20.84 µg/kg.

METHOD BLANKS One method blank was extracted with each batch of samples. No PAH compounds were detected above the MDL; however, 2 of the 17 compounds were detected below the MDL and are flagged with a "J" to indicate the values are estimates. They are pyrene in Batch 1 and naphthalene in Batch 2.

SURROGATES Three isotopically labelled compounds were added prior to extraction to assess the efficiency of the method. These were naphthalene-d₈, acenaphthene-d₁₀, and chrysene-d₁₂. Recoveries of surrogates were within the quality control limits of 30% -150% with one exception. For Batch 1, mean recoveries of naphthalene-d₈, acenaphthene-d₁₀, and chrysene-d₁₂ were 52%, 59%, and 48%, respectively. In one sample, recovery of chrysene-d₁₂ was 28%. For Batch 2, mean recoveries of naphthalene-d₈, acenaphthene-d₁₀, and chrysene-d₁₂ were 62%, 64%, and 57%, respectively.

QA/QC SUMMARY/PAHs (continued)

MATRIX SPIKES

One sample was spiked with all PAH compounds for each batch. Matrix spike recoveries for all analytes in Batch 2 ranged from 57% to 67%. Matrix spike recoveries for all analytes in Batch 1 ranged from 26% to 73%. Six of the analytes in Batch 1 fell outside the acceptable ranges of 50% to 120%. They are 48% for fluoranthene; 47% for pyrene; 44% for benzo[a]anthracene; 38% for chrysene; 26% for benzo[b]fluoranthene; and 32% for benzo[a]pyrene. These PAHs were present at naturally elevated levels in the background sample. A blank spike was prepared with this batch and had acceptable recoveries for all target PAHs. As a result, it appears that the failure of selected PAHs to meet the recovery criteria is related to the sediment sample. The recoveries of PAHs in the MS sample for Batch 2 met the acceptance criteria.

REPLICATES

One sample was extracted in triplicate for each batch. Precision was measured by calculating the relative standard deviation (RSD) between the replicate results. The RSDs ranged from 1% to 20%, within the target precision goal of $\leq 30\%$.

SRMs

One SRM, 1941a, a marine sediment sample obtained from the National Institute of Standards and Technology, was analyzed with each batch of samples. Twelve of the 17 PAH compounds analyzed are certified at levels above the MDLs. Of these, all compounds were detected within 30% of the certified mean, with the exception of chrysene (58% and 73%), benzo[b]fluoranthene (32% and 45%), and dibenz[a,h]anthracene (63% and 40%) in both batches. Percent differences were calculated using SRM concentrations that were corrected for surrogate recovery.

REFERENCES

- Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.
- EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

TABLE A.1. Quality Assurance/Quality Control Data for Grain Size Analysis

Sediment Treatment	Gravimetric Water Content (%)	Batch No.	Total Percent (dry weight)			
			Gravel >2000 μm	Sand 62.4-2000 μm	Silt 3.9-62.4 μm	Clay <3.9 μm
R-CLIS, Replicate 1	109	1	0	6	59	35
R-CLIS, Replicate 2	109	1	0	6	60	34
R-CLIS, Replicate 3	109	1	0	6	60	34
RSD			NA ^(a)	0%	1%	2%
EC-8, Replicate 1	151	2	0	21	39	40
EC-8, Replicate 2	151	2	0	20	40	40
EC-8, Replicate 3	151	2	1	21	38	40
RSD			NA	3%	3%	0%
HU-2, Replicate 1	124	3	1	18	47	34
HU-2, Replicate 2	124	3	0	19	47	34
HU-2, Replicate 3	124	3	2	18	47	33
RSD			NA	3%	0%	2%
HU-22, Replicate 1	139	4	0	16	48	36
HU-22, Replicate 2	139	4	0	16	48	36
HU-22, Replicate 3	139	4	0	15	47	38
RSD			NA	4%	1%	3%
BU-2, Replicate 1	171	5	0	13	42	45
BU-2, Replicate 2	171	5	0	13	40	47
BU-2, Replicate 3	171	5	0	14	41	45
RSD			NA	4%	2%	3%
BC-4, Replicate 1	222	6	0	15	55	30
BC-4, Replicate 2	222	6	0	14	56	30
BC-4, Replicate 3	222	6	0	17	55	28
RSD			NA	10%	1%	4%

(a) NA. Not applicable.

TABLE A.2. Quality Assurance/Quality Control Data for Analysis of Specific Gravity and Bulk Density

Sediment Treatment	Replicate	Sample ID	Batch	Bulk Density		Specific Gravity
				Wet lbs/cu ft	Dry lbs/cu ft	
COMP HU-C	1	NY2-GRA-17	1	92	45	2.61
COMP HU-C	2	NY2-GRA-17	1	ND ^(a)	ND	2.64
RPD				NA ^(b)	NA	1%
I-Stat				NA	NA	0.01
COMP SB-A	1	NY2-GRA-1	1	83	30	2.58
COMP SB-A	2	NY2-GRA-1	1	83	30	2.56
RPD				0%	0%	1%
I-Stat				0.00	0.00	0.00
COMP GR	1	NY2-GRA-9	1	116	94	2.67
COMP GR	2	NY2-GRA-9	1	118	96	ND
RPD				2%	2%	NA
I-Stat				0.01	0.01	NA

(a) ND No data; not tested.

(b) NA Not applicable.

TABLE A.3. Quality Assurance/Quality Control Data for Analysis of TOC and Percentage of Moisture

Sediment Treatment	Batch No.	TOC (% dry wt.)
<u>Method Blanks</u>		
Blank-1	1	0.003
Blank-2	1	0.001
Blank-1	2	0.003
Blank-2	2	0.003
Blank-1	3	0.003
Blank-2	3	0.002
Blank-3	3	0.003
Blank-4	3	0.003
Blank-5	3	0.002
Blank-1	4	0.005
Blank-2	4	0.008
Blank-3	4	0.002
Blank-4	4	0.002
Blank-5	4	0.004
Blank-6	4	0.004
Blank-1	5	0.003
Blank-2	5	0.002
Blank-3	5	0.002
Blank-4	5	0.004
Blank-5	5	0.004
Blank-1	6	0.001
Blank-2	6	0.002
Blank-3	6	0.002
Blank-4	6	0.002
Blank-5	6	0.002
Blank-6	6	0.005
Blank-7	6	0.004
Blank-8	6	0.004
Blank-9	6	0.004
Blank-10	6	0.006
Blank-11	6	0.004
Blank-12	6	0.002
Blank-13	6	0.002
Blank-14	6	0.002

TABLE A.3. (contd)

Sediment Treatment	Batch No.	TOC (% dry wt.)	Percent Difference ^(a)
<u>Standard Reference Material</u>			
Non-certified Value		2.6	
SRM MESS-1	1	2.49	4%
SRM MESS-1	2	2.44	6%
SRM MESS-1	2	2.62	1%
SRM MESS-1	3	2.56	2%
SRM MESS-1	4	2.42	7%
SRM MESS-1	5	2.40	8%
SRM MESS-1	6	2.40	8%
SRM MESS-1	6	2.39	8%
SRM MESS-1	6	2.45	6%
MESS-1Y	6	2.47	
MESS-1Y, Duplicate	6	2.48	
RPD			0%

TABLE A.3. (contd)

Sediment Treatment	Batch No.	TOC (% dry wt.)	Total Percent Solids
<u>Analytical Replicates</u>			
EC-2, Replicate 1	1	1.02	66
EC-2, Replicate 2	1	1.13	66
RPD		10%	0%
GR-1, Replicate 1	1	0.12	80
GR-1, Replicate 2	1	0.13	80
RPD		8%	0%
EC-3, Replicate 1	2	1.26	75
EC-3, Replicate 2	2	1.23	75
EC-3, Replicate 3	2	1.31	75
RSD		3%	0%
HU-1, Replicate 1	3	3.17	53
HU-1, Replicate 2	3	3.13	53
HU-1, Replicate 3	3	3.30	53
RSD		3%	0%
HU-21, Replicate 1	4	3.26	44
HU-21, Replicate 2	4	3.19	44
HU-21, Replicate 3	4	3.15	44
RSD		2%	0%
HU-39, Replicate 1	5	1.95	52
HU-39, Replicate 2	5	1.95	52
HU-39, Replicate 3	5	1.88	52
RSD		2%	0%
BU-4, Replicate 1	6	3.42	37
BU-4, Replicate 2	6	3.44	37
RPD		1%	0%

(a) Percent Difference between results obtained from analysis of SRM MESS-1 and non-certified value of 2.6%. SRM MESS-1 is not certified for TOC, but according to historical analyses from Battelle's records, the estimated value is 2.6% TOC.

TABLE A.4. Quality Assurance/Quality Control Data for Metals in Sediment

Sediment Treatment	Batch	Metals ($\mu\text{g/g}$ dry wt)										
		Ag (ICP/MS)	Ag (ICP/Aqua)	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn	
<u>Method Blanks</u>												
Blank-1	1	0.119 U ^(a)	0.131	0.572 U	0.020	0.401 U	0.525 U	0.001 U	0.849 U	0.14 U	2.55 U	
Blank-2	1	0.119 U	0.119 U	0.572 U	0.020	0.401 U	0.525 U	0.001 U	0.849 U	0.41	2.55 U	
Blank-3	1	NA ^(b)	NA	NA	NA	NA	NA	0.001 U	NA	NA	NA	
Mean blank		NA	NA	NA	NA	NA	NA	NA	NA	0.2	NA	
<u>Standard Reference Material</u>												
A.6	Certified value	NC ^(c)	NC	11.6	0.36	76	18	0.063	32	28.2	138	
	Range	NC	NC	± 1.3	± 0.07	± 3	± 3	± 0.012	± 3	± 1.8	± 6	
	SRM 1646	1	0.119 U	0.275	8.72	0.331	42.7	16.4	0.074	25.4	22.7	93.6
	SRM 1646	1	0.119 U	0.136	8.89	0.350	39.9	16.1	0.079	23.5	22.4	90.6
	SRM 1646	1	NA	NA	NA	NA	NA	NA	0.077	NA	NA	NA
	SRM 1646	1	NA	NA	NA	NA	NA	NA	0.070	NA	NA	NA
	Percent Difference		NA	NA	25% ^(d)	8%	44% ^(d)	9%	17%	21% ^(d)	20%	32% ^(d)
	Percent Difference		NA	NA	23% ^(d)	3%	48% ^(d)	11%	25% ^(d)	27% ^(d)	21% ^(d)	34% ^(d)
	Percent Difference		NA	NA	NA	NA	NA	NA	22% ^(d)	NA	NA	NA
	Percent Difference		NA	NA	NA	NA	NA	NA	11%	NA	NA	NA
<u>Matrix Spike Results</u>												
EC-11/CT COMP EC-B-II	1	2.91	3.38	11.1	4.15	104	250	1.21	44.1	322	379	
EC-11/CT COMP EC-B-II MS	1	4.85	22.0	192	21.4	589	696	11.4	135	840	1140	
Concentration Recovered		1.94	18.6	181	17.3	485	446	10.2	90.9	518	761	
Amount Spiked		20.0	20.0	200	20.0	500	500	10.0	100	500	1000	
Percent Recovery		10% ^(e)	93%	90%	86%	97%	89%	102%	91%	104%	76%	

TABLE A.4. (contd)

Sediment Treatment	Batch	Metals ($\mu\text{g/g}$ dry wt)									
		Ag (ICP/MS)	Ag (ICP/Aqua)	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn
COMP HU-C	1	6.22	7.02	15.2	4.06	169	174	2.55	40.0	194	252
COMP HU-C, MS	1	6.85	25.6	193	21.4	656	612	12.1	125	715	1010
Concentration Recovered		0.63	18.6	178	17.3	487	438	9.55	85.0	521	758
Amount Spiked		20.0	20.0	200	20.0	500	500	10.0	100	500	1000
Percent Recovery		3% ^(e)	93%	89%	87%	97%	88%	96%	85%	104%	76%
<u>Analytical Replicates</u>											
EC-11/CT COMP EC-B-II, Re	1	2.78	3.36	10.9	4.26	102	248	1.27	44.6	322	375
EC-11/CT COMP EC-B-II, Re	1	3.05	3.44	11.3	4.04	107	254	1.18	44.6	333	383
EC-11/CT COMP EC-B-II, Re	1	2.91	3.33	11.1	4.15	103	248	1.19	43.1	312	378
RSD		5%	2%	2%	3%	3%	1%	4%	2%	3%	1%
COMP HU-C, Replicate 1	1	6.10	7.05	15.2	4.05	171	174	2.57	40.3	196	247
COMP HU-C, Replicate 2	1	6.05	7.03	15.5	4.11	167	173	2.66	39.4	193	253
COMP HU-C, Replicate 3	1	6.51	6.98	15.0	4.02	170	175	2.42	40.3	194	257
RSD		4%	1%	2%	1%	1%	1%	5%	1%	1%	2%

(a) U Undetected at or above given concentration.

(b) NA Not applicable.

(c) NC Not certified.

(d) Outside quality control criteria ($\pm 20\%$) for SRMs.

(e) Outside quality control criteria (75-125%) for matrix spike recoveries.

TABLE A.5. Quality Assurance/Quality Control Data for Additional Metals in Sediment

Sediment Treatment	Metals ($\mu\text{g/g}$ dry wt)			
	Be ICP/MS	Sb ICP/MS	Se GFAA	Tl ICP/MS
Method Blanks				
Blank-1	0.5 U ^(a)	0.124	0.13 U	0.024 U
Blank-2	0.5 U	0.030 U	0.13 U	0.024 U
Standard Reference Material				
Certified Value	NC ^(b)	NC	NC	NC
Range	NA ^(c)	NA	NA	NA
Non Certified Value	1.5	0.4	0.6	0.5
1646	1.02	0.300	0.41	0.305
1646	0.912	0.200	0.42	0.322
Percent Difference from Certified value	NA	NA	NA	NA
Percent Difference from Certified value	NA	NA	NA	NA
Matrix Spike Results				
EC-11/CT COMP EC-B-II	NA	0.15	0.21	NA
EC-11/CT COMP EC-B-II MS	NA	2.43	2.89	NA
Amount Recovered	NA	2.28	2.68	NA
Amount Spiked	NS ^(d)	1.00	2.50	NS
Percent Recovery	NA	228% ^(e)	107%	NA
EC-11/CT COMP EC-B-II	0.953	NA	NA	0.461
EC-11/CT COMP EC-B-II MS	4.99	NA	NA	4.68
Amount Recovered	4.04	NA	NA	4.68
Amount Spiked	5.00	NS	NS	5.00
Percent Recovery	81%	NA	NA	94%
Analytical Replicates				
EC-11/CT COMP EC-B-II, Rep 1	0.959	1.52	0.70	0.423
EC-11/CT COMP EC-B-II, Rep 2	0.955	1.46	0.83	0.440
EC-11/CT COMP EC-B-II, Rep 3	0.903	1.48	0.89	0.445
RSD	3%	2%	12%	3%

(a) U Undetected at or above given concentration.

(b) NC Non-certified value.

(c) NA Not applicable.

(d) NS Not spiked.

(e) Outside quality control criteria (75-125%) for matrix spike recoveries.

TABLE A.6. Quality Control Data for 1,4-Dichlorobenzene, Pesticides, and PCB Congeners in Sediment

	Batch: Treatment: Blank	MATRIX SPIKE					
		1	1	1	1	1	1
		EC-10	EC-10, MS	Concentration Recovered	Amount Spiked	Concentration Spiked	Percent Recovery
Sample Size (g)	9.076 ^(a)	6.689	2.289	NA ^(b)	NA	NA	
Units (dry wt) :	µg/kg	µg/kg	µg/kg	µg/kg	ng	µg/kg	
1,4-Dichlorobenzene	1.19 U ^(c)	84.46	510.36	425.91	1425	623	68
2,4-DDD	0.97 U	16.57	18.72	2.15	NS ^(d)	NS	NA
2,4-DDT	0.91 U	NA	NA	NA	NS	NS	NA
4,4-DDD	1.56 U	53.31	154.73	101.42	201.0	88	115
4,4-DDE	2.29 U	38.55	117.11	78.56	200.5	88	90
4,4-DDT	5.19 U	2.19 J ^(e)	74.76	72.56	200.5	88	83
Aldrin	0.87 U	1.18 U	58.05	58.05	200.5	88	66
alpha-Chlordane	1.27 U	14.46	85.02	70.56	200.0	87	81
Dieldrin	1.85 U	8.52	66.86	58.34	200.5	88	67
Endosulfan I /2,4-DDE	2.39 U	3.24 U	73.57	73.57	200.5	88	84
Endosulfan II	1.78 U	2.42 U	72.03	72.03	200.5	88	82
Endosulfan Sulfate	1.68 U	2.28 U	86.48	86.48	200.5	88	99
Endrin ^(f)	3.24 U	4.40 U	78.26	78.26	200.0	87	90
Endrin Aldehyde ^(f)	1.93 U	2.62 U	66.18	66.18	200.5	88	76
Heptachlor	1.96 U	2.65 U	87.96	87.96	200.5	88	100
Heptachlor Epoxide	1.09 U	1.47 U	81.04	81.04	200.5	88	93
alpha-BHC ^(f)	1.21 U	0.28 J	69.22	68.94	200.5	88	79
beta-BHC ^(f)	0.09 J	2.42 U	64.97	64.97	200.5	88	74
delta-BHC ^(f)	1.20 J	2.20 U	68.21	68.21	200.5	88	78
Lindane ^(f)	0.33 J	1.92 U	72.05	72.05	200.5	88	82
Methoxychlor ^(f)	2.03 U	2.75 U	94.68	94.68	200.0	87	108
Toxaphene ^(f)	61.41 U	83.32 U	NA	NA	NS	NS	NA
trans-Nonachlor	1.86 U	7.45	5.57	5.57	NS	NS	NA
CL2(08)	4.38 U	6.47	28.20	21.74	200.00	87	25 ^(g)
CL3(18)	2.78 U	26.86	98.05	71.18	200.00	87	81
CL3(28)	1.83 U	42.91	148.46	105.55	200.00	87	121 ^(g)
CL4(44)	2.65 U	43.52	118.73	75.21	200.00	87	86
CL4(49)	1.66 U	34.91	44.50	9.60	NS	NS	NA
CL4(52)	1.54 U	51.61	122.53	70.92	200.00	87	81
CL4(66)	1.45 U	59.60	158.19	98.58	200.00	87	113
CL5(87)	0.88 U	13.96	15.20	1.24	NS	NS	NA
CL5(101)	0.74 U	33.21	98.14	64.93	200.00	87	74
CL5(105)	0.49 U	12.92	85.99	73.07	200.00	87	84
CL5(118)	1.30 U	28.18	87.87	59.69	200.00	87	68
CL6(128)	1.38 U	5.45	82.99	77.54	200.00	87	89
CL6(138)	1.19 U	31.64	101.08	69.45	200.00	87	79
CL6(153)	5.77 U	26.37	91.20	64.83	200.00	87	74
CL7(170)	1.46 U	17.20	88.02	70.82	200.00	87	81
CL7(180)	0.98 U	31.37	96.83	65.45	200.00	87	75
CL7(183)	1.09 U	4.97	NA	NA	NS	NS	NA
CL7(184)	1.09 U	0.49 J	NA	NA	NS	NS	NA
CL7(187)	0.82 U	15.44	70.69	55.25	200.00	87	63
CL8(195)	1.24 U	6.36	76.77	70.41	200.00	87	81
CL9(206)	1.90 U	14.96	90.94	75.98	200.00	87	87
CL10(209)	1.18 U	9.42	90.27	80.85	200.00	87	93
<u>Surrogate Recoveries (%)</u>							
DBOFB	73	82	86	NA	NA	NA	NA
CL5(112)	64	55	67	NA	NA	NA	NA

TABLE A.6. (contd)

	Batch: 2 Treatment: Blank	MATRIX SPIKE					
		2	2	2	2	2	2
		R-MUD	R-MUD, MS	Concentration	Amount	Concentration	
				Recovered	Spiked	Spiked	Percent
Sample Size (g)	8.542 ^(a)	13.660	13.220	NA	NA	NA	
Units (dry wt) :	µg/kg	µg/kg	µg/kg	µg/kg	ng	µg/kg	Percent Recovery
1,4-Dichlorobenzene	1.27 U	0.79 U	61.78	61.78	1425.00	108	57
2,4-DDD	1.04 U	0.01 J	NA	NA	NS	NS	NA
2,4-DDT	0.97 U	0.60 U	NA	NA	NS	NS	NA
4,4-DDD	1.65 U	0.06 J	11.72	11.66	201.00	15	77
4,4-DDE	2.43 U	0.01 J	10.08	10.07	200.50	15	66
4,4-DDT	5.51 U	3.45 U	10.99	10.99	200.50	15	72
Aldrin	0.93 U	0.58 U	11.35	11.35	200.50	15	75
alpha-Chlordane	1.35 U	0.01 J	11.39	11.39	200.00	15	75
Dieldrin	1.97 U	0.21 J	11.34	11.13	200.50	15	73
Endosulfan I /2,4-DDE	2.54 U	1.59 U	13.52	13.52	200.50	15	89
Endosulfan II	1.89 U	0.05 J	13.24	13.19	200.50	15	87
Endosulfan Sulfate	1.79 U	1.12 U	10.86	10.86	200.50	15	72
Endrin ^(f)	NA	NA	NA	NA	NS	NS	NA
Endrin Aldehyde ^(f)	NA	NA	NA	NA	NS	NS	NA
Heptachlor	2.08 U	1.30 U	10.27	10.27	200.50	15	68
Heptachlor Epoxide	1.15 U	0.72 U	10.60	10.60	200.50	15	70
alpha-BHC ^(f)	NA	NA	NA	NA	NS	NS	NA
beta-BHC ^(f)	NA	NA	NA	NA	NS	NS	NA
delta-BHC ^(f)	NA	NA	NA	NA	NS	NS	NA
Lindane ^(f)	NA	NA	NA	NA	NS	NS	NA
Methoxychlor ^(f)	NA	NA	NA	NA	NS	NS	NA
Toxaphene ^(f)	NA	NA	NA	NA	NS	NS	NA
trans-Nonachlor	1.98 U	0.00 J	NA	NA	NS	NS	NA
CL2(08)	4.65 U	2.91 U	7.05	7.05	200.00	15	47 ^(g)
CL3(18)	2.95 U	1.85 U	8.12	8.12	200.00	15	54
CL3(28)	1.94 U	1.21 U	10.03	10.03	200.00	15	66
CL4(44)	2.82 U	0.22 J	10.29	10.07	200.00	15	67
CL4(49)	1.76 U	0.04 J	NA	NA	NS	NS	NA
CL4(52)	1.63 U	0.06 J	9.91	9.85	200.00	15	65
CL4(66)	1.54 U	0.04 J	10.43	10.39	200.00	15	69
CL5(87)	0.93 U	0.05 J	NA	NA	NS	NS	NA
CL5(101)	0.78 U	0.04 J	10.27	10.23	200.00	15	68
CL5(105)	0.52 U	0.03 J	9.12	9.09	200.00	15	60
CL5(118)	1.38 U	0.02 J	9.25	9.23	200.00	15	61
CL6(128)	1.46 U	0.92 U	9.42	9.42	200.00	15	62
CL6(138)	1.26 U	0.07 J	9.36	9.29	200.00	15	61
CL6(153)	6.13 U	0.03 J	8.56	8.53	200.00	15	56
CL7(170)	1.55 U	0.97 U	9.26	9.26	200.00	15	61
CL7(180)	1.04 U	0.65 U	9.32	9.32	200.00	15	62
CL7(183)	1.15 U	0.72 U	NA	NA	NS	NS	NA
CL7(184)	1.15 U	0.01 J	NA	NA	NS	NS	NA
CL7(187)	0.87 U	0.01 J	9.28	9.27	200.00	15	61
CL8(195)	1.32 U	0.83 U	9.35	9.35	200.00	15	62
CL9(206)	2.02 U	1.26 U	9.13	9.13	200.00	15	60
CL10(209)	1.26 U	0.79 U	9.41	9.41	200.00	15	62
<u>Surrogate Recoveries (%)</u>							
DBOFB	66	65	69	NA	NA	NA	
CL5(112)	72	49	64	NA	NA	NA	

TABLE A.6. (contd)

	STANDARD REFERENCE MATERIAL					
	Batch:	1	1	1	2	2
	Treatment:	SRM	Certified	Percent	SRM	Certified
	Sample Size (g)	5.133	Value	Difference ^(h)	5.057	Value
Units (dry wt) :	µg/kg	µg/kg		µg/kg	µg/kg	Difference
1,4-Dichlorobenzene	NA	NC ⁽ⁱ⁾	NA	NA	NC	NA
2,4-DDD	NA	NC	NA	NA	NC	NA
2,4-DDT	NA	NC	NA	NA	NC	NA
4,4-DDD	2.56 J	5.06	4	4.86	5.06	103
4,4-DDE	3.46 J	6.59	8	3.16 J	6.59	1
4,4-DDT	NA	NC	NA	NA	NC	NA
Aldrin	NA	NC	NA	NA	NC	NA
alpha-Chlordane	1.01 J	2.33	44	1.06 J	2.33	14
Dieldrin	NA	NC	NA	NA	NC	NA
Endosulfan I /2,4-DDE	C ⁽ⁱ⁾	0.73	NA	ND	0.73	NA
Endosulfan II	NA	NC	NA	NA	NC	NA
Endosulfan Sulfate	NA	NC	NA	NA	NC	NA
Endrin ^(f)	NA	NC	NA	NA	NC	NA
Endrin Aldehyde ^(f)	NA	NC	NA	NA	NC	NA
Heptachlor	NA	NC	NA	NA	NC	NA
Heptachlor Epoxide	NA	NC	NA	NA	NC	NA
alpha-BHC ^(f)	NA	NC	NA	NA	NC	NA
beta-BHC ^(f)	NA	NC	NA	NA	NC	NA
delta-BHC ^(f)	NA	NC	NA	NA	NC	NA
Lindane ^(f)	NA	NC	NA	NA	NC	NA
Methoxychlor ^(f)	NA	NC	NA	NA	NC	NA
Toxaphene ^(f)	NA	NC	NA	NA	NC	NA
trans-Nonachlor	0.39 J	1.26	61	0.60 J	1.26	10
CL2(08)	NA	NC	NA	NA	NC	NA
CL3(18)	NA	NC	NA	NA	NC	NA
CL3(28)	NA	NC	NA	NA	NC	NA
CL4(44)	3.88 J	4.80	4	3.92 J	4.80	54
CL4(49)	3.03	9.50	59	3.14 J	9.50	38
CL4(52)	3.20	6.89	40	3.89	6.89	6
CL4(66)	7.11	6.80	34	6.07	6.80	68
CL5(87)	1.45 J	6.70	55	1.72	6.70	46
CL5(101)	9.02	11.00	5	6.94	11.00	19
CL5(105)	1.18	3.65	33	1.05	3.65	39
CL5(118)	3.29	10.00	32	3.55	10.00	25
CL6(128)	3.07	1.87	238	1.82 J	1.87	106
CL6(138)	4.96	13.38	24	6.05	13.38	4
CL6(153)	5.21 J	17.60	39	5.21 J	17.60	37
CL7(170)	4.82	3.00	230	C	3.00	NA
CL7(180)	5.47	5.83	93	5.10	5.83	85
CL7(183)	NA	NC	NA	NA	NC	NA
CL7(184)	NA	NC	NA	NA	NC	NA
CL7(187)	NA	NC	NA	NA	NC	NA
CL8(195)	NA	NC	NA	NA	NC	NA
CL9(206)	C	3.67	NA	2.93 J	3.67	69
CL10(209)	7.52	8.34	85	5.26	8.34	33
<u>Surrogate Recoveries (%)</u>						
DBOFB	78	NA	NA	53	NA	NA
CL5(112)	49	NA	NA	47	NA	NA

TABLE A.6. (contd)

	TRIPPLICATE ANALYSES								
	Batch:	1	1	1		2	2	2	
	Treatment:	EC-15	EC-15	EC-15		GR-10	GR-10	GR-10	
	Sample Size (g)	Replicate 1	Replicate 2	Replicate 3	RSD(%)	Replicate 1	Replicate 2	Replicate 3	RSD(%)
Units (dry wt) :	µg/kg	µg/kg	µg/kg		µg/kg	µg/kg	µg/kg		
1,4-Dichlorobenzene	10.65	8.00	7.52	19	17.73	25.25	19.82	19	
2,4-DDD	10.32	13.52	10.13	17	6.58	9.27	6.64	21	
2,4-DDT	0.84 U	0.87 U	0.88 U	NA	1.01 U	0.96 U	0.95 U	NA	
4,4-DDD	41.51	47.84	42.18	8	5.56	6.05	5.52	5	
4,4-DDE	13.20	12.90	10.14	14	4.58	5.53	5.01	9	
4,4-DDT	2.35 J	4.25 J	2.57 J	34	0.38 J	0.19 J	0.16 J	48	
Aldrin	0.80 U	0.84 U	0.85 U	NA	0.97 U	0.92 U	0.91 U	NA	
alpha-Chlordane	18.62	23.16	22.52	11	1.02 J	1.41	1.09 J	18	
Dieldrin	7.09	7.58	6.22	10	1.27 J	1.35 J	1.46 J	7	
Endosulfan I /2,4-DDE	2.20 U	2.30 U	2.32 U	NA	2.65 U	2.52 U	2.51 U	NA	
Endosulfan II	1.64 U	1.71 U	1.73 U	NA	1.38 J	1.77 J	0.97 J	29	
Endosulfan Sulfate	1.55 U	1.62 U	1.64 U	NA	0.31 J	0.44 J	0.28 J	25	
Endrin ^(f)	2.98 U	3.11 U	3.15 U	NA	NA	NA	NA	NA	
Endrin Aldehyde ^(f)	1.78 U	1.86 U	1.88 U	NA	NA	NA	NA	NA	
Heptachlor	1.80 U	1.88 U	1.90 U	NA	2.17 U	2.07 U	2.05 U	NA	
Heptachlor Epoxide	1.00 U	1.04 U	1.05 U	NA	1.20 U	1.15 U	1.14 U	NA	
alpha-BHC ^(f)	1.11 U	1.16 U	1.17 U	NA	NA	NA	NA	NA	
beta-BHC ^(f)	1.64 U	1.71 U	1.73 U	NA	NA	NA	NA	NA	
delta-BHC ^(f)	1.49 U	1.56 U	1.58 U	NA	NA	NA	NA	NA	
Lindane ^(f)	1.30 U	1.36 U	1.37 U	NA	NA	NA	NA	NA	
Methoxychlor ^(f)	1.87 U	1.95 U	1.97 U	NA	NA	NA	NA	NA	
Toxaphene ^(f)	56.56 U	59.03 U	59.68 U	NA	NA	NA	NA	NA	
trans-Nonachlor	11.31	14.64	14.13	13	0.54 J	0.66 J	0.53 J	12	
CL2(08)	7.98	8.19	6.21	15	2.53 J	2.95 J	2.64 J	8	
CL3(18)	19.18	23.08	22.08	9	3.81	4.43	4.15	7	
CL3(28)	51.14	30.02	31.95	31 ^(k)	13.08	17.79	14.05	17	
CL4(44)	24.24	31.36	29.22	13	5.15	6.44	5.42	12	
CL4(49)	23.21	27.19	24.75	8	5.38	7.00	6.50	13	
CL4(52)	29.20	41.52	36.00	17	6.66	8.07	6.98	10	
CL4(66)	88.09	103.82	92.36	9	10.53	11.61	9.40	10	
CL5(87)	5.33	7.44	6.83	17	1.78	2.11	1.90	8	
CL5(101)	24.93	29.25	28.42	8	5.15	6.22	5.24	11	
CL5(105)	4.86	41.07	7.37	114 ^(k)	2.29	2.35	1.85	13	
CL5(118)	13.11	16.42	15.16	11	4.74	6.11	5.26	13	
CL6(128)	4.50	6.23	7.30	24	2.96	3.47	3.17	8	
CL6(138)	67.37	36.36	24.29	52 ^(k)	5.60	7.00	6.08	11	
CL6(153)	12.25	10.68	12.57	9	4.21 J	5.46 J	5.04 J	13	
CL7(170)	9.06	9.86	8.44	8	2.11	2.81	2.31	15	
CL7(180)	9.43	12.62	10.25	15	3.04	3.82	3.20	12	
CL7(183)	1.45	2.28	2.07	22	0.60 J	0.89 J	0.73 J	19	
CL7(184)	1.19	0.79 J	0.42 J	48	0.38 J	0.36 J	0.45 J	11	
CL7(187)	3.29	4.79	3.73	20	1.61	2.04	1.72	12	
CL8(195)	1.57	2.03	1.59	15	0.35 J	0.41 J	0.37 J	8	
CL9(206)	4.73	5.62	4.95	9	0.74 J	1.07 J	0.86 J	19	
CL10(209)	4.10	5.87	4.75	18	1.27 J	1.49	1.49	9	
<u>Surrogate Recoveries (%)</u>									
DBOFB	84	94	85	NA	50	63	58	NA	
CL5(112)	34	43	34	NA	39	50	44	NA	

TABLE A.6. (contd)

Qualifiers

- (a) Sample concentration of the procedural blank adjusted for the average sample size of the batch.
- (b) NA Not applicable.
- (c) U Undetected at or above given concentration.
- (d) NS Not spiked.
- (e) J Concentration estimated; analyte detected below method detection limit (MDL), but above instrument detection limit (IDL).
- (f) Analyte required only in samples designated for Central Long Island Disposal Testing Site.
- (g) Outside quality control criteria (50-120%) for matrix spike recoveries.
- (h) Percent Difference from certified
= absolute value [(certified value, $\mu\text{g}/\text{kg}$ - value detected corrected for surrogate recovery, $\mu\text{g}/\text{kg}$) / certified value, $\mu\text{g}/\text{kg}$].
- (i) NC No certified value available.
- (j) C Analyte not determined due to co-eluting peak.
- (k) Outside quality control criteria ($\pm 30\%$) for replicates.

TABLE A.7. MDL Verification Study for Analysis of Pesticides and PCBs in Sediment

Battelle ID:	OG99	OH01	OH02	OH03	OH04	OH05	OH06	OH07	Standard Deviation STD (n-1)	Method Detection Limit MDL ^(a) µg/kg	Method Detection Limit MDL (ng)
Sample Size (g):	20.919	19.455	19.201	18.645	19.087	19.434	18.896	18.612			
Units (dry wt):	µg/kg	µg/kg	µg/kg	µg/kg	µg/kg	µg/kg	µg/kg	µg/kg			
1,4-Dichlorobenzene	1.934	1.589	1.642	1.966	1.820	1.483	1.965	2.685	0.372	1.114	21.485
2,4-DDD	NS ^(b)	NS	NS	NS	NS	NS	NS	NS	NA ^(b)	NA	NA
2,4-DDT	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA
4,4-DDD	0.494	0.516	0.533	0.637	0.526	0.570	0.453	0.503	0.055	0.165	3.180
4,4-DDE	0.380	0.433	0.422	0.477	0.464	0.462	0.451	0.456	0.031	0.093	1.791
4,4-DDT	0.853	0.455	0.487	0.474	0.515	0.546	0.498	0.499	0.129	0.387	7.460
Aldrin	0.379	0.460	0.443	0.502	0.459	0.431	0.450	0.477	0.036	0.108	2.077
Alpha-chlordane	0.344	0.427	0.375	0.471	0.435	0.413	0.438	0.440	0.040	0.121	2.328
Dieldrin	0.400	0.451	0.478	0.493	0.456	0.499	0.465	0.441	0.032	0.095	1.836
Endosulfan I	0.423	0.556	0.480	0.562	0.531	0.506	0.517	0.540	0.045	0.136	2.628
Endosulfan II	0.500	0.538	0.544	0.575	0.552	0.558	0.529	0.526	0.023	0.068	1.319
Endosulfan Sulfate	0.416	0.426	0.448	0.476	0.463	0.489	0.473	0.462	0.025	0.076	1.458
Endrin ^(c)	0.381	0.490	0.512	0.557	0.552	0.550	0.540	0.549	0.059	0.178	3.439
Endrin Aldehyde ^(c)	0.425	0.534	0.532	0.619	0.568	0.528	0.558	0.578	0.056	0.169	3.258
Heptachlor	0.445	0.516	0.476	0.581	0.527	0.480	0.528	0.549	0.040	0.119	2.298
Heptachlor epoxide	0.442	0.542	0.495	0.572	0.549	0.514	0.543	0.560	0.042	0.127	2.444
A-BHC ^(d)	0.342	0.415	0.428	0.450	0.433	0.384	0.415	0.433	0.034	0.103	1.985
B-BHC ^(d)	0.442	0.547	0.539	0.541	0.495	0.493	0.513	0.504	0.035	0.104	1.996
D-BHC ^(d)	0.429	0.537	0.489	0.510	0.532	0.473	0.491	0.485	0.034	0.103	1.989
Lindane ^(e)	0.386	0.477	0.457	0.482	0.458	0.431	0.452	0.460	0.030	0.091	1.745
Methoxychlor ^(f)	0.319	0.446	0.497	0.489	0.530	0.553	0.561	0.554	0.081	0.242	4.673
Toxaphene ^(g)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA
Trans-nonachlor	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA

A.14

TABLE A.7. (contd)

Battelle ID:	OG99	OH01	OH02	OH03	OH04	OH05	OH06	OH07	Standard Deviation STD (n-1)	Method Detection Limit MDL ^(a) µg/kg	Method Detection Limit MDL (ng)
Sample Size (g): Units (dry wt):	20.919 µg/kg	19.455 µg/kg	19.201 µg/kg	18.645 µg/kg	19.087 µg/kg	19.434 µg/kg	18.896 µg/kg	18.612 µg/kg			
CL2(08)	0.273	0.302	0.289	0.319	0.244	0.312	0.319	0.378	0.039	0.117	2.265
CL3(18)	0.376	0.447	0.416	0.489	0.452	0.415	0.423	0.445	0.034	0.100	1.937
CL3(28)	0.376	0.491	0.465	0.482	0.439	0.463	0.459	0.486	0.037	0.112	2.155
CL4(44)	0.425	0.529	0.478	0.551	0.506	0.470	0.489	0.511	0.039	0.116	2.243
CL4(49)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA
CL4(52)	0.418	0.491	0.442	0.522	0.471	0.426	0.450	0.473	0.035	0.104	2.000
CL4(66)	0.423	0.526	0.487	0.519	0.493	0.436	0.477	0.490	0.036	0.108	2.087
CL5(87)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA
CL5(101)	0.459	0.587	0.530	0.597	0.551	0.501	0.519	0.532	0.045	0.134	2.581
CL5(105)	0.373	0.381	0.416	0.459	0.405	0.435	0.423	0.421	0.028	0.084	1.618
CL5(118)	0.399	0.479	0.454	0.486	0.463	0.469	0.460	0.467	0.027	0.080	1.534
CL6(128)	0.363	0.414	0.401	0.394	0.400	0.404	0.385	0.401	0.015	0.046	0.887
CL6(138)	0.379	0.422	0.411	0.421	0.418	0.410	0.407	0.417	0.014	0.042	0.806
CL6(153)	0.359	0.416	0.418	0.437	0.430	0.414	0.402	0.414	0.024	0.071	1.378
CL7(170)	0.343	0.402	0.376	0.407	0.394	0.384	0.378	0.380	0.020	0.060	1.149
CL7(180)	0.341	0.384	0.380	0.430	0.426	0.397	0.395	0.390	0.028	0.084	1.622
CL7(183)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA
CL7(184)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA	NA
CL7(187)	0.329	0.384	0.358	0.421	0.400	0.403	0.391	0.378	0.029	0.086	1.654
CL8(195)	0.328	0.367	0.364	0.397	0.390	0.382	0.381	0.371	0.021	0.064	1.227
CL9(206)	0.267	0.303	0.314	0.326	0.328	0.305	0.277	0.299	0.022	0.065	1.256
CL10(209)	0.359	0.399	0.402	0.448	0.447	0.430	0.437	0.425	0.030	0.090	1.738
<u>Surrogate Recoveries (%)</u>											
DBOFB	55	67	58	66	64	61	63	65			
CL5(112)	58	63	61	67	64	67	62	61			

(a) MDL The Method Detection Limit (2.998 x standard deviation).

(b) NS Not spiked.

(c) NA Not applicable.

(d) Analyte required only in samples designated for Central Long Island Disposal Testing Site.

TABLE A.8. Quality Control Data for Polynuclear Aromatic Hydrocarbons (PAH) in Sediment

	BLANKS		MATRIX SPIKE							
	1	2	1	1	2		2			
Batch:	Blank	Blank	EC-10	EC-10, MS	R-MUD		R-MUD, MS			
Treatment:	Blank	Blank	EC-10	EC-10, MS	R-MUD		R-MUD, MS			
Percent Moisture:	NA ^(a)	NA	56.369	19.842	Concentration		Concentration			
Dry Weight (g)	9.076 ^(b)	8.542	6.689	2.289	Spiked	Percent	13.655	13.216	Spiked	Percent
Units (dry wt):	µg/kg	µg/kg	µg/kg	µg/kg	µg/kg	Recovery	µg/kg	µg/kg	µg/kg	Recovery
naphthalene	12.36 U ^(c)	0.73 J ^(d)	293.40	1949.96	2595.02	64	1.13 J	280.30	449	62
1-methylnaphthalene ^(e)	13.00 U	NA	95.73	1781.30	2575.36	65	NA	NA	NS ^(f)	NA
2-methylnaphthalene ^(e)	10.96 U	NA	190.08	1754.99	NS	NA	NA	NA	NS	NA
biphenyl	10.45 U	11.10 U	64.14	1699.62	2588.69	63	6.94 U	285.69	448	64
2,6-dimethylnaphthalene ^(e)	10.21 U	NA	89.93	1798.88	2579.29	66	NA	NA	NS	NA
acenaphthylene	9.94 U	10.57 U	392.81	2109.65	2484.93	69	6.61 U	275.33	430	64
acenaphthene	12.93 U	13.74 U	199.96	1884.07	2681.52	63	8.59 U	299.51	464	64
fluorene	10.69 U	11.36 U	234.41	1876.21	2570.55	64	7.11 U	271.59	445	61
phenanthrene	10.78 U	11.45 U	1129.33	2727.93	2584.10	62	0.72 J	285.68	448	64
anthracene	10.46 U	11.12 U	839.49	2036.08	1956.09	61	6.96 U	211.15	339	62
1-methylphenanthrene ^(e)	9.57 U	NA	343.98	2220.41	2555.70	73	NA	NA	NS	NA
fluoranthene	9.72 U	10.32 U	4118.64	5351.78	2594.15	48 ^(g)	0.53 J	288.99	449	64
pyrene	2.83 J	12.46 U	4171.38	5396.57	2590.65	47 ^(g)	0.55 J	286.47	449	64
benz[a]anthracene	11.56 U	12.29 U	2017.45	3005.59	2245.09	44 ^(g)	0.62 J	230.70	389	59
chrysene	14.17 U	15.06 U	2535.99	3529.16	2602.88	38 ^(g)	9.42 U	291.13	451	65
benzo[b]fluoranthene	10.68 U	11.34 U	3396.16	4074.64	2582.35	26 ^(g)	0.50 J	277.16	447	62
benzo[k]fluoranthene	12.66 U	13.46 U	780.34	2498.31	2572.30	67	8.42 U	298.83	446	67
benzo[e]pyrene ^(e)	7.98 U	NA	1244.09	2852.72	2582.79	62	NA	NA	NS	NA
benzo[a]pyrene	9.90 U	10.52 U	2397.66	3136.38	2332.46	32 ^(g)	6.58 U	231.13	404	57
perylene ^(e)	20.84 U	NA	381.92	1587.57	1953.69	62	NA	NA	NS	NA
indeno[1,2,3-c,d]pyrene	8.55 U	9.08 U	1408.83	2781.05	2292.27	60	5.68 U	239.58	397	60
dibenz[a,h]anthracene	8.68 U	9.22 U	355.49	1583.39	1938.40	63	5.77 U	205.26	336	61
benzo[g,h,i]perylene	7.18 U	7.63 U	1349.43	2656.07	2307.99	57	4.77 U	231.76	400	58
<u>Surrogate Recoveries (%)</u>										
naphthalene-d8	59	69	53	55	NA	NA	54	66	NA	NA
acenaphthene-d10	63	66	60	59	NA	NA	56	63	NA	NA
chrysene-d12	65	63	52	55	NA	NA	58	64	NA	NA

A.16

TABLE A.8. (contd)

Batch: Treatment: Dry Weight (g) Units (dry wt):	STANDARD REFERENCE MATERIAL					
	1			2		
	NIST 1941a	NIST 1941a	Percent Difference ^(h)	NIST 1941a	NIST 1941a	Percent Difference ^(h)
	Certified	Value		Certified	Value	
Value	5.133	µg/kg	5.057	µg/kg		
naphthalene	1010	446.35	2	1010	461.60	10
1-methylnaphthalene ^(e)	NC ^(f)	69.83	NA	NA	NA	NA
2-methylnaphthalene ^(e)	NC	149.85	NA	NA	NA	NA
biphenyl	NC	45.65	NA	NC	45.92	NA
2,6-dimethylnaphthalene ^(e)	NC	33.39	NA	NA	NA	NA
acenaphthylene	NC	50.40	NA	NC	43.38	NA
acenaphthene	NC	23.36	NA	NC	24.71	NA
fluorene	97.3	49.71	18	97	47.87	3
phenanthrene	489	274.57	12	489	275.27	6
anthracene	184	115.14	24	184	114.23	17
1-methylphenanthrene ^(e)	NC	59.14	NA	NA	NA	NA
fluoranthene	981	558.33	13	981	523.89	1
pyrene	811	465.23	14	811	439.33	2
benz[a]anthracene	427	228.99	7	427	208.24	8
chrysene	380	330.74	73 ^(g)	380	318.66	58 ^(g)
benzo[b]fluoranthene	740	540.68	45 ^(g)	740	519.11	32 ^(g)
benzo[k]fluoranthene	361	186.68	3	361	192.57	1
benzo[e]pyrene ^(e)	553	291.70	5	NA	NA	NA
benzo[a]pyrene	628	277.29	12	628	291.97	12
perylene ^(e)	452	202.39	11	NA	NA	NA
indeno[1,2,3-c,d]pyrene	501	264.41	5	501	248.25	6
dibenz[a,h]anthracene	73.9	60.42	63 ^(g)	74	54.65	40 ^(g)
benzo[g,h,i]perylene	525	249.44	6	525	233.31	16
<u>Surrogate Recoveries (%)</u>						
naphthalene-d8	NA	43	NA	NA	51	NA
acenaphthene-d10	NA	50	NA	NA	53	NA
chrysene-d12	NA	51	NA	NA	55	NA

TABLE A.8. (contd)

Batch: Treatment:	ANALYTICAL REPLICATES							
	1	1	1		2	2	2	
	EC-15 Replicate 1	EC-15 Replicate 2	EC-15 Replicate 3		GR-10 Replicate 1	GR-10 Dup. Replicate 2	GR-10 Trip. Replicate 3	
Dry Weight (g)	9.854	9.442	9.339		8.182	8.594	8.657	
Units (dry wt):	µg/kg	µg/kg	µg/kg	RSD(%)	µg/kg	µg/kg	µg/kg	RSD(%)
naphthalene	413.07	383.64	346.57	9	97.15	122.54	106.28	12
1-methylnaphthalene ^(e)	230.13	293.43	294.48	14	NA	NA	NA	NA
2-methylnaphthalene ^(e)	220.96	269.92	256.05	10	NA	NA	NA	NA
biphenyl	67.60	81.01	101.32	20	21.24	27.72	23.89	13
2,6-dimethylnaphthalene ^(e)	141.18	161.94	151.86	7	ND	ND	ND	NA
acenaphthylene	350.59	356.12	360.45	1	72.16	85.43	82.68	9
acenaphthene	393.29	494.18	516.99	14	27.42	37.65	34.18	16
fluorene	496.91	588.49	564.89	9	51.78	69.28	58.58	15
phenanthrene	2775.66	3308.85	2624.88	12	293.40	391.80	305.88	16
anthracene	784.75	917.38	820.41	8	228.03	286.56	241.40	12
1-methylphenanthrene ^(e)	480.83	521.03	513.57	4	NA	NA	NA	NA
fluoranthene	4967.01	5744.20	5225.88	7	809.42	996.86	801.60	13
pyrene	4698.65	5597.13	5124.00	9	877.93	1063.72	851.74	12
benzo[a]anthracene	2158.28	2538.62	2480.41	9	492.12	601.96	493.70	12
chrysene	2530.60	2939.22	2913.86	8	502.85	603.94	493.96	11
benzo[b]fluoranthene	2953.82	3554.01	3284.14	9	572.66	705.11	577.73	12
benzo[k]fluoranthene	678.98	661.98	723.19	5	221.94	269.23	228.73	11
benzo[e]pyrene ^(e)	1586.76	1869.29	1743.18	8	NA	NA	NA	NA
benzo[a]pyrene	2154.13	2586.21	2437.45	9	518.39	627.31	524.98	11
perylene ^(e)	380.77	395.37	445.44	8	NA	NA	NA	NA
indeno[1,2,3-c,d]pyrene	1507.35	1811.51	1634.00	9	276.94	335.32	284.94	11
dibenz[a,h]anthracene	371.68	394.26	398.09	4	71.13	90.76	75.94	13
benzo[g,h,i]perylene	1365.92	1673.81	1530.19	10	249.71	298.49	254.12	10
<u>Surrogate Recoveries (%)</u>								
naphthalene-d8	52	61	55	NA	41	52	46	NA
acenaphthene-d10	57	68	59	NA	47	58	51	NA
chrysene-d12	39	44	41	NA	46	56	49	NA

TABLE A.8. (contd)

Qualifiers

- (a) NA Not applicable.
- (b) Sample concentration of the procedural blank adjusted for the average sample size of the batch.
- (c) U Undetected at or above given concentration.
- (d) J Concentration estimated; analyte detected below method detection limit (MDL), but above instrument detection limit (IDL).
- (e) Analyte required only in samples designated for Central Long Island Disposal Testing Site.
- (f) NS Not spiked.
- (g) Outside quality control criteria (50-120%) for matrix spike recoveries.
- (h) Percent Difference from certified
= absolute value [(certified value, $\mu\text{g}/\text{kg}$ - value detected corrected for surrogate recovery, $\mu\text{g}/\text{kg}$) / certified value, $\mu\text{g}/\text{kg}$].
- (i) NC No certified value available.
- (j) Outside SRM quality control acceptable criteria ($\leq 30\%$).

TABLE A.9. MDL Verification Study for Analysis of Polynuclear Aromatic Hydrocarbons (PAH) in Sediment

Sample Number:	OG99	OH01	OH02	OH03	OH04	OH05	OH06	OH07	Standard	Method Detection Limit	Method Detection Limit
Percent Moisture (%):	38.233	38.160	38.160	38.160	38.098	38.160	38.160	38.161	Deviation	MDL ^(a)	MDL
Sample Dry Weight (g):	20.919	19.455	19.201	18.645	19.087	19.434	18.896	18.612	STD	(µg/kg)	(ng)
Units (dry wt):	(µg/kg)	(µg/kg)	(µg/kg)	(µg/kg)	(µg/kg)	(µg/kg)	(µg/kg)	(µg/kg)			
naphthalene	1.61	1.85	1.88	1.86	1.66	1.72	1.75	1.97	0.12	0.36	7.03
biphenyl	1.30	1.55	1.49	1.61	1.56	1.50	1.57	1.67	0.11	0.33	6.35
acenaphthylene	0.93	1.06	1.09	1.15	1.01	1.18	1.09	1.16	0.08	0.25	4.87
acenaphthene	1.12	1.41	1.16	1.41	1.38	1.21	1.34	1.56	0.15	0.44	8.55
fluorene	1.07	1.31	1.12	1.17	1.09	0.99	1.27	1.25	0.11	0.34	6.48
phenanthrene	1.25	1.41	1.35	1.58	1.42	1.38	1.43	1.59	0.11	0.34	6.52
anthracene	0.73	0.87	0.78	0.88	0.87	0.78	0.77	0.99	0.08	0.25	4.80
fluoranthene	1.10	1.24	1.08	1.24	1.11	1.13	1.13	1.11	0.06	0.19	3.64
pyrene	1.16	1.34	1.21	1.21	1.14	1.15	1.19	1.19	0.06	0.19	3.64
benz[a]anthracene	0.82	1.08	0.94	0.96	0.92	0.89	0.88	0.95	0.08	0.23	4.38
chrysene	0.95	1.12	0.98	1.14	1.01	1.16	0.95	1.02	0.09	0.26	4.98
benzo[b]fluoranthene	0.97	1.02	0.93	1.03	0.89	0.88	0.85	0.86	0.07	0.21	4.03
benzo[k]fluoranthene	0.93	0.92	0.93	1.01	0.89	0.92	1.01	0.69	0.10	0.30	5.72
benzo[a]pyrene	0.67	0.77	0.61	0.79	0.81	0.70	0.71	0.60	0.08	0.24	4.54
indeno[1,2,3-c,d]pyrene	0.85	0.84	0.70	0.75	0.75	0.58	0.66	0.61	0.10	0.30	5.79
dibenz[a,h]anthracene	0.70	0.71	0.53	0.62	0.45	0.53	0.44	0.40	0.12	0.36	6.90
benzo[g,h,i]perylene	0.95	0.87	0.86	0.99	0.73	0.84	0.85	0.76	0.09	0.26	4.99
<u>Surrogate Recoveries (%)</u>											
naphthalene-d8	66	74	69	74	68	74	70	71			
acenaphthene-d10	65	71	69	73	68	73	71	70			
chrysene-d12	58	65	61	65	61	64	62	58			

(a) MDL = STD * 2.998, Average Sample Dry Weight (g) = 19.281.

Appendix B

**Site Water and Elutriate Chemical Analyses and
Quality Assurance/Quality Control Data,
Port Chester Project**

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Metals
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Site Water and Elutriate

QA/QC DATA QUALITY OBJECTIVES

	<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Target Detection Limit (µg/L)</u>
Cadmium	ICP/MS	75-125%	≤20%	≤20%	0.025
Chromium	GFAA	75-125%	≤20%	≤20%	1.0
Copper	ICP/MS	75-125%	≤20%	≤20%	0.35
Lead	ICP/MS	75-125%	≤20%	≤20%	0.35
Mercury	CVAA	75-125%	≤20%	≤20%	0.002
Nickel	ICP/MS	75-125%	≤20%	≤20%	0.3
Silver	ICP/MS	75-125%	≤20%	≤20%	0.25
Zinc	GFAA	75-125%	≤20%	≤20%	0.15

METHOD

A total of eight metals was analyzed in water and elutriate samples: silver (Ag), cadmium (Cd), chromium (Cr), copper (Cu), mercury (Hg), nickel (Ni), lead (Pb) and zinc (Zn). Hg was analyzed using cold-vapor atomic absorption spectroscopy (CVAA) according to the method of Bloom and Crecelius (1983). Cr and Zn were analyzed by Graphite Furnace Atomic Absorption (GFAA) spectrometry following the EPA Method 200.9 (EPA 1991). The remaining metals were analyzed by inductively coupled plasma mass spectrometry (ICP/MS) following a procedure based on EPA Method 200.8 (EPA 1991).

All water and elutriate samples were acidified to pH <2 upon receipt in the laboratory. Five metals, Cd, Cu, Pb, Ni and Ag, were extracted from the water according to a procedure based on EPA Method 218.3 (EPA 1979). This preconcentration involves addition of a chelating agent which results in precipitation of the metals from solution, followed by filtration, and digestion of the filter in concentrated acid in order to achieve low detection limits. The digestates were then analyzed by ICP/MS as described above.

HOLDING TIMES

Twelve site water samples (for triplicate analysis) were received on 3/24/94. Five elutriate samples (for triplicate analysis) were received on 4/11/94, and another five elutriate samples (for triplicate analysis) were received on 4/16/94. All samples were received in good condition, assigned ID numbers according to Battelle's log-in system, acidified to pH<2 with concentrated nitric acid, and held at ambient temperature until analysis.

QA/QC SUMMARY/METALS (continued)

Mercury in water has a holding time of 28 days from collection to analysis. All samples were analyzed within this holding time. Samples were analyzed for the remaining metals within 180 days of collection. Samples were received, digested, and analyzed in two batches, Batch 1a/1b (site waters), and Batch 2 (elutriate). The following table summarizes analysis dates:

<u>Task</u>	<u>Date</u>	
	<u>Batch 1a/1b</u>	<u>Batch 2</u>
APDC Extraction	6/13/94	5/24/94
ICP-MS	7/14/94	7/14/94
CVAA-Hg	4/26-28/94	5/9/94
GFAA-Cr	1a: 5/5/94 1b: 5/6/94	5/9/94
GFAA-Zn	5/16/94	5/16/94

DETECTION LIMITS

Target detection limits were met for all metals except Zn. Detection limits for Zn exceeded the target limits; however, all sample values were well above the detection limits achieved. Method Detection Limits (MDLs) for Ag, Cd, Cu, Hg, Ni and Pb were determined by spiking eight replicates of laboratory deionized water and multiplying the standard deviation of the resulting analysis by the Student's t value for n=8. MDLs reported for Cr and Zn were determined by taking the standard deviation of three replicate analyses of the method blank and multiplying the standard deviation by 3. An MDL verification study was performed within the previous year by spiking four replicates of Sequim Bay seawater and multiplying the standard deviation of the resulting analysis by 4.451. All sample MDLs were lower than the MDL verification values.

METHOD BLANKS

Method blanks were generated during the APDC extraction step and analyzed for the metals that were preconcentrated (Ag, Cd, Cu, Ni and Pb.) The blanks reported for Hg, Cr and Zn (the metals analyzed by direct injection of water samples) consist of a dilute nitric acid solution used to dilute all samples for analysis. For Batch 1a/1b, two APDC procedural blanks were analyzed and no APDC metals were detected in the blanks. Cr and Zn were detected in the blank; Cr at levels less than three times the MD, and Zn at levels greater than three times the MDL. All data were corrected for the blank concentrations, and no data were flagged. For Batch 2, two APDC procedural blanks were analyzed and no APDC metals were detected in the blanks. Zn and Cr were detected in the blank at levels less than three times the MDL. All data were corrected for the blank concentrations.

MATRIX SPIKES

Two samples were spiked in duplicate with all metals except Hg, which was spiked on two single samples. The APDC metals (Ag, Cd, Cu, Ni and Pb) were spiked prior to sample processing and the other metals were spiked just prior to analysis. For Batch 1a/1b, all recoveries were within the QC limits of 75% -125%, with the exception of Ag, Cd, and Cu in some of the spikes. Spike recoveries for these metals ranged from 70% to 74%, just below the lower QC limit. No action was taken. For Batch 2, all recoveries were within the QC limits of 75% -125% with the exception of Pb and Ni in one direct spike. Because Pb and

QA/QC SUMMARY/METALS (continued)

Ni values for the other spikes were acceptable, no further action was taken.

REPLICATES

Each sample was analyzed in triplicate. Precision for triplicate analyses was reported by calculating the relative standard deviation (RSD) of the replicate results. For Batch 1a/1b, RSD values were within the QC limits of $\pm 20\%$, with the exception of Hg, Pb, and Ni on one sample. For Batch 2, RSD values were all within the QC limits of $\pm 20\%$, with the exception of Cd in one sample and Ag in two samples.

SRMs

Standard Reference Material (SRM), CASS-2, a certified seawater sample from the National Institute of Standards and Technology, (NIST), was analyzed for all metals with the exception of Ag and Hg, which are not certified in this SRM. Results for all metals were within $\pm 20\%$ of mean certified value. Cd and Pb are certified below the MDL and were not detected.

A second SRM, 1641b, a freshwater sample from NIST, was analyzed twice for Hg. Results were within $\pm 20\%$ of mean certified value. No salt water SRMs certified for Ag are available.

A third SRM, 1643c, a freshwater sample from NIST, was analyzed for all metals except Hg. All metals were recovered within $\pm 20\%$ of mean certified value.

REFERENCES

Bloom, N. S., and E.A. Crecelius. 1983. "Determination of Mercury in Seawater at Sub-Nanogram per Liter Levels." *Mar. Chem.* 14:49-59.

EPA (U.S. Environmental Protection Agency). 1979. (Revised 1983). *Methods for the Chemical Analysis of Water and Wastes*. EPA-600/4-79-020. Environmental Monitoring Systems Laboratory, Cincinnati, Ohio.

EPA (U.S. Environmental Protection Agency). 1991 *Methods for the Determination of Metals in Environmental Samples*. EPA-600/4-91-010. Environmental Services Division, Monitoring Management Branch, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Chlorinated Pesticides and PCB Congeners
LABORATORY: Battelle Ocean Sciences
MATRIX: Site Water and Elutriate

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>Surrogate Recovery</u>	<u>MS Recovery</u>	<u>Relative Precision</u>	<u>Detection Limit</u>
GC/ECD	30-150%	50-120%	≤30%	2-20 ng/L

SAMPLE CUSTODY Twelve site water samples (in triplicate) were received on 3/31/94. Five elutriate samples (in triplicate) were received on 4/15/94, and another six elutriate samples (in triplicate) were received on 4/19/94. All samples were received in good condition, assigned ID numbers according to Battelle's log-in system, and stored at approximately 4°C until extraction.

METHOD Water samples were extracted with methylene chloride in a separatory funnel under ambient conditions following a procedure based on the National Oceanic and Atmospheric Administration (NOAA) Status and Trends Program method (Krahn et al. 1988). Sample extracts were passed through a silica/alumina (5% deactivated) chromatography column followed by high performance liquid chromatography (HPLC) cleanup (Krahn et al. 1988). Extracts were analyzed for 15 chlorinated pesticides using gas chromatography with electron capture detection (GC/ECD) following a procedure based on EPA Method 8080 (EPA 1986). The GC column used was a J&W DB-17 capillary column (30-m x 0.25-mm I.D.) with confirmatory analysis on a DB-1701 column (also 30-m x 0.25-mm I.D.).

HOLDING TIMES Samples were extracted in four batches: Batches 1 and 2 consisted of site waters; Batches 3 and 4 were elutriate samples. The following table summarizes sample extraction and analysis dates for each batch:

<u>Batch No.</u>	<u>Receipt</u>	<u>Extraction</u>	<u>Analysis</u>
1	3/31/94	4/5/94	4/22-26/94
2	3/31/94	4/5/94	4/26-28/94
3	4/15/94	4/19/94	5/5-7/94
4	4/19/94	4/22/94	5/13-15/94

DETECTION LIMITS Target detection limits (DLs) were met for all pesticides except endosulfan II in some samples (target DL for endosulfan II was 4 ng/L; achieved DL was 11 ng/L).

QA/QC SUMMARY/PESTICIDES AND PCBS (continued)

- METHOD BLANKS** One method blank (Sequim Bay seawater) was extracted with each extraction batch for a total of four method blanks. No pesticides or PCBs were detected in any of the method blanks.
- SURROGATES** Two compounds, dibromooctafluorobiphenyl (DBOBF) and PCB congener 112, were added to all samples to assess the efficiency of the analysis. Sample surrogate recoveries were all within the QC guidelines of 30% -150%.
- MATRIX SPIKES** One water sample in each batch (for a total of four) was spiked with 11 pesticides and 19 PCB congeners. Matrix spike recoveries were within the control limit range of 50-120% with the following exceptions: In the Batch 1, 2, 3, and 4 spike, recovery of PCB 8 was unacceptable due to interference from coelution of the non-target pesticide, alpha-BHC. In the batch 2 matrix spike, recovery of PCB 18 was 48%. In the Batch 3 matrix spike, recovery of endosulfan 1/2,4'DDE was 123% and recovery of heptachlor epoxide was 125%. No action was taken.
- REPLICATES** Each sample was extracted and analyzed in triplicate. Precision was measured by calculating the relative standard deviation (RSD) of the replicate results. The target precision goal was $\leq 30\%$ RSD for analytes > 10 times the Method Detection Limit (MDL). RSDs ranged from 6% to 79%, however, the majority of mean concentrations of all analytes (in each set of triplicate samples) were < 10 times the detection limit. Twenty-five PCB/pesticides had a mean > 10 times the detection limit and had an RSD of $> 30\%$. These RSDs ranged from 31% to 64%.

REFERENCES

- Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.
- EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

TABLE B.1. Metals in Site Water and Elutriate

Sediment Treatment	Replicate	Concentrations in µg/L							
		Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
target detection limit		0.25	0.025	1.0	0.35	0.002	0.30	0.35	0.15
MDL verification ^(a)		0.01	0.025	0.163	0.143	0.0007	0.253	0.035	0.582
PC Site Water	1	0.079	0.325	1.83	8.13	0.0261	2.36	9.83	25.3
PC Site Water	2	0.080	0.360	1.87	8.38	0.0232	2.36	10.1	28.1
PC Site Water	3	0.099	0.336	1.67	8.32	0.0253	2.45	10.5	18.1
PC Elutriate	1	0.0180	0.535	0.76	1.64	0.0236	3.57	1.78	7.81
PC Elutriate	2	0.0220	0.517	0.78	1.60	0.0221	3.48	1.64	6.51
PC Elutriate	3	0.0200	0.539	0.64	1.63	0.0225	3.57	1.76	6.51
Mud Dump Site Water	1	0.023	0.063	0.26	2.09	0.0097	1.29	0.942	9.35
Mud Dump Site Water	2	0.020	0.058	0.32	1.99	0.0093	1.22	0.904	12.2
Mud Dump Site Water	3	0.024	0.060	0.23	2.10	0.0097	1.30	0.947	9.35
Sequim Bay Control	1	0.007 U ^(b)	0.054	0.180	0.468	0.0006 U	0.465	0.035 U	7.88
Sequim Bay Control	2	0.007 U	0.056	0.180	0.452	0.0003	0.456	0.094	8.72
Sequim Bay Control	3	0.007 U	0.057	0.180	0.492	0.0006 U	0.486	0.035 U	11.0

(a) MDL Method detection limit based on standard deviation of 4 replicates of spiked control water x 4.541.

(b) U Not detected at or above concentration shown.

TABLE B.2. Quality Control Data (Method Blanks and Recovery of Matrix Spikes) for Metals in Site Water and Elutriate

Sediment Treatment	Batch	Concentrations in µg/L							
		Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
METHOD BLANKS									
Site Water									
Blank-1	1a	0.007 U ^(a)	0.025 U	0.33	0.143 U	0.0009	0.253 U	0.035 U	7.48
Blank-2	1b	0.007 U	0.025 U	0.41	0.143 U	0.0011	0.253 U	0.035 U	8.42
Blank-3	1b	NS ^(b)	NS	0.45	NS	NS	NS	NS	NS
Elutriate									
Blank-4	2	0.007 U	0.025 U	0.18	0.143 U	0.0009	0.253 U	0.035 U	0.75
Blank-5	2	0.007 U	0.025 U	0.16	0.143 U	0.0009	0.253 U	0.035 U	0.75
MATRIX SPIKES									
PC Site Water	1a	NA ^(c)	NA	1.79	NA	NA	NA	NA	27.2
PC Site Water, MS ^(d)	1a	NA	NA	2.81	NA	NA	NA	NA	67.3
Concentration Recovered		NA	NA	1.02	NA	NA	NA	NA	40.1
Amount Spiked		NS	NS	0.97	NS	NS	NS	NS	44.8
Percent Recovery		NA	NA	105%	NA	NA	NA	NA	90%
PC Site Water	1a	NA	NA	1.79	NA	NA	NA	NA	27.2
PC Site Water, MSD ^(e)	1a	NA	NA	6.47	NA	NA	NA	NA	114
Concentration Recovered		NA	NA	4.68	NA	NA	NA	NA	86.8
Amount Spiked		NS	NS	4.67	NS	NS	NS	NS	89.2
Percent Recovery		NA	NA	100%	NA	NA	NA	NA	97%
RPD ^(f)		NA	NA	5%	NA	NA	NA	NA	8%
SB-A Site Water	1a	0.143	0.112	NA	5.15	0.0165	1.95	2.96	NA
SB-A Site Water, MS	1a	0.945	0.903	NA	5.89	0.0511	2.73	4.19	NA
Concentration Recovered		0.802	0.791	NA	0.74	0.0346	0.78	1.23	NA
Amount Spiked		1.00	1.00	NS	1.00	0.0364	1.00	1.00	NS
Percent Recovery		80%	79%	NA	74% ^(g)	95%	78%	123%	NA
SB-A Site Water	1a	0.143	0.112	NA	5.15	NA	1.95	2.96	NA
SB-A Site Water, MSD	1a	4.49	3.83	NA	9.67	NA	5.94	7.4	NA
Concentration Recovered		4.35	3.72	NA	4.52	NA	3.99	4.44	NA
Amount Spiked		5.00	5.00	NS	5.00	NS	5.00	5.00	NS
Percent Recovery		87%	74% ^(g)	NA	90%	NA	80%	89%	NA
RPD		8%	6%	NA	20%	NA	2%	32%	NA
HU-B Site Water	1b	NA	NA	1.81	NA	NA	NA	NA	NA
HU-B Site Water, MS	1b	NA	NA	2.94	NA	NA	NA	NA	NA
Concentration Recovered		NA	NA	1.13	NA	NA	NA	NA	NA
Amount Spiked		NS	NS	0.97	NS	NS	NS	NS	NS
Percent Recovery		NA	NA	116%	NA	NA	NA	NA	NA
HU-B Site Water	1b	NA	NA	1.81	NA	NA	NA	NA	NA
HU-B Site Water, MSD	1b	NA	NA	6.24	NA	NA	NA	NA	NA
Concentration Recovered		NA	NA	4.43	NA	NA	NA	NA	NA
Amount Spiked		NS	NS	4.67	NS	NS	NS	NS	NS
Percent Recovery		NA	NA	95%	NA	NA	NA	NA	NA
RPD		NA	NA	20%	NA	NA	NA	NA	NA

TABLE B.2. (contd)

Sediment Treatment	Batch	Concentrations in µg/L							
		Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
Mud Dump Site Water	1b	0.022	0.060	NA	2.06	0.0096	1.27	0.931	NA
Mud Dump Site Water, MS	1b	0.743	0.763	NA	3.00	0.0469	20.8	1.86	NA
Concentration Recovered		0.721	0.703	NA	0.94	0.0373	0.810	0.929	NA
Amount Spiked		1.00	1.00	NS	1.00	0.0347	1.00	1.00	NS
Percent Recovery		72% ^(g)	70% ^(g)	NA	94%	107%	81%	93%	NA
Mud Dump Site Water	1b	0.022	0.060	NA	2.06	NA	1.27	0.931	NA
Mud Dump Site Water, MSD	1b	4.13	3.56	NA	6.56	NA	5.3	5.60	NA
Concentration Recovered		4.11	3.50	NA	4.50	NA	4.03	4.67	NA
Amount Spiked		5.00	5.00	NS	5.00	NS	5.00	5.00	NS
Percent Recovery		82%	70% ^(g)	NA	90%	NA	81%	93%	NA
RPD		13%	0.4%	NA	4%	NA	0.5%	1%	NA
PC Elutriate	2	NA	NA	0.78	NA	NA	NA	NA	6.51
PC Elutriate, MS	2	NA	NA	1.70	NA	NA	NA	NA	54.7
Concentration Recovered		NA	NA	0.92	NA	NA	NA	NA	48.2
Amount Spiked		NS	NS	0.97	NS	NS	NS	NS	44.8
Percent Recovery		NA	NA	95%	NA	NA	NA	NA	108%
PC Elutriate	2	NA	NA	0.78	NA	NA	NA	NA	6.51
PC Elutriate, MSD	2	NA	NA	5.44	NA	NA	NA	NA	102
Concentration Recovered		NA	NA	4.66	NA	NA	NA	NA	95.5
Amount Spiked		NS	NS	4.67	NS	NS	NS	NS	89.2
Percent Recovery		NA	NA	100%	NA	NA	NA	NA	107%
RPD		NA	NA	5%	NA	NA	NA	NA	0.5%
SB-B Elutriate	2	0.018	0.025 U	NA	0.741	0.0034	3.02	0.681	NA
SB-B Elutriate, MS	2	0.824	0.856	NA	1.72	0.0245	4.31	2.32	NA
Concentration Recovered		0.806	0.856	NA	0.982	0.0211	1.29	1.64	NA
Amount Spiked		1.00	1.00	NS	1.00	0.0211	1.00	1.00	NS
Percent Recovery		81%	86%	NA	98%	100%	129% ^(g)	164% ^(g)	NA
SB-B Elutriate	2	0.018	0.025 U	NA	0.741	NA	3.02	0.681	NA
SB-B Elutriate, MSD	2	4.34	3.79	NA	5.57	NA	8.10	5.11	NA
Concentration Recovered		4.32	3.79	NA	4.83	NA	5.08	4.43	NA
Amount Spiked		5.00	5.00	NS	5.00	NS	5.00	5.00	NS
Percent Recovery		86%	76%	NA	97%	NA	102%	89%	NA
RPD		7%	12%	NA	2%	NA	24%	60%	NA
EC-B Elutriate	2	NA	NA	NA	NA	0.0275	NA	NA	NA
EC-B Elutriate, MS	2	NA	NA	NA	NA	0.0470	NA	NA	NA
Concentration Recovered		NA	NA	NA	NA	0.0195	NA	NA	NA
Amount Spiked		NS	NS	NS	NS	0.0212	NS	NS	NS
Percent Recovery		NA	NA	NA	NA	92%	NA	NA	NA
HU-B Elutriate	2	NA	NA	0.18	NA	NA	NA	NA	11.0
HU-B Elutriate, MS	2	NA	NA	1.15	NA	NA	NA	NA	59.9
Concentration Recovered		NA	NA	0.97	NA	NA	NA	NA	48.9
Amount Spiked		NS	NS	0.97	NS	NS	NS	NS	44.8
Percent Recovery		NA	NA	100%	NA	NA	NA	NA	109%

TABLE B.2. (contd)

Sediment Treatment	Batch	Concentrations in µg/L							
		Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
HU-B Elutriate	2	NA	NA	0.18	NA	NA	NA	NA	11.0
HU-B Elutriate, MSD	2	NA	NA	5.77	NA	NA	NA	NA	111
Concentration Recovered		NA	NA	5.59	NA	NA	NA	NA	100
Amount Spiked		NS	NS	4.67	NS	NS	NS	NS	89.2
Percent Recovery		NA	NA	120%	NA	NA	NA	NA	112%
RPD		NA	NA	18%	NA	NA	NA	NA	3%
EC-A Elutriate	2	0.007 U	0.025 U	NA	0.661	0.0005	0.771	0.992	NA
EC-A Elutriate, MS	2	0.831	0.805	NA	1.55	0.0319	1.59	1.85	NA
Concentration Recovered		0.831	0.805	NA	0.892	0.0314	0.816	0.857	NA
Amount Spiked		1.00	1.00	NS	1.00	0.0316	1.00	1.00	NS
Percent Recovery		83%	81%	NA	89%	99%	82%	86%	NA
EC-A Elutriate	2	0.004	0.012	NA	0.661	NA	0.771	0.992	NA
EC-A Elutriate, MSD	2	4.34	3.82	NA	5.34	NA	5.11	5.48	NA
Concentration Recovered		4.33	3.81	NA	4.68	NA	4.31	4.49	NA
Amount Spiked		5.00	5.00	NS	5.00	NS	5.00	5.00	NS
Percent Recovery		87%	76%	NA	94%	NA	86%	90%	NA
RPD		4%	6%	NA	5%	NA	5%	5%	NA

(a) U Undetected at or above concentration shown.

(b) NS Not spiked.

(c) NA Not applicable.

(d) MS Matrix spike

(e) MSD Matrix spike duplicate

(f) RPD Relative percent difference.

(g) Outside data quality criteria of 75%-125%.

TABLE B.3. Quality Control Data (Triplicate Analyses) for Metals in Site Water and Elutriate

Sediment Treatment	Repli- cate	Batch	Concentrations in µg/L							
			Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
PC Site Water	1	1a	0.079	0.325	1.83	8.13	0.0261	2.36	9.83	25.3
PC Site Water	2	1a	0.080	0.360	1.87	8.38	0.0232	2.36	10.1	28.1
PC Site Water	3	1a	0.099	0.336	1.67	8.32	0.0253	2.45	10.5	18.1
RSD ^(a)			13%	5%	6%	2%	6%	2%	3%	22% ^(b)
EC-A Site Water	1	1a	0.092	0.503	6.47	13.4	0.0685	4.43	20.5	58.9
EC-A Site Water	2	1a	0.091	0.519	6.71	14.1	0.0640	4.64	22.1	64.5
EC-A Site Water	3	1a	0.087	0.542	6.35	18.6	0.0619	4.43	21.7	64.5
RSD			3%	4%	3%	18%	5%	3%	4%	5%
EC-B Site Water	1	1a	0.152	0.411	4.49	19.0	0.212	4.76	18.7	64.5
EC-B Site Water	2	1a	0.167	0.396	4.61	18.9	0.155	4.58	17.6	69.2
EC-B Site Water	3	1a	0.159	0.419	4.44	18.7	0.182	4.69	18.0	71.1
RSD			5%	3%	2%	1%	16%	2%	3%	5%
HU-A Site Water	1	1a	0.107	0.102	0.83	4.53	0.0178	1.67	3.37	12.2
HU-A Site Water	2	1a	0.082	0.114	0.85	4.59	0.0189	1.79	3.60	14.0
HU-A Site Water	3	1a	0.120	0.114	0.88	4.87	0.0188	1.80	3.78	13.1
RSD			19%	6%	3%	4%	3%	4%	6%	7%
SB-A Site Water	1	1a	0.145	0.108	1.02	5.04	0.0190	1.92	2.85	19.6
SB-A Site Water	2	1a	0.141	0.118	1.15	5.09	0.0160	1.96	3.03	18.7
SB-A Site Water	3	1a	0.142	0.110	1.32	5.33	0.0145	1.97	2.99	21.5
RSD			1%	5%	13%	3%	14%	1%	3%	7%
SB-B Site Water	1	1a	0.075	0.094	0.71	3.53	0.0066	1.67	1.30	9.35
SB-B Site Water	2	1a	0.075	0.093	0.59	3.56	0.0061	1.81	1.32	10.3
SB-B Site Water	3	1a	0.073	0.088	0.68	3.49	0.0062	1.58	1.27	11.2
RSD			2%	4%	9%	1%	4%	7%	2%	9%
BU Site Water	1	1b	0.104	0.090	0.81	4.16	0.0233	1.82	2.79	12.2
BU Site Water	2	1b	0.109	0.080	0.85	4.38	0.0220	1.87	2.79	14.0
BU Site Water	3	1b	0.118	0.096	0.92	4.27	0.0216	1.94	2.85	13.1
RSD			6%	9%	6%	3%	4%	3%	1%	7%
Mud Dump Site Water	1	1b	0.023	0.063	0.26 J ^(c)	2.09	0.0097	1.29	0.942	9.35
Mud Dump Site Water	2	1b	0.020	0.058	0.32 J	1.99	0.0093	1.22	0.904	12.2
Mud Dump Site Water	3	1b	0.024	0.060	0.23 J	2.10	0.0097	1.30	0.947	9.35
RSD			9%	4%	17%	3%	2%	3%	3%	16%
HU-B Site Water	1	1b	0.192	0.105	1.75	6.73	0.0351	2.13	5.34	13.1
HU-B Site Water	2	1b	0.188	0.105	1.92	6.42	0.0369	2.09	4.95	11.2
HU-B Site Water	3	1b	0.182	0.107	1.75	6.57	0.0373	2.07	5.12	13.1
RSD			3%	1%	5%	2%	3%	1%	4%	9%
HU-C Site Water	1	1b	0.144	0.093	0.94	5.52	0.0288	1.85	4.30	30.9
HU-C Site Water	2	1b	0.139	0.087	0.83	5.25	0.0279	1.86	4.15	31.8
HU-C Site Water	3	1b	0.142	0.089	0.90	5.37	0.0296	1.79	4.02	27.1
RSD			2%	3%	6%	3%	3%	2%	3%	8%
HU-D Site Water	1	1b	0.119	0.113	1.43	5.69	0.0263	1.82	4.89	38.3
HU-D Site Water	2	1b	0.119	0.113	1.39	5.59	0.0277	1.65	4.94	37.4
HU-D Site Water	3	1b	0.121	0.111	1.26	5.81	0.0269	4.24	5.17	36.5
RSD			1%	1%	7%	2%	3%	56% ^(b)	3%	2%

TABLE B.3. (Contd)

Sediment Treatment	Replicate	Batch	Concentrations in µg/L							
			Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
PC Elutriate	1	2	0.018	0.535	0.76	1.64	0.0236	3.57	1.78	7.81
PC Elutriate	2	2	0.022	0.517	0.78	1.60	0.0221	3.48	1.64	6.51
PC Elutriate	3	2	0.020	0.539	0.64	1.63	0.0225	3.57	1.76	6.51
RSD			10%	2%	10%	1%	3%	1%	4%	11%
SB-B Elutriate	1	2	0.017	0.025 U ^(d)	0.72	0.755	0.0031	2.95	0.667	3.10
SB-B Elutriate	2	2	0.018	0.025 U	0.58	0.736	0.0032	3.02	0.676	3.47
SB-B Elutriate	3	2	0.018	0.025 U	0.64	0.741	0.0034	3.02	0.681	2.72
RSD			3%	NA ^(e)	11%	1%	5%	1%	1%	12%
SB-A Elutriate	1	2	0.036	0.025 U	1.15	1.28	0.0285	2.61	0.807	3.10
SB-A Elutriate	2	2	0.035	0.025 U	1.21	1.18	0.0290	2.39	0.779	2.63
SB-A Elutriate	3	2	0.030	0.025 U	1.17	1.12	0.0290	2.42	0.772	2.25
RSD			10%	NA	3%	7%	1%	5%	2%	16%
BU Elutriate	1	2	0.021	0.025 U	0.58	0.737	0.0049	2.99	0.586	2.25
BU Elutriate	2	2	0.038	0.025 U	0.62	0.700	0.0051	2.95	0.603	3.28
BU Elutriate	3	2	0.020	0.025 U	0.53	0.709	0.0051	2.85	0.564	2.44
RSD			38% ^(b)	NA	8%	3%	2%	2%	3%	21% ^(b)
EC-B Elutriate	1	2	0.027	0.083	1.62	3.54	0.0263	1.75	5.82	5.35
EC-B Elutriate	2	2	0.023	0.236	1.66	3.57	0.0249	1.73	5.28	5.06
EC-B Elutriate	3	2	0.035	0.121	1.83	3.67	0.0275	1.74	5.34	3.94
RSD			22% ^(b)	54% ^(b)	7%	2%	5%	1%	5%	16%
HU-B Elutriate	1	2	0.075	0.033	2.44	1.90	0.0198	1.39	1.18	1.78
HU-B Elutriate	2	2	0.061	0.034	2.16	1.92	0.0187	1.43	1.11	2.16
HU-B Elutriate	3	2	0.064	0.035	2.42	1.95	0.0179	1.42	1.09	1.88
RSD			11%	3%	7%	1%	5%	1%	4%	10%
HU-A Elutriate	1	2	0.025	0.028	1.44	1.24	0.0130	1.53	0.994	6.19
HU-A Elutriate	2	2	0.022	0.028	1.25	1.22	0.0110	1.50	1.03	6.10
HU-A Elutriate	3	2	0.023	0.025 U	1.17	1.14	0.0108	1.44	0.999	5.91
RSD			7%	NA	11%	4%	10%	3%	2%	2%
EC-A Elutriate	1	2	0.007 U	0.025 U	0.66	0.590	0.0010	0.711	0.971	1.13
EC-A Elutriate	2	2	0.007 U	0.025 U	0.60	0.640	0.0006 U	0.750	0.935	1.41
EC-A Elutriate	3	2	0.007 U	0.025 U	0.55	0.661	0.0005	0.771	0.992	1.41
RSD			NA	NA	9%	6%	NA	4%	3%	12%
HU-C Elutriate	1	2	0.035	0.031	1.73	1.25	0.0152	2.37	1.11	2.25
HU-C Elutriate	2	2	0.030	0.031	1.81	1.14	0.0132	2.24	0.994	2.34
HU-C Elutriate	3	2	0.031	0.033	1.95	1.24	0.0124	2.32	1.09	1.88
RSD			8%	4%	6%	5%	11%	3%	6%	11%
HU-D Elutriate	1	2	0.021	0.025 U	0.84	0.993	0.0125	1.41	0.847	1.69
HU-D Elutriate	2	2	0.016	0.057	0.84	1.06	0.0129	1.39	0.953	1.59
HU-D Elutriate	3	2	0.027	0.045	0.72	1.03	0.0128	1.44	0.846	1.31
RSD			26% ^(b)	NA	9%	3%	2%	2%	7%	13%
Control Site Water	1	2	0.007 U	0.054	0.18	0.468	0.0006 U	0.465	0.035 U	7.88
Control Site Water	2	2	0.007 U	0.056	0.18	0.452	0.0003	0.456	0.094	8.72
Control Site Water	3	2	0.007 U	0.057	0.18	0.492	0.0006 U	0.486	0.035 U	11.0
RSD			NA	3%	0%	4%	NA	3%	NA	18%

(a) RSD Relative standard deviation.

(b) Outside data quality criteria of +/-20% RSD.

(c) J Concentration estimated; analyte detected below detection limit.

(d) U Undetected at or above concentration shown.

(e) NA Not applicable.

TABLE B.4. Quality Control Data (Standard Reference Materials) for Metals in Site Water and Elutriate

Standard Reference Material	Replicate	Batch	Concentrations in µg/L							
			Ag ICP/MS	Cd ICP/MS	Cr GFAA	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn GFAA
Site Water										
SRM CASS-2	1	1a	0.007 U ^(a)	0.025 U	0.32 U	0.695	NA ^(b)	0.301	0.016 J ^(c)	2.04
SRM CASS-2	2	1a	0.007 U	0.025 U	0.32 U	0.730	NA	0.339	0.018 J	2.30
SRM CASS-2	1	1b	NA	NA	0.19 U	NA	NA	NA	NA	NA
Certified Value CASS-2			NC ^(d)	0.019	0.121	0.675	NC	0.298	0.019	1.97
Range			NC	±0.004	±0.016	±0.039	NC	±0.036	±0.006	±0.12
Percent Difference	1		NA	NA	NA	3	NA	1	16	4
Percent Difference	2		NA	NA	NA	8	NA	14	5	17
Percent Difference	1		NA	NA	NA	NA	NA	NA	NA	NA
SRM 1641b	1	1a	NA	NA	NA	NA	1530	NA	NA	NA
SRM 1641b	2	1a	NA	NA	NA	NA	1540	NA	NA	NA
Certified Value 1641b			NC	NC	NC	NC	1520	NC	NC	NC
Range			NC	NC	NC	NC	±40	NC	NC	NC
Percent Difference	1		NA	NA	NA	NA	1	NA	NA	NA
Percent Difference	2		NA	NA	NA	NA	1	NA	NA	NA
SRM 1643c	1	1a	2.09	11.7	20.5	20.6	NA	55.3	33.6	84.2
SRM 1643c	2	1a	2.01	11.0	19.4	19.2	NA	54.2	35.8	84.2
SRM 1643c	1	1b	NA	NA	19.5	NA	NA	NA	NA	NA
Certified Value 1643c			2.21	12.2	19.0	22.3	NC	60.6	35.3	73.9
Range			±0.30	± 1.0	±0.6	±2.8	NC	±7.3	±0.9	±0.9
Percent Difference	1		5	4	8	8	NA	9	5	14
Percent Difference	2		9	10	2	14	NA	11	1	14
Percent Difference	1		NA	NA	3	NA	NA	NA	NA	NA
Elutriate										
SRM CASS-2	1	2	0.003 U	0.025 U	0.103	0.671	NA	0.257	0.035 U	2.10
SRM CASS-2	2	2	0.003 U	0.025 U	0.103	0.668	NA	0.258	0.035 U	1.83
Certified Value CASS-2			NC	0.019	0.118	0.675	NC	0.298	0.019	1.97
Range			NC	±0.004	±0.021	±0.039	NC	±0.036	±0.006	±0.12
Percent Difference	1		NA	NA	13	1	NA	14	NA	7
Percent Difference	2		NA	NA	13	1	NA	13	NA	7
SRM 1641b	1	2	NA	NA	NA	NA	1540	NA	NA	NA
SRM 1641b	2	2	NA	NA	NA	NA	1510	NA	NA	NA
Certified Value 1641b			NC	NC	NC	NC	1520	NC	NC	NC
Range			NC	NC	NC	NC	±40	NC	NC	NC
Percent Difference	1		NA	NA	NA	NA	1	NA	NA	NA
Percent Difference	2		NA	NA	NA	NA	1	NA	NA	NA
SRM 1643c	1	2	1.89	11.3	19.3	20.4	NA	56.7	33.0	76.0
SRM 1643c	2	2	1.80	11.2	21.0	20.0	NA	56.3	32.8	71.9
Certified Value 1643c			2.21	12.2	19.0	22.3	NC	60.6	35.3	73.9
Range			±0.30	± 1.0	±0.6	±2.8	NC	±7.3	±0.9	±0.9
Percent Difference	1		15	7	2	9	NA	6	7	3
Percent Difference	2		19	8	11	10	NA	7	7	3

(a) U Undetected at or above concentration shown.

(b) NA Not applicable.

(c) J Analyte detected below detection limit; concentration estimated.

(d) NC Not certified.

TABLE B.5. Pesticides and PCBs in Site Water and Elutriate

Site/Replicate Matrix	PC Rep. 1 Site Water	PC Rep. 2 Site Water	PC Rep. 3 Site Water	PC Rep 1 Elutriate	PC Rep 2 Elutriate	PC Rep 3 Elutriate
Sample Size (L) Units	1.04 ng/L	1.04 ng/L	1.04 ng/L	0.87 ng/L	0.96 ng/L	0.95 ng/L
2,4-DDD	0.765 U ^(a)	0.765 U	0.765 U	11.1	13.5	17.9
2,4-DDT	0.777 U	0.777 U	0.777 U	5.01	4.62	5.47
4,4-DDD	1.95	1.71	1.90	42.1	48.9	75.1
4,4-DDE	0.626 J ^(b)	0.601 J	0.806 J	11.6	13.8	22.0
4,4-DDT	0.962 U	1.70	0.900 J	1.15 U	1.04 U	1.05 U
Aldrin	0.713 U	0.713 U	0.713 U	0.85 U	0.77 U	0.78 U
<i>alpha</i> -Chlordane	1.80	1.94	1.76	13.4	14.9	21.1
Dieldrin	1.80	1.55	1.56	9.36	11.2	14.8
Endosulfan I/2,4'-DDE	0.813 U	0.813 U	0.813 U	0.97 U	0.88 U	0.89 U
Endosulfan II	1.57 J	10.8 U	10.8 U	4.93 J	4.73 J	6.70 J
Endosulfan sulfate	7.87 U	7.87 U	7.87 U	11.5	13.5	18.0
Heptachlor	0.631 U	0.631 U	0.631 U	0.75 U	0.68 U	0.69 U
Heptachlor epoxide	0.822 U	0.822 U	0.822 U	0.98 U	0.89 U	0.900 U
<i>trans</i> -Nonachlor	0.928 U	0.928 U	0.928 U	6.55	7.38	10.3
CL2(08)	0.841 U	0.841 U	0.841 U	1.01 U	0.91 U	0.92 U
CL3(18)	1.02 U	1.02 U	1.02 U	1.22 U	1.11 U	1.12 U
CL3(28)	4.20	2.69	3.05	5.32	5.88	6.89
CL4(44)	1.17 U	1.17 U	1.17 U	12.2	14.8	19.5
CL4(49)	1.01 U	1.01 U	1.01 U	7.62	7.50	11.4
CL4(52)	1.18 U	1.18 U	1.18 U	24.5	27.5	41.4
CL4(66)	0.917 U	0.917 U	0.917 U	9.78	11.8	21.5
CL5(87)	0.817 J	0.516 J	0.732 J	25.0	26.6	37.1
CL5(101)	1.04 U	1.04 U	1.04 U	67.2	79.1	118
CL5(105)	1.24 U	1.24 U	1.24 U	30.6	34.2	30.0
CL5(118)	0.977 U	0.977 U	0.977 U	47.0	52.5	79.1
CL6(128)	1.10 U	1.10 U	1.10 U	8.85	10.6	14.9
CL6(138)	1.31 U	1.31 U	0.66 J	56.4	66.1	96.5
CL6(153)	1.26 U	1.26 U	0.96 J	35.9	39.0	67.7
CL7(170)	1.12 U	1.12 U	1.12 U	11.3	15.7	22.3
CL7(180)	0.975 U	0.975 U	0.975 U	26.2	29.5	44.9
CL7(183)	1.02 U	1.02 U	1.02 U	5.57	5.91	8.02
CL7(184)	1.02 U	1.02 U	1.02 U	1.22 U	1.11 U	1.12 U
CL7(187)	0.964 U	0.964 U	0.964 U	18.0	20.1	28.0
CL8(195)	1.10 U	1.10 U	1.10 U	3.00	3.41	5.39
CL9(206)	1.08 U	1.08 U	1.08 U	6.07	7.20	11.0
CL10(209)	1.20 U	1.20 U	1.20 U	1.28 J	1.37	1.97
<u>Surrogate Recoveries (%)</u>						
DBOFB	108	105	103	120	120	123
CL5(112)	72	72	71	70.6	65.9	58.2

(a) U Undetected at or above concentration given.

(b) J Concentration estimated; analyte detected is below detection limit.

TABLE B.6. Quality Control Data (Method Blanks and Recovery of Matrix Spikes) for Pesticides and PCBs in Site Water and Elutriate

Sample:	Method Blank	SB-B Rep. 3	SB-B Rep. 3 MS	Amount	Percent
Matrix:	Control Water	Site Water	Site Water	Spiked	Recovery
Sample Size (L):	1.01 ^(a)	0.53	0.51		
Batch:	1	1	1	1	1
Units:	ng/L	ng/L	ng/L	ng	%
2,4-DDD	0.79 U ^(c)	1.52 U	NS ^(c)	NS	NA ^(d)
2,4-DDT	0.80 U	1.54 U	159.31	NS	NA
4,4-DDD	1.15 U	2.21 U	142.46	80.40	90
4,4-DDE	0.98 U	1.88 U	138.23	80.20	88
4,4-DDT	0.99 U	1.90 U	135.93	80.20	86
Aldrin	0.73 U	1.41 U	134.31	80.20	85
<i>alpha</i> -Chlordane	0.92 U	1.77 U	129.31	80.00	82
Dieldrin	0.97 U	2.64	111.18	80.20	69
Endosulfan I/2,4'-DDE	0.84 U	1.61 U	138.52	80.20	88
Endosulfan II	11.07 U	21.33 U	131.51	80.20	84
Endosulfan sulfate	8.09 U	15.59 U	120.25	80.20	76
Heptachlor	0.65 U	1.25 U	117.33	80.20	75
Heptachlor epoxide	0.85 U	1.63 U	118.33	80.20	75
<i>trans</i> -Nonachlor	0.95 U	1.84 U	NS	NS	NA
CL2(08)	0.87 U	1.67 U	C ^(e)	80.00	NC ^(f)
CL3(18)	1.05 U	2.03 U	83.25	80.00	53
CL3(28)	1.18 U	2.27 U	131.73	80.00	84
CL4(44)	1.20 U	2.32 U	114.82	80.00	73
CL4(49)	1.03 U	1.99 U	NS	NS	NA
CL4(52)	1.22 U	2.34 U	108.44	80.00	69
CL4(66)	0.94 U	1.82 U	137.82	80.00	88
CL5(87)	1.06 U	2.04 U	NS	NS	NA
CL5(101)	1.06 U	2.05 U	110.62	80.00	71
CL5(105)	1.28 U	2.46 U	133.30	80.00	85
CL5(118)	1.00 U	1.94 U	121.65	80.00	78
CL6(128)	1.13 U	2.17 U	121.75	80.00	78
CL6(138)	1.35 U	2.60 U	123.58	80.00	79
CL6(153)	1.29 U	2.49 U	108.26	80.00	69
CL7(170)	1.16 U	2.23 U	127.93	80.00	82
CL7(180)	1.00 U	1.93 U	118.14	80.00	75
CL7(183)	1.05 U	2.02 U	NS	NS	NA
CL7(184)	1.05 U	2.02 U	NS	NS	NA
CL7(187)	0.99 U	1.91 U	108.34	80.00	69
CL8(195)	1.14 U	2.19 U	122.94	80.00	78
CL9(206)	1.11 U	2.14 U	117.95	80.00	75
CL10(209)	1.23 U	2.38 U	113.65	80.00	72
<u>Surrogate Recoveries (%)</u>					
DBOFB	86	99	94	NA	NA
CL5(112)	77	74	74	NA	NA

TABLE B.6. (Contd)

Sample:	Method Blank	HU-D Rep. 3	HU-D Rep. 3 MS	Amount	Percent
Matrix:	Control Water	Site Water	Site Water	Spiked	Recovery
Sample Size (L):	1.01 ^(a)	0.52	0.52		
Batch:	2	2	2	2	2
Units:	ng/L	ng/L	ng/L	ng	%
2,4-DDD	0.79 U	1.53 U	NS	NS	NA
2,4-DDT	0.80 U	1.55 U	NS	NS	NA
4,4-DDD	1.15 U	2.23 U	132.72	80.40	86
4,4-DDE	0.98 U	1.90 U	120.53	80.20	78
4,4-DDT	0.99 U	1.92 U	125.17	80.20	81
Aldrin	0.73 U	1.43 U	113.20	80.20	73
<i>alpha</i> -Chlordane	0.92 U	1.72 J ⁽⁹⁾	118.11	80.00	76
Dieldrin	0.98 U	1.53 J	84.92	80.20	54
Endosulfan I/2,4'-DDE	0.84 U	1.63 U	136.31	80.20	88
Endosulfan II	11.08 U	2.71 J	111.86	80.20	71
Endosulfan sulfate	8.10 U	15.74 U	98.59	80.20	64
Heptachlor	0.65 U	1.26 U	103.27	80.20	67
Heptachlor epoxide	0.85 U	1.64 U	117.22	80.20	76
<i>trans</i> -Nonachlor	0.95 U	1.86 U	NS	NS	NA
CL2(08)	0.87 U	1.68 U	C	80.00	NC
CL3(18)	1.05 U	2.05 U	73.37	80.00	48 ⁽ⁿ⁾
CL3(28)	1.18 U	2.29 U	125.42	80.00	82
CL4(44)	1.20 U	2.34 U	109.8	80.00	71
CL4(49)	1.03 U	2.01 U	NS	NS	NA
CL4(52)	1.22 U	2.37 U	103.56	80.00	67
CL4(66)	0.94 U	1.83 U	147	80.00	96
CL5(87)	1.06 U	2.06 U	NS	NS	NA
CL5(101)	1.07 U	2.07 U	118.56	80.00	77
CL5(105)	1.28 U	2.48 U	138.28	80.00	90
CL5(118)	1.00 U	1.95 U	125.01	80.00	81
CL6(128)	1.13 U	2.19 U	122.64	80.00	80
CL6(138)	1.35 U	2.62 U	113.75	80.00	74
CL6(153)	1.29 U	2.52 U	103.09	80.00	67
CL7(170)	1.16 U	2.25 U	130.43	80.00	85
CL7(180)	1.00 U	1.95 U	115.48	80.00	75
CL7(183)	1.05 U	2.04 U	NS	NS	NA
CL7(184)	1.05 U	2.04 U	NS	NS	NA
CL7(187)	0.99 U	1.93 U	94.93	80.00	62
CL8(195)	1.14 U	2.21 U	112.84	80.00	73
CL9(206)	1.11 U	2.16 U	106.60	80.00	69
CL10(209)	1.23 U	2.40 U	96.54	80.00	63
<u>Surrogate Recoveries (%)</u>					
DBOFB	33	32	62	NA	NA
CL5(112)	46	49	64	NA	NA

TABLE B.6. (Contd)

Sample:	Method Blank	EC-B Rep. 3	EC-B Rep. 3 MS	Amount	Percent
Matrix:	Control Water	Elutriate	Elutriate	Spiked	Recovery
Sample Size (L):	0.94 ^(a)	0.50	0.48		
Batch:	3	3	3	3	3
Units:	ng/L	ng/L	ng/L	ng	%
2,4-DDD	0.85 U	3.07	NS	NS	NA
2,4-DDT	0.86 U	0.925 J	NS	NS	NA
4,4-DDD	1.24 U	12.2	185.49	80.40	103
4,4-DDE	1.06 U	6.55	163.88	80.20	94
4,4-DDT	1.07 U	2.00 U	172.90	80.20	103
Aldrin	0.79 U	22.5	199.10	80.20	106
<i>alpha</i> -Chlordane	0.99 U	13.2	189.13	80.00	106
Dieldrin	1.05 U	3.80	122.35	80.20	71
Endosulfan I/2,4'-DDE	0.90 U	1.69 U	205.25	80.20	123 ⁽ⁿ⁾
Endosulfan II	11.97 U	22.4 U	154.59	80.20	93
Endosulfan sulfate	8.75 U	16.4 U	146.38	80.20	88
Heptachlor	0.70 U	1.31 U	179.22	80.20	107
Heptachlor epoxide	0.91 U	1.71 U	209.34	80.20	125 ⁽ⁿ⁾
<i>trans</i> -Nonachlor	1.03 U	7.17	7.24	NS	NA
CL2(08)	0.94 U	1.75 U	C	80.00	NC
CL3(18)	1.14 U	2.13 U	145.89	80.00	88
CL3(28)	1.28 U	15.3	203.61	80.00	113
CL4(44)	1.30 U	12.4	185.74	80.00	104
CL4(49)	1.12 U	8.62	10.64	NS	NA
CL4(52)	1.32 U	66.5	201.24	80.00	81
CL4(66)	1.02 U	17.8	215.42	80.00	119
CL5(87)	1.14 U	4.94	NS	NS	NA
CL5(101)	1.15 U	11.6	181.50	80.00	102
CL5(105)	1.38 U	1.88 J	181.11	80.00	108
CL5(118)	1.09 U	9.71	164.19	80.00	93
CL6(128)	1.22 U	2.54	155.43	80.00	92
CL6(138)	1.46 U	11.1	155.98	80.00	87
CL6(153)	1.40 U	7.32	141.71	80.00	81
CL7(170)	1.25 U	2.34 U	163.91	80.00	98
CL7(180)	1.08 U	2.03 U	152.51	80.00	92
CL7(183)	1.14 U	2.09 J	NS	NS	NA
CL7(184)	1.14 U	2.12 U	NS	NS	NA
CL7(187)	1.07 U	2.01 U	121.21	80.00	73
CL8(195)	1.23 U	2.30 U	143.07	80.00	86
CL9(206)	1.20 U	2.24 U	147.57	80.00	89
CL10(209)	1.33 U	2.49 U	131.96	80.00	79
<u>Surrogate Recoveries (%)</u>					
DBOFB	86	113	111	NA	NA
CL5(112)	79	72	74	NA	NA

TABLE B.6. (Contd)

Sample:	Method Blank	HU-A Rep. 3	HU-A Rep. 3 MS	Amount	Percent
Matrix:	Control Water	Elutriate	Elutriate	Spiked	Recovery
Sample Size (L):	0.94 ^(a)	0.47	0.50	4	4
Batch:	4	4	4	4	4
Units:	ng/L	ng/L	ng/L	ng	%
2,4-DDD	0.85 U	9.81	NS	NS	NA
2,4-DDT	0.86 U	1.62 U	NS	NS	NA
4,4-DDD	1.23 U	9.54	180.43	80.40	100
4,4-DDE	1.05 U	26.82	185.20	80.20	93
4,4-DDT	1.06 U	2.00 U	168.19	80.20	99
Aldrin	0.79 U	1.48 U	145.33	80.20	85
<i>alpha</i> -Chlordane	0.98 U	2.06	152.82	80.00	89
Dieldrin	1.05 U	4.72	129.96	80.20	73
Endosulfan I/2,4'-DDE	0.90 U	10.32	178.82	80.20	99
Endosulfan II	11.89 U	22.40 U	160.96	80.20	94
Endosulfan sulfate	8.69 U	16.37 U	167.71	80.20	98
Heptachlor	0.70 U	1.31 U	176.94	80.20	104
Heptachlor epoxide	0.91 U	0.47 J	176.62	80.20	103
<i>trans</i> -Nonachlor	1.02 U	1.20 J	NS	NS	NA
CL2(08)	0.93 U	1.75 U	C	80.00	NC
CL3(18)	1.13 U	7.52	107.87	80.00	59
CL3(28)	1.27 U	11.32	146.96	80.00	80
CL4(44)	1.29 U	12.98	129.37	80.00	68
CL4(49)	1.11 U	9.72	13.77	NS	NA
CL4(52)	1.31 U	17.50	127.11	80.00	64
CL4(66)	1.01 U	59.92	183.33	80.00	73
CL5(87)	1.14 U	5.12	5.28	NS	NA
CL5(101)	1.14 U	13.99	127.98	80.00	67
CL5(105)	1.37 U	2.31 J	155.08	80.00	90
CL5(118)	1.08 U	8.52	130.92	80.00	72
CL6(128)	1.21 U	4.25	146.69	80.00	84
CL6(138)	1.45 U	15.07	142.49	80.00	75
CL6(153)	1.39 U	10.27	114.82	80.00	61
CL7(170)	1.24 U	5.21	161.93	80.00	92
CL7(180)	1.08 U	8.42	152.31	80.00	85
CL7(183)	1.13 U	3.39	NS	NS	NA
CL7(184)	1.13 U	2.12 U	NS	NS	NA
CL7(187)	1.07 U	2.01 U	118.67	80.00	70
CL8(195)	1.22 U	3.11	163.38	80.00	94
CL9(206)	1.19 U	7.24	171.60	80.00	97
CL10(209)	1.32 U	6.82	153.12	80.00	86
<u>Surrogate Recoveries (%)</u>					
DBOFB	79	83	81	NA	NA
CL5(112)	71	71	65	NA	NA

(a) Sample concentration of the method blank adjusted for the average sample size of the batch.

(b) U Undetected at or above concentration shown.

(c) NS Not spiked.

(d) NA Not applicable.

(e) C PCB congener 08 coeluted with non-target pesticide α -BHC, resulting in unacceptable recovery in matrix spike samples.

(f) NC Percent recovery not calculated due to coeluting peak.

(g) J Concentration estimated; analyte detected below method detection limit (MDL) and above instrument detection limit (IDL).

(h) Outside quality control criteria (50-120%) for matrix spike recovery.

TABLE B.7. Quality Control Data (Triplicate Analyses) for Pesticides and PCBs in Site Water and Elutriate

Matrix	PC Rep. 1 Site Water	PC Rep. 2 Site Water	PC Rep. 3 Site Water	RSD ^(a)	EC-A Rep. 1 Site Water	EC-A Rep. 2 Site Water	EC-A Rep. 3 Site Water	RSD
Sample Size (L)	1.04	1.04	1.04		1.04	1.04	1.04	
Batch	1	1	1		1	1	1	
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.77 U ^(b)	0.77 U	0.77 U	NA ^(c)	0.77 U	0.77 U	0.70 J	NA
2,4-DDT	0.78 U	0.78 U	0.78 U	NA	0.78 U	0.78 U	0.78 U	NA
4,4-DDD	1.95	1.71	1.90	7%	4.99	3.50	3.89	19%
4,4-DDE	0.63 J ^(d)	0.60 J	0.81 J	16%	2.97	1.84	2.64	23%
4,4-DDT	0.96 U	1.70	0.90 J	NA	4.42	3.92	0.96 U	NA
Aldrin	0.71 U	0.71 U	0.71 U	NA	26.7	27.1	0.71 U	NA
<i>alpha</i> -Chlordane	1.80	1.94	1.76	5%	4.35	4.29	5.59	16%
Dieldrin	1.80	1.55	1.56	9%	3.24	1.76	2.53	30%
Endosulfan I/2,4'-DDE	0.81 U	0.81 U	0.81 U	NA	0.81 U	0.81 U	0.81 U	NA
Endosulfan II	1.57 J	10.8 U	10.8 U	NA	10.8 U	10.8 U	10.8 U	NA
Endosulfan sulfate	7.87 U	7.87 U	7.87 U	NA	7.87 U	7.87 U	7.87 U	NA
Heptachlor	0.63 U	0.63 U	0.63 U	NA	0.63 U	0.63 U	0.63 U	NA
Heptachlor epoxide	0.82 U	0.82 U	0.82 U	NA	0.82 U	0.82 U	0.82 U	NA
<i>trans</i> -Nonachlor	0.93 U	0.93 U	0.93 U	NA	1.62	1.60	3.03	39%
CL2(08)	0.84 U	0.84 U	0.84 U	NA	0.84 U	0.84 U	0.84 U	NA
CL3(18)	1.02 U	1.02 U	1.02 U	NA	1.80	1.02 U	1.02 U	NA
CL3(28)	4.20	2.69	3.05	24%	4.25	1.15 U	1.15 U	NA
CL4(44)	1.17 U	1.17 U	1.17 U	NA	2.97	2.59	1.17 U	NA
CL4(49)	1.01 U	1.01 U	1.01 U	NA	1.01 U	1.01 U	1.01 U	NA
CL4(52)	1.18 U	1.18 U	1.18 U	NA	2.96	2.30	1.18 U	NA
CL4(66)	0.92 U	0.92 U	0.92 U	NA	0.92 U	0.92 U	0.92 U	NA
CL5(87)	0.82 J	0.52 J	0.73 J	23%	1.96	0.69 J	1.41	47%
CL5(101)	1.04 U	1.04 U	1.04 U	NA	1.04 U	1.04 U	1.04 U	NA
CL5(105)	1.24 U	1.24 U	1.24 U	NA	0.71 J	0.66 J	1.24 U	NA
CL5(118)	0.98 U	0.98 U	0.98 U	NA	1.50	0.98 U	1.25	NA
CL6(128)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL6(138)	1.31 U	1.31 U	0.66 J	NA	1.41	1.28 J	1.31 U	NA
CL6(153)	1.26 U	1.26 U	0.96 J	NA	1.17 J	1.26	1.26 U	NA
CL7(170)	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	1.12 U	NA
CL7(180)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	0.98 U	NA
CL7(183)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(184)	1.02 U	1.02 U	1.02 U	NA	0.67 J	1.02 U	1.02 U	NA
CL7(187)	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	0.96 U	NA
CL8(195)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL9(206)	1.08 U	1.08 U	1.08 U	NA	1.08 U	1.08 U	1.08 U	NA
CL10(209)	1.20 U	1.20 U	1.20 U	NA	1.20 U	1.20 U	1.20 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	108	105	103	NA	100	112	114	NA
CL5(112)	72	72	71	NA	69	71	69	NA

TABLE B.7. (Contd)

Matrix	EC-B Rep. 1	EC-B Rep. 2	EC-B Rep. 3	RSD	HU-A Rep 1	HU-A Rep 2	HU-A Rep 3	RSD
Sample Size (L)	Site Water	Site Water	Site Water		Site Water	Site Water	Site Water	
Batch	1	1	1	1	1	1	1	
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.77 U	0.77 U	0.77 U	NA	0.77 U	0.77 U	0.77 U	NA
2,4-DDT	0.46 J	0.78 U	0.78 U	NA	0.78 U	0.78 U	0.78 U	NA
4,4-DDD	2.88	2.24	3.07	16%	1.12 U	1.12 U	1.12 U	NA
4,4-DDE	1.03	0.70 J	0.86 J	19%	0.95 U	0.95 U	0.95 U	NA
4,4-DDT	0.96 U	0.96 U	0.88 J	NA	0.96 U	0.96 U	0.96 U	NA
Aldrin	15.5	8.37	7.68	41%	0.71 U	0.71 U	0.71 U	NA
<i>alpha</i> -Chlordane	2.99	2.03	2.57	19%	0.89 U	0.68 J	0.89 U	NA
Dieldrin	1.80	1.14	2.80	44%	2.28	1.42	1.21	35%
Endosulfan I/2,4'-DDE	0.81 U	0.81 U	0.81 U	NA	0.81 U	0.81 U	0.81 U	NA
Endosulfan II	10.8 U	10.8 U	10.8 U	NA	10.8 U	10.8 U	10.8 U	NA
Endosulfan sulfate	7.87 U	7.87 U	7.87 U	NA	7.87 U	7.87 U	7.87 U	NA
Heptachlor	0.63 U	0.63 U	0.63 U	NA	0.63 U	0.63 U	0.63 U	NA
Heptachlor epoxide	0.82 U	0.82 U	0.82 U	NA	0.82 U	0.82 U	0.82 U	NA
<i>trans</i> -Nonachlor	1.00	1.01	1.74	34%	0.93 U	0.93 U	0.93 U	NA
CL2(08)	0.84 U	0.84 U	0.84 U	NA	0.84 U	0.84 U	0.84 U	NA
CL3(18)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL3(28)	7.34	4.16	5.59	28%	1.15 U	1.15 U	1.15 U	NA
CL4(44)	1.17 U	1.17 U	1.94	NA	1.17 U	1.17 U	1.17 U	NA
CL4(49)	1.01 U	1.01 U	1.01 U	NA	1.01 U	1.01 U	1.01 U	NA
CL4(52)	1.18 U	1.18 U	1.18 U	NA	1.18 U	1.18 U	1.18 U	NA
CL4(66)	0.92 U	0.92 U	0.92 U	NA	0.92 U	0.92 U	0.92 U	NA
CL5(87)	0.76 J	0.75 J	1.45	40%	1.56	2.51	2.32	24%
CL5(101)	1.04 U	1.04 U	1.04 U	NA	1.33	0.96 J	1.13	16%
CL5(105)	1.24 U	1.24 U	1.24 U	NA	1.24 U	1.24 U	1.24 U	NA
CL5(118)	0.56 J	0.52 J	0.87 J	29%	0.98 U	0.98 U	0.98 U	NA
CL6(128)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL6(138)	1.31 U	1.31 U	1.45	NA	1.31 U	1.31 U	1.31 U	NA
CL6(153)	0.88 J	0.62 J	0.83 J	18%	1.26 U	1.26 U	1.26 U	NA
CL7(170)	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	1.12 U	NA
CL7(180)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	0.98 U	NA
CL7(183)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(184)	1.02 U	1.02 U	0.50 J	NA	1.02 U	1.02 U	1.02 U	NA
CL7(187)	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	0.96 U	NA
CL8(195)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL9(206)	1.08 U	1.08 U	1.08 U	NA	1.08 U	1.08 U	1.08 U	NA
CL10(209)	1.20 U	1.20 U	1.20 U	NA	1.20 U	1.20 U	1.20 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	108	64	112	NA	86	75	90	NA
CL5(112)	69	42	67	NA	72	69	70	NA

TABLE B.7. (Contd)

Matrix	SB-A Rep 1	SB-A Rep 2	SB-A Rep 3	RSD	SB-B Rep 1	SB-B Rep 2	SB-B Rep 3	RSD
Sample Size (L)	Site Water	Site Water	Site Water		Water	Water	Water	
Batch	1	1	1	1	1	1	1	1
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.77 U	0.77 U	0.77 U	NA	0.77 U	0.77 U	1.52 U	NA
2,4-DDT	0.78 U	0.78 U	0.78 U	NA	0.78 U	0.78 U	1.54 U	NA
4,4-DDD	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	2.21 U	NA
4,4-DDE	0.95 U	0.95 U	0.95 U	NA	0.95 U	0.95 U	1.88 U	NA
4,4-DDT	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	1.90 U	NA
Aldrin	0.71 U	0.71 U	0.71 U	NA	0.71 U	0.71 U	1.41 U	NA
<i>alpha</i> -Chlordane	0.89 U	0.89 U	0.89 U	NA	0.89 U	0.89 U	1.77 U	NA
Dieldrin	0.95 U	1.41	0.95 U	NA	0.95 U	2.18	2.64	NA
Endosulfan I/2,4'-DDE	0.81 U	0.81 U	0.81 U	NA	0.81 U	0.81 U	1.61 U	NA
Endosulfan II	10.8 U	10.8 U	10.8 U	NA	10.8 U	10.8 U	21.3 U	NA
Endosulfan sulfate	7.87 U	7.87 U	7.87 U	NA	7.87 U	7.87 U	15.6 U	NA
Heptachlor	0.63 U	0.63 U	0.63 U	NA	0.63 U	0.63 U	1.25 U	NA
Heptachlor epoxide	0.82 U	0.82 U	0.82 U	NA	0.82 U	0.82 U	1.63 U	NA
<i>trans</i> -Nonachlor	0.93 U	0.93 U	0.93 U	NA	0.93 U	0.93 U	1.84 U	NA
CL2(08)	0.84 U	0.84 U	0.84 U	NA	0.84 U	0.84 U	1.67 U	NA
CL3(18)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	2.03 U	NA
CL3(28)	1.15 U	1.15 U	1.15 U	NA	1.15 U	1.15 U	2.27 U	NA
CL4(44)	1.17 U	1.17 U	1.17 U	NA	1.17 U	1.17 U	2.32 U	NA
CL4(49)	1.01 U	1.01 U	1.01 U	NA	1.01 U	1.01 U	1.99 U	NA
CL4(52)	1.18 U	1.18 U	1.18 U	NA	1.18 U	2.48	2.34 U	NA
CL4(66)	0.92 U	0.92 U	0.92 U	NA	0.92 U	0.92 U	1.82 U	NA
CL5(87)	1.03 U	1.03 U	1.03 U	NA	1.03 U	2.15	2.04 U	NA
CL5(101)	1.04 U	1.23	1.04 U	NA	1.04 U	0.99 J	2.05 U	NA
CL5(105)	1.24 U	1.24 U	1.24 U	NA	1.24 U	1.24 U	2.46 U	NA
CL5(118)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	1.94 U	NA
CL6(128)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	2.17 U	NA
CL6(138)	1.31 U	1.31 U	1.31 U	NA	1.31 U	1.31 U	2.60 U	NA
CL6(153)	1.26 U	1.26 U	1.26 U	NA	1.26 U	1.26 U	2.49 U	NA
CL7(170)	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	2.23 U	NA
CL7(180)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	1.93 U	NA
CL7(183)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	2.02 U	NA
CL7(184)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	2.02 U	NA
CL7(187)	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	1.91 U	NA
CL8(195)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	2.19 U	NA
CL9(206)	1.08 U	1.08 U	1.08 U	NA	1.08 U	1.08 U	2.14 U	NA
CL10(209)	1.20 U	1.20 U	1.20 U	NA	1.20 U	1.20 U	2.38 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	82	94	104	NA	73	97	99	NA
CL5(112)	58	72	74	NA	61	67	74	NA

TABLE B.7. (Contd)

Matrix	BU Rep. 1	BU Rep. 2	BU Rep. 3	RSD	Mud Dump	Mud Dump	Mud Dump	RSD
	Site Water	Site Water	Site Water		Site Rep. 1	Site Rep. 2	Site Rep. 3	
Sample Size (L)	1.04	1.04	1.04		1.04	1.04	1.04	
Batch	2	2	2	2	2	2	2	2
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.77 U	0.77 U	0.77 U	NA	0.77 U	0.77 U	0.77 U	NA
2,4-DDT	0.78 U	0.78 U	0.78 U	NA	0.78 U	0.78 U	0.78 U	NA
4,4-DDD	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	1.12 U	NA
4,4-DDE	0.95 U	0.95 U	0.95 U	NA	0.95 U	0.95 U	0.95 U	NA
4,4-DDT	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	0.96 U	NA
Aldrin	0.71 U	0.71 U	0.71 U	NA	0.71 U	0.71 U	0.71 U	NA
<i>alpha</i> -Chlordane	0.89 U	0.89 U	0.89 U	NA	0.89 U	0.89 U	0.89 U	NA
Dieldrin	0.95 U	0.95 U	0.95 U	NA	0.95 U	0.95 U	0.95 U	NA
Endosulfan I/2,4'-DDE	0.81 U	0.81 U	0.81 U	NA	0.81 U	0.81 U	0.81 U	NA
Endosulfan II	10.8 U	10.8 U	10.8 U	NA	10.8 U	10.8 U	10.8 U	NA
Endosulfan sulfate	7.87 U	7.87 U	7.87 U	NA	7.87 U	7.87 U	7.87 U	NA
Heptachlor	0.63 U	0.63 U	0.63 U	NA	0.63 U	0.63 U	0.63 U	NA
Heptachlor epoxide	0.82 U	0.82 U	0.82 U	NA	0.82 U	0.82 U	0.82 U	NA
<i>trans</i> -Nonachlor	0.93 U	0.93 U	0.93 U	NA	0.93 U	0.93 U	0.93 U	NA
CL2(08)	0.84 U	0.84 U	0.84 U	NA	0.84 U	0.84 U	0.84 U	NA
CL3(18)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL3(26)	1.15 U	1.15 U	1.15 U	NA	1.15 U	1.15 U	1.15 U	NA
CL4(44)	1.17 U	1.17 U	1.17 U	NA	1.17 U	1.17 U	1.17 U	NA
CL4(49)	4.25	1.01 U	1.01 U	NA	1.01 U	1.01 U	1.01 U	NA
CL4(52)	1.18 U	1.18 U	1.18 U	NA	1.18 U	1.18 U	1.18 U	NA
CL4(66)	0.92 U	0.92 U	0.92 U	NA	0.92 U	0.92 U	0.92 U	NA
CL5(87)	1.03 U	1.03 U	1.03 U	NA	1.03 U	1.03 U	1.03 U	NA
CL5(101)	1.04 U	1.04 U	1.04 U	NA	1.04 U	1.04 U	1.04 U	NA
CL5(105)	1.24 U	1.24 U	1.24 U	NA	1.24 U	1.24 U	1.24 U	NA
CL5(118)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	0.98 U	NA
CL6(128)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL6(138)	1.31 U	1.31 U	1.31 U	NA	1.31 U	1.31 U	1.31 U	NA
CL6(153)	1.26 U	1.26 U	1.26 U	NA	1.26 U	1.26 U	1.26 U	NA
CL7(170)	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	1.12 U	NA
CL7(180)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	0.98 U	NA
CL7(183)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(184)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(187)	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	0.96 U	NA
CL8(195)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL9(206)	1.08 U	1.08 U	1.08 U	NA	1.08 U	1.08 U	1.08 U	NA
CL10(209)	1.20 U	1.20 U	1.20 U	NA	1.20 U	1.20 U	1.20 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	30	51	44	NA	45	49	44	NA
CL5(112)	47	57	58	NA	52	56	56	NA

TABLE B.7. (Contd)

Matrix	HU-B Rep. 1	HU-B Rep. 2	HU-B Rep. 3	RSD	HU-C Rep. 1	HU-C Rep. 2	HU-C Rep. 3	RSD
Sample Size (L)	1.04	1.04	1.04		1.04	1.04	1.04	
Batch	2	2	2	2	2	2	2	2
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.77 U	0.77 U	0.77 U	NA	0.77 U	0.77 U	0.77 U	NA
2,4-DDT	0.78 U	0.78 U	0.78 U	NA	0.78 U	0.78 U	0.78 U	NA
4,4-DDD	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	1.12 U	NA
4,4-DDE	0.95 U	0.95 U	0.95 U	NA	0.95 U	0.95 U	0.95 U	NA
4,4-DDT	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	0.96 U	NA
Aldrin	14.7	0.71 U	0.71 U	NA	0.71 U	0.71 U	0.71 U	NA
<i>alpha</i> -Chlordane	0.89 U	0.89 U	0.89 U	NA	0.89 U	0.89 U	0.89 U	NA
Dieldrin	0.95 U	0.95 U	0.95 U	NA	0.95 U	0.95 U	0.95 U	NA
Endosulfan I/2,4'-DDE	0.81 U	0.81 U	0.81 U	NA	0.81 U	0.81 U	0.81 U	NA
Endosulfan II	10.8 U	10.8 U	10.8 U	NA	10.8 U	10.8 U	10.8 U	NA
Endosulfan sulfate	7.87 U	7.87 U	7.87 U	NA	7.87 U	7.87 U	7.87 U	NA
Heptachlor	0.63 U	0.63 U	0.63 U	NA	0.63 U	0.63 U	0.63 U	NA
Heptachlor epoxide	0.82 U	0.82 U	0.82 U	NA	0.82 U	0.82 U	0.82 U	NA
<i>trans</i> -Nonachlor	0.93 U	0.93 U	0.93 U	NA	0.93 U	0.93 U	0.93 U	NA
CL2(08)	0.84 U	0.84 U	0.84 U	NA	0.84 U	0.84 U	0.84 U	NA
CL3(18)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL3(28)	1.15 U	1.15 U	1.15 U	NA	1.15 U	1.15 U	1.15 U	NA
CL4(44)	1.17 U	1.17 U	1.17 U	NA	1.17 U	1.17 U	1.17 U	NA
CL4(49)	1.88	2.22	2.27	10%	1.01 U	1.01 U	1.01 U	NA
CL4(52)	1.18 U	2.08	2.02	NA	1.95	2.10	1.87	6%
CL4(66)	0.92 U	0.81 J	0.92 U	NA	0.92 U	0.92 U	0.92 U	NA
CL5(87)	1.03 U	1.03 U	1.03 U	NA	1.03 U	1.03 U	1.03 U	NA
CL5(101)	1.04 U	1.04 U	1.04 U	NA	1.04 U	1.04 U	1.04 U	NA
CL5(105)	1.24 U	1.24 U	1.24 U	NA	1.24 U	1.24 U	1.24 U	NA
CL5(118)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	0.98 U	NA
CL6(128)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL6(138)	1.31 U	1.31 U	1.31 U	NA	1.31 U	1.31 U	1.31 U	NA
CL6(153)	1.26 U	1.26 U	1.26 U	NA	1.26 U	1.26 U	1.26 U	NA
CL7(170)	1.12 U	1.12 U	1.12 U	NA	1.12 U	1.12 U	1.12 U	NA
CL7(180)	0.98 U	0.98 U	0.98 U	NA	0.98 U	0.98 U	0.98 U	NA
CL7(183)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(184)	1.02 U	1.02 U	1.02 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(187)	0.96 U	0.96 U	0.96 U	NA	0.96 U	0.96 U	0.96 U	NA
CL8(195)	1.10 U	1.10 U	1.10 U	NA	1.10 U	1.10 U	1.10 U	NA
CL9(206)	1.08 U	1.08 U	1.08 U	NA	1.08 U	1.08 U	1.08 U	NA
CL10(209)	1.20 U	1.20 U	1.20 U	NA	1.20 U	1.20 U	1.20 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	47	51	49	NA	49	41	53	NA
CL5(112)	57	63	57	NA	61	57	59	NA

TABLE B.7. (Contd)

Matrix	HU-D Rep. 1	HU-D Rep. 2	HU-D Rep. 3	RSD	GR Rep. 1	GR Rep. 2	GR Rep. 3	RSD
Sample Size (L)	Site Water	Site Water	Site Water		Water	Water	Water	
Batch	2	2	2	2	2	2	2	2
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.77 U	0.77 U	1.53 U	NA	0.77 U	0.77 U	0.77 U	NA
2,4-DDT	0.78 U	0.78 U	1.55 U	NA	0.78 U	0.78 U	0.78 U	NA
4,4-DDD	1.12 U	1.12 U	2.23 U	NA	1.12 U	1.12 U	1.12 U	NA
4,4-DDE	0.95 U	0.95 U	1.90 U	NA	0.95 U	0.95 U	0.95 U	NA
4,4-DDT	0.96 U	0.96 U	1.92 U	NA	0.96 U	0.96 U	0.96 U	NA
Aldrin	0.71 U	0.71 U	1.43 U	NA	0.71 U	0.71 U	0.71 U	NA
<i>alpha</i> -Chlordane	0.89 U	0.89 U	1.72 J	NA	0.89 U	0.89 U	0.89 U	NA
Dieldrin	0.95 U	0.95 U	1.53 J	NA	0.95 U	0.95 U	0.95 U	NA
Endosulfan I/2,4'-DDE	0.81 U	0.81 U	1.63 U	NA	0.81 U	0.81 U	0.81 U	NA
Endosulfan II	10.8 U	10.8 U	2.71 J	NA	10.8 U	10.8 U	10.8 U	NA
Endosulfan sulfate	7.87 U	7.87 U	15.7 U	NA	7.87 U	7.87 U	7.87 U	NA
Heptachlor	0.63 U	0.63 U	1.26 U	NA	0.63 U	0.63 U	0.63 U	NA
Heptachlor epoxide	0.82 U	0.82 U	1.64 U	NA	0.82 U	0.82 U	0.82 U	NA
<i>trans</i> -Nonachlor	0.93 U	0.93 U	1.86 U	NA	0.93 U	0.93 U	0.93 U	NA
CL2(08)	0.84 U	0.84 U	1.68 U	NA	0.84 U	0.84 U	0.84 U	NA
CL3(18)	1.02 U	1.02 U	2.05 U	NA	1.02 U	1.02 U	1.02 U	NA
CL3(28)	1.15 U	1.15 U	2.29 U	NA	1.15 U	1.15 U	1.15 U	NA
CL4(44)	1.17 U	1.17 U	2.34 U	NA	1.17 U	1.17 U	1.17 U	NA
CL4(49)	1.01 U	1.01 U	2.01 U	NA	3.46	2.79	3.21	11%
CL4(52)	1.16 J	1.51	2.37 U	NA	1.18 U	1.18 U	1.18 U	NA
CL4(66)	0.92 U	0.92 U	1.83 U	NA	0.92 U	0.92 U	0.92 U	NA
CL5(87)	1.03 U	1.03 U	2.06 U	NA	1.03 U	1.03 U	1.03 U	NA
CL5(101)	1.04 U	1.04 U	2.07 U	NA	1.04 U	1.04 U	1.04 U	NA
CL5(105)	1.24 U	1.24 U	2.48 U	NA	1.24 U	1.24 U	1.24 U	NA
CL5(118)	0.98 U	0.98 U	1.95 U	NA	0.98 U	0.98 U	0.98 U	NA
CL6(128)	1.10 U	1.10 U	2.19 U	NA	1.10 U	1.10 U	1.10 U	NA
CL6(138)	1.31 U	1.31 U	2.62 U	NA	1.31 U	1.31 U	1.31 U	NA
CL6(153)	1.26 U	1.26 U	2.52 U	NA	1.26 U	1.26 U	1.26 U	NA
CL7(170)	1.12 U	1.12 U	2.25 U	NA	1.12 U	1.12 U	1.12 U	NA
CL7(180)	0.98 U	0.98 U	1.95 U	NA	0.98 U	0.98 U	0.98 U	NA
CL7(183)	1.02 U	1.02 U	2.04 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(184)	1.02 U	1.02 U	2.04 U	NA	1.02 U	1.02 U	1.02 U	NA
CL7(187)	0.96 U	0.96 U	1.93 U	NA	0.96 U	0.96 U	0.96 U	NA
CL8(195)	1.10 U	1.10 U	2.21 U	NA	1.10 U	1.10 U	1.10 U	NA
CL9(206)	1.08 U	1.08 U	2.16 U	NA	1.08 U	1.08 U	1.08 U	NA
CL10(209)	1.20 U	1.20 U	2.40 U	NA	1.20 U	1.20 U	1.20 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	57	70	32	NA	37	36	47	NA
CL5(112)	59	63	49	NA	60	55	60	NA

TABLE B.7. (Contd)

Matrix	PC Rep. 1	PC Rep. 2	PC Rep. 3	RSD	SB-B Rep. 1	SB-B Rep. 2	SB-B Rep. 3	RSD
Sample Size (L)	Elutriate	Elutriate	Elutriate		Elutriate	Elutriate	Elutriate	
Batch	3	3	3	3	3	3	3	3
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	11.1	13.5	17.9	24%	0.82 U	0.81 U	0.81 U	NA
2,4-DDT	5.01	4.62	5.47	8%	0.83 U	0.82 U	0.82 U	NA
4,4-DDD	42.1	48.9	75.1	31% ^(b)	1.20 U	1.18 U	1.18 U	NA
4,4-DDE	11.6	13.8	22.0	35% ^(b)	1.02 U	1.01 U	1.01 U	NA
4,4-DDT	1.15 U	1.04 U	1.05 U	NA	1.03 U	1.02 U	1.02 U	NA
Aldrin	0.85 U	0.77 U	0.78 U	NA	0.76 U	0.76 U	0.76 U	NA
<i>alpha</i> -Chlordane	13.4	14.9	21.1	25%	0.96 U	0.95 U	0.95 U	NA
Dieldrin	9.36	11.2	14.8	24%	1.02 U	1.01 U	1.01 U	NA
Endosulfan 1/2,4'-DDE	0.97 U	0.88 U	0.89 U	NA	0.87 U	0.86 U	0.86 U	NA
Endosulfan-II	4.93 J	4.73 J	6.70 J	20%	11.5 U	11.4 U	11.4 U	NA
Endosulfan sulfate	11.5	13.5	18.0	23%	8.44 U	8.35 U	8.35 U	NA
Heptachlor	0.75 U	0.68 U	0.69 U	NA	0.68 U	0.67 U	0.67 U	NA
Heptachlor epoxide	0.98 U	0.89 U	0.90 U	NA	0.88 U	0.87 U	0.87 U	NA
<i>trans</i> -Nonachlor	6.55	7.38	10.3	25%	0.99 U	0.98 U	0.98 U	NA
CL2(08)	1.01 U	0.91 U	0.92 U	NA	0.90 U	0.89 U	0.89 U	NA
CL3(18)	1.22 U	1.11 U	1.12 U	NA	1.10 U	1.09 U	1.09 U	NA
CL3(28)	5.32	5.88	6.89	13%	1.23 U	1.22 U	1.22 U	NA
CL4(44)	12.2	14.8	19.5	24%	1.25 U	1.24 U	1.24 U	NA
CL4(49)	7.62	7.50	11.4	25%	1.08 U	1.07 U	1.07 U	NA
CL4(52)	24.5	27.5	41.4	29%	1.27 U	1.26 U	1.26 U	NA
CL4(66)	9.78	11.8	21.5	44% ^(b)	0.98 U	0.97 U	0.97 U	NA
CL5(87)	25.0	26.6	37.1	22%	1.10 U	1.09 U	1.09 U	NA
CL5(101)	67.2	79.1	118	30%	1.11 U	1.10 U	1.10 U	NA
CL5(105)	30.6	34.2	30.0	7%	1.33 U	1.32 U	1.32 U	NA
CL5(118)	47.0	52.5	79.1	29%	1.05 U	1.04 U	1.04 U	NA
CL6(128)	8.85	10.6	14.9	27%	1.18 U	1.16 U	1.16 U	NA
CL6(138)	56.4	66.1	96.5	29%	1.41 U	1.39 U	1.39 U	NA
CL6(153)	35.9	39.0	67.7	37% ^(c)	1.35 U	1.33 U	1.33 U	NA
CL7(170)	11.3	15.7	22.3	33% ^(c)	1.21 U	1.19 U	1.19 U	NA
CL7(180)	26.2	29.5	44.9	30%	1.05 U	1.03 U	1.03 U	NA
CL7(183)	5.57	5.91	8.02	20%	1.09 U	1.08 U	1.08 U	NA
CL7(184)	1.22 U	1.11 U	1.12 U	NA	1.09 U	1.08 U	1.08 U	NA
CL7(187)	18.0	20.1	28.0	24%	1.03 U	1.02 U	1.02 U	NA
CL8(195)	3.00	3.41	5.39	32%	1.18 U	1.17 U	1.17 U	NA
CL9(206)	6.07	7.20	11.0	32%	1.16 U	1.14 U	1.14 U	NA
CL10(209)	1.28 J	1.37	1.97	25%	1.29 U	1.27 U	1.27 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	120	120	123	NA	102	101	98	NA
CL5(112)	71	66	58	NA	75	76	82	NA

TABLE B.7. (Contd)

Matrix	SB-A Rep. 1	SB-A Rep. 2	SB-A Rep. 3	RSD	BU Rep. 1	BU Rep. 2	BU Rep. 3	RSD
Sample Size (L)	Elutriate	Elutriate	Elutriate		Elutriate	Elutriate	Elutriate	
Batch	1.00	0.995	0.995		0.95	0.96	0.98	
Units	3	3	3	3	3	3	3	3
	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	0.80 U	0.80 U	0.80 U	NA	0.84 U	0.83 U	0.81 U	NA
2,4-DDT	0.81 U	0.81 U	0.81 U	NA	0.85 U	0.84 U	0.82 U	NA
4,4-DDD	1.16 U	1.17 U	1.17 U	NA	1.22 U	1.21 U	1.18 U	NA
4,4-DDE	0.99 U	0.99 U	0.99 U	NA	1.04 U	1.03 U	1.01 U	NA
4,4-DDT	1.00 U	1.01 U	1.01 U	NA	1.05 U	1.04 U	1.02 U	NA
Aldrin	0.74 U	0.74 U	0.74 U	NA	0.78 U	0.77 U	0.76 U	NA
<i>alpha</i> -Chlordane	0.93 U	0.93 U	0.93 U	NA	0.98 U	0.97 U	0.95 U	NA
Dieldrin	0.99 U	0.99 U	0.99 U	NA	1.04 U	1.03 U	1.01 U	NA
Endosulfan I/2,4'-DDE	0.85 U	0.85 U	0.85 U	NA	0.89 U	0.88 U	0.86 U	NA
Endosulfan II-	11.2 U	11.3 U	11.3 U	NA	11.8 U	11.7 U	11.4 U	NA
Endosulfan sulfate	8.19 U	8.23 U	8.23 U	NA	8.62 U	8.53 U	8.35 U	NA
Heptachlor	0.66 U	0.66 U	0.66 U	NA	0.69 U	0.68 U	0.67 U	NA
Heptachlor epoxide	0.86 U	0.86 U	0.86 U	NA	0.90 U	0.89 U	0.87 U	NA
<i>trans</i> -Nonachlor	0.97 U	0.97 U	0.97 U	NA	1.02 U	1.01 U	0.98 U	NA
CL2(08)	0.88 U	0.88 U	0.88 U	NA	0.92 U	0.91 U	0.89 U	NA
CL3(18)	1.07 U	1.07 U	1.07 U	NA	1.12 U	1.11 U	1.09 U	NA
CL3(28)	1.19 U	1.20 U	1.20 U	NA	1.26 U	1.24 U	1.22 U	NA
CL4(44)	1.22 U	1.22 U	1.22 U	NA	1.28 U	1.27 U	1.24 U	NA
CL4(49)	1.05 U	1.05 U	0.74 J	NA	1.10 U	1.09 U	1.07 U	NA
CL4(52)	1.23 U	1.24 U	2.12	NA	1.29 U	1.28 U	1.26 U	NA
CL4(66)	0.95 U	0.96 U	0.96 U	NA	1.00 U	0.99 U	0.97 U	NA
CL5(87)	1.07 U	1.07 U	1.07 U	NA	1.13 U	1.11 U	1.09 U	NA
CL5(101)	1.08 U	1.08 U	1.22	NA	1.13 U	1.12 U	1.10 U	NA
CL5(105)	1.29 U	1.30 U	1.30 U	NA	1.36 U	1.34 U	1.32 U	NA
CL5(118)	1.02 U	1.02 U	1.02 U	NA	1.07 U	1.06 U	1.04 U	NA
CL6(126)	1.14 U	1.15 U	1.15 U	NA	1.20 U	1.19 U	1.16 U	NA
CL6(136)	1.36 U	1.37 U	1.37 U	NA	1.43 U	1.42 U	1.39 U	NA
CL6(153)	1.31 U	1.31 U	1.31 U	NA	1.38 U	1.36 U	1.33 U	NA
CL7(170)	1.17 U	1.17 U	1.17 U	NA	1.23 U	1.22 U	1.19 U	NA
CL7(180)	1.01 U	1.02 U	1.02 U	NA	1.07 U	1.06 U	1.03 U	NA
CL7(183)	1.06 U	1.07 U	1.07 U	NA	1.12 U	1.11 U	1.08 U	NA
CL7(184)	1.06 U	1.07 U	1.07 U	NA	1.12 U	1.11 U	1.08 U	NA
CL7(187)	1.00 U	1.01 U	1.01 U	NA	1.06 U	1.04 U	1.02 U	NA
CL8(195)	1.15 U	1.15 U	1.15 U	NA	1.21 U	1.20 U	1.17 U	NA
CL9(206)	1.12 U	1.13 U	1.13 U	NA	1.18 U	1.17 U	1.14 U	NA
CL10(209)	1.25 U	1.25 U	1.25 U	NA	1.31 U	1.30 U	1.27 U	NA
<u>Surrogate Recoveries (%)</u>								
DBOFB	101	94	98	NA	96	88	95	NA
CL5(112)	75	80	77	NA	74	75	81	NA

TABLE B.7. (Contd)

Matrix Sample Size (L) Batch Units	EC-B Rep. 1	EC-B Rep. 2	EC-B Rep. 3	RSD	EC-A Rep. 1	EC-A Rep. 2	EC-A Rep. 3	RSD
	Elutriate 0.96 3 ng/L	Elutriate 0.98 3 ng/L	Elutriate 0.50 3 ng/L		Elutriate 0.90 4 ng/L	Elutriate 0.91 4 ng/L	Elutriate 0.92 4 ng/L	
2,4-DDD	3.30	1.82	3.07	29%	2.33	3.20	2.49	17%
2,4-DDT	0.912	0.647 J	0.925 J	19%	0.90 U	0.89 U	0.88 U	NA
4,4-DDD	12.2	6.58	12.2	32%	5.21	4.06	4.49	13%
4,4-DDE	6.27	2.65	6.55	42%	7.99	7.13	6.98	7%
4,4-DDT	1.04 U	1.02 U	2.00 U	NA	1.11 U	1.10 U	1.09 U	NA
Aldrin	14.1	14.9	22.5	27%	0.82 U	0.81 U	0.81 U	NA
<i>alpha</i> -Chlordane	10.0	7.93	13.2	26%	1.43	1.24	1.38	7%
Dieldrin	3.25	2.87	3.80	14%	2.36	2.53	1.66	21%
Endosulfan I/2,4'-DDE	0.88 U	0.86 U	1.69 U	NA	0.94 U	0.93 U	0.92 U	NA
Endosulfan II	11.7 U	11.4 U	22.4 U	NA	12.4 U	12.3 U	12.2 U	NA
Endosulfan sulfate	8.53 U	8.35 U	16.4 U	NA	9.10 U	9.00 U	8.95 U	NA
Heptachlor	0.68 U	0.67 U	1.31 U	NA	0.73 U	0.72 U	0.72 U	NA
Heptachlor epoxide	0.89 U	0.87 U	1.71 U	NA	0.95 U	0.94 U	0.93 U	NA
<i>trans</i> -Nonachlor	6.11	3.94	7.17	29%	0.86 J	0.95 J	0.77 J	10%
CL2(08)	0.91 U	0.89 U	1.75 U	NA	4.26	3.54	4.44	12%
CL3(18)	1.11 U	1.09 U	2.13 U	NA	3.68	4.90	2.30	36%
CL3(28)	6.66	4.10	15.3	68%	9.82	6.22	6.74	26%
CL4(44)	7.88	3.73	12.4	54%	7.46	7.71	5.79	15%
CL4(49)	9.33	4.65	8.62	33%	4.76	3.71	2.83	26%
CL4(52)	39.1	31.06	66.5	41% ^(a)	11.6	10.5	12.5	9%
CL4(66)	19.9	20.11	17.8	7%	35.9	40.5	33.6	10%
CL5(87)	3.13	2.24	4.94	40%	1.82	1.70	1.50	10%
CL5(101)	6.84	5.66	11.6	39%	3.93	3.82	3.90	1%
CL5(105)	1.94	1.81	1.88 J	3%	1.42 J	2.00	1.28 J	24%
CL5(118)	7.55	4.74	9.71	34%	4.42	3.69	3.70	11%
CL6(128)	1.97	1.69	2.54	21%	1.27 U	1.25 U	1.25 U	NA
CL6(138)	9.97	2.83	11.1	56%	5.12	4.29	5.01	9%
CL6(153)	5.18	3.55	7.32	35%	3.42	3.17	2.66	13%
CL7(170)	1.22 U	1.19 U	2.34 U	NA	2.60	2.09	2.19	12%
CL7(180)	1.06 U	1.03 U	2.03 U	NA	2.60	2.08	2.07	13%
CL7(183)	1.39	0.72 J	2.09 J	NA	0.71 J	0.61 J	0.60 J	9%
CL7(184)	1.11 U	1.08 U	2.12 U	NA	1.18 U	1.17 U	1.16 U	NA
CL7(187)	1.04 U	1.02 U	2.01 U	NA	1.79	1.10 U	1.10 U	NA
CL8(195)	1.20 U	1.17 U	2.30 U	NA	0.41 J	0.43 J	0.69 J	31%
CL9(206)	1.17 U	1.14 U	2.24 U	NA	0.87 J	0.61 J	0.61 J	21%
CL10(209)	1.30 U	1.27 U	2.49 U	NA	0.86 J	0.93 J	0.92 J	5%
<u>Surrogate Recoveries (%)</u>								
DBOFB	111	115	113	NA	70	70	64	NA
CL5(112)	72	72	72	NA	56	63	53	NA

TABLE B.7. (Contd)

Matrix	HU-A Rep 1	HU-A Rep 2	HU-A Rep 3	RSD	HU-D Rep. 1	HU-D Rep. 2	HU-D Rep. 3	RSD
	Elutriate	Elutriate	Elutriate		Elutriate	Elutriate	Elutriate	
Sample Size (L)	0.98	0.97	0.50		0.98	0.96	0.96	
Batch	4	4	4	4	4	4	4	4
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	16.6	8.38	9.81	38% ^(a)	3.94	6.65	8.29	35%
2,4-DDT	0.83 U	0.83 U	1.62 U	NA	0.82 U	0.84 U	0.84 U	NA
4,4-DDD	13.4	8.49	9.54	25%	3.50	2.37	5.01	36%
4,4-DDE	52.1	28.4	26.8	40% ^(a)	9.47	5.05	9.47	32%
4,4-DDT	1.03 U	1.03 U	2.00 U	NA	1.02 U	1.04 U	1.04 U	NA
Aldrin	0.76 U	0.76 U	1.48 U	NA	0.76 U	0.77 U	0.77 U	NA
<i>alpha</i> -Chlordane	3.45	1.81	2.06	36%	1.27	0.27 J	1.56	66%
Dieldrin	5.64	4.31	4.72	14%	5.14	2.33	4.13	37%
Endosulfan I/2,4'-DDE	17.0	10.4	10.3	31% ^(a)	0.86 U	0.88 U	0.88 U	NA
Endosulfan II	11.5 U	11.5 U	22.4 U	NA	11.4 U	1.70 J	11.7 U	NA
Endosulfan sulfate	8.40 U	8.44 U	16.4 U	NA	5.37 J	8.53 U	2.88 J	NA
Heptachlor	0.67 U	0.68 U	1.31 U	NA	0.67 U	0.68 U	0.68 U	NA
Heptachlor epoxide	3.25	1.59	0.47 J	79%	0.87 U	0.89 U	0.89 U	NA
<i>trans</i> -Nonachlor	0.85 J	0.83 J	1.20 J	21%	0.65 J	1.01 U	1.00 J	NA
CL2(08)	1.75	1.99	1.75 U	NA	0.89 U	0.91 U	0.91 U	NA
CL3(18)	16.0	9.25	7.52	41%	18.0	8.50	14.9	35% ^(a)
CL3(28)	19.9	11.3	11.3	35% ^(a)	10.7	6.75	11.1	25%
CL4(44)	17.2	11.9	13.0	20%	14.3	8.22	15.0	30%
CL4(49)	16.8	11.0	9.72	30%	13.5	6.39	12.9	36%
CL4(52)	23.4	15.6	17.5	22%	16.9	9.44	19.1	34% ^(a)
CL4(66)	72.7	48.4	59.9	20%	44.1	31.6	49.3	22%
CL5(87)	8.62	5.34	5.12	31%	4.08	2.38	4.89	34%
CL5(101)	21.9	13.6	14.0	28%	9.57	5.72	11.9	34%
CL5(105)	3.56	2.51	2.31 J	24%	1.98	1.36	2.70	33%
CL5(118)	14.9	8.02	8.52	37%	7.57	4.00	8.63	36%
CL6(128)	5.38	3.40	4.25	23%	2.32	0.84 J	2.46	48%
CL6(138)	24.5	14.4	15.1	31% ^(a)	10.3	1.42 U	1.42 U	NA
CL6(153)	19.2	10.3	10.3	39% ^(a)	8.70	4.21	9.28	37%
CL7(170)	7.88	4.82	5.21	28%	3.55	1.52	3.13	39%
CL7(180)	17.4	9.73	8.42	41% ^(a)	5.78	2.58	5.98	40%
CL7(183)	4.43	2.61	3.39	26%	1.89	0.78 J	1.57	41%
CL7(184)	1.09 U	1.09 U	2.12 U	NA	1.08 U	1.11 U	1.11 U	NA
CL7(187)	1.03 U	1.03 U	2.01 U	NA	1.02 U	1.04 U	1.04 U	NA
CL8(195)	6.76	3.81	3.11	42%	2.53	1.07 J	2.55	41%
CL9(206)	16.5	8.70	7.24	46%	5.83	2.19	5.68	45%
CL10(209)	12.8	7.77	6.82	35%	3.50	1.54	3.60	40%
<u>Surrogate Recoveries (%)</u>								
DBOFB	73	64	83	NA	89	70	91	NA
CL5(112)	64	56	71	NA	72	69	80	NA

TABLE B.7. (Contd)

Matrix	HU-B Rep. 1	HU-B Rep. 2	HU-B Rep. 3	RSD	HU-C Rep. 1	HU-C Rep. 2	HU-C Rep. 3	RSD
	Elutriate	Elutriate	Elutriate		Elutriate	Elutriate	Elutriate	
Sample Size (L)	0.98	0.96	0.96		0.96	0.98	1.00	
Batch	4	4	4	4	4	4	4	4
Units	ng/L	ng/L	ng/L		ng/L	ng/L	ng/L	
2,4-DDD	10.3	5.43	6.47	35%	6.49	5.83	5.59	8%
2,4-DDT	0.83 U	0.84 U	0.84 U	NA	0.84 U	0.82 U	0.81 U	NA
4,4-DDD	9.51	4.87	6.98	33%	7.70	6.14	7.89	13%
4,4-DDE	32.2	11.2	14.1	59% ^(e)	26.3	20.6	20.0	16%
4,4-DDT	1.03 U	1.04 U	1.04 U	NA	1.04 U	1.02 U	1.01 U	NA
Aldrin	0.76 U	0.77 U	0.77 U	NA	0.77 U	0.76 U	0.74 U	NA
<i>alpha</i> -Chlordane	3.67	1.31	0.91 J	76%	3.65	3.50	2.79	14%
Dieldrin	6.17	2.38	3.03	53%	5.78	5.50	5.62	2%
Endosulfan I/2,4'-DDE	0.87 U	0.88 U	0.88 U	NA	0.88 U	0.88 U	0.85 U	NA
Endosulfan II	11.5 U	11.7 U	11.7 U	NA	11.7 U	11.4 U	11.3 U	NA
Endosulfan sulfate	10.5	4.68 J	5.43 J	46%	13.5	10.0	10.0	18%
Heptachlor	0.67 U	0.68 U	0.68 U	NA	0.68 U	0.67 U	0.66 U	NA
Heptachlor epoxide	3.35	0.82 J	0.79 J	89%	2.95	3.11	2.72	7%
<i>trans</i> -Nonachlor	1.46	0.81 J	0.88 J	34%	1.39	1.45	1.55	6%
CL2(08)	3.58	4.44	3.85	11%	3.77	3.66	0.88 U	NA
CL3(18)	26.6	10.5	12.0	55% ^(e)	25.1	21.7	16.6	20%
CL3(28)	31.2	11.2	12.1	62% ^(e)	28.6	22.9	22.7	14%
CL4(44)	28.6	11.2	13.7	53% ^(e)	24.9	23.5	21.1	8%
CL4(49)	29.5	9.50	12.0	64% ^(e)	24.9	23.1	21.4	8%
CL4(52)	37.2	18.9	17.8	44% ^(e)	30.3	30.2	27.4	6%
CL4(66)	65.7	33.4	47.5	33% ^(e)	46.2	38.8	20.6	37% ^(e)
CL5(87)	10.2	3.64	5.01	55%	9.99	7.73	7.81	15%
CL5(101)	24.0	10.0	11.5	51% ^(e)	22.7	20.0	18.2	11%
CL5(105)	5.17	2.34	2.37	49%	5.82	4.17	4.82	17%
CL5(118)	1.04 U	7.03	9.63	NA	20.3	15.5	14.7	18%
CL6(128)	4.14	2.15	2.32	38%	3.82	2.92	3.32	13%
CL6(138)	25.2	9.86	12.90	51% ^(e)	27.1	21.7	20.8	15%
CL6(153)	21.3	7.50	10.38	56%	21.2	16.4	16.2	16%
CL7(170)	8.05	3.34	3.80	51%	7.62	5.93	5.75	16%
CL7(180)	16.0	5.53	7.56	57%	14.6	10.8	11.1	17%
CL7(183)	3.88	1.67	2.05	47%	3.94	3.14	3.74	12%
CL7(184)	1.09 U	1.11 U	1.11 U	NA	1.11 U	1.08 U	1.07 U	NA
CL7(187)	1.03 U	1.04 U	1.04 U	NA	1.04 U	1.02 U	1.01 U	NA
CL8(195)	7.19	2.09	2.80	69%	3.89	2.99	3.36	13%
CL9(206)	16.7	4.82	6.65	68%	7.23	4.95	5.10	22%
CL10(209)	9.43	3.60	4.08	57%	6.18	4.99	5.09	12%
<u>Surrogate Recoveries (%)</u>								
DBOFB	79	70	73	NA	74	77	57	NA
CL5(112)	64	63	68	NA	68	71	56	NA

TABLE B.7. (Contd)

Matrix	C-SB Rep. 1	C-SB Rep. 2	C-SB Rep. 3	RSD
Sample Size (L)	1.02	1.02	1.02	
Batch	4	4	4	4
Units	ng/L	ng/L	ng/L	
2,4-DDD	0.78 U	0.78 U	0.78 U	NA
2,4-DDT	0.80 U	0.80 U	0.79 U	NA
4,4-DDD	1.14 U	1.14 U	1.14 U	NA
4,4-DDE	0.97 U	0.97 U	0.97 U	NA
4,4-DDT	0.99 U	0.99 U	0.98 U	NA
Aldrin	0.73 U	0.73 U	0.73 U	NA
<i>alpha</i> -Chlordane	0.91 U	0.91 U	0.91 U	NA
Dieldrin	0.97 U	0.97 U	0.97 U	NA
Endosulfan I/2,4'-DDE	0.83 U	0.83 U	0.83 U	NA
Endosulfan II	11.0 U	11.0 U	11.0 U	NA
Endosulfan sulfate	8.07 U	8.07 U	8.03 U	NA
Heptachlor	2.41	0.65 U	0.64 U	NA
Heptachlor epoxide	0.84 U	0.84 U	0.84 U	NA
<i>trans</i> -Nonachlor	0.95 U	0.95 U	0.95 U	NA
CL2(08)	0.86 U	0.86 U	0.86 U	NA
CL3(18)	1.05 U	1.05 U	1.04 U	NA
CL3(28)	1.18 U	1.18 U	1.17 U	NA
CL4(44)	1.20 U	1.20 U	1.19 U	NA
CL4(49)	1.03 U	1.03 U	1.03 U	NA
CL4(52)	1.21 U	1.21 U	1.21 U	NA
CL4(66)	0.94 U	0.94 U	0.94 U	NA
CL5(87)	1.05 U	1.05 U	1.05 U	NA
CL5(101)	1.06 U	1.06 U	1.06 U	NA
CL5(105)	1.27 U	1.27 U	1.27 U	NA
CL5(118)	1.00 U	1.00 U	1.00 U	NA
CL6(128)	1.12 U	1.12 U	1.12 U	NA
CL6(138)	1.34 U	1.34 U	1.34 U	NA
CL6(153)	1.29 U	1.29 U	1.28 U	NA
CL7(170)	0.30 J	0.14 J	0.13 J	48%
CL7(180)	1.00 U	1.00 U	0.99 U	NA
CL7(183)	1.05 U	1.05 U	1.04 U	NA
CL7(184)	0.42 J	1.05 U	1.04 U	NA
CL7(187)	0.99 U	0.99 U	0.98 U	NA
CL8(195)	1.13 U	1.13 U	1.13 U	NA
CL9(206)	1.10 U	1.10 U	1.10 U	NA
CL10(209)	1.23 U	1.23 U	1.22 U	NA
<u>Surrogate Recoveries (%)</u>				
DBOFB	79	94	84	NA
CL5(112)	75	77	74	NA

(a) % RSD Percent relative standard deviation.

(b) U Undetected at or above concentration shown.

(c) NA Not applicable.

(d) J Concentration estimated; analyte detected below method detection limit (MDL) and above instrument detection limit (IDL).

(e) Outside quality control criteria ($\leq 30\%$ for replicate analysis) for analytes >10 times the achieved MDL.

TABLE B.8. Quality Control Data (Method Detection Limit Verification) for Pesticides and PCBs in Site Water and Elutriate

Sample Matrix	Sequim Bay 1 Control Water	Sequim Bay 1 Control Water	Sequim Bay 1 Control Water	Sequim Bay 1 Control Water	Sequim Bay 2 Control Water	Sequim Bay 2 Control Water	Sequim Bay 2 Control Water	Sequim Bay 2 Control Water	Standard Deviation	Detection Limit
Sample Size (L) Units	1.00 ng/L	1.00 ng/L	1.01 ng/L	0.91 ng/L	1.00 ng/L	1.00 ng/L	1.01 ng/L	0.96 ng/L	STD (n-1)	MDL ^(a) (ng/L)
2,4-DDD	NS ^(a)	NS	NS	NS	NS	NS	NS	NS	NA ^(b)	NA
2,4-DDT	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA
4,4-DDD	9.85	9.95	9.66	11.69	9.90	11.95	10.95	12.23	1.06	3.18
4,4-DDE	9.14	9.34	8.90	9.63	8.78	8.73	9.09	9.75	0.38	1.13
4,4-DDT	10.70	10.49	10.49	12.00	10.65	11.02	11.14	12.74	0.81	2.43
Aldrin	11.33	11.17	11.18	12.03	10.94	10.51	12.02	11.09	0.52	1.55
<i>alpha</i> -Chlordane	9.26	9.72	9.67	10.49	9.22	9.25	9.90	11.44	0.77	2.30
Dieldrin	9.31	9.21	8.87	9.82	8.65	8.61	8.95	9.84	0.44	1.33
Endosulfan I/2,4'-DDE	9.99	10.67	10.31	12.02	10.20	10.70	10.91	13.20	1.09	3.25
Endosulfan II	10.82	10.58	10.45	11.40	10.14	10.30	10.39	11.81	0.58	1.75
Endosulfan sulfate	10.07	9.79	9.74	10.68	9.56	9.73	9.81	10.96	0.50	1.51
Heptachlor	8.85	8.99	8.94	9.80	6.71	8.42	9.38	10.43	0.65	1.96
Heptachlor epoxide	9.30	9.74	9.61	10.27	9.26	9.52	10.01	11.66	0.78	2.34
<i>trans</i> -Nonachlor	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA
CL2(08)	6.04	7.39	6.94	6.50	6.63	6.21	6.63	7.03	0.44	1.32
CL3(18)	7.71	9.10	8.43	8.97	7.60	10.60	9.45	10.69	1.17	3.50
CL3(28)	8.32	8.88	8.78	9.75	8.51	7.95	8.83	9.78	0.64	1.92
CL4(44)	9.38	9.27	9.32	10.59	9.01	8.51	10.03	10.69	0.77	2.30
CL4(49)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA
CL4(52)	8.75	8.25	8.82	9.49	8.31	8.14	9.19	9.84	0.57	1.72
CL4(66)	8.87	9.63	9.58	10.32	9.11	9.67	9.87	11.11	0.70	2.09
CL5(87)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA
CL5(101)	9.26	9.82	10.00	10.08	9.12	8.72	9.67	11.19	0.75	2.25
CL5(105)	9.57	9.50	9.64	10.13	9.34	9.04	9.82	10.35	0.42	1.25
CL5(118)	9.68	10.08	9.64	10.75	9.65	9.64	9.98	10.85	0.50	1.50
CL6(128)	9.68	9.81	8.92	10.19	9.22	9.78	8.96	10.19	0.51	1.52
CL6(138)	9.78	9.78	9.80	11.14	9.52	9.44	10.01	11.57	0.78	2.35
CL6(153)	10.59	10.84	10.46	11.93	10.36	10.62	10.56	12.00	0.66	1.98
CL7(170)	9.15	9.24	9.31	10.07	9.30	9.05	9.50	9.86	0.36	1.07
CL7(180)	9.42	9.40	9.43	10.11	9.01	9.37	9.36	10.57	0.50	1.50
CL7(183)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA
CL7(184)	NS	NS	NS	NS	NS	NS	NS	NS	NA	NA
CL7(187)	9.43	9.34	9.24	10.22	9.03	9.21	9.36	9.69	0.37	1.11
CL8(195)	8.38	8.73	8.33	9.19	8.27	8.21	8.57	8.99	0.36	1.09
CL9(206)	7.86	7.65	7.46	8.03	7.26	7.26	7.55	8.30	0.37	1.11
CL10(209)	8.85	8.63	8.49	8.96	8.02	8.02	8.26	9.14	0.42	1.28
<u>Surrogate Recoveries (%)</u>										
DBQFB	85	89	84	82	81	71	88	71		
CL5(112)	86	88	85	81	82	79	84	84		

(a) MDL Method Detection Limit, calculated as Students-t (2.998 for 8 replicates) x standard deviation.

(b) NS Not spiked.

(c) NA Not applicable.

Appendix C

Water-Column Toxicity Test Data, Port Chester Project

TABLE C.1. Test Results for *M. beryllina* 96-Hour Water Column Toxicity Test

Sediment Treatment	Concentration Percent SPP	Replicate	Live(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	0	1	10	0	1.00		
COMP PC	0	2	10	0	1.00		
COMP PC	0	3	10	0	1.00		
COMP PC	0	4	10	0	1.00		
COMP PC	0	5	10	0	1.00	1.00	0.00
COMP PC	10	1	9	1	0.90		
COMP PC	10	2	10	0	1.00		
COMP PC	10	3	9	1	0.90		
COMP PC	10	4	10	0	1.00		
COMP PC	10	5	10	0	1.00	0.96	0.05
COMP PC	50	1	8	2	0.80		
COMP PC	50	2	9	1	0.90		
COMP PC	50	3	9	1	0.90		
COMP PC	50	4	8	2	0.80		
COMP PC	50	5	8	2	0.80	0.84	0.05
COMP PC	100	1	6	4	0.60		
COMP PC	100	2	2	8	0.20		
COMP PC	100	3	3	7	0.30		
COMP PC	100	4	0	10	0.00		
COMP PC	100	5	2	8	0.20	0.26	0.22

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE C.2. Water Quality Summary for *M. beryllina* 96-Hour Water Column Toxicity Test

Sediment Treatment	Concentration Percent SPP	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
		Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range		18.0	22.0	7.30	8.30	4.0	NA ^(a)	28.0	32.0
COMP PC	0	18.2	19.4	7.94	8.16	7.2	8.9	29.0	30.0
COMP PC	10	18.4	19.4	7.91	8.06	7.2	9.0	29.0	30.0
COMP PC	50	18.4	19.4	7.97	8.17	7.1	8.8	29.5	30.0
COMP PC	100	18.3	19.5	8.06	8.41 ^(b)	5.0	8.8	30.0	30.5

(a) NA Not applicable.

(b) Data point out of range.

TABLE C.3. Test Results for *M. beryllina* 96-Hour Copper Reference Toxicant Test

Copper Concentration ($\mu\text{g/L Cu}$)	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
0	1	10	0	1.00		
0	2	10	0	1.00		
0	3	10	0	1.00	1.00	0.00
16	1	10	0	1.00		
16	2	10	0	1.00		
16	3	10	0	1.00	1.00	0.00
64	1	10	0	1.00		
64	2	8	2	0.80		
64	3	8	2	0.80	0.87	0.12
160	1	1	9	0.10		
160	2	1	9	0.10		
160	3	2	8	0.20	0.13	0.06
400	1	0	10	0.00		
400	2	0	10	0.00		
400	3	0	10	0.00	0.00	0.00

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE C.4. Water Quality Summary for *M. beryllina* 96-Hour Copper Reference Toxicant Test

Copper Concentration ($\mu\text{g/L}$)	Temperature ($^{\circ}\text{C}$)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	4.0	NA ^(a)	28.0	32.0
0	18.5	19.3	7.90	8.09	7.1	7.9	31.0	32.0
16	18.6	19.2	7.98	8.09	7.3	8.0	31.0	32.0
64	18.5	19.2	7.91	8.07	7.4	8.1	31.0	32.0
160	18.6	19.3	7.95	8.08	7.4	8.1	31.0	32.0
400	18.7	19.4	7.85	8.03	7.3	7.6	31.0	31.5

(a) NA Not applicable.

TABLE C.5. Test Results for *M. bahia* 96-Hour Water Column Toxicity Test

Sediment Treatment	Concentration (Percent SPP)	Replicate	Live(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	0	1	9	1	0.90		
COMP PC	0	2	10	0	1.00		
COMP PC	0	3	10	0	1.00		
COMP PC	0	4	10	0	1.00		
COMP PC	0	5	10	0	1.00	0.98	0.04
COMP PC	10	1	9	1	0.90		
COMP PC	10	2	10	0	1.00		
COMP PC	10	3	9	1	0.90		
COMP PC	10	4	10	0	1.00		
COMP PC	10	5	10	0	1.00	0.96	0.05
COMP PC	50	1	8	2	0.80		
COMP PC	50	2	9	1	0.90		
COMP PC	50	3	10	0	1.00		
COMP PC	50	4	8	2	0.80		
COMP PC	50	5	9	1	0.90	0.88	0.08
COMP PC	100	1	8	2	0.80		
COMP PC	100	2	4	6	0.40		
COMP PC	100	3	6	4	0.60		
COMP PC	100	4	6	4	0.60		
COMP PC	100	5	5	5	0.50	0.58	0.15

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE C.6. Water Quality Summary for *M. bahia* 96-Hour Water Column Toxicity Test

Sediment Treatment	Concentration (Percent SPP)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
		Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range		18.0	22.0	7.30	8.30	4.0	NA ^(a)	28.0	32.0
COMP PC	0	18.4	19.4	7.80	8.08	6.5	8.9	29.0	30.0
COMP PC	10	18.6	19.4	7.90	8.11	6.8	8.9	29.0	30.0
COMP PC	50	18.6	19.3	7.99	8.33 ^(b)	6.6	8.2	29.5	30.5
COMP PC	100	18.6	19.4	8.15	8.39 ^(b)	5.2	8.2	30.0	31.0

(a) NA Not applicable.

(b) Data point out of range.

TABLE C.7. Test Results for *M. bahia* 96-Hour Copper Reference Toxicant Test

Copper Concentration (µg/L)	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
0	1	9	1	0.90		
0	2	10	0	1.00		
0	3	10	0	1.00	0.97	0.06
50	1	10	0	1.00		
50	2	9	1	0.90		
50	3	10	0	1.00	0.97	0.06
100	1	8	2	0.80		
100	2	9	1	0.90		
100	3	8	2	0.80	0.83	0.06
150	1	8	2	0.80		
150	2	7	3	0.70		
150	3	7	3	0.70	0.73	0.06
200	1	5	5	0.50		
200	2	5	5	0.50		
200	3	6	4	0.60	0.53	0.06

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE C.8. Water Quality Summary for *M. bahia* 96-Hour Copper Reference Toxicant Tests

Copper Concentration (µg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	4.0	NA ^(a)	28.0	32.0
0	19.3	19.5	7.58	8.08	5.8	8.1	30.5	32.0
50	19.2	19.6	7.81	8.05	7.1	8.0	30.5	32.0
100	19.2	19.5	7.81	8.09	7.0	7.9	30.5	32.0
150	19.2	19.6	7.83	8.08	7.1	7.9	30.5	32.0
200	19.2	19.5	7.85	8.06	7.3	8.0	30.5	32.0

(a) NA Not applicable.

TABLE C.9. Test Results for Larval *M. galloprovincialis* 48-Hour Water Column Toxicity Test

Sediment Treatment	SPP Concentration	Replicate	Mean			Mean			Mean			
			Stocking Density	Number Normal	Number Abnormal	Number Other	Proportion Normal ^(a)	Proportion Normal	Number Surviving	Proportion Surviving ^(a)	Proportion Surviving	Standard Deviation ^(b)
COMP PC	0%	1	261	280	1	6	1.00		287	1.00		
COMP PC	0%	2	261	267	0	4	1.00		271	1.00		
COMP PC	0%	3	261	244	0	3	0.93		247	0.95		
COMP PC	0%	4	261	279	1	9	1.00		289	1.00		
COMP PC	0%	5	261	286	0	9	1.00	0.99	295	1.00	0.99	0.02
COMP PC	10%	1	261	270	2	6	1.00		278	1.00		
COMP PC	10%	2	261	289	3	6	1.00		298	1.00		
COMP PC	10%	3	261	266	0	6	1.00		272	1.00		
COMP PC	10%	4	261	239	0	4	0.92		243	0.93		
COMP PC	10%	5	261	239	0	9	0.92	0.97	248	0.95	0.98	0.03
COMP PC	50%	1	261	204	6	19	0.78		229	0.88		
COMP PC	50%	2	261	214	15	11	0.82		240	0.92		
COMP PC	50%	3	261	217	38	19	0.83		274	1.00		
COMP PC	50%	4	261	190	20	23	0.73		233	0.89		
COMP PC	50%	5	261	138	10	12	0.53	0.74	160	0.61	0.86	0.15
COMP PC	100%	1	261	0	0	213	0.00		213	0.82		
COMP PC	100%	2	261	0	0	209	0.00		209	0.80		
COMP PC	100%	3	261	0	0	231	0.00		231	0.89		
COMP PC	100%	4	261	0	0	236	0.00		236	0.90		
COMP PC	100%	5	261	0	0	208	0.00	0.00	208	0.80	0.84	0.05

(a) When number normal or number surviving exceeded the stocking density, a proportion normal and/or proportion survival of 1.00 was used for mean calculations and statistical analysis.

(b) Standard deviation is based on proportion surviving.

TABLE C.10. Water Quality Summary for *M. galloprovincialis* 48-Hour Water Column Toxicity Test

Sediment Treatment	Percent Concentration	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
		Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range		14.0	18.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
COMP PC	0	16.0	16.6	7.97	8.10	7.5	8.1	30.5	32.0
COMP PC	10	16.0	16.4	7.98	8.18	7.5	8.1	30.5	32.0
COMP PC	50	15.9	16.4	7.99	8.21	7.5	8.0	30.0	31.5
COMP PC	100	15.9	16.4	8.07	8.25	6.7	7.9	29.5	31.5

(a) NA Not applicable.

TABLE C.11. Test Results for Larval *M. galloprovincialis* 48-Hour Copper Reference Toxicant Tests

Copper Concentration (µg/L)	Replicate	Mean Stocking Density	Number			Proportion		Mean Number Surviving	Proportion		Standard Deviation ^(b)
			Normal	Abnormal	Other	Normal ^(a)	Normal		Surviving ^(a)	Surviving	
0.00	1	285	217	0	2	0.76		219	0.77		
0.00	2	285	252	1	15	0.88		268	0.94		
0.00	3	285	232	1	13	0.81		246	0.86		
0.00	4	285	194	0	10	0.68		204	0.72		
0.00	5	285	249	1	14	0.87	0.80	264	0.93	0.84	
1.00	1	285	223	0	19	0.78		242	0.85		
1.00	2	285	248	0	10	0.87		258	0.91		
1.00	3	285	265	2	9	0.93	0.86	276	0.97	0.91	
4.00	1	285	0	0	7	0.00		7	0.02		
4.00	2	285	268	1	10	0.94		279	0.98		
4.00	3	285	264	1	14	0.93	0.62	279	0.98	0.66	
16.00	1	285	16	38	160	0.06		214	0.75		
16.00	2	285	0	13	309	0.00		322	1.00		
16.00	3	285	0	0	242	0.00	0.02	242	0.85	0.87	
64.00	1	285	2	0	1	0.01		3	0.01		
64.00	2	285	254	0	11	0.89		265	0.93		
64.00	3	285	4	0	4	0.01	0.30	8	0.03	0.32	

(a) When number normal or number surviving exceeded the stocking density, a proportion normal and/or proportion survival of 1.00 was used for mean calculations and statistical analysis.

(b) Standard deviation is based on proportion surviving.

TABLE C.12. Water Quality Summary for *M. galloprovincialis* 48-Hour Copper Reference Toxicant Tests

Copper Concentration (µg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	14.0	18.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
0.00	15.9	16.5	8.03	8.14	7.9	8.2	30.5	31.5
1.00	16.0	16.4	8.00	8.15	7.5	8.2	30.5	31.0
4.00	16.0	16.3	7.93	8.06	7.6	8.1	30.5	31.5
16.0	15.8	16.4	8.03	8.15	7.5	8.2	30.5	32.0
64.0	15.9	16.4	8.01	8.18	7.4	8.2	30.5	31.5

(a) NA Not applicable.

Appendix D

Benthic Acute Toxicity Test Data, Port Chester Project

TABLE D.1. Test Results for *A. abdita* 10-Day Static Renewal, Benthic Acute Toxicity Test

Sediment Treatment	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	1	0	20	0.00		
COMP PC	2	0	20	0.00		
COMP PC	3	0	20	0.00		
COMP PC	4	0	20	0.00		
COMP PC	5	0	20	0.00	0.00	0.00
R-MUD	1	17	3	0.85		
R-MUD	2	19	1	0.95		
R-MUD	3	18	2	0.90		
R-MUD	4	19	1	0.95		
R-MUD	5	20	0	1.00	0.93	0.06
R-CLIS	1	19	1	0.95		
R-CLIS	2	20	0	1.00		
R-CLIS	3	19	1	0.95		
R-CLIS	4	20	0	1.00		
R-CLIS	5	19	1	0.95	0.97	0.03
C-AM	1	20	0	1.00		
C-AM	2	20	0	1.00		
C-AM	3	19	1	0.95		
C-AM	4	18	2	0.90		
C-AM	5	20	0	1.00	0.97	0.04

(a) Survival based on initial exposure of 20 organisms per replicate.

TABLE D.2. Water Quality Summary for *A. abdita* 10-Day Static Renewal, Benthic Acute Toxicity Test

Sediment Treatment	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)		Total Ammonia ^(a) (mg/L)	
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	5.0	NA ^(b)	28.0	32.0	NA	30.0
COMP PC	18.2	19.4	8.00	8.80 ^(c)	7.1	8.3	30.5	32.0	<1.00	<1.00
R-MUD	17.9 ^(c)	19.3	7.93	8.14	7.3	8.3	30.5	32.0	<1.00	<1.00
R-CLIS	17.5 ^(c)	19.3	7.95	8.30	6.9	8.4	30.0	32.0	<1.00	<1.00
C-AM	17.9 ^(c)	19.3	7.80	8.16	6.8	8.2	30.0	31.5	<1.00	1.30

(a) Total ammonia measured in overlying water.

(b) NA Not applicable.

(c) Data point out of range.

TABLE D.3. Water Quality Measurements of Porewater for *A. abdita* 10-Day, Static Renewal, Benthic Acute Toxicity Test

Sediment Treatment	Ammonia (mg/L)	Temperature (°C)	pH	Dissolved Oxygen (mg/L)	Salinity (o/oo)
Day 0					
COMP PC	10.7	19.1	8.38	8.1	31.5
R-MUD	0.737	19.2	8.07	7.9	31.5
R-CLIS	2.57	18.2	7.99	8.0	31.0
C-AM	7.12	19.3	8.03	8.1	31.0
Day 10					
COMP PC	4.02	18.8	8.22	8.3	30.5
R-MUD	ND ^(a)	18.9	8.01	8.2	31.0
R-CLIS	1.65	18.7	8.23	8.4	31.0
C-AM	4.61	18.4	8.12	8.1	30.0

(a) ND No data.

TABLE D.4. Test Results for *A. abdita* 96-Hour Cadmium Reference Toxicant Test

Cadmium Concentration (mg/L)	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
0.00	1	20	0	1.00		
0.00	2	19	1	0.95		
0.00	3	20	0	1.00	0.98	0.03
0.25	1	13	7	0.65		
0.25	2	13	7	0.65		
0.25	3	15	5	0.75	0.68	0.06
0.50	1	12	8	0.60		
0.50	2	15	5	0.75		
0.50	3	13	7	0.65	0.67	0.08
1.00	1	4	16	0.20		
1.00	2	5	15	0.25		
1.00	3	5	15	0.25	0.23	0.03
2.00	1	0	20	0.00		
2.00	2	0	20	0.00		
2.00	3	0	20	0.00	0.00	0.00

(a) Survival based on initial exposure of 20 organisms per replicate.

TABLE D.5. Water Quality Summary for 96-Hour *A. abdita* Cadmium Reference Toxicant Test

Cadmium Concentration (mg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
0.00	19.3	19.5	7.97	8.14	7.3	8.0	30.5	31.0
0.25	19.3	19.5	7.92	8.10	7.5	7.9	30.5	31.5
0.50	19.3	19.6	7.91	8.10	7.5	7.8	30.5	31.0
1.00	19.2	19.5	7.90	8.09	7.6	7.9	30.5	31.5
2.00	19.3	19.6	7.85	8.03	7.6	7.9	30.5	31.5

(a) NA Not applicable.

TABLE D.6. Results of *R. abronius* 10-Day, Static Renewal, Benthic Acute Toxicity Test

Sediment Treatment	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	1	14	6	0.70		
COMP PC	2	18	2	0.90		
COMP PC	3	16	4	0.80		
COMP PC	4	15	5	0.75		
COMP PC	5	15	5	0.75	0.78	0.08
R-MUD	1	20	0	1.00		
R-MUD	2	20	0	1.00		
R-MUD	3	20	0	1.00		
R-MUD	4	20	0	1.00		
R-MUD	5	18	2	0.90	0.98	0.04
R-CLIS	1	19	1	0.95		
R-CLIS	2	19	1	0.95		
R-CLIS	3	15	5	0.75		
R-CLIS	4	19	1	0.95		
R-CLIS	5	19	1	0.95	0.91	0.09
C-WB	1	19	1	0.95		
C-WB	2	20	0	1.00		
C-WB	3	21	0	1.00		
C-WB	4	18	2	0.90		
C-WB	5	20	0	1.00	0.97	0.04

(a) Survival based on initial exposure of 20 organisms per replicate.

TABLE D.7. Water Quality Summary for *R. abronius* 10-Day Solid-Phase, Static Renewal, Benthic Acute Toxicity Test

Sediment Treatment	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)		Total Ammonia ^(a) (mg/L)	
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	12.0	16.0	7.30	8.30	5.0	NA ^(b)	28.0	32.0	NA	30.0
COMP PC	14.0	15.0	8.05	8.48 ^(c)	7.4	8.8	30.5	32.0	0.095	1.09
R-MUD	13.8	15.0	7.10	8.12	7.4	8.8	30.5	32.0	0.026	<1.00
R-CLIS	14.0	15.3	7.91	8.13	7.5	8.7	30.0	32.0	0.026	1.72
C-WB	13.8	15.1	7.91	8.40 ^(c)	7.6	8.8	31.0	32.0	0.034	0.219

(a) Total ammonia measured in the overlying water.

(b) NA Not applicable.

(c) Data point out of range.

TABLE D.8. Water Quality Measurements of Porewater for *R. abronius* 10-Day, Static Renewal, Benthic Acute Toxicity Test

Sediment Treatment	Ammonia (mg/L)	Temperature (°C)	pH	Dissolved Oxygen (mg/L)	Salinity (o/oo)
Day 0					
COMP PC	11.0	15.0	8.21	7.9	32.0
R-MUD	0.685	15.0	7.99	8.0	32.0
R-CLIS	2.52	14.5	7.97	7.5	31.5
C-WB	2.74	14.8	7.93	7.7	31.5
Day 10					
COMP PC	2.9	14.4	8.28	8.7	31.0
R-MUD	ND ^(a)	14.5	8.10	8.8	31.0
R-CLIS	1.3	14.4	8.12	8.7	30.5
C-WB	ND	14.3	8.09	8.8	31.0

(a) ND No data.

TABLE D.9. Test Results for *R. abronius* 96-Hour Cadmium Reference Toxicant Test

Cadmium Concentration (mg/L)	Rep	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
0.00	1	18	2	0.90		
0.00	2	20	0	1.00		
0.00	3	20	0	1.00	0.97	0.06
0.38	1	15	5	0.75		
0.38	2	5	5	0.25		
0.38	3	20	0	1.00	0.67	0.38
0.75	1	15	5	0.75		
0.75	2	17	3	0.85		
0.75	3	12	8	0.60	0.73	0.13
1.50	1	8	12	0.40		
1.50	2	2	18	0.10		
1.50	3	9	11	0.45	0.32	0.19
3.00	1	1	19	0.05		
3.00	2	4	16	0.20		
3.00	3	1	19	0.05	0.10	0.09

TABLE D.10. Water Quality Summary for *R. abronius* 96-Hour Cadmium Reference Toxicant Test

Cadmium Concentration (mg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	12.0	16.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
0.00	14.9	15.6	7.91	8.10	7.9	8.3	30.5	32.0
0.38	14.9	15.2	7.90	8.07	8.0	8.4	30.5	32.0
0.75	14.8	15.3	7.90	8.06	8.0	8.3	30.5	31.5
1.50	14.9	15.2	7.87	8.02	8.0	8.3	30.5	32.0
3.00	14.9	15.2	7.66	7.92	7.9	8.2	30.5	32.0

(a) NA Not applicable.

TABLE D.16. Test Results for 10-Day, Static, Benthic Acute Toxicity Test with *M. bahia*

Sediment Treatment	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	1	15	5	0.75		
COMP PC	2	20	0	1.00		
COMP PC	3	15	5	0.75		
COMP PC	4	15	5	0.75		
COMP PC	5	12	8	0.60	0.77	0.14
R-MUD	1	17	3	0.85		
R-MUD	2	17	3	0.85		
R-MUD	3	13	7	0.65		
R-MUD	4	13	7	0.65		
R-MUD	5	16	4	0.80	0.76	0.10
R-CLIS	1	16	4	0.80		
R-CLIS	2	12	8	0.60		
R-CLIS	3	19	1	0.95		
R-CLIS	4	13	7	0.65		
R-CLIS	5	14	6	0.70	0.74	0.14
C-SB ^(b)	1	19	1	0.95		
C-SB	2	16	4	0.80		
C-SB	3	19	1	0.95		
C-SB	4	20	0	1.00		
C-SB	5	19	1	0.95	0.93	0.08

(a) Survival based on initial exposure of 20 organisms per replicate.

(b) Control exposures were run approximately three weeks after the Port Chester sediments were run.

TABLE D.12. Water Quality Summary for 10-Day, Static, Benthic Acute Toxicity Test with *M. bahia*

Sediment Treatment	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)		Ammonia (mg/L)	
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	4.0	NA ^(a)	28.0	32.0	NA	20.0
COMP PC	18.7	19.6	7.87	8.53 ^(b)	6.2	7.8	30.5	32.0	0.201	1.67
R-MUD	18.5	19.5	7.57	7.99	6.0	7.8	30.5	32.0	0.070	3.6
R-CLIS	18.6	19.6	7.64	8.09	5.3	7.7	30.0	32.0	0.069	1.95
C-SB ^(c)	18.6	19.5	7.73	8.24	5.9	7.4	30.0	32.0	3.36	82.0 ^(b)

(a) NA Not applicable.

(b) Data point out of range.

(c) Control exposures were run approximately three weeks after the Port Chester sediments were run.

TABLE D.13. Test Results for 96-Hour, Benthic Acute Toxicity, Copper Reference Toxicant Test with *M. bahia*

Copper Concentration (µg/L)	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
0	1	9	1	0.90		
0	2	10	0	1.00		
0	3	10	0	1.00	0.97	0.06
50	1	10	0	1.00		
50	2	9	1	0.90		
50	3	10	0	1.00	0.97	0.06
100	1	8	2	0.80		
100	2	7	3	0.70		
100	3	8	2	0.80	0.77	0.06
150	1	6	4	0.60		
150	2	5	5	0.50		
150	3	6	4	0.60	0.57	0.06
200	1	1	9	0.10		
200	2	2	8	0.20		
200	3	2	8	0.20	0.17	0.06

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE D.14. Water Quality Summary for 96-Hour, Benthic Acute Toxicity, Copper Reference Toxicant Test with *M. bahia*

Copper Concentration (µg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	4.0	NA ^(a)	28.0	32.0
0.00	18.6	19.3	7.91	8.08	6.5	8.9	30.5	32.0
50.0	18.7	19.3	7.88	8.13	6.6	9.1	30.0	31.5
100	18.7	19.3	7.87	8.08	6.4	9.0	30.5	32.0
150	18.7	19.4	7.86	8.16	6.8	8.9	30.5	32.0
200	18.7	19.4	7.84	8.14	6.7	8.9	30.0	31.5

(a) NA Not applicable.

Appendix E

Bioaccumulation Test Data, Port Chester Project

TABLE E.1. Test Results for 28-Day Bioaccumulation Test with *M. nasuta*

Sediment Treatment	Replicate	Number Live ^(a)	Number Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	1	23	2	0.92		
COMP PC	2	24	1	0.96		
COMP PC	3	21	4	0.84		
COMP PC	4	21	4	0.84		
COMP PC	5	24	1	0.96	0.90	0.06
R-MUD	1	22	3	0.88		
R-MUD	2	20	5	0.80		
R-MUD	3	23	2	0.92		
R-MUD	4	21	4	0.84		
R-MUD	5	24	1	0.96	0.88	0.06
R-CLIS	1	23	2	0.92		
R-CLIS	2	25	0	1.00		
R-CLIS	3	22	3	0.88		
R-CLIS	4	25	0	1.00		
R-CLIS	5	25	0	1.00	0.96	0.06
C-SB	1	25	0	1.00		
C-SB	2	24	1	0.96		
C-SB	3	24	1	0.96		
C-SB	4	24	1	0.96		
C-SB	5	25	0	1.00	0.98	0.02

(a) Survival based on initial exposure of 25 organisms per replicate.

TABLE E.2. Water Quality Summary for 28-day Bioaccumulation Test with *M. nasuta*

Sediment Treatment	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	12.0	16.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
COMP PC	14.3	16.5 ^(b)	7.73	8.00	6.9	8.1	30.0	31.5
R-MUD	14.4	16.4 ^(b)	7.68	8.03	7.4	8.3	30.0	31.0
R-CLIS	14.4	15.9	7.67	8.05	7.2	8.8	30.0	31.0
C-SB	14.3	16.5 ^(b)	7.71	8.01	7.1	8.2	30.5	31.0

(a) NA Not applicable.

(b) Data point out of range.

TABLE E.3. Test Results for 96-Hour Copper Reference Toxicant Test with *M. nasuta*

Copper Concentration (mg/L)	Live ^(a)	Dead or Missing	Proportion Surviving
0.00	10	0	1.00
0.25	10	0	1.00
0.50	10	0	1.00
0.75	8	2	0.80
1.00	10	0	1.00
1.50	8	2	0.80
2.50	4	6	0.40

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE E.4. Water Quality Summary for 96-Hour Copper Reference Toxicant Test with *M. nasuta*

Copper Concentration (mg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	12.0	16.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
0.00	15.1	15.8	7.78	7.96	7.0	8.0	30.5	31.5
0.25	15.0	15.5	7.64	7.94	6.9	8.1	30.5	31.5
0.50	15.0	15.6	7.65	7.94	6.9	8.0	30.5	31.5
0.75	15.0	15.5	7.48	7.93	5.4	8.0	30.5	31.5
1.00	15.1	15.5	7.53	7.88	6.2	8.1	30.5	31.5
1.50	15.0	15.6	7.44	7.88	5.3	8.1	30.5	31.5
2.50	15.0	15.6	7.27 ^(b)	7.86	3.2 ^(b)	8.1	30.5	31.5

(a) NA Not applicable.

(b) Data point out of range.

TABLE E.5. Test Results for 28-Day Bioaccumulation Test with *N. virens*

Sediment Treatment	Replicate	Live ^(a)	Dead or Missing	Proportion Surviving	Mean Proportion Surviving	Standard Deviation
COMP PC	1	16	4	0.80		
COMP PC	2	18	2	0.90		
COMP PC	3	17	3	0.85		
COMP PC	4	10	10	0.50		
COMP PC	5	15	5	0.75	0.76	0.16
R-MUD	1	16	4	0.80		
R-MUD	2	15	5	0.75		
R-MUD	3	18	2	0.90		
R-MUD	4	15	5	0.75		
R-MUD	5	15	5	0.75	0.79	0.07
R-CLIS	1	19	1	0.95		
R-CLIS	2	14	6	0.70		
R-CLIS	3	15	5	0.75		
R-CLIS	4	18	2	0.90		
R-CLIS	5	16	4	0.80	0.82	0.10
C-NR	1	19	1	0.95		
C-NR	2	20	0	1.00		
C-NR	3	16	4	0.80		
C-NR	4	19	1	0.95		
C-NR	5	15	5	0.75	0.89	0.11

(a) Survival based on initial exposure of 20 organisms per replicate.

TABLE E.6. Water Quality Summary for 28-Day Bioaccumulation Test with *N. virens*

Sediment Treatment	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
COMP PC	18.4	20.0	7.57	8.05	5.4	8.2	30.0	31.5
R-MUD	18.0	19.9	7.73	8.88 ^(b)	6.5	8.3	30.5	32.0
R-CLIS	18.1	19.8	7.72	8.01	6.5	8.3	30.0	31.5
C-NR	18.0	19.9	7.70	8.01	6.3	8.2	30.0	31.5

(a) NA Not applicable.

(b) Data point out of range.

TABLE E.7. Test Results for 96-Hour Copper Reference Toxicant Test with *N. virens*

Copper Concentration (mg/L)	Live ^(a)	Dead or Missing	Proportion Surviving
0.00	10	0	1.00
0.05	10	0	1.00
0.075	10	0	1.00
0.15	4	6	0.40
0.20	0	10	0.00
0.25	0	10	0.00
0.30	0	10	0.00

(a) Survival based on initial exposure of 10 organisms per replicate.

TABLE E.8. Water Quality Summary for 96-Hour Copper Reference Toxicant Test with *N. virens*

Copper Concentration (mg/L)	Temperature (°C)		pH		Dissolved Oxygen (mg/L)		Salinity (o/oo)	
	Min	Max	Min	Max	Min	Max	Min	Max
Acceptable Range	18.0	22.0	7.30	8.30	5.0	NA ^(a)	28.0	32.0
0.00	18.6	19.2	7.52	7.94	5.7	7.4	30.5	31.5
0.05	18.6	19.3	7.60	7.95	6.3	7.4	30.5	31.5
0.075	18.6	19.4	7.61	7.91	5.2	7.6	30.5	31.5
0.15	18.6	19.4	7.39	7.93	4.5 ^(b)	7.4	30.5	31.5
0.20	18.7	19.4	7.00 ^(b)	7.82	0.6 ^(b)	7.5	30.5	31.5
0.25	18.6	19.4	7.14 ^(b)	7.86	2.0 ^(b)	7.5	30.5	31.5
0.30	18.6	19.4	7.21 ^(b)	7.90	3.0 ^(b)	7.6	30.5	31.5

(a) NA Not applicable.

(b) Data point out of range.

Appendix F

***Macoma nasuta* Tissue Chemical Analyses and
Quality Assurance/Quality Control Data,
Port Chester Project**

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Metals
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Worm and Clam Tissue

QA/QC DATA QUALITY OBJECTIVES

	<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (µg/g dry wt)</u>
Arsenic	ICP/MS	75-125%	≤20%	≤20%	1.0
Cadmium	ICP/MS	75-125%	≤20%	≤20%	0.1
Chromium	ICP/MS	75-125%	≤20%	≤20%	0.2
Copper	ICP/MS	75-125%	≤20%	≤20%	1.0
Lead	ICP/MS	75-125%	≤20%	≤20%	0.1
Mercury	CVAA	75-125%	≤20%	≤20%	0.02
Nickel	ICP/MS	75-125%	≤20%	≤20%	0.1
Silver	ICP/MS	75-125%	≤20%	≤20%	0.1
Zinc	ICP/MS	75-125%	≤20%	≤20%	1.0

METHOD

A total of nine (9) metals was analyzed for the New York Federal Projects-2 Program: silver (Ag), arsenic (As), cadmium (Cd), chromium (Cr), copper (Cu), mercury (Hg), nickel (Ni), lead (Pb) and zinc (Zn). Hg was analyzed using cold-vapor atomic absorption spectroscopy (CVAA) according to the method of Bloom and Crecelius (1983). The remaining metals were analyzed by inductively coupled plasma mass spectrometry (ICP/MS) following a procedure based on EPA Method 200.8 (EPA 1991).

To prepare tissue for analysis, samples were freeze-dried and blended in a Spex mixer-mill. Approximately 5 g of mixed sample was ground in a ceramic ball mill. For ICP/MS and CVAA analyses, 0.2- to 0.5-g aliquots of dried homogenous sample were digested using a mixture of nitric acid and hydrogen peroxide following EPA Method 200.3 (EPA 1991).

HOLDING TIMES

A total of 68 worm and 68 clam samples was received on 6/15/94 in good condition. Samples were logged into Battelle's log-in system, frozen to -80°C and subsequently freeze dried within approximately 7 days of sample receipt. Samples were analyzed within 180 days of collection. Worms and clams were digested in two separate batches. The following table summarizes the analysis dates:

<u>Task</u>	<u>Clams</u>	<u>Worms</u>
Sample Digestion	8/9/94	9/9/94
ICP-MS	9/15/94	10/6/94
CVAA-Hg	8/17-8/24/94	8/17-8/24/94

QA/QC SUMMARY METALS (continued)

- DETECTION LIMITS** Four aliquots of a background clam tissue were analyzed as four separate replicates. The standard deviation of these results were multiplied by 4.541 to determine a method detection limits (MDL). Target detection limits were exceeded for all metals except Ag, Cd and Hg.
- METHOD BLANKS** One procedural blank was analyzed per 20 samples. No metals were detected in the blanks above the MDLs.
- MATRIX SPIKES** One sample was spiked with all metals at a frequency of 1 per 20 samples. All recoveries were within the QC limits of 75% -125% with the exception of Ag in one spiked worm sample and Zn in three of the four spiked worm samples. Zn was spiked at a level near the level found in the native samples and, in one case, Zn was spiked at a level below that detected in the native sample and no recovery was calculated.
- REPLICATES** One sample was analyzed in triplicate at a frequency of 1 per 20 samples. Precision for triplicate analyses is reported by calculating the relative standard deviation (RSD) between the replicate results. Only the RSDs for Zn in one of the four replicated worm analyses exceeded the QC limits of $\pm 20\%$. RSDs for the rest of the metals were within the QC limits.
- SRMs** Standard Reference Material (SRM), 1566a (Oyster tissue from the National Institute of Standards and Technology, NIST), was analyzed for all metals. Results for all metals were within $\pm 20\%$ of mean certified value with the exception of Cr and Ni. Cr values were below the lower QC limit in two of the five SRMs analyzed with the clams and for three of the four SRMs analyzed with the worms. The SRM certified value for Cr ($1.43 \mu\text{g/g}$) is close to the detection limit ($1.46 \mu\text{g/g}$). Ni was also recovered below or above the control limits in some samples.

REFERENCES

Bloom, N. S., and E.A. Crecelius. 1983. "Determination of Mercury in Seawater at Sub-Nanogram per Liter Levels." *Mar. Chem.* 14:49-59.

EPA (U.S. Environmental Protection Agency). 1991 *Methods for the Determination of Metals in Environmental Samples*. EPA-600/4-91-010. Environmental Services Division, Monitoring Management Branch, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Chlorinated Pesticides/PCB Congeners
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Worm and Clam Tissue

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>Surrogate Recovery</u>	<u>Spike Recovery</u>	<u>Relative Precision</u>	<u>Detection Limit</u>
GC/ECD	30-150%	50-120%	≤30%	0.4 ng/g wet wt.

SAMPLE CUSTODY A total of 68 worm and 68 clam samples was received on 6/15/94 in good condition. Samples were logged into Battelle's log-in system and stored frozen until extraction.

METHOD Tissues were homogenized wet using a stainless steel blade. An aliquot of tissue sample was extracted with methylene chloride using the roller technique under ambient conditions following a procedure which is based on methods used by the National Oceanic and Atmospheric Administration for its Status and Trends Program (Krahn et al. 1988). Samples were then cleaned using silica/alumina (5% deactivated) chromatography followed by HPLC cleanup (Krahn et al. 1988). Extracts were analyzed for 15 chlorinated pesticides and 22 PCB congeners using gas chromatography/electron capture detection (GC/ECD) following a procedure based on EPA Method 8080 (EPA 1986). The column used was a J&W DB-17 and the confirmatory column was a DB-1701, both capillary columns (30m x 0.25mm I.D.). All detections were quantitatively confirmed on the second column.

HOLDING TIMES Samples were extracted in seven batches. All extracts were analyzed by GC/ECD. The following summarizes the extraction and analysis dates:

<u>Batch</u>	<u>Species</u>	<u>Extraction</u>	<u>Analysis</u>
1	<i>M. nasuta</i>	7/28/94	9/9-9/12/94
2	<i>M. nasuta</i>	8/3/94	9/13-9/15/94
3	<i>M. nasuta</i>	8/17/94	9/23-9/25/94
4	<i>N. virens</i>	8/19/95	9/26-9/30/94
5	<i>N. virens</i>	8/26/94	9/8-9/11/94
6	<i>N. virens</i>	9/6/94	9/17-9/19/94
7	<i>M. nasuta/N. virens</i>	9/26/94	9/15-9/17-94
8	<i>M. nasuta</i> MDL study	10/10/94	10/25/94

DETECTION LIMITS Target detection limits of 0.4 ng/g wet weight were met for all pesticides and PCB congeners, with the exception of dieldrin, PCB 8 and PCB 18, and for the samples that were analyzed in triplicate. These elevated detection limits for the replicates were due to the limited amount of tissue available resulting in smaller aliquots used for extraction. Method detection limits (MDLs) reported were determined by multiplying the

QA/QC SUMMARY/PCBs and PESTICIDES (continued)

standard deviation of seven spiked replicates of clam tissue by the Student's t value (99 percentile). Actual pesticide MDLs ranged from approximately 0.1 to 1.1 ng/g wet weight and PCB congener MDLs ranged from approximately 0.1 to 0.9 ng/g wet weight, depending on the compound and the sample weight extracted. MDLs were reported corrected for individual sample wet weight extracted.

Method detection limit verification was performed by analyzing four replicates of a spiked clam sample and multiplying the standard deviation of the results by 3.5. All detection limits calculated in this way were below the target detection limit of 0.4 ng/g wet weight with the exception of 4,4'-DDD which had a DL of 0.467 ng/g.

METHOD BLANKS

One method blank was extracted with each extraction batch. No pesticides or PCBs were detected in any of the method blanks.

SURROGATES

Two compounds, PCB congeners 103 and 198, were added to all samples prior to extraction to assess the efficiency of the analysis. Sample surrogate recoveries were all within the QC guidelines of 30% - 150%, with the exception of one sample in Batch 3 and two samples in Batch 4. All of these incidents involved a high recovery of PCB 198. This was most likely due to matrix interferences with the internal Standard octachloronaphthalene (OCN) which is used to quantify the recovery of surrogate PCB 198. Since no sample data are corrected for the OCN, sample results should not be affected. One sample had low surrogate recoveries for both PCB 103 and 198. This sample was re-extracted once due to surrogate recoveries. Since the recoveries in the reextraction also exceeded control limits, the problem was determined to be matrix interferences and no additional extractions were performed. Sample results were quantified using the surrogate internal standard method.

MATRIX SPIKES

Ten out of the 15 pesticides and 5 of the 22 PCB congeners analyzed were spiked into one sample per extraction batch. Matrix spike recoveries were within the control limit range of 50-120% for all Pesticides and PCBs in Batches 1, 2, 3, 6 and 7 with the exception of PCB 138 in Batch six and three pesticides and 2 PCBs in Batch seven. In all cases, the recoveries were high and are most likely due to matrix interferences. Recoveries for the majority of pesticides and PCBs in Batches four and five exceeded control limits due to high native levels compared with the levels spiked. In most cases, the spiked concentrations were 2 to 10 times lower than the concentrations detected in the samples.

REPLICATES

One sample from each extraction batch was analyzed in triplicate. Precision was measured by calculating the relative standard deviation (RSD) between the replicate results. RSDs for all detectable values were below the target precision goal of $\leq 30\%$ in Batches 1, 2, 3, 4 and 7. The RSD for Endosulfan Sulfate in Batch 5 was high due to comparison of very low concentrations, less than 1 ng/g in the replicates. RSDs for two pesticides and for two PCB congeners in Batch 6 were high due to matrix interferences associated with the first replicate sample.

QA/QC SUMMARY/PCBs and PESTICIDES (continued)

SRMs Not applicable.

MISCELLANEOUS All pesticide and PCB congener results are confirmed using a second dissimilar column. RPDs between the primary and confirmation values must be less than 75% to be considered a confirmed value.

REFERENCES

Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.

EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Polynuclear Aromatic Hydrocarbons (PAH) and 1,4-Dichlorobenzene
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Clam and Worm Tissue

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>MS Recovery</u>	<u>Surrogate Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (wet wt)</u>
GC/MS/SIM	50-120%	30-150%	≤30%	≤30%	4 ng/g

SAMPLE CUSTODY A total of 68 worm and 68 clam samples was received on 6/15/94 in good condition. Samples were logged into Battelle's log-in system and stored frozen until extraction.

METHOD Tissue samples were extracted with methylene chloride using a roller under ambient conditions following a procedure which is based on methods used by the National Oceanic and Atmospheric Administration for its Status and Trends Program (Krahn et al. 1988). Samples were then cleaned using silica/alumina (5% deactivated) chromatography followed by HPLC cleanup.

Extracts were quantified using gas chromatography/mass spectrometry (GC/MS) in the selected ion mode (SIM) following a procedure based on EPA Method 8270 (EPA 1986).

HOLDING TIMES Samples were extracted in seven batches. All extracts were analyzed by GC/MS/SIM. The following summarizes the extraction and analysis dates:

<u>Batch</u>	<u>Species</u>	<u>Extraction</u>	<u>Analysis</u>
1	<i>M. nasuta</i>	7/28/94	9/9-9/12/94
2	<i>M. nasuta</i>	8/3/94	9/13-9/15/94
3	<i>M. nasuta</i>	8/17/94	9/23-9/25/94
4	<i>N. virens</i>	8/19/95	9/26-9/30/94
5	<i>N. virens</i>	8/26/94	9/8-9/11/94
6	<i>N. virens</i>	9/6/94	9/17-9/19/94
7	<i>M. nasuta/N. virens</i>	9/26/94	9/15-9/17-94
8	<i>M. nasuta</i> MDL study	10/10/94	10/25/94

DETECTION LIMITS Target detection limits of 4 ng/g wet weight were met for all PAH compounds except for fluoranthene and pyrene, which had method detection limits (MDL) between 4 and 6 ng/g wet weight. MDLs were determined by multiplying the standard deviation of seven spiked replicates of a background clam sample by the Student's t value (99 percentile). These MDLs were based on a wet weight of 20 g of tissue sample.

QA/QC SUMMARY/PAHs (continued)

Aliquots of samples that were analyzed in triplicate, used for spiking, or were re-extracted, were generally less than 20 g due to limited quantities of tissue available. Because MDLs reported are corrected for sample weight, the MDLs reported for these samples appear elevated and in some cases may exceed the target detection limit.

In addition a method detection limit verification study was performed, which consisted of analyzing four spiked aliquots of a background clam sample received with this project. The standard deviation of the results of these replicate analyses was multiplied by 3.5. Detection limits calculated in this way were all less than the target detection limit of 4 ng/g wet wt.

METHOD BLANKS

One method blank was extracted with each extraction batch. Benz[a]anthracene was detected in blanks from all batches and benzo[b]fluoranthene was detected in the blank from Batch 3. Two method blanks were analyzed with Batch 7 and in addition to benz[a]anthracene, three other compounds were detected in at least one of the two blanks; naphthalene, benzo[a]pyrene and indeno(123-cd)pyrene. All blank levels were less than three times the target MDL of 4 ng/g wet wt. Sample values that were less than five times the value of the method blank associated with that sample were flagged with a "B."

SURROGATES

Five isotopically labeled compounds were added prior to extraction to assess the efficiency of the method. These were d8-naphthalene, d10-acenaphthene, d12-chrysene, d14-dibenz[a,h]anthracene and d4-1,4 dichlorobenzene. Recoveries of all surrogates were within the quality control limits of 30% -150% with the exception of low recoveries for d4-1,4 dichlorobenzene in one sample from Batch 1 and Batch 4 and two samples in Batch seven. In addition, d8-naphthalene recovery was low in two samples in Batch seven.

MATRIX SPIKES

One sample from each batch was spiked with all PAH compounds. Matrix spike recoveries were generally, within QC limits of 50% -120%, with some exceptions. The recoveries for benzo(b)- and benzo(k)fluoranthene were variable due to the poor resolution of these two compounds. Spike recoveries quantified as the sum of these two compounds were within QC limits. Spike recoveries for a number of PAH compounds in Batches 4 and 7 were out of control due to high native levels, relative to the levels spiked. Spike concentrations were from 2 to 20 times lower than native concentrations. Recoveries for a number of compounds in Batches 4 and 6 were slightly above the upper control limit. These recoveries were all between 120% and 140%.

REPLICATES

One sample from each batch was extracted and analyzed in triplicate. Precision was measured by calculating the relative standard deviation (RSD) between the replicate results. All RSDs were within $\pm 30\%$.

SRMs

Not applicable.

QA/QC SUMMARY/PAHs (continued)

MISCELLANEOUS

Some of the compounds are flagged to indicate that the ion ratio for that compound was outside of the QC range. This is due primarily to low levels of the compound of interest. Because the confirmation ion is present at only a fraction of the level of the parent ion, when the native level of the compound is low, the amount of error in the concentration measurement of the confirmation ion goes up. The compound is actually quantified from the parent ion only, so most likely this will not affect the quality of the data. For sample values that are relatively high (>5 times the MDL) it may be an indication of some sort of interference.

REFERENCES

Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.

EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

Table F.1. Metals in Tissue of *M. nasuta* (Wet weight)

Sediment Treatment	Replicate	Batch	% Dry Weight	<i>M. nasuta</i> Metals (wet weight µg/g)								
				Ag ICP/MS	As ICP/MS	Cd ICP/MS	Cr ICP/MS	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn ICP/MS
COMP PC	1	1	12.91%	0.031	2.70	0.169	0.316	2.47	0.014	0.531	0.584	11.1
COMP PC	2	1	13.51%	0.040	2.96	0.235	0.519	2.58	0.016	0.627	0.804	10.4
COMP PC	3	1	13.89%	0.031	3.04	0.206	0.426	2.43	0.016	0.575	0.743	11.7
COMP PC	4	1	14.98%	0.052	3.45	0.255	0.524	2.91	0.015	0.539	0.851	21.9
COMP PC	5	1	13.68%	0.030	3.01	0.178	0.393	2.05	0.013	0.525	0.622	14.9
R-CLIS	1	1	15.08%	0.028	2.96	0.019	0.360	2.37	0.016	0.682	0.841	10.9
R-CLIS	2	2	14.45%	0.031	2.66	0.027	0.523	2.12	0.017	0.542	1.05	9.78
R-CLIS	3	2	14.15%	0.031	2.89	0.020	0.466	1.97	0.015	0.460	0.831	10.6
R-CLIS	4	1	14.06%	0.029	2.84	0.018	0.433	2.31	0.013	0.606	0.725	11.0
R-CLIS	5-1	1	14.57%	0.032	2.49	0.032	0.490	2.78	0.015	0.590	0.816	13.7
R-CLIS	5-2	1	14.57%	0.029	2.68	0.038	0.460	2.83	0.016	0.608	0.798	14.0
R-CLIS	5-3	1	14.57%	0.028	2.45	0.035	0.471	2.70	0.016	0.576	0.771	13.5
R-MUD	1	1	14.08%	0.031	2.13	0.028	0.404	1.48	0.014	0.322	0.282 U ^(a)	11.3
R-MUD	2	1	18.71%	0.058	4.40	0.060	0.400	2.39	0.023	0.608	0.374 U	17.2
R-MUD	3	1	13.02%	0.040	2.75	0.023	0.365	1.39	0.014	0.292	0.261 U	12.1
R-MUD	4	1	11.83%	0.040	2.45	0.027	0.285	1.13	0.012	0.299	0.237 U	9.17
R-MUD	5	1	20.96%	0.035 U	4.07	0.039	0.585	2.49	0.026	0.486	0.419 U	15.6
C-SB	1	1	12.86%	0.024	3.16	0.022	0.404	1.85	0.011	0.579	0.257 U	12.0
C-SB	2	1	12.45%	0.025	2.95	0.020	0.341	1.93	0.012	0.468	0.249 U	8.83
C-SB	3	1	13.90%	0.023 U	3.06	0.030	0.421	1.74	0.012	0.680	0.278 U	8.15
C-SB	4	1	13.16%	0.022 U	2.95	0.019	0.404	1.65	0.012	0.513	0.263 U	9.29
C-SB	5	1	13.21%	0.023	2.92	0.032	0.432	1.99	0.013	0.633	0.264 U	11.4
<i>M. nasuta</i> Background	1	1	15.16%	0.025 U	2.49	0.019	0.249	1.77	0.011	0.303	0.303 U	10.2
<i>M. nasuta</i> Background	2	1	14.86%	0.025 U	2.69	0.034	0.337	1.52	0.012	0.355	0.297 U	11.2
<i>M. nasuta</i> Background	3-1	1	14.87%	0.025 U	2.38	0.021	0.232	1.74	0.011	0.311	0.298 U	10.6
<i>M. nasuta</i> Background	3-2	1	14.87%	0.025 U	2.54	0.025	0.256	1.72	0.013	0.311	0.298 U	10.6
<i>M. nasuta</i> Background	3-3	1	14.87%	0.025 U	2.48	0.026	0.238	1.78	0.011	0.338	0.298 U	10.5

(a) U Undetected at or above given concentration

Table F.2. Metals in Tissue of *M. nasuta* (Dry Weight)

Sed Code ID	Replicate	Batch	% Dry Mass	<i>M. nasuta</i> Metals (dry weight µg/g)								
				Ag ICP/MS	As ICP/MS	Cd ICP/MS	Cr ICP/MS	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn ICP/MS
COMP PC	1	1	12.91%	0.239	20.9	1.31	2.45	19.1	0.112	4.11	4.52	86.3
COMP PC	2	1	13.51%	0.298	21.9	1.74	3.84	19.1	0.115	4.64	5.95	76.7
COMP PC	3	1	13.89%	0.222	21.9	1.48	3.07	17.5	0.116	4.14	5.35	84.0
COMP PC	4	1	14.98%	0.347	23.0	1.70	3.50	19.4	0.097	3.60	5.68	146
COMP PC	5	1	13.68%	0.220	22.0	1.30	2.87	15.0	0.096	3.84	4.55	109
R-CLIS	1	1	15.08%	0.183	19.6	0.126	2.39	15.7	0.103	4.52	5.58	72.1
R-CLIS	2	2	14.45%	0.212	18.4	0.185	3.62	14.7	0.117	3.75	7.24	67.7
R-CLIS	3	2	14.15%	0.216	20.4	0.138	3.29	13.9	0.105	3.25	5.87	74.7
R-CLIS	4	1	14.06%	0.203	20.2	0.130	3.08	16.4	0.096	4.31	5.16	78.3
R-CLIS	5-1	1	14.57%	0.219	17.1	0.217	3.36	19.1	0.103	4.05	5.60	94.1
R-CLIS	5-2	1	14.57%	0.196	18.4	0.259	3.16	19.4	0.108	4.17	5.48	96.1
R-CLIS	5-3	1	14.57%	0.193	16.8	0.238	3.23	18.5	0.111	3.95	5.29	92.7
R-MUD	1	1	14.08%	0.221	15.1	0.196	2.87	10.5	0.099	2.29	2.00 U ^(a)	80.0
R-MUD	2	1	18.71%	0.309	23.5	0.323	2.14	12.8	0.124	3.25	2.00 U	91.9
R-MUD	3	1	13.02%	0.307	21.1	0.180	2.80	10.7	0.111	2.24	2.00 U	93.3
R-MUD	4	1	11.83%	0.336	20.7	0.227	2.41	9.51	0.103	2.53	2.00 U	77.5
R-MUD	5	1	20.96%	0.166 U	19.4	0.186	2.79	11.9	0.126	2.32	2.00 U	74.6
C-SB	1	1	12.86%	0.184	24.6	0.174	3.14	14.4	0.082	4.50	2.00 U	93.4
C-SB	2	1	12.45%	0.203	23.7	0.158	2.74	15.5	0.097	3.76	2.00 U	70.9
C-SB	3	1	13.90%	0.166 U	22.0	0.214	3.03	12.5	0.083	4.89	2.00 U	58.6
C-SB	4	1	13.16%	0.166 U	22.4	0.146	3.07	12.5	0.093	3.90	2.00 U	70.6
C-SB	5	1	13.21%	0.171	22.1	0.242	3.27	15.1	0.102	4.79	2.00 U	86.1
<i>M. nasuta</i> Background	1	1	15.16%	0.166 U	16.4	0.125	1.64	11.7	0.075	2.00	2.00 U	67.4
<i>M. nasuta</i> Background	2	1	14.86%	0.166 U	18.1	0.229	2.27	10.2	0.079	2.39	2.00 U	75.5
<i>M. nasuta</i> Background	3-1	1	14.87%	0.166 U	16.0	0.140	1.56	11.7	0.071	2.09	2.00 U	71.0
<i>M. nasuta</i> Background	3-2	1	14.87%	0.166 U	17.1	0.165	1.72	11.6	0.085	2.09	2.00 U	71.3
<i>M. nasuta</i> Background	3-3	1	14.87%	0.166 U	16.7	0.175	1.60	12.0	0.073	2.27	2.00 U	70.5

(a) U Undetected at or above given concentration

TABLE F.3. Quality Control Summary for Metals in Tissue of *M. nasuta*

Sed Code ID	Replicate	Batch	<i>M. nasuta</i> Metals (µg/g dry weight)								
			Ag	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn
<u>Method Blanks</u>											
Blank-1		1	0.166 U ^(a)	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-2		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-3		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-4		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-5		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
<u>Matrix Spikes</u>											
COMP EC-A	3	1	0.244	19.7	0.276	4.37	20.1	0.113	4.42	10.3	81.3
COMP EC-A, MS	3		1.95	72.7	4.21	14.2	73.9	1.22	14.5	14.8	163
Concentration Recovered			1.71	53.0	3.93	9.83	53.8	1.11	10.1	4.52	81.7
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			82%	102%	94%	95%	103%	106%	97%	108%	82%
COMP HU-C	5	1	0.569	20.9	0.37	8.01	23.5	0.242	5.28	10.4	88.2
COMP HU-C, MS	5	1	2.15	74.0	3.95	17.9	76.3	1.21	15.9	14.5	175
Concentration Recovered			1.58	53.1	3.58	9.89	52.8	0.968	10.6	4.14	86.8
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			76%	102%	86%	95%	101%	93%	102%	99%	87%
R-CLIS	5	1	0.203	17.4	0.238	3.25	19.0	0.107	4.06	5.46	94.3
R-CLIS, MS	5	1	1.91	74.3	4.26	13.9	74.1	1.22	14.8	10.2	190
Concentration Recovered			1.71	56.9	4.02	10.65	55.1	1.11	10.7	4.74	95.7
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			82%	109%	96%	102%	106%	107%	103%	114%	96%
<i>M. nasuta</i> Background	3	1	0.166 U	16.6	0.160	1.63	11.8	0.076	2.15	2.00 U	70.9
<i>M. nasuta</i> Background, MS	3	1	1.78	71.7	3.90	10.9	64.7	1.12	12.6	4.75	163
Concentration Recovered			1.78	55.1	3.74	9.27	52.9	1.04	10.5	4.75	92.1
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			86%	106%	90%	89%	102%	100%	100%	114%	92%

TABLE F.3. (contd)

Sed Code ID	Replicate	Batch	<i>M. nasuta</i> Metals ($\mu\text{g/g}$ dry weight)								
			Ag	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn
<u>Standard Reference Material</u>											
Certified value			1.68	14.0	4.15	1.43	66.3	0.0642	2.25	0.371	830
range			± 0.15	± 1.2	± 0.38	± 0.46	± 4.3	± 0.0067	± 0.44	± 0.014	± 57
SRM 1566a	1	1	1.38	13.6	4.05	1.25	62.6	0.063	1.87	0.372	762
SRM 1566a	2	1	1.41	13.6	4.08	1.23	65.4	0.063	1.61	0.368	808
SRM 1566a	3	1	1.35	13.0	3.99	1.20	64.4	0.060	2.18	0.392	755
SRM 1566a	4	1	1.42	13.8	4.19	0.931	66.9	0.068	2.50	0.382	777
SRM 1566a	5	1	1.44	13.3	3.65	1.04	67.1	0.061	1.51	0.377	765
Percent Difference	1		18	3	2	13	6	2	17	0	8
Percent Difference	2		16	3	2	14	1	2	28 ^(b)	1	3
Percent Difference	3		20	7	4	16	3	7	3	6	9
Percent Difference	4		15	1	1	35 ^(b)	1	6	11	3	6
Percent Difference	5		14	5	12	27 ^(b)	1	5	33 ^(b)	2	8
<u>Analytical Replicates</u>											
COMP EC-A, Replicate 1	3	1	0.246	19.1	0.256	4.66	21.0	0.130	4.80	11.6	81.1
COMP EC-A, Replicate 2	3	1	0.242	18.9	0.305	4.32	20.6	0.105	4.46	9.69	81.9
COMP EC-A, Replicate 3	3	1	0.245	21.0	0.267	4.12	18.8	0.105	4.00	9.54	80.9
RSD			1%	6%	9%	6%	6%	13%	9%	11%	1%
COMP HU-C, Replicate 1	5	1	0.565	20.5	0.396	7.80	24.1	0.242	5.28	10.6	86.3
COMP HU-C, Replicate 2	5	1	0.629	21.8	0.380	8.62	23.4	0.245	5.27	10.7	88.5
COMP HU-C, Replicate 3	5	1	0.514	20.3	0.335	7.60	22.9	0.238	5.28	9.78	89.9
RSD			10%	4%	9%	7%	3%	1%	0%	5%	2%

TABLE F.3. (contd)

Sed Code ID	Replicate	Batch	<i>M. nasuta</i> Metals ($\mu\text{g/g}$ dry weight)								
			Ag	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn
R-CLIS, Replicate 1	5	1	0.219	17.1	0.217	3.36	19.1	0.103	4.05	5.60	94.1
R-CLIS, Replicate 2	5	1	0.196	18.4	0.259	3.16	19.4	0.108	4.17	5.48	96.1
R-CLIS, Replicate 3	5	1	0.193	16.8	0.238	3.23	18.5	0.111	3.95	5.29	92.7
RSD			7%	5%	9%	3%	2%	4%	3%	3%	2%
<i>M. nasuta</i> Background, Rep 1	3	1	0.166 U	16.0	0.140	1.56	11.7	0.071	2.09	2.00 U	71.0
<i>M. nasuta</i> Background, Rep 2	3	1	0.166 U	17.1	0.165	1.72	11.6	0.085	2.09	2.00 U	71.3
<i>M. nasuta</i> Background, Rep 3	3	1	0.166 U	16.7	0.175	1.60	12.0	0.073	2.27	2.00 U	70.5
RSD			NA ^(c)	3%	11%	5%	2%	10%	5%	NA	1%

F.5

- (a) U Undetected at or above given concentration.
 (b) Outside quality control criteria ($\pm 20\%$) for SRMs.
 (c) NA Not applicable.

TABLE F.4. MDL Verification Study for Metals in *M. nasuta* Tissue Chemistry

Sed Code ID	Replicate	Batch	<i>M. nasuta</i> Metals (µg/g dry weight)								
			Ag	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn
COMP SB-B, Replicate 1	3	1	0.462	22.5	0.188	4.32	20.3	0.122	3.86	6.02	90.1
COMP SB-B, Replicate 2	3	1	0.491	22.4	0.242	4.25	21.5	0.122	4.00	6.27	93.4
COMP SB-B, Replicate 3	3	1	0.392	24.5	0.212	3.41	17.5	0.126	3.19	5.00	88.1
COMP SB-B, Replicate 4	3	1	0.494	23.1	0.201	4.10	21.8	0.126	3.94	6.08	91.3
Mean			0.460	23.1	0.211	4.02	20.3	0.124	3.75	5.84	90.7
Standard Deviation			0.0474	0.967	0.0230	0.417	1.96	0.00231	0.376	0.572	2.22
Method Detection Limit (MDL) ^(a)			0.215	4.39	0.105	1.89	8.90	0.0105	1.71	2.60	10.1

(a) MDL calculated by multiplying the standard deviation times Students-t for four replicates (4.541).

TABLE F.5. Pesticides and PCB Congeners (Wet Weight) in Tissue of *M. nasuta*

Treatment	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC	DUP COMP PC	TRIP COMP PC
Replicate	1	2	3	4	5	5	5
Batch	7	7	7	7	7	7	7
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.91	13.51	13.89	14.98	13.68	13.68	13.68
Heptachlor	0.18 U ^(a)	0.19 U	0.62	0.58	0.23 U	0.22 U	0.21 U
Aldrin	0.90	1.45	1.26	1.11	1.14	1.12	1.05
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U	0.13 U	0.16 U	0.16 U	0.15 U
2,4'-DDE	0.25 U	0.26 U	0.26 U	0.26 U	0.32 U	0.31 U	0.29 U
Endosulfan I	0.17 U	0.18 U	0.18 U	0.18 U	0.22 U	0.21 U	0.20 U
α-Chlordane	3.09	4.39	4.54	3.39	3.54	3.06	2.78
Trans Nonachlor	0.52	0.93	0.87	0.59	0.61	0.39	0.32
4,4'-DDE	4.47	5.92	6.56	4.91	5.66	5.28	4.61
Dieldrin	2.94	4.29	4.63	3.24	3.96	3.79	3.43
2,4'-DDD	4.01	5.86	5.92	4.57	5.45	4.75	4.45
2,4'-DDT	0.17 U	0.18 U	0.18 U	0.18 U	0.22 U	0.21 U	0.20 U
4,4'-DDD	8.51	13.1	13.6	10.1	11.4	10.6	9.14
Endosulfan II	0.17 U	0.18 U	0.18 U	0.18 U	0.22 U	0.21 U	0.20 U
4,4'-DDT	0.15 U	0.15 U	0.15 U	0.15 U	0.19 U	0.18 U	0.17 U
Endosulfan Sulfate	0.17 U	0.18 U	0.18 U	0.18 U	0.22 U	0.21 U	0.20 U
PCB 8	0.39 U	0.41 U	0.41 U	0.41 U	0.51 U	0.48 U	0.46 U
PCB 18	0.66	0.43 U	0.43 U	1.29	0.53 U	0.90	0.48 U
PCB 28	0.99	1.31	1.47	1.25	1.33	1.17	1.03
PCB 52	4.18	5.40	6.07	4.90	5.27	4.90	4.38
PCB 49	1.33	1.81	2.03	1.63	1.83	1.58	1.41
PCB 44	0.35	1.18	0.88	0.87	0.50	0.19 U	0.18 U
PCB 66	0.09 U	0.09 U	0.09 U	0.09 U	0.12 U	0.11 U	0.11 U
PCB 101	5.90	7.60	8.08	6.26	7.32	6.83	6.12
PCB 87	2.57	3.43	3.67	2.82	3.21	3.00	2.64
PCB 118	3.67	4.59	5.05	3.83	4.56	4.02	3.83
PCB 184	0.23 U	0.24 U	0.24 U	0.24 U	0.29 U	0.28 U	0.26 U
PCB 153	1.90	2.45	2.52	1.99	2.53	2.19	2.04
PCB 105	1.49	1.98	2.03	1.62	2.11	1.72	1.60
PCB 138	2.42	3.24	3.21	2.46	3.19	2.82	2.59
PCB 187	0.49	0.65	0.59	0.53	0.63	0.50	0.51
PCB 183	0.23 U	0.29	0.39	0.24 U	0.31	0.28 U	0.26 U
PCB 128	0.48	0.74	0.67	0.51	0.73	0.59	0.56
PCB 180	0.57	0.73	0.72	0.56	0.76	0.73	0.64
PCB 170	0.30	0.39	0.17 U	0.17 U	0.39	0.36	0.34
PCB 195	0.10 U	0.10 U	0.10 U	0.10 U	0.12 U	0.12 U	0.11 U
PCB 206	0.11	0.18	0.11 U	0.11 U	0.18	0.18	0.15
PCB 209	1.37	0.09 U	0.09 U	0.09 U	0.12 U	0.11 U	0.11 U
<u>Surrogate Recoveries (%)</u>							
PCB 103 (SIS)	77	87	79	83	95	95	86
PCB 198 (SIS)	72	76	77	79	93	82	75

TABLE F.5. (contd)

Treatment	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	1	2	3	4	5
Batch	2	3	2	3	2
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.08	18.71	13.02	11.83	20.96
Heptachlor	0.19 U	0.19 U	0.19 U	0.19 U	0.17 U
Aldrin	0.13 U	0.73	0.13 U	0.68	0.22
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U	0.13 U	0.12 U
2,4'-DDE	0.26 U	0.26 U	0.26 U	0.37	0.24 U
Endosulfan I	0.18 U	0.18 U	0.18 U	0.18 U	0.17 U
a-Chlordane	0.10 U	0.10 U	0.10 U	0.10 U	0.09 U
Trans Nonachlor	0.15 U	0.15 U	0.15 U	0.15 U	0.13 U
4,4'-DDE	0.30	0.36	0.46	0.36	0.24
Dieldrin	0.52 U	0.52 U	0.52 U	0.52 U	0.47 U
2,4'-DDD	0.25 U	0.25 U	0.25 U	0.25 U	0.23 U
2,4'-DDT	0.18 U	0.18 U	0.18 U	0.18 U	0.16 U
4,4'-DDD	0.26 U	0.26 U	0.26 U	0.26 U	0.24 U
Endosulfan II	0.18 U	0.18 U	0.18 U	0.18 U	0.17 U
4,4'-DDT	0.41	3.51	0.15 U	1.71	0.43
Endosulfan Sulfate	0.18 U	0.18 U	0.18 U	0.18 U	0.17 U
PCB 8	0.41 U	1.76	0.41 U	1.99	0.38 U
PCB 18	0.43 U	0.43 U	0.43 U	0.43 U	0.40 U
PCB 28	0.53	0.67	0.65	0.64	0.60
PCB 52	0.68	0.94	0.78	0.84	0.83
PCB 49	0.24 U	0.24	0.24 U	0.25	0.22 U
PCB 44	0.17 U	0.17 U	0.17 U	0.17 U	0.15 U
PCB 66	0.09 U	0.09 U	0.74	0.09 U	0.09 U
PCB 101	0.33	0.52	0.45	0.42	0.53
PCB 87	0.16 U	0.29	0.16 U	0.27	0.15 U
PCB 118	0.29 U	0.29 U	0.30	0.29 U	0.27 U
PCB 184	0.24 U	0.24 U	0.24 U	0.24 U	0.22 U
PCB 153	0.17	0.14	0.26	0.13	0.11 U
PCB 105	0.11 U	0.11 U	0.13	0.11 U	0.13
PCB 138	0.29 U	0.29 U	0.29 U	0.29 U	0.30
PCB 187	0.13 U	0.13 U	0.13 U	0.13 U	0.12 U
PCB 183	0.24 U	0.24 U	0.24 U	0.24 U	0.22 U
PCB 128	0.15 U	0.15 U	0.15 U	0.15 U	0.14 U
PCB 180	0.18 U	0.18 U	0.18 U	0.18 U	0.17 U
PCB 170	0.18	0.17 U	0.17 U	0.19	0.15 U
PCB 195	0.10 U	0.10 U	0.10 U	0.10 U	0.09 U
PCB 206	0.11 U	0.11 U	0.11 U	0.11 U	0.10 U
PCB 209	0.09 U	0.09 U	0.09 U	0.09 U	0.09 U
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	81	80	83	76	86
PCB 198 (SIS)	66	129	65	121	65

TABLE F.5. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	1	1	1	1	1
Wet Wt.	20.10	20.14	20.18	20.06	20.27
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	15.08	14.45	14.15	14.06	14.57
Heptachlor	0.19 U	0.19 U	0.19 U	0.19 U	0.18 U
Aldrin	0.13 U	0.13 U	0.13 U	0.13 U	0.12 U
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U	0.13 U	0.13 U
2,4'-DDE	0.26 U	0.26 U	0.26 U	0.26 U	0.26 U
Endosulfan I	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
α -Chlordane	0.10 U	0.10 U	0.10 U	0.10 U	0.09 U
Trans-nonachlor	0.15 U	0.15 U	0.15 U	0.15 U	0.14 U
4,4'-DDE	0.97	1.71	1.13	1.38	1.14
Dieldrin	0.52 U	0.59	0.52 U	0.52 U	0.51 U
2,4'-DDD	0.25 U	0.25 U	0.25 U	0.25 U	0.25 U
2,4'-DDT	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
4,4'-DDD	0.26 U	0.29	0.26 U	0.26 U	0.26 U
Endosulfan II	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
4,4'-DDT	7.73	5.24	8.54	12.3	2.21
Endosulfan Sulfate	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
PCB 8	0.41 U	0.41 U	0.41 U	0.41 U	0.40 U
PCB 18	0.43 U	0.43 U	0.43 U	0.43 U	0.42 U
PCB 28	0.66	0.83	0.73	0.98	0.67
PCB 52	0.63	0.87	0.62	0.95	0.65
PCB 49	0.56	0.72	0.52	0.78	0.53
PCB 44	0.17 U	0.43	0.17 U	0.17 U	0.16 U
PCB 66	1.12	1.33	1.17	0.09 U	1.15
PCB 101	0.88	1.03	0.85	1.16	0.91
PCB 87	0.16 U	0.47	0.16 U	0.16 U	0.25
PCB 118	0.29 U	0.83	0.29 U	0.29 U	0.77
PCB 184	0.24 U	0.24 U	0.24 U	0.24 U	0.23 U
PCB 153	0.98	1.16	0.95	1.16	1.07
PCB 105	0.11 U	0.14	0.12	0.11 U	0.12
PCB 138	0.54	0.64	0.53	0.66	0.59
PCB 187	1.03	0.83	0.81	0.25	2.11
PCB 183	0.24 U	0.24 U	0.24 U	0.24 U	0.23 U
PCB 128	0.16	0.16	0.15 U	0.18	0.15 U
PCB 180	0.29	0.30	0.24	0.28	0.21
PCB 170	0.17 U	0.17	0.17 U	0.17 U	0.16 U
PCB 195	0.10 U	0.10 U	0.10 U	0.10 U	0.10 U
PCB 206	0.11 U	0.11 U	0.11 U	0.11 U	0.11 U
PCB 209	0.09 U	0.09 U	0.09 U	0.09 U	0.09 U
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	75	73	74	70	52
PCB 198 (SIS)	61	58	62	73	42

TABLE F.5. (contd)

Treatment	C-SB	C-SB, Dup	C-SB, Trip	C-SB	C-SB
Replicate	1	1	1	2	3
Batch	3	3	3	2	3
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.86	12.86	12.86	12.45	13.9
Heptachlor	0.36 U	0.36 U	0.37 U	0.19 U	0.18 U
Aldrin	0.25 U	0.25 U	0.25 U	0.13 U	0.12 U
Heptachlor Epoxide	0.26 U	0.26 U	0.26 U	0.13 U	0.13 U
2,4'-DDE	0.51 U	0.51 U	0.52 U	0.26 U	0.26 U
Endosulfan I	0.35 U	0.35 U	0.36 U	0.18 U	0.18 U
α -Chlordane	0.19 U	0.19 U	0.19 U	0.10 U	0.09 U
Trans Nonachlor	0.28 U	0.28 U	0.29 U	0.15 U	0.14 U
4,4'-DDE	0.81	0.37 U	0.37 U	0.36	0.52
Dieldrin	1.01 U	1.01 U	1.02 U	0.52 U	0.51 U
2,4'-DDD	0.50 U	0.50 U	0.50 U	0.25 U	0.25 U
2,4'-DDT	0.35 U	0.35 U	0.35 U	0.18 U	0.18 U
4,4'-DDD	0.51 U	0.51 U	0.52 U	0.26 U	0.26 U
Endosulfan II	0.35 U	0.35 U	0.36 U	0.18 U	0.18 U
4,4'-DDT	0.30 U	0.30 U	0.30 U	0.37	1.24
Endosulfan Sulfate	0.35 U	0.35 U	0.36 U	0.18 U	0.18 U
PCB 8	0.82	1.26	0.94	0.41 U	0.54
PCB 18	0.84 U	0.84 U	0.85 U	0.43 U	0.42 U
PCB 28	0.40 U	0.40 U	0.40 U	0.20 U	0.23
PCB 52	0.70 U	0.70 U	0.71 U	0.36 U	0.35 U
PCB 49	0.46 U	0.46 U	0.47 U	0.24 U	0.23 U
PCB 44	0.32 U	0.32 U	0.33 U	0.17 U	0.16 U
PCB 66	0.19 U	0.30	0.32	0.90 U	0.09 U
PCB 101	0.29 U	0.29 U	0.29 U	0.15 U	0.19
PCB 87	0.31 U	0.31 U	0.32 U	0.16 U	0.16 U
PCB 118	0.58 U	0.58 U	0.58 U	0.29 U	0.29 U
PCB 184	0.46 U	0.46 U	0.47 U	0.24 U	0.23 U
PCB 153	0.24 U	0.24 U	0.24 U	0.12 U	0.12 U
PCB 105	0.22 U	0.22 U	0.22 U	0.11 U	0.11 U
PCB 138	0.57 U	0.57 U	0.57 U	0.29 U	0.28 U
PCB 187	0.25 U	0.25 U	0.25 U	0.13 U	0.12 U
PCB 183	0.46 U	0.46 U	0.47 U	0.24 U	0.23 U
PCB 128	0.30 U	0.30 U	0.31 U	0.15 U	0.15 U
PCB 180	0.36 U	0.36 U	0.37 U	0.18 U	0.18 U
PCB 170	0.33 U	0.34	0.33 U	0.17 U	0.16 U
PCB 195	0.20 U	0.20 U	0.20 U	0.10 U	0.10 U
PCB 206	0.22 U	0.22 U	0.22 U	0.11 U	0.11 U
PCB 209	0.19 U	0.19 U	0.19 U	0.09 U	0.09 U
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	89	79	88	77	94
PCB 198 (SIS)	144	125	141	59	162 ^(b)

TABLE F.5. (contd)

Treatment	C-SB	C-SB	C-SB, Dup	C-SB, Trip
Replicate	4	5	5	5
Batch	2	2	2	2
Units	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.16	13.21	13.21	13.21
Heptachlor	0.19 U	0.36 U	0.37 U	0.36 U
Aldrin	0.13 U	0.25 U	0.25 U	0.25 U
Heptachlor Epoxide	0.13 U	0.26 U	0.26 U	0.26 U
2,4'-DDE	0.26 U	0.51 U	0.52 U	0.51 U
Endosulfan I	0.18 U	0.35 U	0.36 U	0.25 U
α -Chlordane	0.10 U	0.19 U	0.19 U	0.19 U
Trans Nonachlor	0.15 U	0.28 U	0.29 U	0.28 U
4,4'-DDE	0.45	0.54	0.37 U	0.36 U
Dieldrin	0.52 U	1.01 U	1.02 U	1.00 U
2,4'-DDD	0.25 U	0.50 U	0.50 U	0.49 U
2,4'-DDT	0.18 U	0.35 U	0.35 U	0.35 U
4,4'-DDD	0.26 U	0.51 U	0.52 U	0.51 U
Endosulfan II	0.18 U	0.35 U	0.36 U	0.35 U
4,4'-DDT	0.39	0.91	0.30 U	0.34
Endosulfan Sulfate	0.18 U	0.35 U	0.36 U	0.35 U
PCB 8	0.41 U	0.81 U	0.81 U	0.80 U
PCB 18	0.43 U	0.84 U	0.85 U	0.83 U
PCB 28	0.20 U	0.40 U	0.40 U	0.40 U
PCB 52	0.36 U	0.70 U	0.71 U	0.69 U
PCB 49	0.24 U	0.46 U	0.47 U	0.46 U
PCB 44	0.17 U	0.32 U	0.33 U	0.32 U
PCB 66	0.09 U	0.19 U	0.19 U	0.18 U
PCB 101	0.15 U	0.29 U	0.29 U	0.28 U
PCB 87	0.16 U	0.31 U	0.32 U	0.31 U
PCB 118	0.29 U	0.58 U	0.58 U	0.57 U
PCB 184	0.24 U	0.46 U	0.47 U	0.46 U
PCB 153	0.12 U	0.24 U	0.24 U	0.24 U
PCB 105	0.11 U	0.22 U	0.22 U	0.21 U
PCB 138	0.29 U	0.57 U	0.57 U	0.56 U
PCB 187	0.13 U	0.25 U	0.25 U	0.24 U
PCB 183	0.24 U	0.46 U	0.47 U	0.46 U
PCB 128	0.15 U	0.30 U	0.31 U	0.30 U
PCB 180	0.18 U	0.36 U	0.37 U	0.36 U
PCB 170	0.17 U	0.33 U	0.45	0.32 U
PCB 195	0.10 U	0.20 U	0.20 U	0.19 U
PCB 206	0.11 U	0.22 U	0.22 U	0.22 U
PCB 209	0.09 U	0.19 U	0.19 U	0.18 U
<u>Surrogate Recoveries (%)</u>				
PCB 103 (SIS)	84	82	76	75
PCB 198 (SIS)	66	61	57	58

TABLE F.5. (contd)

Treatment	<i>M. nasuta</i>	<i>M. nasuta</i>	<i>M. nasuta</i>
Replicate	Background	Background	Background
Batch	1	2	3
Units	7	7	7
Percent Dry Weight	ng/g	ng/g	ng/g
	15.16	14.86	14.87
Heptachlor	0.18 U	0.19 U	0.19 U
Aldrin	0.12 U	0.13 U	0.13 U
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U
2,4'-DDE	0.26 U	0.26 U	0.26 U
Endosulfan I	0.18 U	0.18 U	0.18 U
a-Chlordane	0.09 U	0.10 U	0.10 U
Trans Nonachlor	0.14 U	0.15 U	0.15 U
4,4'-DDE	0.58	0.19 U	0.19 U
Dieldrin	0.51 U	0.52 U	0.52 U
2,4'-DDD	0.25 U	0.25 U	0.25 U
2,4'-DDT	0.18 U	0.18 U	0.18 U
4,4'-DDD	0.26 U	0.26 U	0.26 U
Endosulfan II	0.18 U	0.18 U	0.18 U
4,4'-DDT	0.15 U	0.15 U	0.15 U
Endosulfan Sulfate	0.55	0.47	0.39
PCB 8	0.40 U	0.41 U	0.41 U
PCB 18	0.42 U	0.43 U	0.43 U
PCB 28	0.50	0.77	0.20 U
PCB 52	0.35 U	0.36 U	0.36 U
PCB 49	0.23 U	0.24 U	0.24 U
PCB 44	0.16 U	0.17 U	0.17 U
PCB 66	0.09 U	0.09 U	0.09 U
PCB 101	0.14 U	0.15 U	0.15 U
PCB 87	0.16 U	0.16 U	0.16 U
PCB 118	0.29 U	0.29 U	0.29 U
PCB 184	0.23 U	0.24 U	0.24 U
PCB 153	0.12 U	0.12 U	0.12 U
PCB 105	0.11 U	0.11 U	0.11 U
PCB 138	0.28 U	0.29 U	0.29 U
PCB 187	0.12 U	0.13 U	0.13 U
PCB 183	0.23 U	0.24 U	0.24 U
PCB 128	0.15 U	0.15 U	0.15 U
PCB 180	0.18 U	0.18 U	0.18 U
PCB 170	0.16 U	0.17 U	0.17 U
PCB 195	0.10 U	0.10 U	0.10 U
PCB 206	0.11 U	0.11 U	0.11 U
PCB 209	0.09 U	0.09 U	0.09 U
<u>Surrogate Recoveries (%)</u>			
PCB 103 (SIS)	61	61	62
PCB 198 (SIS)	74	76	80

(a) U Undetected at or above given concentration.

(b) Result is outside quality control range (30-150%) for surrogate internal standard.

TABLE F.6. Pesticides and PCB Congeners (Dry Weight) in Tissue of *M. nasuta*

Treatment	COMP PC					DUP	TRIP
	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC
Replicate	1	2	3	4	5	5	5
Batch	7	7	7	7	7	7	7
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.91	13.51	13.89	14.98	13.68	13.68	13.68
Heptachlor	1.4 U ^(a)	1.4 U	4.5	3.9	1.7 U	1.6 U	1.5 U
Aldrin	7.0	10.73	9.07	7.41	8.33	8.19	7.68
Heptachlor Epoxide	1.0 U	1.0 U	0.9 U	0.9 U	1.2 U	1.2 U	1.1 U
2,4'-DDE	1.9 U	1.9 U	1.9 U	1.7 U	2.3 U	2.3 U	2.1 U
Endosulfan I	1.3 U	1.3 U	1.3 U	1.2 U	1.6 U	1.5 U	1.5 U
α -Chlordane	23.9	32.5	32.7	22.6	25.9	22.4	20.3
Trans-nonachlor	4.0	6.9	6.3	3.9	4.5	2.9	2.3
4,4'-DDE	34.6	43.8	47.2	32.8	41.4	38.6	33.7
Dieldrin	22.8	31.8	33.3	21.6	28.9	27.7	25.1
2,4'-DDD	31.1	43.4	42.6	30.5	39.8	34.7	32.5
2,4'-DDT	1.3 U	1.3 U	1.3 U	1.2 U	1.6 U	1.5 U	1.5 U
4,4'-DDD	65.9	97.0	97.9	67.4	83.3	77.6	66.8
Endosulfan II	1.3 U	1.3 U	1.3 U	1.2 U	1.6 U	1.5 U	1.5 U
4,4'-DDT	1.2 U	1.1 U	1.1 U	1.0 U	1.4 U	1.3 U	1.2 U
Endosulfan Sulfate	1.3 U	1.3 U	1.3 U	1.2 U	1.6 U	1.5 U	1.5 U
PCB 8	3.0 U	3.0 U	3.0 U	2.7 U	3.7 U	3.5 U	3.4 U
PCB 18	5.1	3.2 U	3.1 U	8.61	3.9 U	6.6	3.5 U
PCB 28	7.7	9.70	10.6	8.34	9.72	8.55	7.53
PCB 52	32.4	40.0	43.7	32.7	38.5	35.8	32.0
PCB 49	10.3	13.4	14.6	10.9	13.4	11.5	10.3
PCB 44	2.7	8.73	6.3	5.8	3.7	1.4 U	1.3 U
PCB 66	0.70 U	0.67 U	0.65 U	0.60 U	0.88 U	0.80 U	0.80 U
PCB 101	45.7	56.3	58.2	41.8	53.5	49.9	44.7
PCB 87	19.9	25.4	26.4	18.8	23.5	21.9	19.3
PCB 118	28.4	34.0	36.4	25.6	33.3	29.4	28.0
PCB 184	1.8 U	1.8 U	1.7 U	1.6 U	2.1 U	2.0 U	1.9 U
PCB 153	14.7	18.1	18.1	13.3	18.5	16.0	14.9
PCB 105	11.5	14.7	14.6	10.8	15.4	12.6	11.7
PCB 138	18.7	24.0	23.1	16.4	23.3	20.6	18.9
PCB 187	3.8	4.8	4.2	3.5	4.6	3.7	3.7
PCB 183	1.8 U	2.1	2.8	1.6 U	2.3	2.0 U	1.9 U
PCB 128	3.7	5.5	4.8	3.4	5.3	4.3	4.1
PCB 180	4.4	5.4	5.2	3.7	5.6	5.3	4.7
PCB 170	2.3	2.9	1.2 U	1.1 U	2.9	2.6	2.5
PCB 195	0.77 U	0.74 U	0.72 U	0.67 U	0.88 U	0.88 U	0.80 U
PCB 206	0.85	1.3	0.79 U	0.73 U	1.3	1.3	1.1
PCB 209	10.6	0.67 U	0.65 U	0.60 U	0.88 U	0.80 U	0.80 U

TABLE F.6. (contd)

Treatment	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	1	2	3	4	5
Batch	2	3	2	3	2
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.08	18.71	13.02	11.83	20.96
Heptachlor	1.3 U	1.0 U	1.5 U	1.6 U	0.81 U
Aldrin	0.92 U	3.9	1.0 U	5.7	1.0
Heptachlor Epoxide	0.92 U	0.69 U	1.0 U	1.1 U	0.57 U
2,4'-DDE	1.8 U	1.4 U	2.0 U	3.1	1.1 U
Endosulfan I	1.3 U	0.96 U	1.4 U	1.5 U	0.81 U
a-Chlordane	0.71 U	0.53 U	0.77 U	0.85 U	0.4 U
Trans Nonachlor	1.1 U	0.80 U	1.2 U	1.3 U	0.62 U
4,4'-DDE	2.1	1.9	3.5	3.0	1.1
Dieldrin	3.7 U	2.8 U	4.0 U	4.4 U	2.2 U
2,4'-DDD	1.8 U	1.3 U	1.9 U	2.1 U	1.1 U
2,4'-DDT	1.3 U	1.0 U	1.4 U	1.5 U	0.76 U
4,4'-DDD	1.8 U	1.4 U	2.0 U	2.2 U	1.1 U
Endosulfan II	1.3 U	1.0 U	1.4 U	1.5 U	0.81 U
4,4'-DDT	2.9	18.8	1.2 U	14.5	2.1
Endosulfan Sulfate	1.3 U	0.96 U	1.4 U	1.5 U	0.81 U
PCB 8	2.9 U	9.41	3.1 U	16.8	1.8 U
PCB 18	3.1 U	2.3 U	3.3 U	3.6 U	1.9 U
PCB 28	3.8	3.6	5.0	5.4	2.9
PCB 52	4.8	5.0	6.0	7.1	4.0
PCB 49	1.7 U	1.3	1.8 U	2.1	1.0 U
PCB 44	1.2 U	0.91 U	1.3 U	1.4 U	0.72 U
PCB 66	0.6 U	0.5 U	5.7	0.8 U	0.4 U
PCB 101	2.3	2.8	3.5	3.6	2.5
PCB 87	1.1 U	1.5	1.2 U	2.3	0.72 U
PCB 118	2.1 U	1.5 U	2.3	2.5 U	1.3 U
PCB 184	1.7 U	1.3 U	1.8 U	2.0 U	1.0 U
PCB 153	1.2	0.75	2.0	1.1	0.52 U
PCB 105	0.78 U	0.59 U	1.0	0.93 U	0.62
PCB 138	2.1 U	1.5 U	2.2 U	2.5 U	1.4
PCB 187	0.92 U	0.69 U	1.0 U	1.1 U	0.57 U
PCB 183	1.7 U	1.3 U	1.8 U	2.0 U	1.0 U
PCB 128	1.1 U	0.80 U	1.2 U	1.3 U	0.67 U
PCB 180	1.3 U	0.96 U	1.4 U	1.5 U	0.81 U
PCB 170	1.3	0.91 U	1.3 U	1.6	0.72 U
PCB 195	0.71 U	0.53 U	0.77 U	0.85 U	0.4 U
PCB 206	0.78 U	0.59 U	0.84 U	0.93 U	0.48 U
PCB 209	0.6 U	0.5 U	0.7 U	0.8 U	0.4 U

TABLE F.6. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	1	1	1	1	1
Wet Wt.	20.10	20.14	20.18	20.06	20.27
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	15.08	14.45	14.15	14.06	14.57
Heptachlor	1.3 U	1.3 U	1.3 U	1.4 U	1.2 U
Aldrin	0.86 U	0.90 U	0.92 U	0.92 U	0.82 U
Heptachlor Epoxide	0.86 U	0.90 U	0.92 U	0.92 U	0.89 U
2,4'-DDE	1.7 U	1.8 U	1.8 U	1.8 U	1.8 U
Endosulfan I	1.2 U	1.2 U	1.3 U	1.3 U	1.2 U
a-Chlordane	0.66 U	0.69 U	0.71 U	0.71 U	0.62 U
Trans-nonachlor	0.99 U	1.0 U	1.1 U	1.1 U	1.0 U
4,4'-DDE	6.4	11.8	7.99	9.82	7.82
Dieldrin	3.4 U	4.1	3.7 U	3.7 U	3.5 U
2,4'-DDD	1.7 U	1.7 U	1.8 U	1.8 U	1.7 U
2,4'-DDT	1.2 U	1.2 U	1.3 U	1.3 U	1.2 U
4,4'-DDD	1.7 U	2.0	1.8 U	1.8 U	1.8 U
Endosulfan II	1.2 U	1.2 U	1.3 U	1.3 U	1.2 U
4,4'-DDT	51.3	36.3	60.4	87.5	15.2
Endosulfan Sulfate	1.2 U	1.2 U	1.3 U	1.3 U	1.2 U
PCB 8	2.7 U	2.8 U	2.9 U	2.9 U	2.7 U
PCB 18	2.9 U	3.0 U	3.0 U	3.1 U	2.9 U
PCB 28	4.4	5.7	5.2	7.0	4.6
PCB 52	4.2	6.0	4.4	6.8	4.5
PCB 49	3.7	5.0	3.7	5.5	3.6
PCB 44	1.1 U	3.0	1.2 U	1.2 U	1.1 U
PCB 66	7.43	9.20	8.27	0.6 U	7.89
PCB 101	5.8	7.13	6.0	8.25	6.2
PCB 87	1.1 U	3.3	1.1 U	1.1 U	1.7
PCB 118	1.9 U	5.7	2.0 U	2.1 U	5.3
PCB 184	1.6 U	1.7 U	1.7 U	1.7 U	1.6 U
PCB 153	6.5	8.03	6.7	8.25	7.34
PCB 105	0.73 U	0.97	0.85	0.78 U	0.82
PCB 138	3.6	4.4	3.7	4.7	4.0
PCB 187	6.83	5.7	5.7	1.8	14.5
PCB 183	1.6 U	1.7 U	1.7 U	1.7 U	1.6 U
PCB 128	1.1	1.1	1.1 U	1.3	1.0 U
PCB 180	1.9	2.1	1.7	2.0	1.4
PCB 170	1.1 U	1.2	1.2 U	1.2 U	1.1 U
PCB 195	0.66 U	0.69 U	0.71 U	0.71 U	0.69 U
PCB 206	0.73 U	0.76 U	0.78 U	0.78 U	0.75 U
PCB 209	0.6 U	0.6 U	0.6 U	0.6 U	0.6 U

TABLE F.6. (contd)

Treatment	C-SB	C-SB, Dup	C-SB, Trip	C-SB	C-SB
Replicate	1	1	1	2	3
Batch	3	3	3	2	3
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.86	12.86	12.86	12.45	13.9
Heptachlor	2.8 U	2.8 U	2.9 U	1.5 U	1.3 U
Aldrin	1.9 U	1.9 U	1.9 U	1.0 U	0.86 U
Heptachlor Epoxide	2.0 U	2.0 U	2.0 U	1.0 U	0.94 U
2,4'-DDE	4.0 U	4.0 U	4.0 U	2.1 U	1.9 U
Endosulfan I	2.7 U	2.7 U	2.8 U	1.4 U	1.3 U
a-Chlordane	1.5 U	1.5 U	1.5 U	0.80 U	0.65 U
Trans Nonachlor	2.2 U	2.2 U	2.3 U	1.2 U	1.0 U
4,4'-DDE	6.3	2.9 U	2.9 U	2.9	3.7
Dieldrin	7.85 U	7.85 U	7.93 U	4.2 U	3.7 U
2,4'-DDD	3.9 U	3.9 U	3.9 U	2.0 U	1.8 U
2,4'-DDT	2.7 U	2.7 U	2.7 U	1.4 U	1.3 U
4,4'-DDD	4.0 U	4.0 U	4.0 U	2.1 U	1.9 U
Endosulfan II	2.7 U	2.7 U	2.8 U	1.4 U	1.3 U
4,4'-DDT	2.3 U	2.3 U	2.3 U	3.0	8.92
Endosulfan Sulfate	2.7 U	2.7 U	2.8 U	1.4 U	1.3 U
PCB 8	6.4	9.80	7.3	3.3 U	3.9
PCB 18	6.5 U	6.5 U	6.6 U	3.5 U	3.0 U
PCB 28	3.1 U	3.1 U	3.1 U	1.6 U	1.7
PCB 52	5.4 U	5.4 U	5.5 U	2.9 U	2.5 U
PCB 49	3.6 U	3.6 U	3.7 U	1.9 U	1.7 U
PCB 44	2.5 U	2.5 U	2.6 U	1.4 U	1.2 U
PCB 66	1.5 U	2.3	2.5	7.2 U	0.6 U
PCB 101	2.3 U	2.3 U	2.3 U	1.2 U	1.4
PCB 87	2.4 U	2.4 U	2.5 U	1.3 U	1.2 U
PCB 118	4.5 U	4.5 U	4.5 U	2.3 U	2.1 U
PCB 184	3.6 U	3.6 U	3.7 U	1.9 U	1.7 U
PCB 153	1.9 U	1.9 U	1.9 U	0.96 U	0.86 U
PCB 105	1.7 U	1.7 U	1.7 U	0.88 U	0.79 U
PCB 138	4.4 U	4.4 U	4.4 U	2.3 U	2.0 U
PCB 187	1.9 U	1.9 U	1.9 U	1.0 U	0.86 U
PCB 183	3.6 U	3.6 U	3.7 U	1.9 U	1.7 U
PCB 128	2.3 U	2.3 U	2.4 U	1.2 U	1.1 U
PCB 180	2.8 U	2.8 U	2.9 U	1.4 U	1.3 U
PCB 170	2.6 U	2.6	2.6 U	1.4 U	1.2 U
PCB 195	1.6 U	1.6 U	1.6 U	0.80 U	0.72 U
PCB 206	1.7 U	1.7 U	1.7 U	0.88 U	0.79 U
PCB 209	1.5 U	1.5 U	1.5 U	0.7 U	0.6 U

TABLE F.6. (contd)

Treatment	C-SB	C-SB	C-SB, Dup	C-SB, Trip
Replicate	4	5	5	5
Batch	2	2	2	2
Units	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.16	13.21	13.21	13.21
Heptachlor	1.4 U	2.7 U	2.8 U	2.7 U
Aldrin	0.99 U	1.9 U	1.9 U	1.9 U
Heptachlor Epoxide	0.99 U	1.97 U	1.97 U	1.97 U
2,4'-DDE	2.0 U	3.9 U	3.9 U	3.9 U
Endosulfan I	1.4 U	2.6 U	2.7 U	1.9 U
α -Chlordane	0.76 U	1.4 U	1.4 U	1.4 U
Trans Nonachlor	1.1 U	2.1 U	2.2 U	2.1 U
4,4'-DDE	3.4	4.1	2.8 U	2.7 U
Dieldrin	4.0 U	7.65 U	7.72 U	7.57 U
2,4'-DDD	1.9 U	3.8 U	3.8 U	3.7 U
2,4'-DDT	1.4 U	2.6 U	2.6 U	2.6 U
4,4'-DDD	2.0 U	3.9 U	3.9 U	3.9 U
Endosulfan II	1.4 U	2.6 U	2.7 U	2.6 U
4,4'-DDT	3.0	6.9	2.3 U	2.6
Endosulfan Sulfate	1.4 U	2.6 U	2.7 U	2.6 U
PCB 8	3.1 U	6.1 U	6.1 U	6.1 U
PCB 18	3.3 U	6.4 U	6.4 U	6.3 U
PCB 28	1.5 U	3.0 U	3.0 U	3.0 U
PCB 52	2.7 U	5.3 U	5.4 U	5.2 U
PCB 49	1.8 U	3.5 U	3.6 U	3.5 U
PCB 44	1.3 U	2.4 U	2.5 U	2.4 U
PCB 66	0.7 U	1.4 U	1.4 U	1.4 U
PCB 101	1.1 U	2.2 U	2.2 U	2.1 U
PCB 87	1.2 U	2.3 U	2.4 U	2.3 U
PCB 118	2.2 U	4.4 U	4.4 U	4.3 U
PCB 184	1.8 U	3.5 U	3.6 U	3.5 U
PCB 153	0.91 U	1.8 U	1.8 U	1.8 U
PCB 105	0.84 U	1.7 U	1.7 U	1.6 U
PCB 138	2.2 U	4.3 U	4.3 U	4.2 U
PCB 187	1.0 U	1.9 U	1.9 U	1.8 U
PCB 183	1.8 U	3.5 U	3.6 U	3.5 U
PCB 128	1.1 U	2.3 U	2.3 U	2.3 U
PCB 180	1.4 U	2.7 U	2.8 U	2.7 U
PCB 170	1.3 U	2.5 U	3.4	2.4 U
PCB 195	0.76 U	1.5 U	1.5 U	1.4 U
PCB 206	0.84 U	1.7 U	1.7 U	1.7 U
PCB 209	0.7 U	1.4 U	1.4 U	1.4 U

TABLE F.6. (contd)

Treatment	<i>M. nasuta</i> Background	<i>M. nasuta</i> Background	<i>M. nasuta</i> Background
Replicate	1	2	3
Batch	7	7	7
Units	ng/g	ng/g	ng/g
Percent Dry Weight	15.16	14.86	14.87
Heptachlor	1.2 U	1.3 U	1.3 U
Aldrin	0.79 U	0.87 U	0.87 U
Heptachlor Epoxide	0.86 U	0.87 U	0.87 U
2,4'-DDE	1.7 U	1.7 U	1.7 U
Endosulfan I	1.2 U	1.2 U	1.2 U
a-Chlordane	0.59 U	0.67 U	0.67 U
Trans Nonachlor	0.9 U	1.0 U	1.0 U
4,4'-DDE	3.8	1.3 U	1.3 U
Dieldrin	3.4 U	3.5 U	3.5 U
2,4'-DDD	1.6 U	1.7 U	1.7 U
2,4'-DDT	1.2 U	1.2 U	1.2 U
4,4'-DDD	1.7 U	1.7 U	1.7 U
Endosulfan II	1.2 U	1.2 U	1.2 U
4,4'-DDT	1.0 U	1.0 U	1.0 U
Endosulfan Sulfate	3.6	3.2	2.6
PCB 8	2.6 U	2.8 U	2.8 U
PCB 18	2.8 U	2.9 U	2.9 U
PCB 28	3.3	5.2	1.3 U
PCB 52	2.3 U	2.4 U	2.4 U
PCB 49	1.5 U	1.6 U	1.6 U
PCB 44	1.1 U	1.1 U	1.1 U
PCB 66	0.6 U	0.6 U	0.6 U
PCB 101	0.92 U	1.0 U	1.0 U
PCB 87	1.1 U	1.1 U	1.1 U
PCB 118	1.9 U	2.0 U	2.0 U
PCB 184	1.5 U	1.6 U	1.6 U
PCB 153	0.79 U	0.81 U	0.81 U
PCB 105	0.73 U	0.74 U	0.74 U
PCB 138	1.8 U	2.0 U	2.0 U
PCB 187	0.79 U	0.87 U	0.87 U
PCB 183	1.5 U	1.6 U	1.6 U
PCB 128	1.0 U	1.0 U	1.0 U
PCB 180	1.2 U	1.2 U	1.2 U
PCB 170	1.1 U	1.1 U	1.1 U
PCB 195	0.66 U	0.67 U	0.67 U
PCB 206	0.73 U	0.74 U	0.74 U
PCB 209	0.6 U	0.6 U	0.6 U

(a) U Undetected at or above given concentration.

TABLE F.7. Quality Control Summary for Pesticides and PCB Congeners in Tissue of *M. nasuta* (Wet Weight)

<u>Matrix Spike Results</u>										
Treatment	Matrix Spike		Matrix Spike		Matrix Spike		Matrix Spike		Amount Spiked	Percent Recovery
	COMP	HU-A	COMP	HU-A	COMP	HU-C	COMP	HU-C		
Replicate	1	1	1	1	5	5	5	5		
Batch	1	1	1	1	2	2	2	2		
Wet Wt Units	20.12 ng/g	20.12 ng/g	20.12 ng/g	20.12 ng/g	10.14 ng/g	10.25 ng/g	10.14 ng/g	10.25 ng/g		
Heptachlor	0.19 U ^(a)	2.62	2.50	105	0.37 U	4.69	4.90	96		
Aldrin	1.66	4.28	2.50	105	3.40	5.96	4.90	52		
Heptachlor Epoxide	0.13 U	2.13	2.50	85	0.26 U	3.53	4.90	72		
2,4'-DDE	0.26 U	NA ^(b)	NS ^(c)	NA	0.52 U	NA	NS	NA		
Endosulfan I	0.18 U	2.28	2.50	91	0.36 U	3.31	4.90	68		
α-Chlordane	0.10 U	NA	NS	NA	0.85	NA	NS	NA		
Trans Nonachlor	0.15 U	NA	NS	NA	0.29 U	NA	NS	NA		
4,4'-DDE	5.48	7.48	2.50	80	10.1	13.9	4.90	78		
Dieldrin	0.91	3.12	2.50	88	2.13	5.15	4.90	62		
2,4'-DDD	0.77	NS	NS	NS	1.49	NA	NS	NA		
2,4'-DDT	0.18 U	NS	NS	NS	0.35 U	NA	NS	NA		
4,4'-DDD	2.67	5.24	2.50	103	4.61	8.58	4.90	81		
Endosulfan II	0.18 U	2.92	2.50	117	0.36 U	4.49	4.90	92		
4,4'-DDT	12.6	14.1	2.50	60	0.96	6.16	4.90	106		
Endosulfan Sulfate	0.18 U	2.00	2.50	80	0.65	4.51	4.90	79		
PCB 8	0.41 U	NA	NS	NA	0.81 U	NA	NS	NA		
PCB 18	4.09	NA	NS	NA	17.0	NA	NS	NA		
PCB 28	4.92	8.51	3.19	113	24.6	30.9	6.25	101		
PCB 52	4.65	10.5	6.65	88	21.1	33.0	13.0	92		
PCB 49	3.33	NS	NS	NS	16.7	NA	NS	NA		
PCB 44	1.37	NA	NS	NA	9.51	NA	NS	NA		
PCB 66	4.11	NA	NS	NA	19.6	NA	NS	NA		
PCB 101	2.54	6.73	4.51	93	9.97	17.9	8.84	90		
PCB 87	0.86	NA	NS	NA	3.11	NA	NS	NA		
PCB 118	1.62	NA	NS	NA	7.68	NA	NS	NA		
PCB 184	0.24 U	NA	NS	NA	0.47 U	NA	NS	NA		
PCB 153	1.26	3.31	2.64	78	4.43	8.76	5.17	84		
PCB 105	0.63	NA	NS	NA	2.85	NA	NS	NA		
PCB 138	1.02	2.75	2.04	85	3.68	7.29	3.99	90		
PCB 187	1.18	NA	NS	NA	0.25 U	NA	NS	NA		
PCB 183	0.24 U	NA	NS	NA	0.54	NA	NS	NA		
PCB 128	0.27	NA	NS	NA	0.90	NA	NS	NA		
PCB 180	0.40	NA	NS	NA	1.25	NA	NS	NA		
PCB 170	0.17 U	NA	NS	NA	0.33 U	NA	NS	NA		
PCB 195	0.10 U	NA	NS	NA	0.20 U	NA	NS	NA		
PCB 206	0.24	NA	NS	NA	0.41	NA	NS	NA		
PCB 209	0.11	NA	NS	NA	0.29	NA	NS	NA		
<u>Surrogate Recoveries (%)</u>										
PCB 103 (SIS)	65	65	NA	NA	81	77	NA	NA		
PCB 198 (SIS)	63	69	NA	NA	59	59	NA	NA		

TABLE F.7. (contd)

Matrix Spike Results									
Treatment	COMP SB-A				COMP PC				
	COMP SB-A	MS	Amount Spiked	Percent Recovery	COMP PC	MS	Amount Spiked	Percent Recovery	
Replicate	3	3			1	1			
Batch	3	3			7	7			
Wet Wt Units	10.06 ng/g	10.32 ng/g			20.84 ng/g	20.18 ng/g			
Heptachlor	0.37 U	4.35	4.85	90	0.18 U	2.41	2.50	96	
Aldrin	1.45	5.18	4.85	77	0.90	2.96	2.50	82	
Heptachlor Epoxide	0.26 U	3.97	4.85	82	0.13 U	2.58	2.50	103	
2,4'-DDE	0.52 U	NA	NS	NA	0.25 U	NA	NS	NA	
Endosulfan I	0.36 U	3.62	4.85	75	0.17 U	2.11	2.50	84	
a-Chlordane	0.75	NA	NS	NA	3.09	NA	NS	NA	
Trans Nonachlor	0.29 U	NA	NS	NA	0.52	NA	NS	NA	
4,4'-DDE	4.00	7.91	4.85	81	4.47	7.19	2.50	109	
Dieldrin	1.50	4.84	4.85	69	2.94	5.83	2.50	116	
2,4'-DDD	0.55	NA	NS	NA	4.01	NA	NS	NA	
2,4'-DDT	0.35 U	NA	NS	NA	0.17 U	NA	NS	NA	
4,4'-DDD	2.22	7.25	4.85	104	8.51	13.3	2.50	192 ^(e)	
Endosulfan II	0.36 U	3.77	4.85	78	0.17 U	2.72	2.50	109	
4,4'-DDT	2.12	7.55	4.85	112	0.15 U	3.22	2.50	129 ^(e)	
Endosulfan Sulfate	0.36 U	4.57	4.85	94	0.17 U	3.04	2.50	122 ^(e)	
PCB 8	1.54	NA	NS	NA	0.39 U	NA	NS	NA	
PCB 18	1.63	NA	NS	NA	0.66	NA	NS	NA	
PCB 28	3.31	9.60	6.18	102	0.99	4.93	3.19	124 ^(e)	
PCB 52	3.35	14.8	12.9	89	4.18	10.9	6.65	101	
PCB 49	2.63	NA	NS	NA	1.33	NA	NS	NA	
PCB 44	0.84	NA	NS	NA	0.35	NA	NS	NA	
PCB 66	4.44	NA	NS	NA	0.09 U	NA	NS	NA	
PCB 101	3.34	11.8	8.75	97	5.90	11.0	4.51	113	
PCB 87	1.12	NA	NS	NA	2.57	NA	NS	NA	
PCB 118	1.71	NA	NS	NA	3.67	NA	NS	NA	
PCB 184	0.47 U	NA	NS	NA	0.23 U	NA	NS	NA	
PCB 153	1.61	4.95	5.12	65	1.90	4.21	2.64	88	
PCB 105	0.57	NA	NS	NA	1.49	NA	NS	NA	
PCB 138	1.30	4.93	3.95	92	2.42	4.63	2.04	108	
PCB 187	0.37	NA	NS	NA	0.49	NA	NS	NA	
PCB 183	0.47 U	NA	NS	NA	0.23 U	NA	NS	NA	
PCB 128	0.31 U	NA	NS	NA	0.48	NA	NS	NA	
PCB 180	0.94	NA	NS	NA	0.57	NA	NS	NA	
PCB 170	0.63	NA	NS	NA	0.30	NA	NS	NA	
PCB 195	0.20 U	NA	NS	NA	0.10 U	NA	NS	NA	
PCB 206	0.22 U	NA	NS	NA	0.11	NA	NS	NA	
PCB 209	0.19 U	NA	NS	NA	1.37	NA	NS	NA	
<u>Surrogate Recoveries (%)</u>									
PCB 103 (SIS)	86	82	NA	NA	77	82	NA	NA	
PCB 198 (SIS)	154 ^(d)	147	NA	NA	72	67	NA	NA	

TABLE F.7. (contd)

Analytical Replicate Results

Treatment	DUP		TRIP		Control-SB	DUP		TRIP	
	COMP	EC-B	COMP	EC-B		Control-SB	Control-SB	Control-SB	Control-SB
Replicate	5	5	5	5	5	5	5	5	
Batch	1	1	1	1	2	2	2	2	
Wet Wt	10.04	10.02	10.11		10.16	10	10	10	NA
Units	ng/g	ng/g	ng/g	RSD%	ng/g	ng/g	ng/g	ng/g	RSD%
Heptachlor	0.37 U	0.37 U	0.37 U	NA	0.36 U	0.37 U	0.36 U	0.36 U	NA
Aldrin	1.15	1.23	1.21	3	0.25 U	0.25 U	0.25 U	0.25 U	NA
Heptachlor Epoxide	0.27 U	0.27 U	0.26 U	NA	0.26 U	0.26 U	0.26 U	0.26 U	NA
2,4'-DDE	0.52 U	0.52 U	0.52 U	NA	0.51 U	0.52 U	0.51 U	0.51 U	NA
Endosulfan I	0.36 U	0.36 U	0.36 U	NA	0.35 U	0.36 U	0.25 U	0.25 U	NA
a-Chlordane	2.58	2.98	2.92	8	0.19 U	0.19 U	0.19 U	0.19 U	NA
Trans Nonachlor	0.75	1.06	1.01	18	0.28 U	0.29 U	0.28 U	0.28 U	NA
4,4'-DDE	3.65	3.82	3.91	3	0.54	0.37 U	0.37 U	0.36 U	NA
Dieldrin	1.77	1.95	1.92	5	1.01 U	1.02 U	1.00 U	1.00 U	NA
2,4'-DDD	1.62	1.50	1.59	4	0.50 U	0.50 U	0.49 U	0.49 U	NA
2,4'-DDT	0.36 U	0.36 U	0.35 U	NA	0.35 U	0.35 U	0.35 U	0.35 U	NA
4,4'-DDD	5.35	5.63	5.96	5	0.51 U	0.52 U	0.51 U	0.51 U	NA
Endosulfan II	0.36 U	0.36 U	0.36 U	NA	0.35 U	0.36 U	0.35 U	0.35 U	NA
4,4'-DDT	1.86	2.54	3.15	26	0.91	0.30 U	0.34	0.34	NA
Endosulfan Sulfate	0.36 U	0.36 U	0.36 U	NA	0.35 U	0.36 U	0.35 U	0.35 U	NA
PCB 8	0.82 U	0.82 U	0.82 U	NA	0.81 U	0.81 U	0.80 U	0.80 U	NA
PCB 18	6.73	6.77	6.82	1	0.84 U	0.85 U	0.83 U	0.83 U	NA
PCB 28	7.35	7.93	7.85	4	0.40 U	0.40 U	0.40 U	0.40 U	NA
PCB 52	7.26	7.29	7.44	1	0.70 U	0.71 U	0.69 U	0.69 U	NA
PCB 49	4.78	4.89	4.99	2	0.46 U	0.47 U	0.46 U	0.46 U	NA
PCB 44	2.17	2.65	2.54	10	0.32 U	0.33 U	0.32 U	0.32 U	NA
PCB 66	6.75	7.12	7.26	4	0.19 U	0.19 U	0.18 U	0.18 U	NA
PCB 101	3.35	3.42	3.73	6	0.29 U	0.29 U	0.28 U	0.28 U	NA
PCB 87	1.23	1.35	1.41	7	0.31 U	0.32 U	0.31 U	0.31 U	NA
PCB 118	2.48	2.49	2.70	5	0.58 U	0.58 U	0.57 U	0.57 U	NA
PCB 184	0.47 U	0.47 U	0.47 U	NA	0.46 U	0.47 U	0.46 U	0.46 U	NA
PCB 153	1.38	1.39	1.46	3	0.24 U	0.24 U	0.24 U	0.24 U	NA
PCB 105	0.93	0.97	1.03	5	0.22 U	0.22 U	0.21 U	0.21 U	NA
PCB 138	1.19	1.23	1.31	5	0.57 U	0.57 U	0.56 U	0.56 U	NA
PCB 187	3.47	3.11	3.41	6	0.25 U	0.25 U	0.24 U	0.24 U	NA
PCB 183	0.47 U	0.47 U	0.47 U	NA	0.46 U	0.47 U	0.46 U	0.46 U	NA
PCB 128	0.33	0.31 U	0.34	NA	0.30 U	0.31 U	0.30 U	0.30 U	NA
PCB 180	0.68	0.65	0.62	5	0.36 U	0.37 U	0.36 U	0.36 U	NA
PCB 170	0.33 U	0.33 U	0.33 U	NA	0.33 U	0.45	0.32 U	0.32 U	NA
PCB 195	0.20 U	0.20 U	0.20 U	NA	0.20 U	0.20 U	0.19 U	0.19 U	NA
PCB 206	0.23 U	0.23 U	0.23 U	NA	0.22 U	0.22 U	0.22 U	0.22 U	NA
PCB 209	0.19 U	0.19 U	0.19 U	NA	0.19 U	0.19 U	0.18 U	0.18 U	NA
<u>Surrogate Recoveries (%)</u>									
PCB 103 (SIS)	67	80	74	NA	82	76	75	75	NA
PCB 198 (SIS)	54	74	62	NA	61	57	58	58	NA

TABLE F.7. (contd)

Analytical Replicate Results

Treatment	DUP		TRIP		COMP PC	DUP		TRIP	
	C-SB	C-SB	C-SB	C-SB		COMP PC	COMP PC	COMP PC	COMP PC
Replicate	1	1	1	1	5	5	5	5	
Batch	3	3	3	3	7	7	7	7	
Wet Wt	10.22	10.18	10.08	NA	16.10	16.99	17.88		
Units	ng/g	ng/g	ng/g	RSD%	ng/g	ng/g	ng/g	RSD%	
Heptachlor	0.36 U	0.36 U	0.37 U	NA	0.23 U	0.22 U	0.21 U	NA	
Aldrin	0.25 U	0.25 U	0.25 U	NA	1.14	1.12	1.05	4	
Heptachlor Epoxide	0.26 U	0.26 U	0.26 U	NA	0.16 U	0.16 U	0.15 U	NA	
2,4'-DDE	0.51 U	0.51 U	0.52 U	NA	0.32 U	0.31 U	0.29 U	NA	
Endosulfan I	0.35 U	0.35 U	0.36 U	NA	0.22 U	0.21 U	0.20 U	NA	
α -Chlordane	0.19 U	0.19 U	0.19 U	NA	3.54	3.06	2.78	12	
Trans Nonachlor	0.28 U	0.28 U	0.29 U	NA	0.61	0.39	0.32	34	
4,4'-DDE	0.81	0.37 U	0.37 U	NA	5.66	5.28	4.61	10	
Dieldrin	1.01 U	1.01 U	1.02 U	NA	3.96	3.79	3.43	7	
2,4'-DDD	0.50 U	0.50 U	0.50 U	NA	5.45	4.75	4.45	11	
2,4'-DDT	0.35 U	0.35 U	0.35 U	NA	0.22 U	0.21 U	0.20 U	NA	
4,4'-DDD	0.51 U	0.51 U	0.52 U	NA	11.4	10.6	9.14	11	
Endosulfan II	0.35 U	0.35 U	0.36 U	NA	0.22 U	0.21 U	0.20 U	NA	
4,4'-DDT	0.30 U	0.30 U	0.30 U	NA	0.19 U	0.18 U	0.17 U	NA	
Endosulfan Sulfate	0.35 U	0.35 U	0.36 U	NA	0.22 U	0.21 U	0.20 U	NA	
PCB 8	0.82	1.26	0.94	23	0.51 U	0.48 U	0.46 U	NA	
PCB 18	0.84 U	0.84 U	0.85 U	NA	0.53 U	0.90	0.48 U	NA	
PCB 28	0.40 U	0.40 U	0.40 U	NA	1.33	1.17	1.03	13	
PCB 52	0.70 U	0.70 U	0.71 U	NA	5.27	4.90	4.38	9	
PCB 49	0.46 U	0.46 U	0.47 U	NA	1.83	1.58	1.41	13	
PCB 44	0.32 U	0.32 U	0.33 U	NA	0.50	0.19 U	0.18 U	NA	
PCB 66	0.19 U	0.30	0.32	NA	0.12 U	0.11 U	0.11 U	NA	
PCB 101	0.29 U	0.29 U	0.29 U	NA	7.32	6.83	6.12	9	
PCB 87	0.31 U	0.31 U	0.32 U	NA	3.21	3.00	2.64	10	
PCB 118	0.58 U	0.58 U	0.58 U	NA	4.56	4.02	3.83	9	
PCB 184	0.46 U	0.46 U	0.47 U	NA	0.29 U	0.28 U	0.26 U	NA	
PCB 153	0.24 U	0.24 U	0.24 U	NA	2.53	2.19	2.04	11	
PCB 105	0.22 U	0.22 U	0.22 U	NA	2.11	1.72	1.60	15	
PCB 138	0.57 U	0.57 U	0.57 U	NA	3.19	2.82	2.59	11	
PCB 187	0.25 U	0.25 U	0.25 U	NA	0.63	0.50	0.51	13	
PCB 183	0.46 U	0.46 U	0.47 U	NA	0.31	0.28 U	0.26 U	NA	
PCB 128	0.30 U	0.30 U	0.31 U	NA	0.73	0.59	0.56	14	
PCB 180	0.36 U	0.36 U	0.37 U	NA	0.76	0.73	0.64	9	
PCB 170	0.33 U	0.34	0.33 U	NA	0.39	0.36	0.34	7	
PCB 195	0.20 U	0.20 U	0.20 U	NA	0.12 U	0.12 U	0.11 U	NA	
PCB 206	0.22 U	0.22 U	0.22 U	NA	0.18	0.18	0.15	10	
PCB 209	0.19 U	0.19 U	0.19 U	NA	0.12 U	0.11 U	0.11 U	NA	
<u>Surrogate Recoveries (%)</u>									
PCB 103 (SIS)	89	79	88	NA	95	95	86	NA	
PCB 198 (SIS)	144	125	141	NA	93	82	75	NA	

(a) U Undetected at or above given concentration.

(b) NA Not applicable.

(c) NS Not spiked.

(d) Outside quality control range (30-150%) for SIS.

(e) Outside quality control criteria (50-120%) for matrix spike recovery.

TABLE F.8. Polynuclear Aromatic Hydrocarbons (PAH) and 1,4-Dichlorobenzene (Wet Weight) in Tissue of *M. nasuta*

Treatment	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC	DUP COMP PC
Replicate	1	2	3	4	5-1	5-2
Batch	7	7	7	7	7	7
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.91%	13.51%	13.89%	14.98%	13.68%	13.68%
1,4-Dichlorobenzene	1.79 U ^(a)	1.86 U	1.86 U	1.86 U	2.31 U	2.20 U
Naphthalene	3.19 ^(b)	4.16	2.98 ^(b)	3.67	4.65	4.68
Acenaphthylene	0.70 U	0.77 ^(b)	0.73 U	0.73 U	0.93 ^(b)	0.86 U
Acenaphthene	14.3	21.9	22.4	11.4	20.2	18.4
Fluorene	5.12 ^(b)	6.39 ^(b)	6.66 ^(b)	5.31 ^(b)	6.90	6.56
Phenanthrene	23.9	31.3	38.3	23.5	34.0	30.5
Anthracene	27.2	37.2	47.1	29.1	36.7	34.0
Fluoranthene	495	661	779	561	627	587
Pyrene	364	494	574	422	453	425
Benzo(a)anthracene	80.6	109	119	85.3	106	96.8
Chrysene	96.0	125	138	104	122	112
Benzo(b)fluoranthene	69.4	69.4	94.7	71.5	69.3	81.1
Benzo(k)fluoranthene	1.60 U	17.7	1.67 U	1.67 U	17.6	1.97 U
Benzo(a)pyrene	25.6	32.5	34.1	26.1	32.8	30.5
Indeno(123-cd)pyrene	9.45	11.3	12.0	9.33	12.2	11.4
Dibenzo(a,h)anthracene	2.97	3.35	3.64	2.99	3.88	3.64
Benzo(g,h,i)perylene	9.36	11.1	11.6	9.55	12.1	11.4
<u>Surrogate Internal Standards (%)</u>						
d4 1,4-Dichlorobenzene	49	58	45	53	62	68
d8 Naphthalene	63	69	60	67	74	80
d10 Acenaphthene	73	79	76	79	88	91
d12 Chrysene	79	87	81	83	95	94
d14 Dibenzo(a,h,i)anthracene	96	107	100	101	118	114

TABLE F.8. (contd)

Treatment	TRIP					
	COMP PC	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	5-3	1	2	3	4	5
Batch	7	2	3	2	3	2
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.68%	14.08%	18.71%	13.02%	11.83%	20.96%
1,4-Dichlorobenzene	2.09 U	1.86 U	1.86 U	1.86 U	1.86 U	1.71 U
Naphthalene	4.39	1.86 U	1.86 U	1.86 U	1.86 U	1.87 ^(b)
Acenaphthylene	0.82 ^(b)	0.72 U	0.72 U	0.72 U	0.72 U	0.67 U
Acenaphthene	17.5	1.30 U	1.30 U	1.30 U	1.30 U	1.20 U
Fluorene	5.99	1.24 U	1.24 U	1.24 U	1.24 U	1.14 U
Phenanthrene	28.1	2.56 U	2.56 U	2.56 U	2.56 U	2.35 U
Anthracene	30.8	2.24 U	2.24 U	2.24 U	2.24 U	2.06 U
Fluoranthene	533	5.36 U	5.36 U	5.36 U	5.36 U	4.94 U
Pyrene	383	4.57 U	4.57 U	4.57 U	4.57 U	4.20 U
Benzo(a)anthracene	85.5	2.16 ^(b) B ^(c)	2.38 ^(b) B	2.73 ^(b) B	2.34 ^(b) B	2.20 ^(b) B
Chrysene	99.5	2.27 U	2.27 U	2.27 U	2.27 U	2.09 U
Benzo(b)fluoranthene	57.6	2.98 ^(b)	3.25 ^(b) B	4.14 ^(d)	2.95 ^(b) B	3.54
Benzo(k)fluoranthene	13.7	2.05 ^(b)	2.12 ^(b)	1.67 U	2.17 ^(b)	1.96
Benzo(a)pyrene	26.6	1.49 U	1.49 U	1.54 ^(b)	1.62 ^(b)	1.41
Indeno(123-cd)pyrene	10.1	1.76 U	1.76 U	1.76 U	1.76 U	1.62 U
Dibenzo(a,h)anthracene	3.25	1.26 U	1.26 U	1.26 U	1.26 U	1.16 U
Benzo(g,h,i)perylene	10.0	1.40 U	1.40 U	1.46 ^(b)	1.40 U	1.41 ^(b)
<u>Surrogate Internal Standards (%)</u>						
d4 1,4-Dichlorobenzene	50	58	51	55	43	60
d8 Naphthalene	63	66	60	65	51	71
d10 Acenaphthene	79	68	63	70	56	73
d12 Chrysene	83	73	61	72	61	73
d14 Dibenzo(a,h,i)anthracene	102	88	70	86	71	86

TABLE F.8. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	1	1	1	1	1
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	15.08%	14.45%	14.15%	14.06%	14.57%
1,4-Dichlorobenzene	1.86 U	1.86 U	1.86 U	1.86 U	1.86 U
Naphthalene	1.86 U	1.86 U	1.86 U	1.86 U	1.86 U
Acenaphthylene	1.19 ^(b)	0.955 ^(b)	0.877 ^(b)	0.845 ^(b)	1.08 ^(b)
Acenaphthene	1.30 U	1.30 U	1.30 U	1.30 U	1.30 U
Fluorene	1.24 U	1.24 U	1.24 U	1.24 U	1.24 U
Phenanthrene	3.32	4.53	2.56 U	3.66	3.67
Anthracene	3.13 ^(b)	3.28	2.83 ^(b)	3.05 ^(b)	2.95 ^(b)
Fluoranthene	9.13	11.2	7.20	9.82	8.54
Pyrene	10.4	14.2	9.46	12.2	11.8
Benzo(a)anthracene	5.66 B	5.93 B	4.25 B	5.52 B	4.78 B
Chrysene	5.50	5.92	3.87	5.75	4.91
Benzo(b)fluoranthene	13.2	14.6	11.0	14.0	13.3
Benzo(k)fluoranthene	5.91	5.91	4.97	5.94	5.49
Benzo(a)pyrene	6.41	6.96	4.88	6.48	5.17
Indeno(123-cd)pyrene	4.28	4.77	4.00	4.32	4.55
Dibenzo(a,h)anthracene	1.26 U	1.27	1.26 U	1.26 U	1.26 U
Benzo(g,h,i)perylene	4.39	4.97	3.88	4.35	4.49
<u>Surrogate Internal Standards (%)</u>					
d4 1,4-Dichlorobenzene	53	53	58	58	29 ^(e)
d8 Naphthalene	65	65	71	72	36
d10 Acenaphthene	65	66	71	73	41
d12 Chrysene	76	75	81	80	51
d14 Dibenzo(a,h,i)anthracene	92	92	101	103	63

TABLE F.8. (contd)

Treatment	DUP			TRIP		
	C-SB	C-SB	C-SB	C-SB	C-SB	C-SB
Replicate	1-1	1-2	1-3	2	3	4
Batch	3	3	3	2	3	2
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.86%	12.86%	12.86%	12.45%	13.90%	13.16%
1,4-Dichlorobenzene	3.65 U	3.65 U	3.69 U	1.86 U	1.86 U	1.86 U
Naphthalene	3.65 U	3.65 U	3.69 U	1.86 U	1.86 U	1.86 U
Acenaphthylene	1.42 U	1.42 U	1.44 U	0.72 U	0.72 U	0.72 U
Acenaphthene	2.56 U	2.56 U	2.58 U	1.30 U	1.30 U	1.30 U
Fluorene	2.42 U	2.42 U	2.45 U	1.24 U	1.24 U	1.24 U
Phenanthrene	5.02 U	5.02 U	5.07 U	2.56 U	2.56 U	2.56 U
Anthracene	4.39 U	4.39 U	4.43 U	2.24 U	2.74 ^(b)	2.24 U
Fluoranthene	10.5 U	10.5 U	10.6 U	5.36 U	5.76	5.92
Pyrene	8.95 U	8.95 U	9.05 U	4.57 U	4.57 U	4.57 U
Benzo(a)anthracene	4.54 ^{(b)B}	4.95 ^{*B}	4.65 ^{(b)B}	2.52 ^{(b)B}	2.57 ^{(b)B}	2.46 ^{(b)B}
Chrysene	4.45 U	4.45 U	4.49 U	2.27 U	2.27 U	2.27 U
Benzo(b)fluoranthene	6.41 ^{(b)B}	5.72 ^{(b)B}	6.18 ^{(b)B}	3.54	4.11 ^{(b)B}	4.35 ^(d)
Benzo(k)fluoranthene	3.27 U	3.93 ^(b)	3.31 U	2.09 ^(b)	1.67 U	1.67 U
Benzo(a)pyrene	2.92 U	2.93 U	2.96 U	1.49 U	1.49 U	1.49 U
Indeno(123-cd)pyrene	3.45 U	3.45 U	3.49 U	1.76 U	1.76 U	1.76 U
Dibenzo(a,h)anthracene	2.47 U	2.47 U	2.50 U	1.26 U	1.26 U	1.26 U
Benzo(g,h,i)perylene	2.75 U	2.75 U	2.78 U	1.40 U	1.40 U	1.48

Surrogate Internal Standards (%)

d4 1,4-Dichlorobenzene	54	57	59	57	65	53
d8 Naphthalene	64	65	71	62	74	65
d10 Acenaphthene	67	66	76	64	73	69
d12 Chrysene	80	75	87	65	78	75
d14 Dibenzo(a,h,i)anthracene	83	77	91	76	89	87

TABLE F.8. (contd)

Treatment	C-SB	DUP C-SB	TRIP C-SB	M. nasuta Background	M. nasuta Background	M. nasuta Background
Replicate	5-1	5-2	5-3	1	2	3
Batch	2	2	2	7	7	7
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.21%	13.21%	13.21%	15.16%	14.86%	14.87%
1,4-Dichlorobenzene	3.65 U	3.69 U	3.62 U	1.83 U	1.86 U	1.86 U
Naphthalene	3.65 U	3.69 U	3.62 U	2.31	2.51	3.18 ^(b)
Acenaphthylene	1.42 U	1.44 U	1.41 U	0.71 U	0.73 U	0.73 U
Acenaphthene	2.56 U	2.58 U	2.53 U	1.28 U	1.3 U	1.3 U
Fluorene	2.42 U	2.45 U	2.40 U	1.21 U	2.82 ^(b)	2.86 ^(b)
Phenanthrene	5.02 U	5.07 U	4.96 U	5.25	3.74	3.96
Anthracene	4.39 U	4.43 U	4.34 U	2.19 U	2.24 U	2.24 U
Fluoranthene	10.5 U	10.6 U	10.4 U	6.49 ^(b)	7.05 ^(b)	7.42 ^(b)
Pyrene	8.95 U	9.05 U	8.86 U	4.61 ^(b)	5.10	5.49
Benzo(a)anthracene	4.73	4.80 ^{(b)B}	4.53 ^{(b)B}	4.00 ^(b)	4.04 ^(b)	4.06 ^(b)
Chrysene	4.45 U	4.49 U	4.40 U	2.22 U	2.27 U	2.27 U
Benzo(b)fluoranthene	5.67	5.81 ^(b)	6.38	4.90	4.67 ^(b)	4.97 ^(b)
Benzo(k)fluoranthene	3.98	4.08 ^(b)	3.24 U	2.51 ^(b)	2.65 ^(b)	2.62 ^(b)
Benzo(a)pyrene	4.70	2.96 U	2.90 U	2.85 ^(b)	2.26 ^(b)	2.64 ^(b)
Indeno(123-cd)pyrene	3.45 U	3.49 U	3.42 U	3.31 ^(b)	3.48 ^(b)	3.44 ^(b)
Dibenzo(a,h)anthracene	2.47 U	2.50 U	2.45 U	1.24 U	1.26 U	1.26 U
Benzo(g,h,i)perylene	2.75 U	2.78 U	2.72 U	3.12 ^(b)	1.4 U	1.4 U

Surrogate Internal Standards (%)

d4 1,4-Dichlorobenzene	58	59	53	11 ^(e)	45	31
d8 Naphthalene	67	67	61	18 ^(e)	59	44
d10 Acenaphthene	68	66	62	27 ^(e)	76	66
d12 Chrysene	68	63	63	70	75	75
d14 Dibenzo(a,h,i)anthracene	79	71	74	88	71	92

(a) U Undetected at or above given concentration.

(b) Ion ratio out or confirmation ion not detected.

(c) B Value is < 5 times concentration in blank.

(d) Benzo(b)fluoranthene is the sum of benzo(b)fluoranthene and benzo(k)fluoranthene.

Benzo(k)fluoranthene is present but could not be quantified due to poor resolution.

(e) Outside quality control criteria (30-150%) for SIS.

TABLE F.9. Polynuclear Aromatic Hydrocarbons (PAH) and 1,4-Dichlorobenzene (Dry Weight) in Tissue of *M. nasuta*

Treatment	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC
Replicate	1	2	3	4	5-1	5-2
Batch	7	7	7	7	7	7
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.91%	13.51%	13.89%	14.98%	13.68%	13.68%
1,4-Dichlorobenzene	13.9 U ^(a)	13.8 U	13.4 U	12.4 U	16.9 U	16.1 U
Naphthalene	24.7 ^(b)	30.8	21.5 ^(b)	24.5	34.0	34.2
Acenaphthylene	5.4 U	5.7 ^(b)	5.3 U	4.9 U	6.8 ^(b)	6.3 U
Acenaphthene	111	162	161	76.1	148	135
Fluorene	39.7 ^(b)	47.3 ^(b)	47.9 ^(b)	35.4 ^(b)	50.4	48.0
Phenanthrene	185	232	276	157	249	223
Anthracene	211	275	339	194	268	249
Fluoranthene	3830	4890	5610	3750	4580	4290
Pyrene	2820	3660	4130	2820	3310	3110
Benzo(a)anthracene	624	807	857	569	775	708
Chrysene	744	925	994	694	892	819
Benzo(b)fluoranthene	538	514	682	477	507	593
Benzo(k)fluoranthene	12.4 U	131	12.0 U	11.1 U	129	14.4 U
Benzo(a)pyrene	198	241	246	174	240	223
Indeno(123-cd)pyrene	73.2	83.6	86.4	62.3	89.2	83.3
Dibenzo(a,h)anthracene	23.0	24.8	26.2	20.0	28.4	26.6
Benzo(g,h,i)perylene	72.5	82.2	83.5	63.8	88.5	83.3

TABLE F.9. (contd)

Treatment	COMP PC	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	5-3	1	2	3	4	5
Batch	7	2	3	2	3	2
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.68%	14.08%	18.71%	13.02%	11.83%	20.96%
1,4-Dichlorobenzene	15.3 U	13.2 U	9.94 U	14.3 U	15.7 U	8.16 U
Naphthalene	32.1	13.2 U	9.94 U	14.3 U	15.7 U	8.92 ^(b)
Acenaphthylene	6.0 ^(b)	5.1 U	3.8 U	5.5 U	6.1 U	3.2 U
Acenaphthene	128	9.23 U	6.95 U	9.98 U	11.0 U	5.73 U
Fluorene	43.8	8.81 U	6.63 U	9.52 U	10.5 U	5.44 U
Phenanthrene	205	18.2 U	13.7 U	19.7 U	21.6 U	11.2 U
Anthracene	225	15.9 U	12.0 U	17.2 U	18.9 U	9.83 U
Fluoranthene	3900	38.1 U	28.6 U	41.2 U	45.3 U	23.6 U
Pyrene	2800	32.5 U	24.4 U	35.1 U	38.6 U	20.0 U
Benzo(a)anthracene	625	15.3 ^(b) B ^(c)	12.7 ^(b) B	21.0 ^(b) B	19.8 ^(b) B	10.5 ^(b) B
Chrysene	727	16.1 U	12.1 U	17.4 U	19.2 U	9.97 U
Benzo(b)fluoranthene	421	21.2 ^(b)	17.4 ^(b) B	31.8 ^(d)	24.9 ^(b) B	16.9
Benzo(k)fluoranthene	100	14.6 ^(b)	11.3 ^(b)	12.8 U	18.3 ^(b)	9.35
Benzo(a)pyrene	194	10.6 U	7.96 U	11.8 ^(b)	13.7 ^(b)	6.73
Indeno(123-cd)pyrene	73.8	12.5 U	9.41 U	13.5 U	14.9 U	7.73 U
Dibenzo(a,h)anthracene	23.8	8.95 U	6.73 U	9.68 U	10.7 U	5.53 U
Benzo(g,h,i)perylene	73.1	9.94 U	7.48 U	11.2 ^(b)	11.8 U	6.73 ^(b)

TABLE F.9. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	1	1	1	1	1
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	15.08%	14.45%	14.15%	14.06%	14.57%
1,4-Dichlorobenzene	12.3 U	12.9 U	13.1 U	13.2 U	12.8 U
Naphthalene	12.3 U	12.9 U	13.1 U	13.2 U	12.8 U
Acenaphthylene	7.89 ^(b)	6.61 ^(b)	6.20 ^(b)	6.01 ^(b)	7.41 ^(b)
Acenaphthene	8.62 U	9.00 U	9.19 U	9.25 U	8.92 U
Fluorene	8.22 U	8.58 U	8.76 U	8.82 U	8.51 U
Phenanthrene	22.0	31.3	18.1 U	26.0	25.2
Anthracene	20.8 ^(b)	22.7	20.0 ^(b)	21.7 ^(b)	20.2 ^(b)
Fluoranthene	60.5	77.5	50.9	69.8	58.6
Pyrene	69.0	98.3	66.9	86.8	81.0
Benzo(a)anthracene	37.5 B	41.0 B	30.0 B	39.3 B	32.8 B
Chrysene	36.5	41.0	27.3	40.9	33.7
Benzo(b)fluoranthene	87.5	101	77.7	99.6	91.3
Benzo(k)fluoranthene	39.2	40.9	35.1	42.2	37.7
Benzo(a)pyrene	42.5	48.2	34.5	46.1	35.5
Indeno(123-cd)pyrene	28.4	33.0	28.3	30.7	31.2
Dibenzo(a,h)anthracene	8.36 U	8.79	8.90 U	8.96 U	8.65 U
Benzo(g,h,i)perylene	29.1	34.4	27.4	30.9	30.8

TABLE F.9. (contd)

Treatment	Control-SB	Control-SB	Control-SB	Control-SB	Control-SB	Control-SB
Replicate	1-1	1-2	1-3	2	3	4
Batch	3	3	3	2	3	2
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	12.86%	12.86%	12.86%	12.45%	13.90%	13.16%
1,4-Dichlorobenzene	28.4 U	28.4 U	28.7 U	14.9 U	13.4 U	14.1 U
Naphthalene	28.4 U	28.4 U	28.7 U	14.9 U	13.4 U	14.1 U
Acenaphthylene	11.0 U	11.0 U	11.2 U	5.78 U	5.18 U	5.47 U
Acenaphthene	19.9 U	19.9 U	20.1 U	10.4 U	9.35 U	9.88 U
Fluorene	18.8 U	18.8 U	19.1 U	9.96 U	8.92 U	9.42 U
Phenanthrene	39.0 U	39.0 U	39.4 U	20.6 U	18.4 U	19.5 U
Anthracene	34.1 U	34.1 U	34.4 U	18.0 U	19.7 (b)	17.0 U
Fluoranthene	81.6 U	81.6 U	82.4 U	43.1 U	41.4	45.0
Pyrene	69.6 U	69.6 U	70.4 U	36.7 U	32.9 U	34.7 U
Benzo(a)anthracene	35.3 ^(b) B	38.5 ^(b) B	36.2 ^(b) B	20.2 ^(b) B	18.5 ^(b) B	18.7 ^(b) B
Chrysene	34.6 U	34.6 U	34.9 U	18.2 U	16.3 U	17.2 U
Benzo(b)fluoranthene	49.8 ^(b) B	44.5 ^(b) B	48.1 ^(b) B	28.4	29.6 ^(b) B	33.1 ^(d)
Benzo(k)fluoranthene	25.4 U	30.6 (b)	25.7 U	16.8 (b)	12.0 U	12.7 U
Benzo(a)pyrene	22.7 U	22.8 U	23.0 U	12.0 U	10.7 U	11.3 U
Indeno(123-cd)pyrene	26.8 U	26.8 U	27.1 U	14.1 U	12.7 U	13.4 U
Dibenzo(a,h)anthracene	19.2 U	19.2 U	19.4 U	10.1 U	9.06 U	9.57 U
Benzo(g,h,i)perylene	21.4 U	21.4 U	21.6 U	11.2 U	10.1 U	11.2

TABLE F.9. (contd)

Treatment	Control-SB	Control-SB	Control-SB	<i>M. nasuta</i>	<i>M. nasuta</i>	<i>M. nasuta</i>
	Replicate	Replicate	Replicate	Background	Background	Background
Batch	5-1	5-2	5-3	1	2	3
Units	2	2	2	7	7	7
Percent Dry Weight	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g
	13.21%	13.21%	13.21%	15.16%	14.86%	14.87%
1,4-Dichlorobenzene	27.6 U	27.9 U	27.4 U	12.1 U	12.5 U	12.5 U
Naphthalene	27.6 U	27.9 U	27.4 U	15.2	16.9	21.4 ^(b)
Acenaphthylene	10.7 U	10.9 U	10.7 U	4.68 U	4.91 U	4.91 U
Acenaphthene	19.4 U	19.5 U	19.2 U	8.44 U	8.75 U	8.74 U
Fluorene	18.3 U	18.5 U	18.2 U	7.98 U	19.0 ^(b)	19.2 ^(b)
Phenanthrene	38.0 U	38.4 U	37.5 U	34.6	25.2	26.6
Anthracene	33.2 U	33.5 U	32.9 U	14.4 U	15.1 U	15.1 U
Fluoranthene	79.5 U	80.2 U	78.7 U	42.8 ^(b)	47.4 ^(b)	49.9 ^(b)
Pyrene	67.8 U	68.5 U	67.1 U	30.4 ^(b)	34.3	36.9
Benzo(a)anthracene	35.8	36.3 ^(b) B	34.3 ^(b) B	26.4 ^(b)	27.2 ^(b)	27.3 ^(b)
Chrysene	33.7 U	34.0 U	33.3 U	14.6 U	15.3 U	15.3 U
Benzo(b)fluoranthene	42.9	44.0 ^(b)	48.3	32.3	31.4 ^(b)	33.4 ^(b)
Benzo(k)fluoranthene	30.1	30.9 ^(b)	24.5 U	16.6 ^(b)	17.8 ^(b)	17.6 ^(b)
Benzo(a)pyrene	35.6	22.4 U	22.0 U	18.8 ^(b)	15.2 ^(b)	17.8 ^(b)
Indeno(123-cd)pyrene	26.1 U	26.4 U	25.9 U	21.8 ^(b)	23.4 ^(b)	23.1 ^(b)
Dibenzo(a,h)anthracene	18.7 U	18.9 U	18.5 U	8.18 U	8.48 U	8.47 U
Benzo(g,h,i)perylene	20.8 U	21.0 U	20.6 U	20.6 ^(b)	9.4 U	9.41 U

(a) U Undetected at or above given concentration.

(b) Ion ratio out or confirmation ion not detected.

(c) B Value is < 5 times concentration in blank.

(d) Benzo(b)fluoranthene is the sum of benzo(b)fluoranthene and benzo(k)fluoranthene.

Benzo(K)fluoranthene is present but could not be quantified due to poor resolution.

TABLE F.10. Quality Control Summary for Polynuclear Aromatic Hydrocarbons (PAHs) and 1,4-Dichlorobenzene in Tissue of *M. nasuta* (Wet Weight)

Matrix Spike Results

Treatment Replicate Batch Wet Weight Units	COMP PC 1 7 20.84 ng/g	Matrix Spike		Amount Spiked ng/g	Percent Recovery
		COMP PC 1 7 20.18 ng/g	COMP PC(MS)		
1,4-Dichlorobenzene	1.79 U ^(a)	22.3	24.8	90	
Naphthalene	3.19 ^(b)	30.6	24.8	111	
Acenaphthylene	0.70 U	26.0	24.8	105	
Acenaphthene	14.3	44.1	24.8	120	
Fluorene	5.12 ^(b)	32.5	24.8	110	
Phenanthrene	23.9	54.5	24.8	123 ^(c)	
Anthracene	27.2	62.2	24.8	141 ^(c)	
Fluoranthene	495	555	24.8	242 ^(c)	
Pyrene	364	414	24.8	202 ^(c)	
Benzo(a)anthracene	80.6	118	24.8	151 ^(c)	
Chrysene	96.0	128	24.8	129 ^(c)	
Benzo(b)fluoranthene	69.4	83.3	24.8	56	
Benzo(k)fluoranthene	1.60 U	47.1	24.8	190 ^(c)	
Benzo(a)pyrene	25.6	55.7	24.8	121 ^(c)	
Indeno(123-cd)pyrene	9.45	34.9	24.8	103	
Dibenzo(a,h)anthracene	2.97	30.9	24.8	113	
Benzo(g,h,i)perylene	9.36	33.5	24.8	97	
Surrogate Internal Standards (%)					
d4 1,4-Dichlorobenzene	49	57	NA ^(d)	NA	
d8 Naphthalene	63	67	NA	NA	
d10 Acenaphthene	73	74	NA	NA	
d12 Chrysene	79	76	NA	NA	
d14 Dibenzo(a,h,i)anthracene	96	93	NA	NA	

TABLE F.10. (contd)

<u>Matrix Spike Results</u>				
Treatment	COMP HU-A	<i>Matrix Spike</i>		
Replicate	1	COMP HU-A(MS)		
Batch	1	1		Amount
Wet Weight	20.12	20.12	Spiked	Percent
Units	ng/g	ng/g	ng/g	Recovery
1,4-Dichlorobenzene	1.86 U	37.1	37.8	98
Naphthalene	3.34	25.8	24.9	90
Acenaphthylene	2.20 ^(b)	24.4	24.9	89
Acenaphthene	7.45	31.8	24.9	98
Fluorene	8.07	31.9	24.9	96
Phenanthrene	90.2	112	24.9	92
Anthracene	42.8	68.2	24.9	102
Fluoranthene	232	251	24.9	76
Pyrene	278	291	24.9	52
Benzo(a)anthracene	144	167	24.9	92
Chrysene	155	173	24.9	72
Benzo(b)fluoranthene	86.6	110	24.9	94
Benzo(k)fluoranthene	24.1	49.8	24.9	103
Benzo(a)pyrene	69.7	94.1	24.9	98
Indeno(123-cd)pyrene	13.9	34.2	24.9	82
Dibenzo(a,h)anthracene	4.22	25.5	24.9	85
Benzo(g,h,i)perylene	14.4	34.8	24.9	82
<u>Surrogate Internal Standards (%)</u>				
d4 1,4-Dichlorobenzene	43	53	NA	NA
d8 Naphthalene	53	65	NA	NA
d10 Acenaphthene	62	69	NA	NA
d12 Chrysene	76	84	NA	NA
d14 Dibenzo(a,h,i)anthracene	84	95	NA	NA

TABLE F.10. (contd)

Analytical Replicate Results

	COMP PC	<i>Dup</i> COMP PC	<i>Trip</i> COMP PC	
Treatment	5-1	5-2	5-3	
Replicate	7	7	7	
Batch	16.10	16.99	17.88	
Wet Weight	ng/g	ng/g	ng/g	RSD%
Units				
1,4-Dichlorobenzene	2.31 U	2.20 U	2.09 U	NA
Naphthalene	4.65	4.68	4.39	3
Acenaphthylene	0.93 ^(b)	0.86 U	0.82 ^(b)	NA
Acenaphthene	20.2	18.4	17.5	7
Fluorene	6.90	6.56	5.99	7
Phenanthrene	34.0	30.5	28.1	10
Anthracene	36.7	34.0	30.8	9
Fluoranthene	627	587	533	8
Pyrene	453	425	383	8
Benzo(a)anthracene	106	96.8	85.5	11
Chrysene	122	112	99.5	10
Benzo(b)fluoranthene	69.3	81.1	57.6	17
Benzo(k)fluoranthene	17.6	1.97 U	13.7	NA
Benzo(a)pyrene	32.8	30.5	26.6	10
Indeno(123-cd)pyrene	12.2	11.4	10.1	9
Dibenzo(a,h)anthracene	3.88	3.64	3.25	9
Benzo(g,h,i)perylene	12.1	11.4	10.0	10
<u>Surrogate Internal Standards (%)</u>				
d4 1,4-Dichlorobenzene	62	68	50	NA
d8 Naphthalene	74	80	63	NA
d10 Acenaphthene	88	91	79	NA
d12 Chrysene	95	94	83	NA
d14 Dibenzo(a,h,i)anthracene	118	114	102	NA

TABLE F.10. (contd)

Analytical Replicate Results

Treatment Replicate Batch	COMP EC-B 5-1 1	<i>Dup</i> COMP EC-B 5-2 1	<i>Trip</i> COMP EC-B 5-3 1	RSD%
1,4-Dichlorobenzene	3.73 U	3.73 U	3.73 U	NA
Naphthalene	5.99	4.80	5.64	11
Acenaphthylene	3.26 ^(b)	3.21 ^(b)	3.24 ^(b)	1
Acenaphthene	40.0	41.5	41.8	2
Fluorene	25.8	26.2	25.9	1
Phenanthrene	210	213	213	1
Anthracene	103	106	106	2
Fluoranthene	453	464	475	2
Pyrene	466	476	484	2
Benzo(a)anthracene	183	188	190	2
Chrysene	226	233	234	2
Benzo(b)fluoranthene	139	139	146	3
Benzo(k)fluoranthene	31.7	34.1	32.7	4
Benzo(a)pyrene	88.9	91.4	94.4	3
Indeno(123-cd)pyrene	22.2	22.3	22.9	2
Dibenzo(a,h)anthracene	4.77	5.06	5.17	4
Benzo(g,h,i)perylene	24.1	24.4	25.0	2
<u>Surrogate Internal Standards (%)</u>				
d4 1,4-Dichlorobenzene	44	52	53	NA
d8 Naphthalene	54	65	64	NA
d10 Acenaphthene	58	74	70	NA
d12 Chrysene	69	89	78	NA
d14 Dibenzo(a,h,i)anthracene	79	102	89	NA

TABLE F.10. (contd)

<u>Analytical Replicate Results</u>				
Treatment	C-SB	<i>Dup</i> C-SB	<i>Trip</i> C-SB	
Replicate	5-1	5-2	5-3	
Batch	2	2	2	
Wet Weight	10.16	10.14	10.34	
Units	ng/g	ng/g	ng/g	RSD%
1,4-Dichlorobenzene	3.65 U	3.69 U	3.62 U	NA
Naphthalene	3.65 U	3.69 U	3.62 U	NA
Acenaphthylene	1.42 U	1.44 U	1.41 U	NA
Acenaphthene	2.56 U	2.58 U	2.53 U	NA
Fluorene	2.42 U	2.45 U	2.40 U	NA
Phenanthrene	5.02 U	5.07 U	4.96 U	NA
Anthracene	4.39 U	4.43 U	4.34 U	NA
Fluoranthene	10.5 U	10.6 U	10.4 U	NA
Pyrene	8.95 U	9.05 U	8.86 U	NA
Benzo(a)anthracene	4.73	4.80 ^(b) B ^(e)	4.53 ^(b) B	3
Chrysene	4.45 U	4.49 U	4.40 U	NA
Benzo(b)fluoranthene	5.67	5.81 ^(b)	6.38	7
Benzo(k)fluoranthene	3.98	4.08 ^(b)	3.24 U	NA
Benzo(a)pyrene	4.70	2.96 U	2.90 U	NA
Indeno(123-cd)pyrene	3.45 U	3.49 U	3.42 U	NA
Dibenzo(a,h)anthracene	2.47 U	2.50 U	2.45 U	NA
Benzo(g,h,i)perylene	2.75 U	2.78 U	2.72 U	NA
<u>Surrogate Internal Standards (%)</u>				
d4 1,4-Dichlorobenzene	58	59	53	NA
d8 Naphthalene	67	67	61	NA
d10 Acenaphthene	68	66	62	NA
d12 Chrysene	68	63	63	NA
d14 Dibenzo(a,h,i)anthracene	79	71	74	NA

TABLE F.10. (contd)

<u>Analytical Replicate Results</u>				
Treatment	C-SB	<i>Dup</i> C-SB	<i>Trip</i> C-SB	
Replicate	1-1	1-2	1-3	
Batch	3	3	3	
Wet Weight	10.22	10.18	10.08	
Units	ng/g	ng/g	ng/g	RSD%
1,4-Dichlorobenzene	3.65 U	3.65 U	3.69 U	NA
Naphthalene	3.65 U	3.65 U	3.69 U	NA
Acenaphthylene	1.42 U	1.42 U	1.44 U	NA
Acenaphthene	2.56 U	2.56 U	2.58 U	NA
Fluorene	2.42 U	2.42 U	2.45 U	NA
Phenanthrene	5.02 U	5.02 U	5.07 U	NA
Anthracene	4.39 U	4.39 U	4.43 U	NA
Fluoranthene	10.5 U	10.5 U	10.6 U	NA
Pyrene	8.95 U	8.95 U	9.05 U	NA
Benzo(a)anthracene	4.54 ^(b) B	4.95 ^(b) B	4.65 ^(b) B	5
Chrysene	4.45 U	4.45 U	4.49 U	NA
Benzo(b)fluoranthene	6.41 ^(b) B	5.72 ^(b) B	6.18 ^(b) B	6
Benzo(k)fluoranthene	3.27 U	3.93 ^(b)	3.31 U	NA
Benzo(a)pyrene	2.92 U	2.93 U	2.96 U	NA
Indeno(123-cd)pyrene	3.45 U	3.45 U	3.49 U	NA
Dibenzo(a,h)anthracene	2.47 U	2.47 U	2.50 U	NA
Benzo(g,h,i)perylene	2.75 U	2.75 U	2.78 U	NA
<u>Surrogate Internal Standards (%)</u>				
d4 1,4-Dichlorobenzene	54	57	59	NA
d8 Naphthalene	64	65	71	NA
d10 Acenaphthene	67	66	76	NA
d12 Chrysene	80	75	87	NA
d14 Dibenzo(a,h,i)anthracene	83	77	91	NA

(a) U Undetected at or above given concentration.

(b) Ion ratio out or confirmation ion not detected.

(c) Outside quality control range (50-120%) for matrix spike recovery.

(d) NA Not applicable.

(e) B Value is less than 5 times concentration in associated blank.

TABLE F.11. Lipids in Tissue of *M. nasuta*

Sediment Treatment	Replicate	Sample Weight	% Dry Weight	% Lipids (wet weight)	% Lipids (dry weight)
<i>Macoma</i> Background	1	5.18	15.16	0.58	3.83
<i>Macoma</i> Background	2	5.07	14.86	0.59	3.97
<i>Macoma</i> Background	3	5.04	14.87	0.60	4.03

Appendix G

***Nereis virens* Tissue Chemical Analyses and
Quality Assurance/Quality Control Data,
Port Chester Project**

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Metals
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Worm and Clam Tissue

QA/QC DATA QUALITY OBJECTIVES

	<u>Reference Method</u>	<u>Range of Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (µg/g dry wt)</u>
Arsenic	ICP/MS	75-125%	≤20%	≤20%	1.0
Cadmium	ICP/MS	75-125%	≤20%	≤20%	0.1
Chromium	ICP/MS	75-125%	≤20%	≤20%	0.2
Copper	ICP/MS	75-125%	≤20%	≤20%	1.0
Lead	ICP/MS	75-125%	≤20%	≤20%	0.1
Mercury	CVAA	75-125%	≤20%	≤20%	0.02
Nickel	ICP/MS	75-125%	≤20%	≤20%	0.1
Silver	ICP/MS	75-125%	≤20%	≤20%	0.1
Zinc	ICP/MS	75-125%	≤20%	≤20%	1.0

METHOD

A total of nine (9) metals was analyzed for the New York Federal Projects-2 Program: silver (Ag), arsenic (As), cadmium (Cd), chromium (Cr), copper (Cu), mercury (Hg), nickel (Ni), lead (Pb) and zinc (Zn). Hg was analyzed using cold-vapor atomic absorption spectroscopy (CVAA) according to the method of Bloom and Creelius (1983). The remaining metals were analyzed by inductively coupled plasma mass spectrometry (ICP/MS) following a procedure based on EPA Method 200.8 (EPA 1991).

To prepare tissue for analysis, samples were freeze-dried and blended in a Spex mixer-mill. Approximately 5 g of mixed sample was ground in a ceramic ball mill. For ICP/MS and CVAA analyses, 0.2- to 0.5-g aliquots of dried homogenous sample were digested using a mixture of nitric acid and hydrogen peroxide following EPA Method 200.3 (EPA 1991).

HOLDING TIMES

A total of 68 worm and 68 clam samples was received on 6/15/94 in good condition. Samples were logged into Battelle's log-in system, frozen to -80°C and subsequently freeze dried within approximately 7 days of sample receipt. Samples were analyzed within 180 days of collection. Worms and clams were digested in two separate batches. The following table summarizes the analysis dates:

<u>Task</u>	<u>Clams</u>	<u>Worms</u>
Sample Digestion	8/9/94	9/9/94
ICP-MS	9/15/94	10/6/94
CVAA-Hg	8/17-8/24/94	8/17-8/24/94

QA/QC SUMMARY METALS (continued)

- DETECTION LIMITS** Four aliquots of a background clam tissue were analyzed as four separate replicates. The standard deviation of these results were multiplied by 4.541 to determine a method detection limits (MDL). Target detection limits were exceeded for all metals except Ag, Cd and Hg.
- METHOD BLANKS** One procedural blank was analyzed per 20 samples. No metals were detected in the blanks above the MDLs.
- MATRIX SPIKES** One sample was spiked with all metals at a frequency of 1 per 20 samples. All recoveries were within the QC limits of 75% -125% with the exception of Ag in one spiked worm sample and Zn in three of the four spiked worm samples. Zn was spiked at a level near the level found in the native samples and, in one case, Zn was spiked at a level below that detected in the native sample and no recovery was calculated.
- REPLICATES** One sample was analyzed in triplicate at a frequency of 1 per 20 samples. Precision for triplicate analyses is reported by calculating the relative standard deviation (RSD) between the replicate results. Only the RSDs for Zn in one of the four replicated worm analyses exceeded the QC limits of $\pm 20\%$. RSDs for the rest of the metals were within the QC limits.
- SRMs** Standard Reference Material (SRM), 1566a (Oyster tissue from the National Institute of Standards and Technology, NIST), was analyzed for all metals. Results for all metals were within $\pm 20\%$ of mean certified value with the exception of Cr and Ni. Cr values were below the lower QC limit in two of the five SRMs analyzed with the clams and for three of the four SRMs analyzed with the worms. The SRM certified value for Cr ($1.43 \mu\text{g/g}$) is close to the detection limit ($1.46 \mu\text{g/g}$). Ni was also recovered below or above the control limits in some samples.

REFERENCES

Bloom, N. S., and E.A. Crecelius. 1983. "Determination of Mercury in Seawater at Sub-Nanogram per Liter Levels." *Mar. Chem.* 14:49-59.

EPA (U.S. Environmental Protection Agency). 1991 Methods for the Determination of Metals in Environmental Samples. EPA-600/4-91-010. Environmental Services Division, Monitoring Management Branch, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Chlorinated Pesticides/PCB Congeners
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Worm and Clam Tissue

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>Surrogate Recovery</u>	<u>Spike Recovery</u>	<u>Relative Precision</u>	<u>Detection Limit</u>
GC/ECD	30-150%	50-120%	≤30%	0.4 ng/g wet wt.

SAMPLE CUSTODY A total of 68 worm and 68 clam samples was received on 6/15/94 in good condition. Samples were logged into Battelle's log-in system and stored frozen until extraction.

METHOD Tissues were homogenized wet using a stainless steel blade. An aliquot of tissue sample was extracted with methylene chloride using the roller technique under ambient conditions following a procedure which is based on methods used by the National Oceanic and Atmospheric Administration for its Status and Trends Program (Krahn et al. 1988). Samples were then cleaned using silica/alumina (5% deactivated) chromatography followed by HPLC cleanup (Krahn et al. 1988). Extracts were analyzed for 15 chlorinated pesticides and 22 PCB congeners using gas chromatography/electron capture detection (GC/ECD) following a procedure based on EPA Method 8080 (EPA 1986). The column used was a J&W DB-17 and the confirmatory column was a DB-1701, both capillary columns (30m x 0.25mm I.D.). All detections were quantitatively confirmed on the second column.

HOLDING TIMES Samples were extracted in seven batches. All extracts were analyzed by GC/ECD. The following summarizes the extraction and analysis dates:

<u>Batch</u>	<u>Species</u>	<u>Extraction</u>	<u>Analysis</u>
1	<i>M. nasuta</i>	7/28/94	9/9-9/12/94
2	<i>M. nasuta</i>	8/3/94	9/13-9/15/94
3	<i>M. nasuta</i>	8/17/94	9/23-9/25/94
4	<i>N. virens</i>	8/19/95	9/26-9/30/94
5	<i>N. virens</i>	8/26/94	9/8-9/11/94
6	<i>N. virens</i>	9/6/94	9/17-9/19/94
7	<i>M. nasuta/N. virens</i>	9/26/94	9/15-9/17-94
8	<i>M. nasuta</i> MDL study	10/10/94	10/25/94

DETECTION LIMITS Target detection limits of 0.4 ng/g wet weight were met for all pesticides and PCB congeners, with the exception of dieldrin, PCB 8 and PCB 18, and for the samples that were analyzed in triplicate. These elevated detection limits for the replicates were due to the limited amount of tissue available resulting in smaller aliquots used for extraction. Method detection limits (MDLs) reported were determined by multiplying the

QA/QC SUMMARY/PCBs and PESTICIDES (continued)

standard deviation of seven spiked replicates of clam tissue by the Student's t value (99 percentile). Actual pesticide MDLs ranged from approximately 0.1 to 1.1 ng/g wet weight and PCB congener MDLs ranged from approximately 0.1 to 0.9 ng/g wet weight, depending on the compound and the sample weight extracted. MDLs were reported corrected for individual sample wet weight extracted.

Method detection limit verification was performed by analyzing four replicates of a spiked clam sample and multiplying the standard deviation of the results by 3.5. All detection limits calculated in this way were below the target detection limit of 0.4 ng/g wet weight with the exception of 4,4'-DDD which had a DL of 0.467 ng/g.

METHOD BLANKS

One method blank was extracted with each extraction batch. No pesticides or PCBs were detected in any of the method blanks.

SURROGATES

Two compounds, PCB congeners 103 and 198, were added to all samples prior to extraction to assess the efficiency of the analysis. Sample surrogate recoveries were all within the QC guidelines of 30% - 150%, with the exception of one sample in Batch 3 and two samples in Batch 4. All of these incidents involved a high recovery of PCB 198. This was most likely due to matrix interferences with the internal Standard octachloronaphthalene (OCN) which is used to quantify the recovery of surrogate PCB 198. Since no sample data are corrected for the OCN, sample results should not be affected. One sample had low surrogate recoveries for both PCB 103 and 198. This sample was re-extracted once due to surrogate recoveries. Since the recoveries in the reextraction also exceeded control limits, the problem was determined to be matrix interferences and no additional extractions were performed. Sample results were quantified using the surrogate internal standard method.

MATRIX SPIKES

Ten out of the 15 pesticides and 5 of the 22 PCB congeners analyzed were spiked into one sample per extraction batch. Matrix spike recoveries were within the control limit range of 50-120% for all Pesticides and PCBs in Batches 1, 2, 3, 6 and 7 with the exception of PCB 138 in Batch six and three pesticides and 2 PCBs in Batch seven. In all cases, the recoveries were high and are most likely due to matrix interferences. Recoveries for the majority of pesticides and PCBs in Batches four and five exceeded control limits due to high native levels compared with the levels spiked. In most cases, the spiked concentrations were 2 to 10 times lower than the concentrations detected in the samples.

REPLICATES

One sample from each extraction batch was analyzed in triplicate. Precision was measured by calculating the relative standard deviation (RSD) between the replicate results. RSDs for all detectable values were below the target precision goal of $\leq 30\%$ in Batches 1, 2, 3, 4 and 7. The RSD for Endosulfan Sulfate in Batch 5 was high due to comparison of very low concentrations, less than 1 ng/g in the replicates. RSDs for two pesticides and for two PCB congeners in Batch 6 were high due to matrix interferences associated with the first replicate sample.

QA/QC SUMMARY/PCBs and PESTICIDES (continued)

SRMs Not applicable.

MISCELLANEOUS All pesticide and PCB congener results are confirmed using a second dissimilar column. RPDs between the primary and confirmation values must be less than 75% to be considered a confirmed value.

REFERENCES

Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.

EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

QA/QC SUMMARY

PROGRAM: New York/New Jersey Federal Projects-2
PARAMETER: Polynuclear Aromatic Hydrocarbons (PAH) and 1,4-Dichlorobenzene
LABORATORY: Battelle/Marine Sciences Laboratory, Sequim, Washington
MATRIX: Clam and Worm Tissue

QA/QC DATA QUALITY OBJECTIVES

<u>Reference Method</u>	<u>MS Recovery</u>	<u>Surrogate Recovery</u>	<u>SRM Accuracy</u>	<u>Relative Precision</u>	<u>Detection Limit (wet wt)</u>
GC/MS/SIM	50-120%	30-150%	≤30%	≤30%	4 ng/g

SAMPLE CUSTODY A total of 68 worm and 68 clam samples was received on 6/15/94 in good condition. Samples were logged into Battelle's log-in system and stored frozen until extraction.

METHOD Tissue samples were extracted with methylene chloride using a roller under ambient conditions following a procedure which is based on methods used by the National Oceanic and Atmospheric Administration for its Status and Trends Program (Krahn et al. 1988). Samples were then cleaned using silica/alumina (5% deactivated) chromatography followed by HPLC cleanup.

Extracts were quantified using gas chromatography/mass spectrometry (GC/MS) in the selected ion mode (SIM) following a procedure based on EPA Method 8270 (EPA 1986).

HOLDING TIMES Samples were extracted in seven batches. All extracts were analyzed by GC/MS/SIM. The following summarizes the extraction and analysis dates:

<u>Batch</u>	<u>Species</u>	<u>Extraction</u>	<u>Analysis</u>
1	<i>M. nasuta</i>	7/28/94	9/9-9/12/94
2	<i>M. nasuta</i>	8/3/94	9/13-9/15/94
3	<i>M. nasuta</i>	8/17/94	9/23-9/25/94
4	<i>N. virens</i>	8/19/95	9/26-9/30/94
5	<i>N. virens</i>	8/26/94	9/8-9/11/94
6	<i>N. virens</i>	9/6/94	9/17-9/19/94
7	<i>M. nasuta/N. virens</i>	9/26/94	9/15-9/17-94
8	<i>M. nasuta</i> MDL study	10/10/94	10/25/94

DETECTION LIMITS Target detection limits of 4 ng/g wet weight were met for all PAH compounds except for fluoranthene and pyrene, which had method detection limits (MDL) between 4 and 6 ng/g wet weight. MDLs were determined by multiplying the standard deviation of seven spiked replicates of a background clam sample by the Student's t value (99 percentile). These MDLs were based on a wet weight of 20 g of tissue sample.

QA/QC SUMMARY/PAHs (continued)

Aliquots of samples that were analyzed in triplicate, used for spiking, or were re-extracted, were generally less than 20 g due to limited quantities of tissue available. Because MDLs reported are corrected for sample weight, the MDLs reported for these samples appear elevated and in some cases may exceed the target detection limit.

In addition a method detection limit verification study was performed, which consisted of analyzing four spiked aliquots of a background clam sample received with this project. The standard deviation of the results of these replicate analyses was multiplied by 3.5. Detection limits calculated in this way were all less than the target detection limit of 4 ng/g wet wt.

METHOD BLANKS

One method blank was extracted with each extraction batch. Benz[a]anthracene was detected in blanks from all batches and benzo[b]fluoranthene was detected in the blank from Batch 3. Two method blanks were analyzed with Batch 7 and in addition to benz[a]anthracene, three other compounds were detected in at least one of the two blanks; naphthalene, benzo[a]pyrene and indeno(123-cd)pyrene. All blank levels were less than three times the target MDL of 4 ng/g wet wt. Sample values that were less than five times the value of the method blank associated with that sample were flagged with a "B."

SURROGATES

Five isotopically labeled compounds were added prior to extraction to assess the efficiency of the method. These were d8-naphthalene, d10-acenaphthene, d12-chrysene, d14-dibenz[a,h]anthracene and d4-1,4 dichlorobenzene. Recoveries of all surrogates were within the quality control limits of 30% -150% with the exception of low recoveries for d4-1,4 dichlorobenzene in one sample from Batch 1 and Batch 4 and two samples in Batch seven. In addition, d8-naphthalene recovery was low in two samples in Batch seven.

MATRIX SPIKES

One sample from each batch was spiked with all PAH compounds. Matrix spike recoveries were generally, within QC limits of 50% -120%, with some exceptions. The recoveries for benzo(b)- and benzo[k]fluoranthene were variable due to the poor resolution of these two compounds. Spike recoveries quantified as the sum of these two compounds were within QC limits. Spike recoveries for a number of PAH compounds in Batches 4 and 7 were out of control due to high native levels, relative to the levels spiked. Spike concentrations were from 2 to 20 times lower than native concentrations. Recoveries for a number of compounds in Batches 4 and 6 were slightly above the upper control limit. These recoveries were all between 120% and 140%.

REPLICATES

One sample from each batch was extracted and analyzed in triplicate. Precision was measured by calculating the relative standard deviation (RSD) between the replicate results. All RSDs were within $\pm 30\%$.

SRMs

Not applicable.

QA/QC SUMMARY/PAHs (continued)

MISCELLANEOUS

Some of the compounds are flagged to indicate that the ion ratio for that compound was outside of the QC range. This is due primarily to low levels of the compound of interest. Because the confirmation ion is present at only a fraction of the level of the parent ion, when the native level of the compound is low, the amount of error in the concentration measurement of the confirmation ion goes up. The compound is actually quantified from the parent ion only, so most likely this will not affect the quality of the data. For sample values that are relatively high (>5 times the MDL) it may be an indication of some sort of interference.

REFERENCES

Krahn, M.M., C.A. Wigren, R.W. Pearce, L.K. Moore, R.G. Bogar, W.D. MacLeod, Jr., S-L Chan, and D.W. Brown. 1988. *New HPLC Cleanup and Revised Extraction Procedures for Organic Contaminants*. NOAA Technical Memorandum NMFS F/NWC-153. National Oceanic and Atmospheric Administration, National Marine Fisheries, Seattle, Washington.

EPA (U.S. Environmental Protection Agency). 1986. *Test Methods for Evaluating Solid Waste: Physical/Chemical Methods*. SW-846. U.S. Document No. 955-001-00000, U.S. Environmental Protection Agency, Washington D.C.

Table G.1. Metals in Tissue of *N. virens* (Wet Weight)

Sediment Treatment	Replicate	Batch	% Dry Weight	<i>N. virens</i> Metals (µg/g wet weight)								
				Ag ICP/MS	As ICP/MS	Cd ICP/MS	Cr ICP/MS	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn ICP/MS
COMP PC	1	1	14.07%	0.023 U ^(a)	1.89	0.088	0.205 U	1.20	0.006	0.277	0.335	36.9
COMP PC	2	1	16.08%	0.027 U	1.96	0.133	0.235 U	1.72	0.011	0.277	0.563	10.1
COMP PC	3	1	13.88%	0.023 U	1.72	0.092	0.203 U	1.29	0.004	0.480	0.344	85.9
COMP PC	4	1	15.67%	0.026 U	1.86	0.119	0.229 U	2.16	0.008	0.235	0.548	8.26
COMP PC	5	1	15.11%	0.025 U	1.81	0.079	0.220 U	1.69	0.007	0.239	0.302 U	16.0
R-CLIS	1	1	13.70%	0.023 U	1.99	0.053	0.200 U	1.51	0.012	0.180 U	0.319	8.52
R-CLIS	2	1	16.08%	0.027 U	2.27	0.060	0.235 U	1.66	0.010	0.212 U	0.486	9.29
R-CLIS	3	1	15.15%	0.025 U	1.80	0.049	0.221 U	1.35	0.008	0.199 U	0.333	40.6
R-CLIS	4	1	14.02%	0.023 U	2.27	0.055	0.205 U	1.37	0.011	0.222	0.331	52.6
R-CLIS	5	1	14.53%	0.024 U	2.06	0.057	0.212 U	1.73	0.012	0.248	0.334	20.2
R-MUD	1	1	13.12%	0.022	1.86	0.063	0.191 U	1.64	0.011	0.173 U	0.321	8.46
R-MUD	2	1	14.94%	0.029	2.29	0.079	0.218 U	10.8	0.013	0.197 U	0.647	11.6
R-MUD	3	1	15.21%	0.025 U	2.18	0.053	0.222 U	1.11	0.010	0.200 U	0.304 U	10.4
R-MUD	4	1	14.00%	0.026	2.11	0.062	0.204 U	1.58	0.011	0.184 U	0.280 U	8.08
R-MUD	5	1	13.24%	0.022	1.91	0.053	0.193 U	1.34	0.015	0.174 U	0.297	17.7
C-NV	1	1	14.84%	0.025 U	2.37	0.056	0.217 U	1.23	0.011	0.195 U	0.297 U	7.73
C-NV	2	1	12.32%	0.020 U	1.71	0.048	0.180 U	1.02	0.010	0.162 U	0.247 U	27.1
C-NV	3	1	14.51%	0.024 U	2.02	0.077	0.212 U	1.51	0.016	0.191 U	0.315	8.20
C-NV	4	1	13.67%	0.023 U	2.16	0.062	0.199 U	1.35	0.012	0.180 U	0.325	16.4
C-NV	5	1	14.91%	0.025 U	2.03	0.085	0.218 U	1.76	0.014	0.196 U	0.416	9.87
<i>N. virens</i> Background	1	1	12.86%	0.021 U	1.84	0.051	0.247	1.61	0.011	0.240	0.257 U	9.75
<i>N. virens</i> Background	2	1	12.94%	0.021 U	2.02	0.045	0.189 U	1.24	0.016	0.170 U	0.259 U	8.14
<i>N. virens</i> Background	3	1	12.05%	0.020 U	1.57	0.055	0.180	1.78	0.018	0.172	0.241 U	9.97

(a) U Undetected at or above given concentration.

Table G.2. Metals in Tissue of *N. virens* (Dry Weight)

Sediment Treatment	Replicate	Batch	% Dry Weight	<i>N. virens</i> Metals (µg/g dry weight)								
				Ag ICP/MS	As ICP/MS	Cd ICP/MS	Cr ICP/MS	Cu ICP/MS	Hg CVAF	Ni ICP/MS	Pb ICP/MS	Zn ICP/MS
COMP PC	1	1	14.07%	0.166 U ^(a)	13.4	0.626	1.46 U	8.54	0.045	1.97	2.38	262
COMP PC	2	1	16.08%	0.166 U	12.2	0.828	1.46 U	10.7	0.071	1.72	3.50	63.0
COMP PC	3	1	13.88%	0.166 U	12.4	0.665	1.46 U	9.30	0.026	3.46	2.48	619
COMP PC	4	1	15.67%	0.166 U	11.9	0.760	1.46 U	13.8	0.050	1.50	3.50	52.7
COMP PC	5	1	15.11%	0.166 U	12.0	0.523	1.46 U	11.2	0.044	1.58	2.00 U	106
R-CLIS	1	1	13.70%	0.166 U	14.5	0.385	1.46 U	11.0	0.085	1.32 U	2.33	62.2
R-CLIS	2	1	16.08%	0.166 U	14.1	0.372	1.46 U	10.3	0.061	1.32 U	3.02	57.8
R-CLIS	3	1	15.15%	0.166 U	11.9	0.324	1.46 U	8.88	0.050	1.32 U	2.20	268
R-CLIS	4	1	14.02%	0.166 U	16.2	0.395	1.46 U	9.80	0.078	1.58	2.36	375
R-CLIS	5	1	14.53%	0.166 U	14.2	0.393	1.46 U	11.9	0.082	1.71	2.30	139
R-MUD	1	1	13.12%	0.168	14.2	0.478	1.46 U	12.5	0.086	1.32 U	2.45	64.5
R-MUD	2	1	14.94%	0.196	15.3	0.531	1.46 U	72.6	0.089	1.32 U	4.33	77.5
R-MUD	3	1	15.21%	0.166 U	14.3	0.347	1.46 U	7.27	0.067	1.32 U	2.00 U	68.3
R-MUD	4	1	14.00%	0.186	15.1	0.444	1.46 U	11.3	0.075	1.32 U	2.00 U	57.7
R-MUD	5	1	13.24%	0.166	14.4	0.397	1.46 U	10.1	0.116	1.32 U	2.24	134
C-NV	1	1	14.84%	0.166 U	16.0	0.376	1.46 U	8.26	0.074	1.32 U	2.00 U	52.1
C-NV	2	1	12.32%	0.166 U	13.9	0.387	1.46 U	8.28	0.082	1.32 U	2.00 U	220
C-NV	3	1	14.51%	0.166 U	13.9	0.530	1.46 U	10.4	0.112	1.32 U	2.17	56.5
C-NV	4	1	13.67%	0.166 U	15.8	0.454	1.46 U	9.86	0.086	1.32 U	2.38	120
C-NV	5	1	14.91%	0.166 U	13.6	0.573	1.46 U	11.8	0.097	1.32 U	2.79	66.2
<i>N. virens</i> Background	1	1	12.86%	0.166 U	14.3	0.398	1.92	12.5	0.089	1.87	2.00 U	75.8
<i>N. virens</i> Background	2	1	12.94%	0.166 U	15.6	0.349	1.46 U	9.58	0.120	1.32 U	2.00 U	62.9
<i>N. virens</i> Background	3	1	12.05%	0.166 U	13.0	0.459	1.49	14.8	0.148	1.43	2.00 U	82.7

(a) U Undetected at or above given concentration.

TABLE G.3. Quality Control Summary for Metals in Tissue of *N. virens*

Sediment Treatment	Replicate	Batch	<i>N. virens</i> Metals (µg/g dry weight)								
			Ag	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn
Method Blanks											
Blank-1		1	0.166 U ^(a)	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-2		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-3		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Blank-4		1	0.166 U	3.39 U	0.081 U	1.46 U	6.86 U	0.001 U	1.32 U	2.00 U	10.8 U
Matrix Spikes											
COMP BU	2	1	0.166 U	13.9	0.404	1.46 U	10.6	0.059	1.32 U	2.42	69
COMP BU, MS	2	1	1.90	61.6	4.34	9.63	57.6	1.02	10.3	6.66	183
Concentration Recovered			1.90	47.7	3.94	9.63	47.0	0.96	10.3	4.24	114.0
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			91%	92%	94%	93%	90%	92%	99%	102%	114%
COMP BU	4	1	0.191	14.3	0.385	1.46 U	8.4	0.068	1.32 U	2.19	93.8
COMP BU, MS	4	1	2.06	63.4	4.45	10.2	57.4	1.18	10.4	6.13	153
Concentration Recovered			1.87	49.1	4.07	10.2	49.0	1.11	10.4	4.75	59.2
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			90%	94%	97%	98%	94%	107%	100%	114%	59% ^(b)
COMP EC-A	3	1	0.178 U	14.7	0.476	1.46 U	10.2	0.059	1.32 U	2.79	NA ^(c)
COMP EC-A, MS	3	1	0.968	61.3	4.28	9.84	56.8	1.04	10.1	6.95	NA
Concentration Recovered			0.968	46.6	3.80	9.84	46.6	0.98	10.1	4.16	NA
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	NS ^(d)
Percent Recovery			47% ^(b)	89%	91%	95%	89%	94%	97%	100%	NA
COMP HU-A	5	1	0.173 U	15.8	0.5313	1.46 U	11.0	0.077	1.32 U	2.77	98.7
COMP HU-A, MS	5	1	1.91	63.8	4.56	9.78	58.7	1.05	10.3	7.13	160
Concentration Recovered			1.91	48.0	4.03	9.78	47.7	0.973	10.3	4.36	61.3
Amount Spiked			2.08	52.1	4.17	10.4	52.1	1.04	10.4	4.17	100
Percent Recovery			92%	92%	97%	94%	91%	94%	99%	105%	61% ^(b)

G 3

TABLE G.3. (contd)

Sed Code ID	Replicate	Batch	<i>N. virens</i> Metals ($\mu\text{g/g}$ dry weight)									
			Ag	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn	
<u>Standard Reference Material</u>												
Certified value			1.68	14.0	4.15	1.43	66.3	0.0642	2.25	0.371	830	
Range			± 0.15	± 1.2	± 0.38	± 0.46	± 4.3	± 0.0067	± 0.44	± 0.014	± 57	
SRM 1566a	1	1	1.62	13.2	4.25	1.23	63.6	0.064	2.13	0.369	854	
SRM 1566a	2	1	1.54	12.5	4.01	1.00	58.3	0.057	3.05	0.389	778	
SRM 1566a	3	1	1.47	11.9	4.00	0.921	57.9	0.058	1.86	0.369	764	
SRM 1566a	4	1	1.51	11.9	4.01	0.948	60.4	0.061	1.65	0.363	792	
Percent Difference	1		4	6	2	14	4	0	5	1	3	
Percent Difference	2		8	11	3	30 ^(a)	12	11	36 ^(e)	5	6	
Percent Difference	3		13	15	4	36 ^(e)	13	10	17	1	8	
Percent Difference	4		10	15	3	34 ^(e)	9	5	27 ^(e)	2	5	
<u>Analytical Replicates</u>												
COMP BU, Replicate 1	4	1	0.195	14.4	0.388	1.459 U	8.30	0.065	1.32 U	2.18	60.2	
COMP BU, Replicate 2	4	1	0.195	14.0	0.362	1.459 U	8.34	0.074	1.32 U	2.19	59.1	
COMP BU, Replicate 3	4	1	0.182	14.6	0.404	1.459 U	8.55	0.066	1.32 U	2.19	162	
RSD			4%	2%	6%	NA	2%	7%	NA	0%	63% ^(f)	
COMP EC-A, Replicate 1	3	1	0.166 U	13.6	0.472	1.459 U	9.66	0.059	1.32 U	2.58	156	
COMP EC-A, Replicate 2	3	1	0.166 U	15.4	0.466	1.459 U	10.8	0.061	1.32 U	2.88	155	
COMP EC-A, Replicate 3	3	1	0.166 U	15.1	0.491	1.459 U	10.3	0.058	1.32 U	2.90	165	
RSD			NA	7%	3%	NA	6%	3%	NA	6%	3%	
COMP BU, Replicate 1	2	1	0.166 U	13.5	0.396	1.459 U	10.3	0.055	1.32 U	2.30	87.2	
COMP BU, Replicate 2	2	1	0.166 U	14.1	0.401	1.459 U	10.8	0.064	1.32 U	2.43	61.8	
COMP BU, Replicate 3	2	1	0.166 U	14.0	0.416	1.459 U	10.7	0.058	1.32 U	2.54	58.1	
RSD			NA	2%	3%	NA	2%	8%	NA	5%	23% ^(f)	
COMP HU-A, Replicate 1	5	1	0.166 U	16.3	0.568	1.459 U	11.4	0.071	1.32 U	2.84	98.9	
COMP HU-A, Replicate 2	5	1	0.166 U	15.7	0.490	1.459 U	11.1	0.090	1.32 U	2.76	80.1	
COMP HU-A, Replicate 3	5	1	0.166 U	15.5	0.536	1.459 U	10.6	0.069	1.32 U	2.70	117	
RSD			NA	3%	7%	NA	4%	15%	NA	3%	19%	

(a) U Undetected at or above given concentration.

(b) Outside quality control criteria (75-125%) for matrix spike recovery.

(c) NA Not applicable.

(d) NS Not spiked.

(e) Outside quality control criteria ($\pm 20\%$) for SRMs.(f) Outside quality control criteria ($\pm 20\%$) for RSD.

TABLE G.4. Pesticides and PCB Congeners (Wet Weight) in Tissue of *N. virens*

Treatment	PC	PC	PC	PC	PC
Replicate	1	2	3	4	5
Batch	6	7	4	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.07	16.08	13.88	15.67	15.11
Heptachlor	0.19 U ^(a)	0.19 U	0.19 U	0.20 U	0.19 U
Aldrin	1.48	1.24	0.99	0.84	1.55
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U	0.15 U	0.13 U
2,4'-DDE	0.26 U	0.26 U	0.26 U	0.29 U	0.26 U
Endosulfan I	0.18 U	0.18 U	0.18 U	0.20 U	0.18 U
α -Chlordane	4.10	5.92	2.89	7.22	5.60
Trans Nonachlor	2.95	2.96	2.61	2.96	4.01
4,4'-DDE	6.38	5.04	4.15	6.69	8.35
Dieldrin	11.1	8.34	5.42	9.18	9.54
2,4'-DDD	15.2	7.36	5.40	10.0	10.4
2,4'-DDT	0.18 U	0.18 U	0.18 U	0.20 U	0.18 U
4,4'-DDD	71.0	18.4	16.5	27.6	30.7
Endosulfan II	0.18 U	0.18 U	0.18 U	0.20 U	0.18 U
4,4'-DDT	0.15 U	0.15 U	0.15 U	0.17 U	0.15 U
Endosulfan Sulfate	3.43	0.18 U	0.18 U	0.20 U	5.09
PCB 8	0.41 U	0.41 U	0.41 U	0.45 U	0.41 U
PCB 18	1.09	0.43 U	0.43 U	0.62	1.12
PCB 28	1.38	1.01	0.39	1.12	1.39
PCB 52	11.9	7.17	5.68	9.52	12.8
PCB 49	2.69	1.59	0.99	2.07	2.87
PCB 44	1.71	1.19	0.54	1.62	1.92
PCB 66	0.09 U	0.09 U	0.09 U	0.10 U	0.09 U
PCB 101	18.4	11.7	11.4	16.4	24.1
PCB 87	2.37	1.56	1.26	2.27	3.46
PCB 118	12.3	6.65	5.28	9.54	13.8
PCB 184	0.24 U	0.24 U	0.24 U	0.26 U	0.24 U
PCB 153	15.7	10.2	7.81	13.0	17.6
PCB 105	7.42	5.27	4.33	6.95	9.16
PCB 138	15.9	11.3	8.39	14.5	18.4
PCB 187	4.38	3.34	2.22	3.72	5.00
PCB 183	2.64	1.73	1.78	2.02	3.89
PCB 128	3.40	0.15 U	1.98	0.17 U	4.40
PCB 180	7.04	4.05	4.13	4.88	8.68
PCB 170	3.29	1.95	2.17	2.35	3.90
PCB 195	0.63	0.10 U	0.41	0.46	0.76
PCB 206	1.20	0.98	0.80	1.02	1.25
PCB 209	0.31	0.26	0.22	0.26	0.22
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	100	72	69	81	97
PCB 198 (SIS)	79	82	115	81	74

TABLE G.4. (contd)

Treatment	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	1	2	3	4	5
Batch	4	5	6	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.12	14.94	15.21	14.00	13.24
Heptachlor	0.19 U	0.18 U	0.19 U	0.19 U	0.23 U
Aldrin	0.13 U	0.12 U	0.13 U	0.13 U	0.16 U
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U	0.13 U	0.16 U
2,4'-DDE	0.26 U	0.26 U	0.26 U	0.26 U	0.32 U
Endosulfan I	0.18 U	0.18 U	0.18 U	0.18 U	0.22 U
α-Chlordane	0.10 U	0.09 U	0.10 U	0.10 U	0.12 U
Trans Nonachlor	0.43	0.61	0.67	0.39	0.61
4,4'-DDE	0.19 U	0.18 U	0.35	0.19 U	0.23 U
Dieldrin	0.94	0.71	0.52 U	0.66	0.64 U
2,4'-DDD	0.25 U	0.35	0.25 U	0.25 U	0.31 U
2,4'-DDT	0.18 U	0.18 U	0.18 U	0.18 U	0.22 U
4,4'-DDD	1.00	0.39	0.26 U	0.85	0.32 U
Endosulfan II	0.18 U	0.18 U	0.18 U	0.18 U	0.22 U
4,4'-DDT	0.15 U	0.15 U	0.15 U	0.15 U	0.19 U
Endosulfan Sulfate	0.18 U	0.18 U	0.18 U	0.18 U	0.22 U
PCB 8	0.41 U	0.40 U	0.41 U	0.41 U	0.51 U
PCB 18	0.43 U	0.42 U	0.43 U	0.43 U	0.53 U
PCB 28	0.20 U	0.20 U	0.20 U	0.20 U	0.25 U
PCB 52	0.36 U	0.35 U	0.43	0.36 U	0.64
PCB 49	0.24 U	0.23 U	0.24 U	0.24 U	0.29 U
PCB 44	0.17 U	0.16 U	0.17 U	0.17 U	0.20 U
PCB 66	0.09 U	0.09 U	0.09 U	0.09 U	0.12 U
PCB 101	0.15 U	0.81	0.44	0.45	0.54
PCB 87	0.16 U	0.16 U	0.23	0.16 U	0.20 U
PCB 118	0.29 U	0.29 U	0.29 U	0.29 U	0.37 U
PCB 184	0.24 U	0.23 U	0.24 U	0.24 U	0.29 U
PCB 153	1.76	2.35	2.20	2.08	1.66
PCB 105	0.11 U	0.11 U	0.24	0.28	0.27
PCB 138	0.92	1.44	1.17	1.36	1.03
PCB 187	0.38	0.53	0.60	0.58	0.43
PCB 183	0.24 U	0.24	0.24	0.24 U	0.29 U
PCB 128	0.19	0.22	0.20	0.20	0.90 U
PCB 180	0.45	0.69	0.60	0.56	0.59
PCB 170	0.17 U	0.37	0.33	0.27	0.34
PCB 195	0.10 U	0.10 U	0.10 U	0.10 U	0.12 U
PCB 206	0.30	0.23	0.23	0.11 U	0.31
PCB 209	0.16	0.15	0.16	0.17	0.15
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	77	93	83	58	84
PCB 198 (SIS)	118	82	66	57	64

TABLE G.4. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	6	5	4	6	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.70	16.08	15.15	14.02	14.53
Heptachlor	0.19 U	0.19 U	0.18 U	0.18 U	0.19 U
Aldrin	1.04	0.79	0.77	0.80	0.68
Heptachlor Epoxide	0.27	0.13 U	0.13 U	0.13 U	0.13 U
2,4'-DDE	0.26 U	0.26 U	0.26 U	0.26 U	0.26 U
Endosulfan I	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
a-Chlordane	0.10 U	0.17	0.11	0.20	0.10 U
Trans Nonachlor	0.76	0.69	0.59	0.80	0.23
4,4'-DDE	1.25	0.70	0.60	0.44	0.19 U
Dieldrin	1.62	0.92	1.08	0.51 U	0.61
2,4'-DDD	3.00	1.24	0.50	0.66	0.25 U
2,4'-DDT	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
4,4'-DDD	6.12	1.95	1.18	0.26 U	0.26 U
Endosulfan II	0.18 U	0.18 U	0.18 U	0.18 U	0.18 U
4,4'-DDT	0.15 U	0.15 U	0.15 U	0.15 U	0.15 U
Endosulfan Sulfate	0.18 U	0.18 U	0.18 U	0.22	0.18 U
PCB 8	0.41 U	0.41 U	0.40 U	0.40 U	0.41 U
PCB 18	0.43 U	0.43 U	0.42 U	0.42 U	0.43 U
PCB 28	0.48	0.37	0.28	0.20 U	0.20 U
PCB 52	5.31	1.65	0.99	0.94	0.36 U
PCB 49	1.41	0.47	0.34	0.31	0.24 U
PCB 44	0.22	0.17 U	0.16 U	0.16 U	0.17 U
PCB 66	0.09 U	0.09 U	0.09 U	0.09 U	0.09 U
PCB 101	8.13	3.32	1.62	1.47	0.43
PCB 87	0.75	0.16 U	0.17	0.16 U	0.16 U
PCB 118	5.67	2.06	0.99	0.89	0.29 U
PCB 184	0.24 U	0.24 U	0.23 U	0.23 U	0.24 U
PCB 153	7.38	4.36	2.92	3.45	0.84
PCB 105	2.12	1.13	0.45	0.45	0.13
PCB 138	6.11	3.64	1.88	2.22	0.50
PCB 187	1.76	0.91	0.88	1.06	0.23
PCB 183	0.88	0.41	0.34	0.43	0.24 U
PCB 128	1.21	0.68	0.36	0.42	0.15 U
PCB 180	2.39	1.20	0.95	0.92	0.41
PCB 170	1.11	0.67	0.56	0.51	0.19
PCB 195	0.10 U	0.10 U	0.10 U	0.10 U	0.10 U
PCB 206	0.38	0.34	0.38	0.35	0.14
PCB 209	0.24	0.23	0.19	0.22	0.09 U
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	89	97	52	80	89
PCB 198 (SIS)	70	75	85	65	155

TABLE G.4. (contd)

Treatment	C-NV	C-NV	C-NV	C-NV	C-NV
Replicate	1	2	3	4	5
Batch	6	6	7	4	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.84	12.32	14.51	13.67	14.91
Heptachlor	0.19 U	0.19 U	0.31 U	0.19 U	0.19 U
Aldrin	0.13 U	0.13 U	0.21 U	0.80	0.13 U
Heptachlor Epoxide	0.13 U	0.13 U	0.22 U	0.13 U	0.13 U
2,4'-DDE	0.26 U	0.26 U	0.43 U	0.26 U	0.26 U
Endosulfan I	0.18 U	0.18 U	0.30 U	0.18 U	0.18 U
a-Chlordane	0.10 U	0.10 U	0.26	0.10 U	0.10 U
Trans Nonachlor	0.61	0.60	0.24 U	0.48	0.38
4,4'-DDE	0.22	0.29	0.31 U	0.47	0.19 U
Dieldrin	0.92	0.93	1.37	0.52 U	0.52 U
2,4'-DDD	0.42	0.40	3.25	1.67	0.25 U
2,4'-DDT	0.18 U	0.18 U	0.30 U	0.18 U	0.18 U
4,4'-DDD	0.71	0.83	10.5	5.21	0.26 U
Endosulfan II	0.18 U	0.18 U	0.30 U	0.18 U	0.18 U
4,4'-DDT	0.15 U	0.15 U	0.38	0.15 U	0.15 U
Endosulfan Sulfate	0.18 U	0.18 U	0.30 U	0.18 U	0.18 U
PCB 8	0.41 U	0.41 U	0.68 U	0.41 U	0.41 U
PCB 18	0.43 U	0.43 U	0.71 U	0.43 U	0.43 U
PCB 28	0.20 U	0.20 U	0.34 U	0.20 U	0.20 U
PCB 52	0.69	0.52	0.59 U	2.45	0.40
PCB 49	0.24 U	0.24 U	0.39 U	0.26	0.24 U
PCB 44	0.17 U	0.17 U	0.27 U	0.17 U	0.17 U
PCB 66	0.09 U	0.09 U	0.16 U	0.09 U	0.09 U
PCB 101	0.80	0.78	2.53	3.69	0.15 U
PCB 87	0.16 U	0.16 U	0.26 U	0.16 U	0.16 U
PCB 118	0.47	0.45	0.95	1.95	0.47
PCB 184	0.24 U	0.24 U	0.39 U	0.24 U	0.24 U
PCB 153	2.19	2.20	4.48	3.73	1.93
PCB 105	0.34	0.33	1.02	1.09	0.28
PCB 138	1.47	1.42	3.46	3.05	1.19
PCB 187	0.64	0.62	0.88	0.86	0.51
PCB 183	0.28	0.25	0.41	0.44	0.24 U
PCB 128	0.26	0.25	0.63	0.61	0.22
PCB 180	0.71	0.72	1.19	1.44	0.57
PCB 170	0.43	0.38	0.58	0.75	0.38
PCB 195	0.10 U	0.10 U	0.17 U	0.10 U	0.10 U
PCB 206	0.29	0.27	0.29	0.41	0.21
PCB 209	0.16	0.16	0.83	0.21	0.12
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)	83	87	81	71	41
PCB 198 (SIS)	68	69	84	124	63

TABLE G.4. (contd)

Treatment Replicate Batch Units Percent Dry Weight	<i>N. virens</i>	<i>N. virens</i>	<i>N. virens</i>
	Background	Background	Background
	1	2	3
	7	7	7
	ng/g	ng/g	ng/g
	12.86	12.94	12.05
Heptachlor	0.19 U	0.19 U	0.19 U
Aldrin	0.73	0.13 U	0.13 U
Heptachlor Epoxide	0.13 U	0.13 U	0.13 U
2,4'-DDE	0.26 U	0.26 U	0.26 U
Endosulfan I	0.18 U	0.18 U	0.18 U
α -Chlordane	0.10 U	0.10 U	0.10 U
Trans Nonachlor	0.44	0.15 U	0.46
4,4'-DDE	0.19 U	0.99	0.19 U
Dieldrin	0.52 U	1.01	0.65
2,4'-DDD	0.25 U	0.25 U	0.25 U
2,4'-DDT	0.18 U	0.18 U	0.18 U
4,4'-DDD	0.26 U	0.26 U	0.56
Endosulfan II	0.18 U	0.18 U	0.18 U
4,4'-DDT	0.18	0.15 U	0.15 U
Endosulfan Sulfate	0.18 U	0.18 U	0.18 U
PCB 8	0.41 U	0.41 U	0.41 U
PCB 18	0.43 U	0.43 U	0.43 U
PCB 28	0.21	0.20 U	0.20 U
PCB 52	0.36 U	0.36 U	0.36 U
PCB 49	0.24 U	0.24 U	0.24 U
PCB 44	0.17 U	0.17 U	0.17 U
PCB 66	0.73	0.09 U	0.55
PCB 101	0.58	0.45	0.44
PCB 87	0.16 U	0.62	0.16 U
PCB 118	0.29 U	0.29 U	0.29 U
PCB 184	0.24 U	0.24 U	0.24 U
PCB 153	2.24	1.97	1.72
PCB 105	0.26	0.23	0.25
PCB 138	1.60	1.35	1.19
PCB 187	0.63	0.54	0.41
PCB 183	0.24	0.24 U	0.24 U
PCB 128	0.24	0.20	0.17
PCB 180	0.49	0.43	0.43
PCB 170	0.17 U	0.21	0.19
PCB 195	0.10 U	0.10 U	0.10 U
PCB 206	0.11 U	0.11 U	0.11 U
PCB 209	0.10	0.09 U	0.09 U
<u>Surrogate Recoveries (%)</u>			
PCB 103 (SIS)	96	84	75
PCB 198 (SIS)	84	80	81

(a) U Undetected at or above given concentration.

TABLE G.5. Pesticides and PCB Congeners (Dry Weight) in Tissue of *N. virens*

Treatment	PC	PC	PC	PC	PC
Replicate	1	2	3	4	5
Batch	6	7	4	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.07	16.08	13.88	15.67	15.11
Heptachlor	1.35 U	1.18 U	1.37 U	1.28 U	1.26 U
Aldrin	10.5	7.71	7.13	5.36	10.3
Heptachlor Epoxide	0.92 U	0.81 U	0.94 U	0.96 U	0.86 U
2,4'-DDE	1.85 U	1.62 U	1.87 U	1.85 U	1.72 U
Endosulfan I	1.28 U	1.12 U	1.30 U	1.28 U	1.19 U
α -Chlordane	29.1	36.8	20.8	46.1	37.1
Trans Nonachlor	21.0	18.4	18.8	18.9	26.5
4,4'-DDE	45.3	31.3	29.9	42.7	55.3
Dieldrin	78.9	51.9	39.0	58.6	63.1
2,4'-DDD	108	45.8	38.9	63.8	68.8
2,4'-DDT	1.28 U	1.12 U	1.30 U	1.28 U	1.19 U
4,4'-DDD	505	114	119	176	203
Endosulfan II	1.28 U	1.12 U	1.30 U	1.28 U	1.19 U
4,4'-DDT	1.07 U	0.93 U	1.08 U	1.08 U	0.99 U
Endosulfan Sulfate	24.4	1.12 U	1.30 U	1.28 U	33.7
PCB 8	2.91 U	2.55 U	2.95 U	2.87 U	2.71 U
PCB 18	7.75	2.67 U	3.10 U	3.96	7.41
PCB 28	9.81	6.28	2.81	7.15	9.20
PCB 52	84.6	44.6	40.9	60.8	84.7
PCB 49	19.1	9.89	7.13	13.2	19.0
PCB 44	12.2	7.40	3.89	10.3	12.7
PCB 66	0.64 U	0.56 U	0.65 U	0.64 U	0.60 U
PCB 101	131	72.8	82.1	105	159
PCB 87	16.8	9.70	9.08	14.5	22.9
PCB 118	87.4	41.4	38.0	60.9	91.3
PCB 184	1.71 U	1.49 U	1.73 U	1.66 U	1.59 U
PCB 153	112	63.4	56.3	83.0	116
PCB 105	52.7	32.8	31.2	44.4	60.6
PCB 138	113	70.3	60.4	92.5	122
PCB 187	31.1	20.8	16.0	23.7	33.1
PCB 183	18.8	10.8	12.8	12.9	25.7
PCB 128	24.2	0.93 U	14.3	1.08 U	29.1
PCB 180	50.0	25.2	29.8	31.1	57.4
PCB 170	23.4	12.1	15.6	15.0	25.8
PCB 195	4.48	0.62 U	2.95	2.94	5.03
PCB 206	8.53	6.09	5.76	6.51	8.27
PCB 209	2.20	1.62	1.59	1.66	1.46

TABLE G.5. (contd)

Treatment	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	1	2	3	4	5
Batch	4	5	6	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.12	14.94	15.21	14.00	13.24
Heptachlor	1.45 U	1.20 U	1.25 U	1.36 U	1.74 U
Aldrin	0.99 U	0.80 U	0.85 U	0.93 U	1.21 U
Heptachlor Epoxide	0.99 U	0.87 U	0.85 U	0.93 U	1.21 U
2,4'-DDE	1.98 U	1.74 U	1.71 U	1.86 U	2.42 U
Endosulfan I	1.37 U	1.20 U	1.18 U	1.29 U	1.66 U
α -Chlordane	0.76 U	0.60 U	0.66 U	0.71 U	0.91 U
Trans Nonachlor	3.28	4.08	4.40	2.79	4.61
4,4'-DDE	1.45 U	1.20 U	2.30	1.36 U	1.74 U
Dieldrin	7.16	4.75	3.42 U	4.71	4.83 U
2,4'-DDD	1.91 U	2.34	1.64 U	1.79 U	2.34 U
2,4'-DDT	1.37 U	1.20 U	1.18 U	1.29 U	1.66 U
4,4'-DDD	7.62	2.61	1.71 U	6.07	2.42 U
Endosulfan II	1.37 U	1.20 U	1.18 U	1.29 U	1.66 U
4,4'-DDT	1.14 U	1.00 U	0.99 U	1.07 U	1.44 U
Endosulfan Sulfate	1.37 U	1.20 U	1.18 U	1.29 U	1.66 U
PCB 8	3.13 U	2.68 U	2.70 U	2.93 U	3.85 U
PCB 18	3.28 U	2.81 U	2.83 U	3.07 U	4.00 U
PCB 28	1.52 U	1.34 U	1.31 U	1.43 U	1.89 U
PCB 52	2.74 U	2.34 U	2.83	2.57 U	4.83
PCB 49	1.83 U	1.54 U	1.58 U	1.71 U	2.19 U
PCB 44	1.30 U	1.07 U	1.12 U	1.21 U	1.51 U
PCB 66	0.69 U	0.60 U	0.59 U	0.64 U	0.91 U
PCB 101	1.14 U	5.42	2.89	3.21	4.08
PCB 87	1.22 U	1.07 U	1.51	1.14 U	1.51 U
PCB 118	2.21 U	1.94 U	1.91 U	2.07 U	2.79 U
PCB 184	1.83 U	1.54 U	1.58 U	1.71 U	2.19 U
PCB 153	13.4	15.7	14.5	14.9	12.5
PCB 105	0.84 U	0.74 U	1.58	2.00	2.04
PCB 138	7.01	9.64	7.69	9.71	7.78
PCB 187	2.90	3.55	3.94	4.14	3.25
PCB 183	1.83 U	1.61	1.58	1.71 U	2.19 U
PCB 128	1.45	1.47	1.31	1.43	6.80 U
PCB 180	3.43	4.62	3.94	4.00	4.46
PCB 170	1.30 U	2.48	2.17	1.93	2.57
PCB 195	0.76 U	0.67 U	0.66 U	0.71 U	0.91 U
PCB 206	2.29	1.54	1.51	0.79 U	2.34
PCB 209	1.22	1.00	1.05	1.21	1.13

TABLE G.5. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	6	5	4	6	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.70	16.08	15.15	14.02	14.53
Heptachlor	1.39 U	1.18 U	1.19 U	1.28 U	1.31 U
Aldrin	7.59	4.91	5.08	5.71	4.68
Heptachlor Epoxide	1.97	0.81 U	0.86 U	0.93 U	0.89 U
2,4'-DDE	1.90 U	1.62 U	1.72 U	1.85 U	1.79 U
Endosulfan I	1.31 U	1.12 U	1.19 U	1.28 U	1.24 U
a-Chlordane	0.73 U	1.06	0.73	1.43	0.69 U
Trans Nonachlor	5.55	4.29	3.89	5.71	1.58
4,4'-DDE	9.12	4.35	3.96	3.14	1.31 U
Dieldrin	11.8	5.72	7.13	3.64 U	4.20
2,4'-DDD	21.9	7.71	3.30	4.71	1.72 U
2,4'-DDT	1.31 U	1.12 U	1.19 U	1.28 U	1.24 U
4,4'-DDD	44.7	12.1	7.79	1.85 U	1.79 U
Endosulfan II	1.31 U	1.12 U	1.19 U	1.28 U	1.24 U
4,4'-DDT	1.09 U	0.93 U	0.99 U	1.07 U	1.03 U
Endosulfan Sulfate	1.31 U	1.12 U	1.19 U	1.57	1.24 U
PCB 8	2.99 U	2.55 U	2.64 U	2.85 U	2.82 U
PCB 18	3.14 U	2.67 U	2.77 U	3.00 U	2.96 U
PCB 28	3.50	2.30	1.85	1.43 U	1.38 U
PCB 52	38.8	10.3	6.53	6.70	2.48 U
PCB 49	10.3	2.92	2.24	2.21	1.65 U
PCB 44	1.61	1.06 U	1.06 U	1.14 U	1.17 U
PCB 66	0.66 U	0.56 U	0.59 U	0.64 U	0.62 U
PCB 101	59.3	20.6	10.7	10.5	2.96
PCB 87	5.47	1.00 U	1.12	1.14 U	1.10 U
PCB 118	41.4	12.8	6.53	6.35	2.00 U
PCB 184	1.75 U	1.49 U	1.52 U	1.64 U	1.65 U
PCB 153	53.9	27.1	19.3	24.6	5.78
PCB 105	15.5	7.03	2.97	3.21	0.89
PCB 138	44.6	22.6	12.4	15.8	3.44
PCB 187	12.8	5.66	5.81	7.56	1.58
PCB 183	6.42	2.55	2.24	3.07	1.65 U
PCB 128	8.83	4.23	2.38	3.00	1.03 U
PCB 180	17.4	7.46	6.27	6.56	2.82
PCB 170	8.10	4.17	3.70	3.64	1.31
PCB 195	0.73 U	0.62 U	0.66 U	0.71 U	0.69 U
PCB 206	2.77	2.11	2.51	2.50	0.96
PCB 209	1.75	1.43	1.25	1.57	0.62 U

TABLE G.5. (contd)

Treatment	C-NV	C-NV	C-NV	C-NV	C-NV
Replicate	1	2	3	4	5
Batch	6	6	7	4	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.84	12.32	14.51	13.67	14.91
Heptachlor	1.28 U	1.54 U	2.14 U	1.39 U	1.27 U
Aldrin	0.88 U	1.06 U	1.45 U	5.85	0.87 U
Heptachlor Epoxide	0.88 U	1.06 U	1.52 U	0.95 U	0.87 U
2,4'-DDE	1.75 U	2.11 U	2.96 U	1.90 U	1.74 U
Endosulfan I	1.21 U	1.46 U	2.07 U	1.32 U	1.21 U
α -Chlordane	0.67 U	0.81 U	1.79	0.73 U	0.67 U
Trans Nonachlor	4.11	4.87	1.65 U	3.51	2.55
4,4'-DDE	1.48	2.35	2.14 U	3.44	1.27 U
Dieldrin	6.20	7.55	9.44	3.80 U	3.49 U
2,4'-DDD	2.83	3.25	22.4	12.2	1.68 U
2,4'-DDT	1.21 U	1.46 U	2.07 U	1.32 U	1.21 U
4,4'-DDD	4.78	6.74	72.6	38.1	1.74 U
Endosulfan II	1.21 U	1.46 U	2.07 U	1.32 U	1.21 U
4,4'-DDT	1.01 U	1.22 U	2.62	1.10 U	1.01 U
Endosulfan Sulfate	1.21 U	1.46 U	2.07 U	1.32 U	1.21 U
PCB 8	2.76 U	3.33 U	4.69 U	3.00 U	2.75 U
PCB 18	2.90 U	3.49 U	4.89 U	3.15 U	2.88 U
PCB 28	1.35 U	1.62 U	2.34 U	1.46 U	1.34 U
PCB 52	4.65	4.22	4.07 U	17.9	2.68
PCB 49	1.62 U	1.95 U	2.69 U	1.90	1.61 U
PCB 44	1.15 U	1.38 U	1.86 U	1.24 U	1.14 U
PCB 66	0.61 U	0.73 U	1.10 U	0.66 U	0.60 U
PCB 101	5.39	6.33	17.4	27.0	1.01 U
PCB 87	1.08 U	1.30 U	1.79 U	1.17 U	1.07 U
PCB 118	3.17	3.65	6.55	14.3	3.15
PCB 184	1.62 U	1.95 U	2.69 U	1.76 U	1.61 U
PCB 153	14.8	17.9	30.9	27.3	12.9
PCB 105	2.29	2.68	7.03	7.97	1.88
PCB 138	9.91	11.5	23.8	22.3	7.98
PCB 187	4.31	5.03	6.06	6.29	3.42
PCB 183	1.89	2.03	2.83	3.22	1.61 U
PCB 128	1.75	2.03	4.34	4.46	1.48
PCB 180	4.78	5.84	8.20	10.5	3.82
PCB 170	2.90	3.08	4.00	5.49	2.55
PCB 195	0.67 U	0.81 U	1.17 U	0.73 U	0.67 U
PCB 206	1.95	2.19	2.00	3.00	1.41
PCB 209	1.08	1.30	5.72	1.54	0.80

TABLE G.5. (contd)

Treatment	<i>N. virens</i> Background	<i>N. virens</i> Background	<i>N. virens</i> Background
Replicate	1	2	3
Batch	7	7	7
Units	ng/g	ng/g	ng/g
Percent Dry Weight	12.86	12.94	12.05
Heptachlor	1.5 U	1.5 U	1.6 U
Aldrin	5.7	1.0 U	1.1 U
Heptachlor Epoxide	1.0 U	1.0 U	1.1 U
2,4'-DDE	2.0 U	2.0 U	2.2 U
Endosulfan I	1.4 U	1.4 U	1.5 U
α -Chlordane	0.78 U	0.77 U	0.83 U
Trans Nonachlor	3.4	1.2 U	3.8
4,4'-DDE	1.5 U	7.7	1.6 U
Dieldrin	4.0 U	7.81	5.4
2,4'-DDD	1.9 U	1.9 U	2.1 U
2,4'-DDT	1.4 U	1.4 U	1.5 U
4,4'-DDD	2.0 U	2.0 U	4.6
Endosulfan II	1.4 U	1.4 U	1.5 U
4,4'-DDT	1.4	1.2 U	1.2 U
Endosulfan Sulfate	1.4 U	1.4 U	1.5 U
PCB 8	3.2 U	3.2 U	3.4 U
PCB 18	3.3 U	3.3 U	3.6 U
PCB 28	1.6	1.5 U	1.7 U
PCB 52	2.8 U	2.8 U	3.0 U
PCB 49	1.9 U	1.9 U	2.0 U
PCB 44	1.3 U	1.3 U	1.4 U
PCB 66	5.7	0.7 U	4.6
PCB 101	4.5	3.5	3.7
PCB 87	1.2 U	4.8	1.3 U
PCB 118	2.3 U	2.2 U	2.4 U
PCB 184	1.9 U	1.9 U	2.0 U
PCB 153	17.4	15.2	14.3
PCB 105	2.0	1.8	2.1
PCB 138	12.4	10.4	9.88
PCB 187	4.9	4.2	3.4
PCB 183	1.9	1.9 U	2.0 U
PCB 128	1.9	1.5	1.4
PCB 180	3.8	3.3	3.6
PCB 170	1.3 U	1.6	1.6
PCB 195	0.78 U	0.77 U	0.83 U
PCB 206	0.86 U	0.85 U	0.91 U
PCB 209	0.78	0.7 U	0.7 U

(a) U Undetected at or above given concentration.

**TABLE G.6. Quality Control Summary for Pesticides and PCB Congeners
in Tissue of *N. virens* (Wet Weight)**

<u>Blanks</u>	Treatment Replicate Batch Wet Wt. Units	Blank 1 4 NA ng/g	Blank 1 5 NA ng/g	Blank 1 6 NA ng/g	Blank 1 7 NA ng/g
Heptachlor		0.20 U ^(a)	0.19 U	0.19 U	0.21 U
Aldrin		0.13 U	0.13 U	0.13 U	0.15 U
Heptachlor epoxide		0.14 U	0.14 U	0.14 U	0.15 U
2,4'-DDE		0.28 U	0.27 U	0.27 U	0.30 U
Endosulfan I		0.19 U	0.18 U	0.19 U	0.21 U
α -Chlordane		0.10 U	0.10 U	0.10 U	0.11 U
Trans Nonachlor		0.15 U	0.15 U	0.15 U	0.17 U
4,4'-DDE		0.20 U	1.90 U	0.20 U	0.22 U
Dieldrin		0.55 U	0.53 U	0.54 U	0.60 U
2,4'-DDD		0.27 U	0.26 U	0.26 U	0.29 U
2,4'-DDT		0.19 U	0.18 U	0.19 U	0.21 U
4,4'-DDD		0.28 U	0.27 U	0.27 U	0.30 U
Endosulfan II		0.19 U	0.18 U	0.19 U	0.21 U
4,4'-DDT		0.16 U	0.15 U	0.16 U	0.18 U
Endosulfan Sulfate		0.19 U	0.18 U	0.19 U	0.21 U
PCB 8		0.44 U	0.42 U	0.43 U	0.48 U
PCB 18		0.46 U	0.44 U	0.45 U	0.50 U
PCB 28		0.22 U	0.21 U	0.21 U	0.24 U
PCB 52		0.38 U	0.37 U	0.37 U	0.42 U
PCB 49		0.25 U	0.24 U	0.25 U	0.27 U
PCB 44		0.17 U	0.17 U	0.17 U	0.19 U
PCB 66		0.10 U	0.10 U	0.10 U	0.11 U
PCB 101		0.15 U	0.15 U	0.15 U	0.17 U
PCB 87		0.17 U	0.16 U	0.17 U	0.19 U
PCB 118		0.31 U	0.30 U	0.31 U	0.34 U
PCB 184		0.25 U	0.24 U	0.25 U	0.27 U
PCB 153		0.13 U	0.12 U	0.13 U	0.14 U
PCB 105		0.12 U	0.11 U	0.12 U	0.13 U
PCB 138		0.31 U	0.30 U	0.30 U	0.34 U
PCB 187		0.13 U	0.13 U	0.13 U	0.15 U
PCB 183		0.25 U	0.24 U	0.25 U	0.27 U
PCB 128		0.16 U	0.16 U	0.16 U	0.18 U
PCB 180		0.20 U	0.19 U	0.19 U	0.21 U
PCB 170		0.18 U	0.17 U	0.17 U	0.19 U
PCB 195		0.11 U	0.10 U	0.10 U	0.12 U
PCB 206		0.12 U	0.12 U	0.12 U	0.13 U
PCB 209		0.10 U	0.10 U	0.10 U	0.11 U
<u>Surrogate Recoveries (%)</u>					
PCB 103 (SIS)		68	82	86	104
PCB 198 (SIS)		106	79	79	110

TABLE G.6. (contd)

Matrix Spike Results

Treatment	Matrix Spike				Matrix Spike			
	COMP SB-A		COMP SB-A		COMP EC-A		COMP EC-A	
	1	1	Amount Spiked ng/g	Percent Recovery	1	1	Amount Spiked ng/g	Percent Recovery
	Replicate	Batch			5	5		
Wet Wt. Units	20.08 ng/g	20.02 ng/g	20.08 ng/g	20.05 ng/g	20.08 ng/g	20.05 ng/g		
Heptachlor	1.39	2.45	2.50	42 ^(b)	0.19 U	3.10	2.50	124 ^(b)
Aldrin	1.57	3.16	2.50	64	2.08	2.72	2.50	116
Heptachlor epoxide	0.13 U	2.10	2.50	84	0.13 U	2.33	2.50	93
2,4'-DDE	0.26 U	NA ^(c)	NS ^(d)	NA	0.26 U	NA	NS	NA
Endosulfan I	0.18 U	1.96	2.50	78	0.18 U	2.23	2.50	89
α-Chlordane	0.84	NA	NS	NA	1.29	NA	NS	NA
Trans Nonachlor	0.83	NA	NS	NA	1.40	NA	NS	NA
4,4'-DDE	5.68	8.14	2.50	98	2.68	7.38	2.50	188 ^(b)
Dieldrin	2.56	4.63	2.50	83	1.58	6.23	2.50	186 ^(b)
2,4'-DDD	2.52	NA	NS	NA	0.25 U	NA	NS	NA
2,4'-DDT	0.18 U	NA	NS	NA	0.18 U	NA	NS	NA
4,4'-DDD	14.4	19.3	2.50	196 ^(b)	2.16	13.2	2.50	442 ^(b)
Endosulfan II	0.18 U	1.50	2.50	60	0.18 U	1.52	2.50	61
4,4'-DDT	0.15 U	2.59	2.50	104	0.15 U	2.55	2.50	102
Endosulfan Sulfate	0.18 U	1.95	2.50	78	0.18 U	1.72	2.50	69
PCB 8	0.41 U	NA	NS	NA	0.41 U	NA	NS	NA
PCB 18	11.8	NA	NS	NA	1.58	NA	NS	NA
PCB 28	14.5	21.1	3.18	208 ^(b)	3.24	9.65	3.18	202 ^(b)
PCB 52	17.0	30.4	6.65	202 ^(b)	5.08	19.5	6.65	217 ^(b)
PCB 49	10.0	NA	NS	NA	3.10	NA	NS	NA
PCB 44	6.29	NA	NS	NA	1.28	NA	NS	NA
PCB 66	14.3	NA	NS	NA	0.09 U	NA	NS	NA
PCB 101	10.6	17.7	4.51	157 ^(b)	5.24	18.2	4.51	287 ^(b)
PCB 87	1.71	NA	NS	NA	0.48	6.62	5.70	108
PCB 118	5.18	NA	NS	NA	2.84	NA	NS	NA
PCB 184	0.24 U	NA	NS	NA	0.24 U	NA	NS	NA
PCB 153	6.10	9.64	2.64	134 ^(b)	5.61	12.0	2.64	242 ^(b)
PCB 105	2.52	NS	NS	NS	1.33	NS	NS	NS
PCB 138	5.36	9.10	2.04	183 ^(b)	4.40	14.6	2.04	500 ^(b)
PCB 187	1.79	NA	NS	NA	1.56	NA	NS	NA
PCB 183	0.90	NA	NS	NA	0.74	NA	NS	NA
PCB 128	1.05	NA	NS	NA	0.69	NA	NS	NA
PCB 180	3.21	NA	NS	NA	2.34	NA	NS	NA
PCB 170	1.55	NA	NS	NA	1.13	NA	NS	NA
PCB 195	0.31	NA	NS	NA	0.10 U	NA	NS	NA
PCB 206	1.85	NA	NS	NA	0.50	NA	NS	NA
PCB 209	0.92	NA	NS	NA	0.21	NA	NS	NA
<u>Surrogate Recoveries (%)</u>								
PCB 103 (SIS)	73	49	NA	NA	86	94	NA	NA
PCB 198 (SIS)	131	83	NA	NA	78	87	NA	NA

TABLE G.6. (contd)

Matrix Spike Results								
Treatment Replicate Batch Wet Wt. Units	Matrix Spike				Matrix Spike			
	C-NV	C-NV			COMP HU-C	COMP HU-C	Amount Spiked ng/g	Percent Recovery
	2	2	Amount	Percent	1	1		
	6	6	Spiked	Recovery	7	7		
20.08 ng/g	20.17 ng/g	12.96 ng/g	12.71 ng/g					
Heptachlor	0.19 U	2.71	2.50	108	0.28 U	4.76	3.95	121 ^(b)
Aldrin	0.13 U	2.23	2.50	89	1.77	4.88	3.95	79
Heptachlor epoxide	0.13 U	2.48	2.50	99	0.20 U	3.45	3.95	87
2,4'-DDE	0.26 U	NA	NS	NA	0.40 U	NA	NS	NA
Endosulfan I	0.18 U	2.40	2.50	96	0.28 U	2.93	3.95	74
α -Chlordane	0.10 U	NA	NS	NA	2.21	NA	NS	NA
Trans Nonachlor	0.60	NA	NS	NA	0.68	NA	NS	NA
4,4'-DDE	0.29	2.11	2.50	73	3.87	7.30	3.95	87
Dieldrin	0.93	2.96	2.50	81	2.50	6.10	3.95	91
2,4'-DDD	0.40	NA	NS	NA	0.39 U	NA	NS	NA
2,4'-DDT	0.18 U	NA	NS	NA	0.28 U	NA	NS	NA
4,4'-DDD	0.83	3.5	2.50	105	4.66	10.1	3.95	138
Endosulfan II	0.18 U	1.71	2.50	68	0.28 U	3.00	3.95	76
4,4'-DDT	0.15 U	2.31	2.50	92	0.23 U	4.23	3.95	107
Endosulfan Sulfate	0.18 U	2.23	2.50	89	0.28 U	3.71	3.95	94
PCB 8	0.41 U	NA	NS	NA	0.63 U	NA	NS	NA
PCB 18	0.43 U	NA	NS	NA	9.95	NA	NS	NA
PCB 28	0.20 U	3.98	3.19	118	14.30	21.78	5.04	148 ^(b)
PCB 52	0.52	7.4	6.65	104	19.31	31.6	10.51	117
PCB 49	0.24 U	NA	NS	NA	10.00	NA	NS	NA
PCB 44	0.17 U	NA	NS	NA	4.98	NA	NS	NA
PCB 66	0.09 U	NA	NS	NA	15.27	NA	NS	NA
PCB 101	0.78	5.7	4.51	109	9.92	19.7	7.13	137 ^(b)
PCB 87	0.16 U	NA	NS	NA	0.88	NA	NS	NA
PCB 118	0.45	NA	NS	NA	5.30	NA	NS	NA
PCB 184	0.24 U	NA	NS	NA	0.36 U	NA	NS	NA
PCB 153	2.20	4.5	2.64	88	7.80	11.3	4.17	83
PCB 105	0.33	NA	NS	NA	3.38	NA	NS	NA
PCB 138	1.42	5.6	2.04	202 ^(b)	7.19	10.4	3.22	99
PCB 187	0.62	NA	NS	NA	2.51	NA	NS	NA
PCB 183	0.25	NA	NS	NA	1.21	NA	NS	NA
PCB 128	0.25	NA	NS	NA	1.28	NA	NS	NA
PCB 180	0.72	NA	NS	NA	3.05	NA	NS	NA
PCB 170	0.38	NA	NS	NA	1.45	NA	NS	NA
PCB 195	0.10 U	NA	NS	NA	0.22	NA	NS	NA
PCB 206	0.27	NA	NS	NA	1.23	NA	NS	NA
PCB 209	0.16	NA	NS	NA	0.82	NA	NS	NA
<u>Surrogate Recoveries (%)</u>								
PCB 103 (SIS)	87	83	NA	NA	64	77	NA	NA
PCB 198 (SIS)	69	61	NA	NA	68	80	NA	NA

TABLE G.6. (contd)

Analytical Replicate Results

Treatment Replicate Batch Wet Wt. Units	<i>DUP</i>		<i>TRIP</i>		<i>DUP</i>		<i>TRIP</i>	
	COMP HU-A	COMP HU-A	COMP HU-A	RSD%	COMP SB-B	COMP SB-B	COMP SB-B	RSD%
	5	5	5		2	2	2	
	4	4	4		5	5	5	
	14.57	13.76	13.79		17.11	17.25	17.13	
	ng/g	ng/g	ng/g		ng/g	ng/g	ng/g	
Heptachlor	1.02	0.89	1.00	7	0.21 U	0.21 U	0.21 U	NA
Aldrin	3.64	3.48	3.65	3	1.67	1.72	1.64	2
Heptachlor epoxide	0.18 U	0.19 U	0.19 U	NA	0.15 U	0.24	0.15 U	NA
2,4'-DDE	0.36 U	0.38 U	0.38 U	NA	0.3 U	0.3 U	0.3 U	NA
Endosulfan I	0.25 U	0.26 U	0.26 U	NA	0.21 U	0.21 U	0.21 U	NA
α -Chlordane	0.13 U	0.14 U	0.14 U	NA	0.8	0.89	0.85	5
Trans Nonachlor	0.54	0.21 U	0.21 U	NA	0.86	0.96	0.94	6
4,4'-DDE	6.42	6.41	6.43	0	1.9	2.05	1.95	4
Dieldrin	2.00	1.69	1.85	8	1.80	1.9	1.81	3
2,4'-DDD	0.93	1.12	1.38	20	5.42	5.91	5.86	5
2,4'-DDT	0.25 U	0.26 U	0.26 U	NA	0.21 U	0.21 U	0.21 U	NA
4,4'-DDD	6.97	6.32	6.62	5	10.30	11.7	12	8
Endosulfan II	0.25 U	0.26 U	0.26 U	NA	0.21 U	0.21 U	0.21 U	NA
4,4'-DDT	0.21 U	0.22 U	0.22 U	NA	0.18 U	2.33	0.18 U	NA
Endosulfan Sulfate	0.25 U	0.26 U	0.44	34 ^(e)	0.65	0.45	0.3	38 ^(e)
PCB 8	0.57 U	0.60 U	0.60 U	NA	0.48 U	0.48 U	0.48 U	NA
PCB 18	8.28	8.45	8.44	1	1.18	1.34	1.21	7
PCB 28	8.87	8.92	9.03	1	2.39	2.46	2.30	3
PCB 52	9.39	9.06	9.43	2	4.22	4.32	3.85	6
PCB 49	5.31	5.21	5.38	2	2.23	2.27	2.07	5
PCB 44	3.08	3.02	3.05	1	0.79	0.86	0.86	5
PCB 66	0.13 U	0.14 U	0.14 U	NA	0.11 U	0.11 U	0.11 U	NA
PCB 101	5.04	4.93	5.10	2	4.37	4.52	4.09	5
PCB 87	0.91	0.99	0.82	9	0.19 U	0.28	0.33	27
PCB 118	2.51	2.44	2.54	2	2.79	2.72	2.23	12
PCB 184	0.33 U	0.34 U	0.34 U	NA	0.27 U	0.27 U	0.27 U	NA
PCB 153	4.40	4.40	4.47	1	5.28	5.19	4.11	13
PCB 105	1.25	1.11	1.18	6	1.42	1.41	1.16	11
PCB 138	2.92	2.91	2.91	0	4.06	4.1	3.41	10
PCB 187	1.39	1.32	1.36	3	1.32	1.29	1.03	13
PCB 183	0.65	0.54	0.60	9	0.62	0.6	0.48	13
PCB 128	0.60	0.50	0.56	9	0.69	0.69	0.56	12
PCB 180	1.71	1.69	1.65	2	1.94	2.01	1.78	6
PCB 170	0.23 U	0.24 U	0.24 U	NA	0.98	1.01	0.88	7
PCB 195	0.17	0.17	0.15 U	NA	0.17	0.12 U	0.12 U	NA
PCB 206	1.25	1.29	1.24	2	0.49	0.51	0.42	10
PCB 209	0.87	0.77	0.83	6	0.32	0.31	0.25	13
<u>Surrogate Recoveries (%)</u>								
PCB 103 (SIS)	75	74	66	NA	65	81	72	NA
PCB 198 (SIS)	116	115	102	NA	61	73	66	NA

TABLE G.6. (contd)

Analytical Replicate Results

Treatment	DUP		TRIP		DUP		TRIP	
	COMP HU-C	COMP HU-C	COMP HU-C	COMP HU-C	COMP BU	COMP BU	COMP BU	COMP BU
Replicate	4	4	4		3	3	3	
Batch	6	6	6		7	7	7	
Wet Wt.	17.18	17.51	16.38		8.6	8.47	8.21	
Units	ng/g	ng/g	ng/g	RSD%	ng/g	ng/g	ng/g	RSD%
Heptachlor	2.5	2.43	2.33	4	0.43 U	0.44 U	0.45 U	NA
Aldrin	2.42	2.25	2.29	4	2.42	2.74	2.2	11
Heptachlor epoxide	0.15 U	0.15 U	0.16 U	NA	0.31 U	0.31 U	0.32 U	NA
2,4'-DDE	0.3 U	0.3 U	0.32 U	NA	0.61 U	0.62 U	0.64 U	NA
Endosulfan I	0.21 U	0.21 U	0.22 U	NA	0.42 U	0.42 U	0.44 U	NA
a-Chlordane	1.83	1.78	1.66	5	1.13	1.46	1.11	16
Trans Nonachlor	1.65	1.61	1.52	4	0.54	0.77	0.35 U	NA
4,4'-DDE	16.8	7.5	6.89	53 ^(e)	2.01	2.54	2.23	12
Dieldrin	0.60 U	4.31	4.16	69 ^(e)	1.43	1.84	1.58	13
2,4'-DDD	7.71	7.61	7.11	4	0.59 U	0.60 U	0.62 U	NA
2,4'-DDT	0.21 U	0.2 U	0.22 U	NA	0.42 U	0.42 U	0.44 U	NA
4,4'-DDD	26.00	22.5	21.3	10	2.24	2.56	1.85	16
Endosulfan II	0.21 U	0.21 U	0.22 U	NA	0.42 U	0.42 U	0.44 U	NA
4,4'-DDT	0.18 U	0.17 U	0.18 U	NA	0.35 U	0.36 U	0.37 U	NA
Endosulfan Sulfate	0.21 U	0.21 U	0.22 U	NA	0.42 U	0.75	0.44 U	NA
PCB 8	0.48 U	0.47 U	0.50 U	3	0.95 U	0.97 U	1.00 U	NA
PCB 18	19.8	19.3	18.5	3	1 U	1.01 U	1.05 U	NA
PCB 28	25.70	24.30	23.80	4	2.34	3.19	2.54	17
PCB 52	37.10	34.00	31.8	8	3.94	5.27	4.37	15
PCB 49	17.80	16.7	16.5	4	2.09	2.79	2.14	17
PCB 44	11.60	10.6	9.58	10	1.07	1.44	1.18	15
PCB 66	27.20	25.10	24.1	6	0.22 U	0.22 U	0.23 U	NA
PCB 101	20.80	19.3	18.70	6	3.09	4.17	3.26	17
PCB 87	20.60	2.04	1.82	132 ^(e)	0.37 U	0.41	0.39 U	NA
PCB 118	18.40	10.5	9.87	37 ^(e)	1.51	2.05	1.68	16
PCB 184	0.27 U	0.27 U	0.29 U	NA	0.55 U	0.56 U	0.58 U	NA
PCB 153	17.90	13.60	12.8	19	3.89	5.28	4.33	16
PCB 105	6.30	5.72	5.38	8	0.95	1.33	1.08	17
PCB 138	13.30	12	11.5	8	3.06	4.33	3.44	18
PCB 187	3.62	3.2	3	10	0.99	1.51	1.13	22
PCB 183	1.85	1.68	1.57	8	0.55 U	0.65	0.58 U	NA
PCB 128	2.64	2.46	2.27	8	0.52	0.68	0.56	14
PCB 180	3.77	4.79	4.46	12	1.39	1.97	1.55	18
PCB 170	2.44	2.44	2.25	5	0.73	0.96	0.79	14
PCB 195	0.25	0.39	0.12 U	NA	0.23 U	0.24 U	0.24 U	NA
PCB 206	1.53	1.24	1.14	16	0.42	0.57	0.45	17
PCB 209	0.92	0.90	0.88	2	0.23	0.31	0.26	15
<u>Surrogate Recoveries (%)</u>								
PCB 103 (SIS)	89	82	88	NA	81	66	74	NA
PCB 198 (SIS)	81	67	70	NA	83	67	79	NA

(a) U Undetected at or above given concentration.

(b) Outside Spike QC range (50-120%) for matrix spike recoveries

(c) NA Not applicable.

(d) NS Not spiked.

(e) Exceeds quality control criteria ($\pm 30\%$) for replicates.

TABLE G.7. MDL Verification Study for Pesticide/PCB Tissue Chemistry

Treatment	MDL	MDL	MDL	MDL	
Replicate	R1	R2	R3	R4	
Batch	8	8	8	8	
Wet Wt.	20.12	20.40	20.09	20.03	
Units	ng/g	ng/g	ng/g	ng/g	MDL ^(a)
Heptachlor	1.01	1.08	1.09	1.04	0.129
Aldrin	0.82	0.79	0.83	0.82	0.061
Heptachlor Epoxide	1.32	1.27	1.33	1.28	0.103
2,4'-DDE	1.18	1.2	1.24	1.19	0.092
Endosulfan I	NA ^(b)	NA	NA	NA	NA
α -Chlordane	0.94	0.96	0.95	1.1	0.264
Trans Nonachlor	1.43	1.49	1.46	1.61	0.276
4,4'-DDE	1.87	1.62	1.77	1.78	0.363
Dieldrin	2.27	2.38	2.39	2.32	0.196
2,4'-DDD	1.40	1.52	1.52	1.52	0.210
2,4'-DDT	1.07	1.02	1.17	1.18	0.273
4,4'-DDD	1.40	1.52	1.67	1.68	0.467
Endosulfan II	NA	NA	NA	NA	NA
4,4'-DDT	1.04	1.18	1.13	1.25	0.309
Endosulfan Sulfate	NA	NA	NA	NA	NA
PCB 8	0.56	0.57	0.54	0.56	0.044
PCB 18	0.84	0.80	0.85	0.84	0.078
PCB 28	1.04	1.01	1.07	1.10	0.136
PCB 52	1.20	1.20	1.27	1.31	0.191
PCB 49	0.24 U ^(c)	0.23 U	0.24 U	0.24 U	NA
PCB 44	0.96	0.90	0.93	0.94	0.088
PCB 66	1.47	1.42	1.47	1.44	0.086
PCB 101	1.59	1.54	1.62	1.55	0.129
PCB 87	0.79	0.81	0.79	0.97	0.305
PCB 118	1.02	1.00	1.05	1.10	0.152
PCB 184	0.24 U	0.23 U	0.24 U	0.24 U	NA
PCB 153	2.54	2.46	2.61	2.60	0.241
PCB 105	1.00	0.95	1.03	1.04	0.141
PCB 138	1.91	1.89	1.89	1.96	0.116
PCB 187	1.24	1.23	1.24	1.35	0.199
PCB 183	0.24 U	0.23 U	0.24 U	0.24 U	NA
PCB 128	0.87	0.87	0.88	0.92	0.083
PCB 180	1.18	1.34	1.22	1.17	0.273
PCB 170	0.98	0.93	1.01	1.03	0.152
PCB 195	0.82	0.80	0.84	0.89	0.135
PCB 206	1.03	1.01	1.09	1.13	0.193
PCB 209	1.00	0.95	1.03	1.06	0.164

(a) MDL Calculated by multiplying the standard deviation of the four replicates by Students-t (4.54).

(b) NA Not applicable.

(c) U Undetected at or above given concentration.

TABLE G.8. Polynuclear Aromatic Hydrocarbons (PAH) and 1,4-Dichlorobenzene (Wet Weight) in Tissue of *N. virens*

Treatment	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC
Replicate	1	2	3	4	5
Batch	6	7	4	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.07%	16.08%	13.88%	15.67%	15.11%
1,4-Dichlorobenzene	1.86 U ^(a)	1.86 U	1.86 U	2.05 U	1.86 U
Naphthalene	2.14 ^(b)	6.48 B	2.38 ^(b)	6.78 B	2.92
Acenaphthylene	2.20 ^(b)	1.20 ^(b)	0.84 ^(b)	1.09 ^(b)	2.52 ^(b)
Acenaphthene	13.4	16.5	6.99	16.5	19.5
Fluorene	3.38 ^(b)	4.20	1.24 U	3.91 ^(b)	3.43
Phenanthrene	4.26	7.08	2.82	6.16	4.80
Anthracene	4.57 ^(b)	5.06	3.84	5.08	5.19
Fluoranthene	124	96.6	72.0	91.7	117
Pyrene	84.0	68.6	48.0	62.2	86.2
Benzo(a)anthracene	8.05 B ^(c)	8.01 B	4.94 ^(b) B	7.73 B	8.39 B
Chrysene	30.3	24.5	19.7	24.9	32.8
Benzo(b)fluoranthene	8.25 ^(b)	8.28 ^(b)	5.39 ^(b)	7.76 ^(b)	9.25
Benzo(k)fluoranthene	5.46	5.17 ^(b)	3.97	4.85	5.58
Benzo(a)pyrene	5.13	5.05 ^(b) B	2.73 ^(b)	4.59	5.50
Indeno(123-cd)pyrene	3.86 ^(b)	4.34 B	1.76 U	3.94 ^(b) B	3.92
Dibenzo(a,h)anthracene	1.88 ^(b)	2.19 ^(b)	1.26 U	2.29 ^(b) B	2.02 ^(b)
Benzo(g,h,i)perylene	3.95	4.18	2.26	4.15 ^(b)	4.74
<u>Surrogate Internal Standards (</u>					
d4 1,4-Dichlorobenzene	72	39	48	42	69
d8 Naphthalene	86	53	65	57	80
d10 Acenaphthene	93	72	76	74	87
d12 Chrysene	86	70	72	78	83
d14 Dibenzo(a,h,i)anthracene	95	90	80	98	92

TABLE G.8. (contd)

Treatment	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	1	2	3	4	5
Batch	4	5	6	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.12%	14.94%	15.21%	14.00%	13.24%
1,4-Dichlorobenzene	1.86 U	1.83 U	1.86 U	1.86 U	2.31 U
Naphthalene	1.86 U	1.83 U	2.71 ^(b)	6.00 ^{(b)B}	11.9
Acenaphthylene	0.73 U	0.71 U	0.73 U	0.73 U	2.93 ^(b)
Acenaphthene	1.30 U	1.28 U	2.28 ^(b)	3.24	3.29
Fluorene	1.24 U	1.21 U	1.24 U	3.31	4.07
Phenanthrene	2.56 U	2.51 U	2.56 U	4.04	7.21
Anthracene	2.24 U	2.19 U	2.24 U	2.24 U	2.77 U
Fluoranthene	5.36 U	5.26 U	5.36 U	5.36 U	6.65 U
Pyrene	4.57 U	4.48 U	4.57 U	5.54 ^(b)	6.97 ^(b)
Benzo(a)anthracene	2.43 ^{(b)B}	2.47 B	3.68 ^{(b)B}	4.05 ^{(b)B}	4.51 ^{(b)B}
Chrysene	2.27 U	2.22 U	2.27 U	2.27 U	2.81 U
Benzo(b)fluoranthene	2.51 ^(b)	1.61 U	4.09 ^(b)	1.64 U	5.09 ^(b)
Benzo(k)fluoranthene	1.92 ^(b)	1.64 U	1.67 U	1.67 U	2.07 U
Benzo(a)pyrene	1.49 U	1.46 U	1.49 U	1.49 U	1.85 U
Indeno(123-cd)pyrene	1.76 U	1.73 U	1.76 U	1.76 U	3.66 ^(b)
Dibenzo(a,h)anthracene	1.26 U	1.24 U	1.26 U	1.26 U	1.56 U
Benzo(g,h,i)perylene	1.40 U	1.37 U	1.40 U	1.40 U	3.57 ^(b)
<u>Surrogate Internal Standards (%)</u>					
d4 1,4-Dichlorobenzene	69	63	64	12 ^(d)	66
d8 Naphthalene	82	85	76	28 ^(d)	76
d10 Acenaphthene	83	92	81	47	79
d12 Chrysene	72	93	77	54	78
d14 Dibenzo(a,h,i)anthracene	82	102	86	70	87

TABLE G.8. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	6	5	4	6	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.70%	16.08%	15.15%	14.02%	14.53%
1,4-Dichlorobenzene	1.86 U	1.86 U	1.83 U	1.83 U	1.86 U
Naphthalene	2.33 ^(b)	1.86 U	2.46	2.59 ^(b)	1.86 U
Acenaphthylene	0.73 U	0.73 U	0.71 U	0.71 U	0.73 U
Acenaphthene	2.47	1.30 U	1.28 U	2.60 ^(b)	1.30 U
Fluorene	1.24 U	1.24 U	1.21 U	1.21 U	1.24 U
Phenanthrene	2.56 U	2.56 U	2.51 U	2.64 ^(b)	2.56 U
Anthracene	2.24 U	2.24 U	2.19 U	2.19 U	2.24 U
Fluoranthene	5.36 U	5.36 U	5.26 U	5.26 U	5.36 U
Pyrene	6.36	4.57 U	4.48 U	5.54 ^(b)	4.57 U
Benzo(a)anthracene	3.32 ^{(b)B}	1.09 U	2.15 ^{(b)B}	1.07 U	2.11 ^{(b)B}
Chrysene	2.62	2.27 U	2.22	2.42	2.27 U
Benzo(b)fluoranthene	4.53 ^(b)	2.61 ^(b)	2.75	4.32	2.42 ^(b)
Benzo(k)fluoranthene	3.14 ^(b)	1.97 ^(b)	2.06 ^(b)	2.81 ^(b)	1.83 ^(b)
Benzo(a)pyrene	2.29 ^(b)	1.49 U	1.46 U	1.46 U	1.49 U
Indeno(123-cd)pyrene	3.01 ^(b)	1.76 U	1.73 U	2.86 ^(b)	1.76 U
Dibenzo(a,h)anthracene	1.26 U	1.26 U	1.24 U	1.24 U	1.26 U
Benzo(g,h,i)perylene	2.91 ^(b)	1.40 U	1.37 U	2.75 ^(b)	1.40 U
<u>Surrogate Internal Standards (%)</u>					
d4 1,4-Dichlorobenzene	66	71	40	63	63
d8 Naphthalene	81	93	51	74	84
d10 Acenaphthene	88	99	55	79	95
d12 Chrysene	81	98	52	78	102
d14 Dibenzo(a,h,i)anthracene	93	103	55	85	103

TABLE G.8. (contd)

Treatment	C-NV	C-NV	C-NV	C-NV	C-NV
Replicate	1	2	3	4	5
Batch	6	6	4	4	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.84%	12.32%	14.51%	13.67%	14.91%
1,4-Dichlorobenzene	1.86 U	1.86 U	1.86 U	1.86 U	1.86 U
Naphthalene	2.16 ^(b)	2.72 ^(b)	2.49	2.80	2.09 ^(b)
Acenaphthylene	2.04 ^(b)	0.73 U	0.73 U	0.73 U	0.73 U
Acenaphthene	1.30 U	2.34 ^(b)	1.30 U	1.40 ^(b)	1.30 U
Fluorene	1.24 U	2.76	1.24 U	1.24 U	1.24 U
Phenanthrene	2.56 ^(b)	2.76 ^(b)	2.56 U	2.56 U	2.56 U
Anthracene	2.24 U	2.24 U	2.24 U	2.24 U	2.24 U
Fluoranthene	7.87 ^(b)	6.80	11.1	5.46	5.36 U
Pyrene	9.30	7.20	14.7	4.95	5.01 ^(b)
Benzo(a)anthracene	3.95 B	1.09 U	2.45 ^(b) B	2.26 ^(b) B	1.09 U
Chrysene	3.21	2.87	3.77	2.27 U	2.27 U
Benzo(b)fluoranthene	5.00	4.44 ^(b)	3.53	2.60	2.70 ^(b)
Benzo(k)fluoranthene	3.19 ^(b)	2.81 ^(b)	2.48 ^(b)	2.02 ^(b)	2.05 ^(b)
Benzo(a)pyrene	2.64 ^(b)	1.49 U	1.49 U	1.49	1.49 U
Indeno(123-cd)pyrene	3.07 ^(b)	2.87 ^(b)	1.76 U	1.76 ^(b)	1.76 U
Dibenzo(a,h)anthracene	1.26 U	1.26 U	1.26 U	1.26	1.26 U
Benzo(g,h,i)perylene	2.96 ^(b)	2.78 ^(b)	1.40 U	1.40 ^(b)	1.40 U
<u>Surrogate Internal Standards (%)</u>					
d4 1,4-Dichlorobenzene	68	71	46	55	27 ^(d)
d8 Naphthalene	82	85	58	71	35
d10 Acenaphthene	89	88	63	76	38
d12 Chrysene	78	80	58	71	41
d14 Dibenzo(a,h,i)anthracene	85	92	61	77	38

TABLE G.8. (contd)

Treatment	<i>N. virens</i> Background	<i>N. virens</i> Background	<i>N. virens</i> Background
Replicate	1	2	3
Batch	7	7	7
Units	ng/g	ng/g	ng/g
Percent Dry Weight	12.86%	12.94%	12.05%
1,4-Dichlorobenzene	1.86 U	1.86 U	1.86 U
Naphthalene	2.79	2.67	2.98
Acenaphthylene	0.73 U	2.79 U	0.73 U
Acenaphthene	2.12	2.24 ^(b)	2.09 ^(b)
Fluorene	1.24 U	1.24 U	1.24 U
Phenanthrene	2.56 U	2.56 U	2.67 ^(b)
Anthracene	3.49	2.24 U	2.24 U
Fluoranthene	5.36 U	5.36 U	5.36 U
Pyrene	4.57 U	4.57 U	4.57 U
Benzo(a)anthracene	4.22	3.86 ^(b)	3.77 ^(b)
Chrysene	2.27 U	2.27 U	2.27 U
Benzo(b)fluoranthene	1.64 U	1.64 U	4.49 ^(b)
Benzo(k)fluoranthene	1.67 U	1.67 U	1.67 U
Benzo(a)pyrene	1.49 U	2.59	1.49 U
Indeno(123-cd)pyrene	1.76 U	1.76 U	1.76 U
Dibenzo(a,h)anthracene	1.26 U	1.26 U	1.26 U
Benzo(g,h,i)perylene	1.40 U	1.40 U	1.40 U
<u>Surrogate Internal Standards (%)</u>			
d4 1,4-Dichlorobenzene	72	68	51
d8 Naphthalene	85	82	67
d10 Acenaphthene	91	89	84
d12 Chrysene	84	81	82
d14 Dibenzo(a,h,i)anthracene	105	103	104

(a) U Undetected at or above given concentration.

(b) Ion ratio out or confirmation ion not detected.

(c) B Value is < 5 times concentration in blank.

(d) Outside quality control criteria (30-150%) for surrogate internal standards.

TABLE G.9. Polynuclear Aromatic Hydrocarbons (PAH) and 1,4-Dichlorobenzene (Dry Weight) in Tissue of *N. virens*

Treatment	COMP PC	COMP PC	COMP PC	COMP PC	COMP PC
Replicate	1	2	3	4	5
Batch	6	7	4	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.07%	16.08%	13.88%	15.67%	15.11%
1,4-Dichlorobenzene	13.2 U ^(a)	11.6 U	13.4 U	13.1 U	12.3 U
Naphthalene	15.2 ^(b)	40.3 B	17.1 ^(b)	43.3 B	19.3
Acenaphthylene	15.6 ^(b)	7.46 ^(b)	6.05 ^(b)	6.96 ^(b)	16.7 ^(b)
Acenaphthene	95.2	103	50.4	105	129
Fluorene	24.0 ^(b)	26.1	8.93 U	25.0 ^(b)	22.7
Phenanthrene	30.3	44.0	20.3	39.3	31.8
Anthracene	32.5 ^(b)	31.5	27.7	32.4	34.3
Fluoranthene	881	601	519	585	774
Pyrene	597	427	346	397	570
Benzo(a)anthracene	57.2 B ^(c)	49.8 B	35.6 ^(b) B	49.3 B	55.5 B
Chrysene	215	152	142	159	217
Benzo(b)fluoranthene	58.6 ^(b)	51.5 ^(b)	38.8 ^(b)	49.5 ^(b)	61.2
Benzo(k)fluoranthene	38.8	32.2 ^(b)	28.6	31.0	36.9
Benzo(a)pyrene	36.5	31.4 ^(b) B	19.7 ^(b)	29.3	36.4
Indeno(123-cd)pyrene	27.4 ^(b)	27.0 B	12.7 U	25.1 ^(b) B	25.9
Dibenzo(a,h)anthracene	13.4 ^(b)	13.6 ^(b)	9.08 U	14.6 ^(b) B	13.4 ^(b)
Benzo(g,h,i)perylene	28.1	26.0	16.3	26.5 ^(b)	31.4

TABLE G.9. (contd)

Treatment	R-MUD	R-MUD	R-MUD	R-MUD	R-MUD
Replicate	1	2	3	4	5
Batch	4	5	6	7	6
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.12%	14.94%	15.21%	14.00%	13.24%
1,4-Dichlorobenzene	14.2 U	12.2 U	12.2 U	13.3 U	17.4 U
Naphthalene	14.2 U	12.2 U	17.8 ^(b)	42.9 ^{(b)B}	89.9
Acenaphthylene	5.56 U	4.8 U	4.8 U	5.2 U	22.1 ^(b)
Acenaphthene	9.91 U	8.57 U	15.0 ^(b)	23.1	24.8
Fluorene	9.45 U	8.10 U	8.15 U	23.6	30.7
Phenanthrene	19.5 U	16.8 U	16.8 U	28.9	54.5
Anthracene	17.1 U	14.7 U	14.7 U	16.0 U	20.9 U
Fluoranthene	40.9 U	35.2 U	35.2 U	38.3 U	50.2 U
Pyrene	34.8 U	30.0 U	30.0 U	39.6 ^(b)	52.6 ^(b)
Benzo(a)anthracene	18.5 ^{(b)B}	16.5 B	24.2 ^{(b)B}	28.9 ^{(b)B}	34.1 ^{(b)B}
Chrysene	17.3 U	14.9 U	14.9 U	16.2 U	21.2 U
Benzo(b)fluoranthene	19.1 ^(b)	10.8 U	26.9 ^(b)	11.7 U	38.4 ^(b)
Benzo(k)fluoranthene	14.6 ^(b)	11.0 U	11.0 U	11.9 U	15.6 U
Benzo(a)pyrene	11.4 U	9.77 U	9.80 U	10.6 U	14.0 U
Indeno(123-cd)pyrene	13.4 U	11.6 U	11.6 U	12.6 U	27.6 ^(b)
Dibenzo(a,h)anthracene	9.60 U	8.30 U	8.28 U	9.00 U	11.8 U
Benzo(g,h,i)perylene	10.7 U	9.17 U	9.20 U	10.0 U	27.0 ^(b)

TABLE G.9. (contd)

Treatment	R-CLIS	R-CLIS	R-CLIS	R-CLIS	R-CLIS
Replicate	1	2	3	4	5
Batch	6	5	4	6	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	13.70%	16.08%	15.15%	14.02%	14.53%
1,4-Dichlorobenzene	13.6 U	11.6 U	12.1 U	13.1 U	12.8 U
Naphthalene	17.0 ^(b)	11.6 U	16.2	18.5 ^(b)	12.8 U
Acenaphthylene	5.3 U	4.5 U	4.7 U	5.1 U	5.0 U
Acenaphthene	18.0	8.08 U	8.45 U	18.5 ^(b)	8.95 U
Fluorene	9.05 U	7.71 U	7.99 U	8.63 U	8.53 U
Phenanthrene	18.7 U	15.9 U	16.6 U	18.8 ^(b)	17.6 U
Anthracene	16.4 U	13.9 U	14.5 U	15.6 U	15.4 U
Fluoranthene	39.1 U	33.3 U	34.7 U	37.5 U	36.9 U
Pyrene	46.4	28.4 U	29.6 U	39.5 ^(b)	31.5 U
Benzo(a)anthracene	24.2 ^(b) B	6.78 U	14.2 ^(b) B	7.63 U	14.5 ^(b) B
Chrysene	19.1	14.1 U	14.7	17.3	15.6 U
Benzo(b)fluoranthene	33.1 ^(b)	16.2 ^(b)	18.2	30.8	16.7 ^(b)
Benzo(k)fluoranthene	22.9 ^(b)	12.3 ^(b)	13.6 ^(b)	20.0 ^(b)	12.6 ^(b)
Benzo(a)pyrene	16.7 ^(b)	9.27 U	9.64 U	10.4 U	10.3 U
Indeno(123-cd)pyrene	22.0 ^(b)	10.9 U	11.4 U	20.4 ^(b)	12.1 U
Dibenzo(a,h)anthracene	9.20 U	7.84 U	8.18 U	8.84 U	8.67 U
Benzo(g,h,i)perylene	21.2 ^(b)	8.7 U	9.04 U	19.6 ^(b)	9.64 U

TABLE G.9. (contd)

Treatment	C-NV	C-NV	C-NV	C-NV	C-NV
Replicate	1	2	3	4	5
Batch	6	6	4	4	4
Units	ng/g	ng/g	ng/g	ng/g	ng/g
Percent Dry Weight	14.84%	12.32%	14.51%	13.67%	14.91%
1,4-Dichlorobenzene	12.5 U	15.1 U	12.8 U	13.6 U	12.5 U
Naphthalene	14.6 ^(b)	22.1 ^(b)	17.2	20.5	14.0 ^(b)
Acenaphthylene	13.7 ^(b)	5.9 U	5.0 U	5.3 U	4.9 U
Acenaphthene	8.76 U	19.0 ^(b)	9.0 U	10.2 ^(b)	8.72 U
Fluorene	8.36 U	22.4	8.55 U	9.07 U	8.32 U
Phenanthrene	17.3 ^(b)	22.4 ^(b)	17.6 U	18.7 U	17.2 U
Anthracene	15.1 U	18.2 U	15.4 U	16.4 U	15.0 U
Fluoranthene	53.0 ^(b)	55.2	76.5	39.9	35.9 U
Pyrene	62.7	58.4	101	36.2	33.6 ^(b)
Benzo(a)anthracene	26.6 B	8.85 U	16.9 ^(b) B	16.5 ^(b) B	7.31 U
Chrysene	21.6	23.3	26.0	16.6 U	15.2 U
Benzo(b)fluoranthene	33.7	36.0 ^(b)	24.3	19.0	18.1 ^(b)
Benzo(k)fluoranthene	21.5 ^(b)	22.8 ^(b)	17.1 ^(b)	14.8 ^(b)	13.7 ^(b)
Benzo(a)pyrene	17.8 ^(b)	12.1 U	10.3 U	10.9	9.99 U
Indeno(123-cd)pyrene	20.7 ^(b)	23.3 ^(b)	12.1 U	12.9 ^(b)	11.8 U
Dibenzo(a,h)anthracene	8.49 U	10.2 U	8.68 U	9.22	8.45 U
Benzo(g,h,i)perylene	19.9 ^(b)	22.6 ^(b)	9.65 U	10.2 ^(b)	9.39 U

TABLE G.9. (contd)

Treatment	<i>N. virens</i> Background	<i>N. virens</i> Background	<i>N. virens</i> Background
Replicate	1	2	3
Batch	7	7	7
Units	ng/g	ng/g	ng/g
Percent Dry Weight	12.86%	12.94%	12.05%
1,4-Dichlorobenzene	14.5 U	14.4 U	15.4 U
Naphthalene	21.7	20.6	24.7
Acenaphthylene	5.7 U	21.6 U	6.1 U
Acenaphthene	16.5	17.3 ^(b)	17.3 ^(b)
Fluorene	9.64 U	9.58 U	10.3 U
Phenanthrene	19.9 U	19.8 U	22.2 ^(b)
Anthracene	27.1	17.3 U	18.6 U
Fluoranthene	41.7 U	41.4 U	44.5 U
Pyrene	35.5 U	35.3 U	37.9 U
Benzo(a)anthracene	32.8	29.8 ^(b)	31.3 ^(b)
Chrysene	17.7 U	17.5 U	18.8 U
Benzo(b)fluoranthene	12.8 U	12.7 U	37.3 ^(b)
Benzo(k)fluoranthene	13.0 U	12.9 U	13.9 U
Benzo(a)pyrene	11.6 U	20.0	12.4 U
Indeno(123-cd)pyrene	13.7 U	13.6 U	14.6 U
Dibenzo(a,h)anthracene	9.80 U	9.74 U	10.5 U
Benzo(g,h,i)perylene	10.9 U	10.8 U	11.6 U

(a) U Undetected at or above given concentration.

(b) Ion ratio out or confirmation ion not detected.

(c) B Value is < 5 times concentration in blank.

**TABLE G.10. Quality Control Summary for Polynuclear Aromatic Hydrocarbons (PAHs)
in Tissue of *N. virens* (Wet Weight)**

METHOD BLANKS					
Treatment	BLANK	BLANK	BLANK	BLANK	BLANK
Replicate	1	1	1	1	2
Batch	4	5	6	7	7
Wet Wt.	NA	NA	NA	NA	NA
Units	ng/g	ng/g	ng/g	ng/g	ng/g
1,4-Dichlorobenzene	1.98 U ^(a)	1.90 U	1.94 U	2.24 U	2.16 U
Naphthalene	1.98 U	1.90 U	1.94 U	2.24 U	2.24 ^(b)
Acenaphthylene	0.77 U	0.74 U	0.75 U	0.87 U	0.84 U
Acenaphthene	1.38 U	1.33 U	1.36 U	1.56 U	1.51 U
Fluorene	1.31 U	1.26 U	1.29 U	1.48 U	1.43 U
Phenanthrene	2.71 U	2.61 U	2.66 U	3.07 U	2.97 U
Anthracene	2.37 U	2.28 U	2.33 U	2.69 U	6.22 U
Fluoranthene	5.69 U	5.47 U	5.58 U	6.44 U	5.30 U
Pyrene	4.84 U	4.66 U	4.75 U	5.48 U	5.30 U
Benzo(a)anthracene	2.29	2.13 ^(b)	3.50 ^(b)	4.40 ^(b)	4.41 ^(b)
Chrysene	2.40 U	2.31 U	2.36 U	2.72 U	2.63 U
Benzo(b)fluoranthene	1.74 U	1.67 U	1.71 U	1.97 U	1.90 U
Benzo(k)fluoranthene	1.77 U	1.70 U	1.74 U	2.00 U	1.94 U
Benzo(a)pyrene	1.58 U	1.52 U	1.55 U	2.75	1.73 U
Indeno(123-cd)pyrene	1.87 U	1.80 U	1.83 U	4.02 ^(b)	2.04 U
Dibenzo(a,h)anthracene	1.34 U	1.29 U	1.31 U	1.51 U	1.46 U
Benzo(g,h,i)perylene	1.49 U	1.43 U	1.46 U	1.68 U	1.63 U
<u>Surrogate Internal Standards (%)</u>					
d4 1,4-Dichlorobenzene	59 ^(b)	76	78	89	59
d8 Naphthalene	70	91	84	91	65
d10 Acenaphthene	72	87	81	94	72
d12 Chrysene	81	75	83	105	77
d14 Dibenzo(a,h,i)anthracene	66	78	76	108	97

TABLE G.10. (contd)

Treatment	MATRIX SPIKES							
	COMP	COMP			COMP	COMP		
	EC-A	EC-A, MS			HU-C	HU-C, MS		
	1	1			1	1		
Batch	5	5	Amount		7	7	Amount	
Wet Wt.	20.08	20.05	Spiked	Percent	12.96	12.71	pike	Percent
Units	ng/g	ng/g	ng/g	Recover	ng/g	ng/g	ng/g	Recovery
1,4-Dichlorobenzene	1.86 U	21.5	24.9	86	2.87 U	36.1	39.3	92
Naphthalene	1.86 U	23.5	24.9	94	7.42	47.9	39.3	103
Acenaphthylene	1.58 ^(b)	21.4	24.9	80	1.59	39.3	39.3	100
Acenaphthene	6.17	27.8	24.9	87	3.75	47.6	39.3	112
Fluorene	1.90 ^(b)	23.2	24.9	86	1.90 U	46.1	39.3	117
Phenanthrene	6.07	25.1	24.9	76	5.24	52.6	39.3	121 ^(c)
Anthracene	4.07	27.1	24.9	92	3.45 U	51.3	39.3	131 ^(c)
Fluoranthene	45.0	133	24.9	353 ^(c)	19.0	73.9	39.3	140 ^(c)
Pyrene	65.0	134	24.9	277 ^(c)	22.7	69.9	39.3	120
Benzo(a)anthracene	6.87	30.0	24.9	93	6.61 ^(b)	55.6	39.3	125 ^(c)
Chrysene	25.7	46.0	24.9	82	10.3	54.0	39.3	111
Benzo(b)fluoranthene	7.13	32.6	24.9	102	8.74	54.5	39.3	116
Benzo(k)fluoranthene	4.61	28.4	24.9	96	4.77 ^(b)	54.7	39.3	127 ^(c)
Benzo(a)pyrene	6.27 ^(b)	27.9	24.9	87	5.14	53.8	39.3	124 ^(c)
Indeno(123-cd)pyrene	1.76 U	23.0	24.9	85	5.85 ^(b)	47.6	39.3	106
Dibenzo(a,h)anthracene	1.26 U	22.8	24.9	87	1.94 U	47.8	39.3	122 ^(c)
Benzo(g,h,i)perylene	2.91	22.1	24.9	77	5.28 ^(b)	43.5	39.3	97
<u>Surrogate Internal Standards (%)</u>								
d4 1,4-Dichlorobenzene	56	70	NA	NA	41	52	NA	NA
d8 Naphthalene	75	90	NA	NA	53	63	NA	NA
d10 Acenaphthene	86	97	NA	NA	66	77	NA	NA
d12 Chrysene	92	96	NA	NA	67	81	NA	NA
d14 Dibenzo(a,h,i)anthracene	101	103	NA	NA	85	102	NA	NA

TABLE G.10. (contd)

Treatment Replicate Batch Wet Wt. Units	MATRIX SPIKES							
	COMP	COMP			C-NV	C-NV, MS		
	SB-A	SB-A, MS			2	2		
	1	1	mount	Percent	6	6	Amount	Percent
	20.08	20.02	Spiked	Recovery	20.08	20.17	Spiked	Percent
	ng/g	ng/g	ng/g		ng/g	ng/g	ng/g	Recovery
1,4-Dichlorobenzene	1.86 U	20.2	25.0	81	1.86 U	24.1	24.8	97
Naphthalene	3.79	27.5	25.0	95	2.72 ^(b)	30.5	24.8	112
Acenaphthylene	1.92 ^(b)	23.0	25.0	84	0.73 U	27.1	24.8	109
Acenaphthene	23.2	52.2	25.0	116	2.34 ^(b)	31.1	24.8	116
Fluorene	11.1	36.9	25.0	103	2.76	28.1	24.8	102
Phenanthrene	62.7	101	25.0	153 ^(c)	2.76 ^(b)	30.4	24.8	111
Anthracene	14.4	42.8	25.0	114	2.24 U	30.2	24.8	122 ^(c)
Fluoranthene	152	218	25.0	264 ^(c)	6.80	40.1	24.8	134 ^(c)
Pyrene	146	208	25.0	248 ^(c)	7.20	35.8	24.8	115
Benzo(a)anthracene	12.6	38.8	25.0	105	1.09 U	33.9	24.8	137 ^(c)
Chrysene	33.8	63.8	25.0	120	2.87	31.0	24.8	113
Benzo(b)fluoranthene	10.3 ^(b)	33.7	25.0	94	4.44 ^(b)	32.5	24.8	113
Benzo(k)fluoranthene	4.84	29.4	25.0	98	2.81 ^(b)	32.5	24.8	120
Benzo(a)pyrene	7.74	32.4	25.0	99	1.49 U	31.3	24.8	126 ^(c)
Indeno(123-cd)pyrene	2.45	24.1	25.0	87	2.87 ^(b)	29.1	24.8	106
Dibenzo(a,h)anthracene	1.26 U	24.1	25.0	96	1.26 U	29.8	24.8	120
Benzo(g,h,i)perylene	3.53	25.4	25.0	87	2.78 ^(b)	27.4	24.8	99
<u>Surrogate Internal Standards (%)</u>								
d4 1,4-Dichlorobenzene	60	37	NA	NA	71	59	NA	NA
d8 Naphthalene	76	46	NA	NA	85	69	NA	NA
d10 Acenaphthene	82	50	NA	NA	88	77	NA	NA
d12 Chrysene	80	49	NA	NA	80	73	NA	NA
d14 Dibenzo(a,h,i)anthracene	87	53	NA	NA	92	83	NA	NA

TABLE G.10. (contd)

Treatment	ANALYTICAL REPLICATES							RSD	RSD%
	COMP HU-A	COMP HU-A Dup	COMP HU-A Trip	COMP HU-C	COMP HU-C Dup	COMP HU-C Trip			
Replicate	5-1	5-2	5-3	4-1	4-2	4-3			
Batch	4	4	4	6	6	6			
Wet Wt.	14.57	13.76	13.79	17.18	17.51	16.38			
Units	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g	RSD%	
1,4-Dichlorobenzene	2.57 U	2.72 U	2.72 U	NA	2.16 U	2.12 U	2.27 U	NA	
Naphthalene	4.51	3.53	3.67	14	3.01 ^(b)	3.22	3.50 ^(b)	8	
Acenaphthylene	2.97 ^(b)	3.18 ^(b)	2.79 ^(b)	7	2.59 ^(b)	2.84 ^(b)	2.71 ^(b)	5	
Acenaphthene	23.5	22.8	23.6	2	4.77	4.59	4.75	2	
Fluorene	9.15	9.0	9.20	1	3.39 ^(b)	3.40 ^(b)	3.96	9	
Phenanthrene	53.3	53.7	55.1	2	6.43	5.66	5.74	7	
Anthracene	17.6	17.4	18.0	2	4.34 ^(b)	4.12 ^(b)	3.75 ^(b)	7	
Fluoranthene	263	258	264	1	46.1	44.8	43.5	3	
Pyrene	295	289	292	1	59.7	57.6	56.3	3	
Benzo(a)anthracene	34.7	34.4	34.6	0	7.37 B	7.18 B	7.30 B	1	
Chrysene	79.1	76.9	79.2	2	20.7	19.8	19.2	4	
Benzo(b)fluoranthene	24.5	34.1	24.6	20	9.45	9.35	9.07	2	
Benzo(k)fluoranthene	10.1 ^(b)	2.44 U	11.1	NA	5.05	4.69	5.29	6	
Benzo(a)pyrene	19.2	19.5	20.1	2	5.87	5.72	5.79	1	
Indeno(123-cd)pyrene	5.01	5.09	5.03	1	3.95	3.77 ^(b)	4.12	4	
Dibenzo(a,h)anthracene	1.98 ^(b)	1.84 U	2.07	NA	2.14 ^(b)	2.14 ^(b)	2.23 ^(b)	2	
Benzo(g,h,i)perylene	6.20	6.44	6.52	3	4.23	4.09	4.28	2	
<u>Surrogate Internal Standards (%)</u>									
d4 1,4-Dichlorobenzene	63	60	52	NA	63	62	68	NA	
d8 Naphthalene	77	77	67	NA	74	77	81	NA	
d10 Acenaphthene	80	82	70	NA	79	81	86	NA	
d12 Chrysene	73	75	65	NA	76	79	81	NA	
d14 Dibenzo(a,h,i)anthracene	82	85	73	NA	82	88	90	NA	

TABLE G.10. (contd)

Treatment	ANALYTICAL REPLICATES							
	COMP SB-B	COMP SB-B Dup	COMP SB-B Trip		COMP BU	COMP BU Dup	COMP BU Trip	
Replicate	2-1	2-2	2-3		3-1	3-2	3-3	
Batch	5	5	5		7	7	7	
Wet Wt.	17.11	17.25	17.13		8.60	8.47	8.21	
Units	ng/g	ng/g	ng/g	RSD%	ng/g	ng/g	ng/g	RSD%
1,4-Dichlorobenzene	2.24 U	2.24 U	2.24 U	NA	4.32 U	4.40 U	4.55 U	NA
Naphthalene	2.33 ^(b)	2.31 ^(b)	2.33	0	10.8	11.2	10.2	5
Acenaphthylene	1.76 ^(b)	1.62 ^(b)	1.40 ^(b)	11	1.68 U	1.85 ^(b)	1.77 U	NA
Acenaphthene	7.39	6.96	6.72	5	5.01	5.63	5.95 ^(b)	9
Fluorene	2.21	2.02 ^(b)	1.83	9	6.39	2.92 U	6.84 ^(b)	NA
Phenanthrene	6.73	7.08	6.61	4	7.61	8.28	7.52	5
Anthracene	4.76	4.92	4.99	2	7.93 ^(b)	5.28 U	5.46 U	NA
Fluoranthene	49.4	50.7	45.6	5	16.3	19.6	17.6	9
Pyrene	69.5	70.2	63.8	5	21.1	24.8	22.1	8
Benzo(a)anthracene	7.72 B	7.14 B	6.68 B	7	2.54 U	9.61 ^(b)	2.67 U	NA
Chrysene	21.1	21.7	19.1	7	10.2	10.8	10.9	4
Benzo(b)fluoranthene	7.70	7.49 ^(b)	6.76	7	11.9	12.6	12.5	3
Benzo(k)fluoranthene	4.59	4.44	3.98	7	6.60 ^(b)	6.85 ^(b)	6.78 ^(b)	2
Benzo(a)pyrene	6.38 ^(b)	5.52 ^(b)	5.18	11	6.06	6.67	6.38	5
Indeno(123-cd)pyrene	2.11 U	2.11 U	2.11 U	NA	8.11 ^(b)	8.18	8.54 ^(b)	3
Dibenzo(a,h)anthracene	1.51 U	1.51 U	1.51 U	NA	2.92 U	2.97 U	3.08 U	NA
Benzo(g,h,i)perylene	2.82	2.68	2.53	5	7.71	8.09	7.98	2
<u>Surrogate Internal Standards (%)</u>								
d4 1,4-Dichlorobenzene	44	61	53	NA	50	41	50	NA
d8 Naphthalene	60	80	71	NA	60	50	60	NA
d10 Acenaphthene	64	83	76	NA	78	65	74	NA
d12 Chrysene	64	83	75	NA	83	67	77	NA
d14 Dibenzo(a,h,i)anthracene	71	92	82	NA	104	85	99	NA

(a) U Undetected at or above given concentration.

(b) Ion ratio out or confirmation ion not detected.

(c) Outside quality control range (50-120%) for matrix spike recovery.

(d) NA Not applicable.

TABLE G.11. Lipids in Tissue of *N. virens*

<u>Sediment Treatment</u>	<u>Replicate</u>	<u>Sample Weight</u>	<u>% Dry Weight</u>	<u>% Lipids (wet weight)</u>	<u>% Lipids (dry weight)</u>
<i>Nereis</i> Background	1	5.04	12.86	1.98	15.4
<i>Nereis</i> Background	2	5.07	12.94	2.17	16.8
<i>Nereis</i> Background	3	5.13	12.05	2.14	17.8