



**Fermi National Accelerator Laboratory**

**FERMILAB Conf-96/274-E  
CDF & DØ**

# **Prospects for Mixing and CP Violation at the Tevatron**

**Michael Rijssenbeek  
For the CDF and DØ Collaborations**

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510-0500*

*State University of New York at Stony Brook  
Stony Brook, NY 11794-3800*

**August 1996**

Published Proceedings at the *XI Topical Workshop on  $p\bar{p}$  Collider Physics*, Padova, Italy, May 27-June 1, 1996.

## **Disclaimer**

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process or service by trade name, trademark, manufacturer or otherwise, does not necessarily constitute or imply its endorsement, recommendation or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

## **Distribution**

*Approved for public release: further dissemination unlimited.*

## PROSPECTS FOR MIXING AND CP VIOLATION AT THE TEVATRON

Michael Rijssenbeek  
*State University of New York at Stony Brook*  
*Stony Brook, NY 11794-3800*

The Fermilab  $p\bar{p}$  Tevatron collider, operating at  $\sqrt{s} = 1.8$  TeV, is a proficient source of  $B$  hadrons. The first collider run, Run I, has clearly proven that  $B$  states can be observed cleanly and can be studied in great detail in a hadron collider environment. In this presentation the future of the study of  $B\bar{B}$  mixing and of CP violation at the Tevatron, with an upgraded machine and detectors, is discussed.

### 1 Introduction

The Tevatron Run I ended in May '96. The data,  $\simeq 115 \text{ pb}^{-1}$  per detector, led to the long-awaited discovery of the top quark with combined mass  $175 \pm 6$  GeV, to the current precision measurement of the  $W$  mass with a combined uncertainty of 130 MeV, and to numerous new and improved measurements in electroweak physics, QCD physics, and searches for phenomena beyond the Minimal Standard Model (MSM).

In the field of  $B$  physics, CDF has presented a number of pioneering studies – e.g.  $B \rightarrow J/\psi K_s^0$ ,  $\Lambda_b$ ,  $B$  hadron cross section, lifetime and branching fraction measurements, measurement of mixing parameter  $x_d$  – that have solidly proven the feasibility of detailed  $B$  physics studies at hadron colliders<sup>1</sup>. The DØ Collaboration, with large muon coverage and excellent muon id, but without the benefits of a central magnetic field and of micro vertexing, has contributed with measurements of inclusive  $\mu$  and  $b$  quark production cross sections, inclusive central and forward  $J/\psi$  production, and  $B\bar{B}$  mixing<sup>2</sup>.

### 2 The Tevatron Collider – Past and Future

At Fermilab the Main Injector ring, adjacent to the Tevatron ring, is under construction. The Main Injector is a conventional high intensity 150 GeV proton synchrotron, intended to improve the  $\bar{p}$  production rate, and to increase the number of  $p$ 's and  $\bar{p}$ 's available for collisions in the Tevatron. With improvements at the  $\bar{p}$  production target, in the cooling and accumulator rings, and with 36 on 36 bunches in the Tevatron, the luminosity  $\mathcal{L}$  will increase by a factor 5–10 in Run II (early 1999).

Under the name TeV33, further improvements are proposed: the addition of a permanent magnet antiproton storage and recycling ring. This would

increase the luminosity by a factor five in the period before the LHC era.

The luminosity evolution for the various run periods is summarized in Table 1, with crossing interval and machine improvements<sup>3</sup>.

Table 1: Typical luminosity, crossing interval, and machine improvements for the past and future Tevatron run periods.

Run Period	Typical $\mathcal{L}$	Bunch Interval	Improvements
1989	$1.6 \times 10^{30}$	$3.0 \mu\text{s}$	original design
Ia 1991–1993	$5.4 \times 10^{30}$	$3.0 \mu\text{s}$	Separators, $\bar{p}$ improvements
Ibc 1993–1996	$1.9 \times 10^{31}$	$3.0 \mu\text{s}$	Linac upgrade
IIa early 1999–	$1 \times 10^{32}$	396 ns	Main Injector,
IIb	$2 \times 10^{32}$	132 ns	$\bar{p}$ improvements
TeV33 2002– (?)	$1 \times 10^{33}$	132 ns	Recycler ring, electron cooling

### 3 The Detector Upgrades

Both detector collaborations have ambitious upgrade plans (fully approved for DØ, partially approved for CDF as of June).

The CDF collaboration plans to extend its  $B$ -physics capability with a larger and improved 3-d, 5-layer silicon vertex detector (SVX II), an intermediate fiber tracker (IFT), an outer small-cell drift chamber (COT) with  $dE/dx$ , followed by a time-of-flight counter (TOF). Muon coverage will be increased (CMU, FMU), and the present plug calorimeter (PCAL) improved<sup>5</sup>. The flexible trigger system will use information on  $p_T$ , decay-length,  $dE/dx$ , and TOF, and will have a 40 kHz accept capability at the first (deadtime-less) level.

Extrapolations based on current performance predict that CDF will be able to address all CKM topics of interest with great precision and within a single experiment<sup>5</sup>.

The DØ Collaboration is replacing its central and forward tracking systems completely. For Run II, a central/forward silicon barrels and disks micro-vertex tracker with  $10 \mu\text{m}$  hit resolution will be installed, followed by 16 doublet-layers of scintillating fiber detection (axial and stereo views) with doublet  $r\phi$  resolution of  $130 \mu\text{m}$ . This tracking is surrounded by a 2 T,  $1 X_0$  superconducting solenoid, located inside the bore of the existing central cryostat of the liquid Argon-Uranium calorimeter. The resolution  $\delta p_T/p_T$  is about 2% at

$p_T = 1 \text{ GeV}$ , and 4% at  $p_T = 10 \text{ GeV}$ . After the solenoidal coil, and also in front of the end calorimeters, a preshower detector is placed to help in  $e/\pi$  separation and to sample the energy lost in the solenoid coil.

The calorimeter and muon system electronics are overhauled in order to cope with the reduced bunch interval. The muon system will be upgraded in areas of trigger, rejection of cosmics, and shielding. A  $p_T$  track trigger has been designed for level 1, and faster trigger processors will be installed.

DØ will concentrate on leptonic decay modes for its  $B$  physics program, making full use of its large  $|\eta| \leq 2.5$  coverage for both  $\mu$  and  $e$  triggers in combination with the fast  $p_T$  trigger at several thresholds. Because of cost, no impact parameter trigger is planned at the start of Run II.

#### 4 $B\bar{B}$ Mixing and CP Violation at the Tevatron

The Tevatron is a high energy  $B$  factory: at 1.8 TeV the cross section for  $b\bar{b}$  production is  $\simeq 100 \mu\text{b}$ , i.e. about  $10^{11}$   $b$ 's are produced per  $\text{fb}^{-1}$  ( $\simeq 1$  year), versus  $10^7$  for BaBar or Belle. Moreover, all  $B$  species are produced copiously:  $B_u$ ,  $B_d$ ,  $B_s$ ,  $B_c$ , and  $b$ -baryons.

The increased luminosity and the shortened bunch interval require substantial upgrades to detectors, triggers, and DAQ. Although high  $p_T$  physics remains a priority of both collaborations, the availability of very large numbers of  $b\bar{b}$  pairs has focussed strong interest onto the field of  $B$ -physics.

Both detectors plan to have acceptance with full momentum determination up to pseudorapidity  $|\eta| \simeq 2-3$ , i.e. a large fraction of the  $b$ -quark phase-space. Because many  $B$  decay modes of interest have small branching fractions, very selective triggers are required, with identification of one or more of the decay products. With the decay products being relatively soft, it is important to have low ( $p_T \simeq 2 \text{ GeV}$ ) trigger thresholds. At such thresholds the raw trigger rates are huge (up to 40 kHz for a central di-hadron trigger), and seriously challenge the trigger system and DAQ capabilities.

The study of  $B\bar{B}$  mixing and CP violation aims to accurately determine elements of the CKM mixing matrix: their magnitudes and phases. Using both exclusive and semi-exclusive  $B$  decays, it is possible to over-constrain the CKM matrix, thereby testing the MSM. In the Wolfenstein approximation the CKM matrix elements are written in powers of  $\lambda$ ,  $\lambda \simeq V_{us} = \sin \theta_C$ . Unitarity provides six non-trivial relationships between products of CKM elements. For the  $B$  system the most interesting relationship is:

$$\begin{aligned} 0 &= V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* \\ &\simeq A\lambda^3[(\rho + i\eta) - 1 + (1 - \rho - i\eta)] \end{aligned} \quad (1)$$

These types of relationships in the SM are requirements of closure of “triangles” in the complex plane, where the sides are the product terms in Eq. 1.

The CKM quantities of interest are measurable with various  $B$  decay channels, and generally rely on interference between two dominant amplitudes contributing to the decay mode, e.g. interference between  $B^0 \rightarrow f$  and  $B^0 \rightarrow \bar{B}^0 \rightarrow \bar{f}$  in cases where  $f = \bar{f}$ . In Table 2 a list of CKM parameters is given with the process that will (probably) measure it best. These, and many other processes have been investigated; beautiful and detailed overviews, both theoretical and experimental, can be found e.g. in the many contributions to the Snowmass '93  $B$  Physics Workshop<sup>4</sup>.

Table 2: Physics CKM quantities, “best” measurement mode, and measurement technique and critical detection issues.

Physics	Modes	Technique/Issues
$\beta$	CP asym. in $B \rightarrow J/\psi K_s$	$\mu$ trigger; low background
$\alpha$	CP asym. in $B_d \rightarrow \pi\pi$	Di-hadron, decay-length trig. $\pi X$ backgrounds, penguins?
$\gamma$	CP asym. in $B_s \rightarrow D_s K$	Trigger? Resolve oscillations? Fully reconstruct final states.
$V_{td}/V_{ts}$	$B_d$ mixing, $B_s$ mixing	Trigger? Resolve oscillations?
$V_{ub}/V_{cb}$	$B \rightarrow \rho\ell\nu, D^* \ell\nu$	Lepton trigger; rates?
QCD	$B_c$ decays ( $B_c \rightarrow J/\psi\pi$ )	Lepton trigger; rates?
rare	$B \rightarrow \mu\mu, B^{\pm/0} \rightarrow \mu\mu K^{\pm}/K^*$	Lepton trigger; rates?

In the following we present an – incomplete – overview of the approaches by the collaborations for measurement of the CKM parameters.

$\sin(2\beta)$ :

In the “golden” decay mode  $B^0, \bar{B}^0 \rightarrow J/\psi K_s^0$ , a CP eigenstate, the CP asymmetry  $A_{CP}$  is directly related to  $\sin(2\beta)$ :  $A_{CP}(t) = \sin(2\beta) \sin(\Delta M_d t)$ , where  $\Delta M_d$  is the mass difference between the  $B$  mass eigenstates. The *measured* asymmetry is “diluted” by a factor  $D$  due to mis-tagging, and also due to mixing ( $D_{mix} = x/(1+x^2) \simeq 0.47$ ) if the time-integrated asymmetry is measured.

The uncertainty  $\delta A_{CP}$ , depends on the number of events  $N$ , the  $B$  flavor tagging efficiency  $\epsilon_{tag}$  and the “dilution” factor  $D$  from mis-tagging ( $D \equiv (N_r - N_w)/(N_r + N_w)$ , where  $N_r(N_w)$  is the number of right(wrong)  $B$  tags) as follows:  $\delta A_{CP}^2 = N \times \epsilon_{tag} D^2$ . In Table 3 the product  $\epsilon_{tag} D^2$  estimated for CDF is listed for several flavor tagging methods. The resulting uncertainty

from CDF on  $\sin(2\beta)$  expected in  $1 \text{ fb}^{-1}$  is between 0.14 and 0.08, competitive with results predicted for future  $B$  factories.

Table 3: Values of  $\epsilon_{tag}D^2$  as measured in the present data, extrapolations based on predicted upgrade performance, and detection elements involved in the flavor tagging technique.

Tagging Technique	$\epsilon_{tag}D^2$ (measured)	$\epsilon_{tag}D^2$ (expected)	Upgrade
jet charge	$1.3 \pm 0.3$	4.0	SVX II, IFT
central muon	$0.7 \pm 0.2$	0.7	CMU
non-central muon	—	0.3	FMU, IFT
electron	—	1.0	PCAL, IFT
same-side $\pi$	$1.5 \pm 0.9$	2.0	SVX II, IFT
total	3.5	8.0(?)	correlations?

DØ expects a  $\sim 40 \text{ MeV}$  resolution in the  $J/\psi$ -mass. Using a di-muon trigger, DØ expects a  $\delta \sin(2\beta)$  of about 0.12 with  $1 \text{ fb}^{-1}$ .

$\sin(2\alpha)$ :

The decay mode  $B_d \rightarrow \pi^+\pi^-$ , a CP eigenstate, appears to be best for measurement of  $\sin(2\alpha)$ . The trigger requires di-hadrons of  $p_T > 1.5 \text{ GeV}$  plus a separated vertex. The method is similar to that above apart from two caveats. First, backgrounds from  $B_s \rightarrow K\pi, KK$ , and combinatorics will further dilute  $A_{CP}$ , and thus excellent mass resolution and  $\pi/K$  separation are essential. Second, the theoretical side of CP violation in  $B_d \rightarrow \pi\pi$  is not clean, as spoiling contributions from penguin diagrams exist and are difficult to estimate.

In CDF the expected mass resolution is 20 MeV, whereas the COT plus TOF will provide  $\pi/K$  separation. With present theoretical estimates and assuming a very conservative signal-over-background ratio of 25%, CDF expects an uncertainty on  $\sin(2\alpha) \simeq 0.10$  with this decay mode.

Angle  $\gamma$ :

The angle  $\gamma$  can be accessed with the decay modes  $B_s(t) \rightarrow D_s^\pm K^\mp$  and  $B^\pm \rightarrow DK^\pm$ , with  $D = D^0, \bar{D}^0$ , or  $D_{CP+}^0$  the CP-even combination of  $D^0$  and  $\bar{D}^0$ . Only all-hadronic decay modes can be used, e.g.  $D_s^\pm \rightarrow \phi\pi^\pm, \phi \rightarrow K^+K^-$ . In view of the large value of  $x_s \geq 10$  ( $x \equiv \Delta M/\Gamma$ ), the  $B_s$  decay modes have to be measured as function of proper time. The decays measure angle  $\gamma$  directly and are free from spoiling penguin contributions.

Alternatively the six decay modes  $B^\pm \rightarrow DK^\pm$ ,  $D = D^0, \bar{D}^0$ , or  $D_{CP}$  maybe used. However, some of these branching fractions are very small. Also,

these decays measure  $\gamma$  up to a hadronic phase from final state effects, which can be as large as 0.1 radians. Ideally, several processes and methods should be used to constrain and measure angle  $\gamma$ .

$|V_{td}/V_{ts}|$ :

A measurement of this ratio, combined with the measurement of  $|V_{ub}|$  and  $|V_{cb}|$  from e.g. CLEO, yields an alternative measurement of the unitarity triangle, from its sides rather than from its angles.  $|V_{td}/V_{ts}|$  can be derived from a measurement of  $\Delta m_{B_s}/\Delta m_{B_d} = (B_{B_s} f_{B_s}^2)/(B_{B_d} f_{B_d}^2) \times |V_{ts}/V_{td}|^2$ , where enters the ratio of theoretical form factors  $(B_{B_s} f_{B_s}^2)$ . Lattice QCD calculations presently determine the proportionality constant as  $1.2 \pm 0.1$  and from LEP  $\Delta m_{B_d}$  is measured to  $\pm 4\%$ . ALEPH sets a lower limit on  $\Delta m_{B_s} > 5.9 \text{ ps}^{-1}$  (equivalent to  $x_s > 9.5$ ) at 95% CL.

Thus oscillations will be rapid and the  $x_s$  reach will crucially depend on resolution in proper time, i.e. in decay length and  $B$ -momentum. For semi-leptonic decay modes, where the missing decay neutrino degrades the  $B$  momentum resolution, CDF expects to measure  $x_s$  up to 15, while fully reconstructed decays (e.g.  $B_s \rightarrow D_s \pi$ ,  $D_s 3\pi$ , or  $\psi\phi$ ) might reach to beyond 20. For the latter, Fig. 1 shows that with 3-D reconstruction of primary and decay vertices with the SVX, oscillations can be resolved well for  $x_s = 20$ .




Figure 1: Simulation of the mixed  $B_s$  fraction for fully reconstructed  $B_s$  hadrons with the upgraded CDF detector in Run II




Figure 2: Simulation of difference between unmixed and mixed event samples of fully reconstructed  $B_s \rightarrow D_s^\pm \pi^\mp (\pi^+ \pi^-)$ ,  $D_s \rightarrow \phi \pi (\pi \pi)$  for DØ in Run II

CP violation in fully “visible” hadronic modes  $B_s \rightarrow D_s \pi (\pi \pi)$ ,  $D_s^\pm \rightarrow \phi \pi (\pi \pi)$  decays is used to explore the capability of DØ to measure the mixing parameter  $x_s$ , with a simulation of the upgraded DØ detector. In Table 4 the



estimated branching fractions and efficiencies for  $D\bar{0}$  are listed, indicating a yield of about 4000 fully reconstructed  $B_s$  in  $1\text{ fb}^{-1}$ . In the Fig. 2 the CP asymmetry is shown as function of proper decay time for  $x_s = 12$ . The oscillations are well resolved, and a reach up to 20 seems possible.

Table 4: Rates, Branching fractions, and efficiencies for fully reconstructed  $B_s$  for  $D\bar{0}$

$B$ -pairs per $1\text{ fb}^{-1}$	$4 \times 10^{10}$
Fraction $B_s$	0.3 /pair
$\text{BR}(B_s \rightarrow D_s \pi(\pi\pi))$ (all charged second.)	0.03 (est.)
$\text{BR}(D_s^\pm \rightarrow \phi\pi^\pm, \phi \rightarrow K^+K^-)$	0.018 (meas.)
$\text{BR}(D_s^\pm \rightarrow \phi\pi^\pm\pi^+\pi^-, \phi \rightarrow K^+K^-)$	0.009 (meas.)
Tagging efficiency ( $e, \mu$ )	$0.3 \times 0.22$ (meas.)
Trigger efficiency ( $p_T > 4$ )	0.06 (meas.)
Geometry, vertex, & track efficiency	0.1 (simul.)

## Acknowledgments

The author would like to thank Ron Lipton of  $D\bar{0}$ , and the conveners of the  $B$  physics groups, P. Spiccas and M. Schmidt in CDF, and A. Zieminski in  $D\bar{0}$ , for their generous help and information.

## References

1. Proc. XI<sup>th</sup> Topical Workshop on  $p\bar{p}$  Collider Physics, Padova, Italy, May 1996, and the CDF  $B$  physics webpage [www-cdf.fnal.gov](http://www-cdf.fnal.gov).
2. Proc. XI<sup>th</sup> Topical Workshop on  $p\bar{p}$  Collider Physics, Padova, Italy, May 1996, and the  $D\bar{0}$   $B$  physics webpage [www-d0sg10.fnal.gov](http://www-d0sg10.fnal.gov).
3. Proc. XI<sup>th</sup> Topical Workshop on  $p\bar{p}$  Collider Physics, Padova, Italy, May 1996.
4. Proc. of the Workshop on  $B$  Physics at Hadron Accelerators, eds. P. McBride and C. Mishra, Snowmass, Colorado (1993); Proc. 10<sup>th</sup> Topical Workshop on  $p\bar{p}$  Collider Physics, eds. R. Raja and J. Yoh, Fermilab (1995); Proc. 6<sup>th</sup> International Symposium on Heavy Flavor Physics, (1995).
5. CDF Upgrade Proposal to the Fermilab PAC, June 1996.