

CONF-961005--32

GA-A22467

# IMPLICATIONS FROM DIMENSIONLESS PARAMETER SCALING EXPERIMENTS

by  
T.C. LUCE, C.C. PETTY, B. BALET, and J.G. CORDEY

RECEIVED  
MAR 17 1997  
OSTI

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

*J*

OCTOBER 1996

MASTER

 **GENERAL ATOMICS**

## DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DISCLAIMER**

**Portions of this document may be illegible  
in electronic image products. Images are  
produced from the best available original  
document.**

GA-A22467

# IMPLICATIONS FROM DIMENSIONLESS PARAMETER SCALING EXPERIMENTS

by

T.C. LUCE, C.C. PETTY, B. BALET,\* and J.G. CORDEY\*

This is a preprint of a paper to be presented at the Sixteenth IAEA International Conference on Plasma Physics and Controlled Nuclear Research, October 7-11, 1996, Montreal, Canada, and to be published in *The Proceedings*.

\*JET Joint Undertaking, Abingdon, United Kingdom.

Work supported by  
the U.S. Department of Energy  
under Contract No. DE-AC03-89ER51114

GA PROJECT 3466  
OCTOBER 1996



F1-CN-64/AP1-3

## IMPLICATIONS FROM DIMENSIONLESS PARAMETER SCALING EXPERIMENTS

### ABSTRACT

The dimensionless parameter scaling approach is increasingly useful for predicting future tokamak performance and guiding theoretical models of energy transport. Experiments to determine the  $\rho_*$  (gyroradius normalized to plasma size) scaling have been carried out in many regimes. The electron  $\rho_*$  scaling is always "gyro-Bohm" ( $\chi \propto \chi_B \rho_*$ ), while the ion  $\rho_*$  scaling varies with regime. The ion variation is correlated with both density scale length (L mode, H mode) and current profile. The ion  $\rho_*$  scaling in the low- $q$ , H-mode regime (where fusion power plants are expected to operate) is gyro-Bohm, which is the most favorable confinement scaling observed. New experiments in  $\beta$  scaling and collisionality scaling have been carried out in low- $q$  discharges in both L mode and H mode. In L mode, global analysis shows that there is a slightly unfavorable  $\beta$  dependence ( $\beta^{-0.1}$ ) and no  $v_*$  dependence. In H-mode, global analysis finds a weak  $\beta$  dependence ( $\beta^{0.1}$ ) and an unfavorable dependence on  $v_*$  ( $v_*^{-0.35}$ ). The lack of significant  $\beta$  scaling spans the range of  $\beta_N$  from 0.25 to 2.0. The very small  $\beta$  dependence in L mode and H mode is in contradiction with the standard global scaling relations (ITER-89P:  $\tau \propto \beta^{-0.52}$ , ITER-93H:  $\tau \propto \beta^{-1.23}$ ). This contradiction in H mode may be indicative of the impact on the H-mode database of low- $n$  tearing instabilities which are observed at slightly higher  $\beta_N$  in the  $\beta$  scaling experiments. For the low- $q$ , H-mode experiments, the observed scalings can be combined to yield a global scaling law  $\tau_E \propto I^{6z/11} B^{(2-z)6/11} P^{-5/11} n^{3/11} a^{(31-6z)/11}$ , where  $z$  is the yet unmeasured  $q$  scaling experiment. The measured  $\beta$  and  $v_*$  scalings explain the weak density dependence observed in engineering parameter scans. It also points to the power of the dimensionless parameter approach, since it is possible to obtain a definitive size scaling from experiments on a single tokamak.

### 1. INTRODUCTION

Significant progress has been made in the last two years on applying dimensionless parameter scaling techniques to the problem of predicting and understanding tokamak energy transport. The value of this approach with respect to prediction of future machine performance is that present-day devices can operate at ignition-relevant values of the standard dimensionless parameters with the exception of  $\rho_*$ , which is the gyroradius normalized to the plasma linear size [1]. Therefore, the performance extrapolation is reduced to knowledge of a single parameter scaling which can be validated independently on many machines.

Experiments to measure the  $\rho_*$  scaling of local transport in various operating regimes have been carried out on the DIII-D tokamak. In low- $q$  H-mode, the  $\rho_*$  scaling is "gyro-Bohm" [2], which means the net diffusivity  $\chi$  scales like  $\chi_B \rho_*$  where  $\chi_B$  is the Bohm diffusivity. This gyro-Bohm scaling is qualitatively consistent with the H-mode global regression scaling (ITER-93H) used by the ITER EDA for its confinement projections. For L-mode plasmas, the measured global scaling lies near Bohm scaling (consistent with  $\chi \propto \chi_B \rho_*^0$ ). This is consistent with the standard L-mode regression scaling (ITER-89P). Two-fluid analysis showed the origin of the

derivation from the expected gyro-Bohm scaling to be exclusively in the ion scaling [3,4]. Recently, high- $q$  H-mode discharges have also been shown to exhibit Bohm scaling in the ion channel [5]. The data clearly show that the deviation from gyro-Bohm scaling in the ion channel is correlated with shorter density scale length (L-mode) or a current profile scale length (high- $q$ ). The physical mechanisms for this behavior are a topic of active research. Combining the most favorable confinement scaling regime (low- $q$  H-mode) and the H-mode power threshold scaling, it is possible to construct a paradigm for minimizing the size of the plasma core for an ignition power balance point [6]. Extrapolating from DIII-D discharges, a plasma with 500 MW of fusion power could be sustained in the H mode against transport losses at  $B = 5.7$  T,  $I = 9.9$  MA,  $\bar{n} = 2.0 \times 10^{20} \text{ m}^{-3}$ ,  $R = 2.7$  m, and  $\beta_N = 3.3$ . This ignition power balance point is below the Troyon  $\beta$  limit and the Greenwald density limit.

In addition to  $\rho_*$  scaling experiments, scaling studies for the dimensionless parameters  $\beta$  and collisionality have also been carried out in both L-mode and H-mode. These experiments help clarify what physical processes are involved in the energy transport. In principle, a complete set of scaling experiments would define an empirical local scaling relation for transport. The experiments reported here represent a significant step in that direction, including the first reported  $\beta$  and collisionality scaling experiments in H-mode.

## 2. PROOF-OF-PRINCIPLE EXPERIMENTS

A critical test of the dimensionless parameter scaling approach to transport studies is the realization of "identity" discharges — discharges on two tokamaks with widely different physical parameters, but exact matches of the dimensionless parameters. The first steps in this direction have been successfully carried out by comparing ELMing H-mode discharges on DIII-D and JET. The magnetic geometry of both machines was identical to 1% in aspect ratio, elongation, and  $q$ . The fluid dimensionless parameters  $\rho_*$ ,  $\beta$ , and collisionality  $\nu_*$  will be constant if  $na^2$ ,  $Wa^{-1/2}$ , and  $Ba^{5/4}$  are constant. The experimental match is 2%, 5%, and 3%, respectively. The proof of principle for the global confinement is whether the thermal confinement normalized to the cyclotron frequency is constant. This figure of merit agrees to within 2% for these discharges [5]. The next step is to show that the diffusivities scale appropriately ( $\chi a^{-3/4}$  constant) across the plasma for both electrons and ions. Unfortunately,  $T_i$  is unavailable for the JET discharge, so it is assumed that  $T_i = T_e$  and a one-fluid diffusivity  $\chi \equiv (\kappa_e + \kappa_i) / (n_e + n_i)$  is used. The comparison is shown in Fig. 1. The scaled diffusivities agree to within 20% everywhere from  $\rho = 0.2$ – $0.8$ . Recent JET discharges with ion temperatures will be analyzed in the near future. The present degree of agreement between the scaled JET and DIII-D plasmas provides confidence in the dimensionless scaling approach.

## 3. $\beta$ SCALING

Experimental results from the scaling of energy transport with  $\beta$  should differentiate between various proposed instability mechanisms. Most drift wave models show little enhancement or perhaps even slight reduction in transport with increasing  $\beta$ . On the other hand, transport models which invoke resistive MHD or magnetic fluctuations are generally expected to have strong  $\beta$  degradation. Standard regression analysis of global confinement databases would favor the latter. The ITER-89P L-mode scaling gives  $\beta^{-0.52}$ , while the ITER-93H H-mode scaling gives a very strong  $\beta^{-1.2}$  dependence. In order to keep  $\rho_*$  and  $\nu_*$  fixed while  $\beta$  varies, the




Fig. 1. The normalized one-fluid diffusivity for JET and DIII-D is shown vs. normalized radius.

following relations must be held for fixed plasma geometry:  $n \propto B^4$ ,  $T \propto B^2$ , and  $I \propto B$ . This results in  $\Delta\beta \propto (\Delta B)^4$ . In the experiments reported here, a factor of 2 scan in  $\beta$  is made.

For the L-mode experiment, the normalized confinement scales like  $\beta^{-0.1}$ . The global engineering parameters and the appropriate combinations proportional to the dimensionless parameters are shown in Table I. The density and stored energy matches are better than 10%. The normalized confinement time varies only slightly as  $\beta_N$  goes from 0.25 to 0.5. This is significantly weaker than the ITER-89P scaling. Preliminary two-fluid analysis shows that both fluids scale similarly.

In H-mode, the normalized confinement time scales like  $\beta^{0.1}$  as  $\beta_N$  is scanned from 1.0 to 2.0. Table I also shows the engineering and dimensionless variables for the H-mode scan. The parameters were chosen to be in a regime where gyro-Bohm  $\rho_*$  scaling would be expected. The density and stored energy matches are better than 10%. This is significantly different from the  $\beta$  scaling in the ITER-93H scaling relation.

One speculation on the source of this difference for H-mode plasmas is the effect of tearing modes at higher  $\beta_N$ . For the shots in the same sequence with slightly higher  $\beta_N$ , a  $m=3/n=2$  tearing mode is destabilized and the confinement can drop by up to 30%. This destabilization is attributed to neoclassical effects rather than classical  $\Delta' > 0$  destabilization [7]. Since this effect depends on collisionality and aspect ratio, it would not be discriminated in the ITER regression database by windowing on  $\beta_N$ . A similar effect is unlikely in the L-mode database. The source of the apparent  $\beta$  scaling there may be fast ion losses due to beam-driven instabilities (e.g., fishbones) in the highest  $\beta_N$  shots.

Table I: Comparison of global plasma parameters ( $\beta$ )

Parameter	L-mode		H-mode	
	Low- $\beta$	High- $\beta$	Low- $\beta$	High- $\beta$
$B$ (T)	1.63	1.91	1.62	1.93
$I_p$ (MA)	1.13	1.35	1.13	1.35
$\bar{n}$ ( $10^{19} \text{ m}^{-3}$ )	1.8	3.7	3.4	7.2
$W_{th}$ (kJ)	93	246	260	830
$P_{tot}$ (MW)	0.91	2.91	1.75	6.2
$\tau$ (s)	0.102	0.084	0.150	0.134
$R/a$	2.68	2.67	2.76	2.77
$\kappa$	1.72	1.73	1.81	1.83
$q_{95}$	3.66	3.64	3.67	3.85
$\bar{n}/B^4$	0.25	0.28	0.49	0.52
$W_{th}/B^6$	5.0	5.1	14.4	16.1
$\beta$ (%)	0.28	0.54	0.87	1.90
$B\tau$	0.17	0.16	0.24	0.26

#### 4. COLLISIONALITY SCALING

The theoretical expectations for scaling with collisionality are less clear. Neoclassical theory gives normalized confinement inversely proportional to collisionality in the banana regime. Drift wave transport is expected to be roughly independent of collisionality in the banana regime. Both the ITER-89P L-mode scaling and the ITER-93H-mode scaling have a  $v_*^{-0.28}$  dependence. In order to keep  $\rho_*$  and  $\beta$  fixed as collisionality varies, the following relations must hold:  $n \propto B^0$ ,  $T \propto B^2$ , and  $I \propto B$ . This implies  $\Delta v_* \propto (\Delta B)^{-4}$ . The experiments reported here vary  $v_*$  by a factor of 8.

For the L-mode scan, the normalized confinement is almost independent of  $v_*$ . The engineering parameters and the effective global dimensionless parameters are shown in Table II. Because of the large scan in  $v_*$ , the effect of mismatches in  $\rho_*$  and  $\beta$  are minimized. The density and stored energy matches are within 10%.

For the H-mode  $v_*$  scan, the normalized confinement time drops as the collisionality increases, consistent with a  $v_*^{-0.35}$  scaling. The H-mode  $v_*$  parameters are also given in Table II. The experiments were carried out in the low- $q$  H-mode regime where gyro-Bohm  $\rho_*$  scaling would be expected. The  $\beta$  values are similar to the ITER demonstration discharges. It is not yet clear whether this  $v_*$  scaling is observed in both channels. The observed scaling is close to the  $\rho_*$  scaling in the ITER-93H scaling relation.



Table II: Comparison of global plasma parameters ( $v_*$ )

Parameter	L-mode		H-mode	
	Low- $v_*$	High- $v_*$	Low- $v_*$	High- $v_*$
$B$ (T)	1.91	1.14	1.91	1.14
$I_p$ (MA)	1.37	0.81	1.35	0.8
$\bar{n}$ ( $10^{19} \text{ m}^{-3}$ )	2.7	2.4	6.3	6.2
$W_{th}$ (kJ)	228	82	930	340
$P_{tot}$ (MW)	3.55	0.72	3.6	1.61
$\tau$ (s)	0.064	0.113	0.26	0.21
$R/a$	2.69	2.70	2.74	2.77
$\kappa$	1.73	1.73	1.7	1.7
$q_{95}$	3.56	3.60	3.84	4.055
$\sqrt{W_{th}/\bar{n}}/B$	4.9	5.1	6.4	6.5
$\beta$ (%)	0.52	0.52	2.2	2.2
$B\tau$	0.12	0.13	0.50	0.24

## 5. DISCUSSION

Converting from dimensionless variables back to engineering variables for a global confinement scaling is a straightforward algebraic manipulation. Assuming a standard power law form for the scaling relation, the power degradation and density scaling are completely determined by the  $\rho_*$ ,  $\beta$ , and  $v_*$  scaling. [In this paper, the collisionality  $v_*$  is defined without  $q$ ; this is the collision frequency normalized to the transit time rather than the bounce time.] For the H-mode scans, the conversion to a global scaling law in engineering variables is clear because the experiments are carried out in a regime where both the electron and ion  $\rho_*$  scaling are expected to be the same. Taking the observed scalings:  $\rho_*^{-3}$  (gyro-Bohm),  $\beta^0$ , and  $v_*^{-1/3}$ , the following relation is obtained:

$$\tau_E \propto I^{6z/11} B^{(2-z)/6} P^{-5/11} n^{3/11} a^{(31-6z)/11}, \quad (1)$$

where  $-z$  is the  $q$  scaling in the dimensionless parameter formalism. Note that the weak density dependence observed in engineering variable scans [8] is recovered from the combination of three different scans. The rather weak power degradation is surprising. A weaker collisionality scaling than  $v_*^{-1/3}$  would give more density scaling and more power degradation. It is important to point out that Eq. (1) also demonstrates that the dimensionless parameter scaling approach yields a definitive prediction for the size scaling from single machine experiments. By verifying the dimensionless parameter scalings independently on several machines, confidence in the size scaling can be gained while avoiding the inevitable systematic effects in multiple machine databases. If  $z \equiv 11/6$  to give the standard linear current scaling, then Eq. (1) becomes

$$\tau_E \propto I^{1.0} B^{0.09} P^{-0.45} n^{0.27} a^{1.82}. \quad (2)$$

The next step is to compare these inferred engineering scaling relations with the larger set of discharges taken to find the dimensionless parameter matches to see if the experimental deviations can be described by the derived scaling relation.

The same exercise for the L-mode scans is not very straightforward, due to the variable ion  $\rho_*$  scaling. Without knowing the physical mechanism for this variation or, at least an empirical characterization of it, scaling of ion transport is uncertain. Also, various heating schemes deposit power in electrons and ions in differing proportions and the exchange term is not "dimensionally correct," so the fraction of power lost in electrons and ions is uncertain.

In conclusion, great progress toward the prediction and understanding of energy transport has been made by applying the dimensionless parameter scaling approach. In the next few years, it may be possible to derive a complete scaling rule by means of this approach.

## 6. ACKNOWLEDGMENTS

This is a report of work supported by the U.S. Department of Energy under Contract No. DE-AC03-89ER51114. We gratefully acknowledge the work of R.J. Groebner, P. Gohil, D.M. Thomas, and D.R. Baker in analyzing the CER data.

## 7. REFERENCES

- [1] WALTZ, R.E., DeBOO, J.C., ROSENBLUTH, M.N., Phys. Rev. Lett. **65** 2390 (1990).
- [2] PETTY, C.C., et al., Phys. Plasmas **2** 2342 (1995).
- [3] PETTY, C.C., et al., Phys. Rev. Lett. **74** 1763 (1995).
- [4] LUCE, T.C., PETTY, C.C., in *Controlled Fusion and Plasma Physics* (Proc. 22nd EPS Conf. Bournemouth, 1995) European Physical Society, Petit-Lancy, Switzerland, 1995) Vol. 19C, Part III, p. 25.
- [5] PETTY, C.C., et al., to be published in *Controlled Fusion and Plasma Physics* (Proc. 23rd EPS Conf. Kiev, 1996).
- [6] PETTY, C.C., LUCE, T.C., "Projections of Gyroradius Scaling Experiments to an Ignition Tokamak," General Atomics Report GA-A22271 (1996), to be published in Nucl. Fusion **36** (1996).
- [7] LA HAYE, R.J., this conference.
- [8] SCHISSEL, D.P., et al., Nucl. Fusion **34** 1401 (1994).