

UCRL-JRNL-221969



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Potential bias of model projected greenhouse warming in irrigated regions

D. Lobell, G. Bala, C. Bonfils, P. Duffy

June 9, 2006

Geophysical Research Letters

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Potential bias of model projected greenhouse warming in irrigated regions

D.B. Lobell<sup>1\*</sup>, G. Bala<sup>1</sup>, C. Bonfils<sup>2</sup>, P.B. Duffy<sup>1,2</sup>

<sup>1</sup> Energy and Environment Directorate, Lawrence Livermore National Laboratory,  
Livermore, CA 94550

<sup>2</sup> School of Natural Sciences, University of California, Merced, CA 95346

\* Corresponding Author:

David B. Lobell

Lawrence Livermore National Laboratory

P.O. Box 808 L-103

Livermore, CA 94550

Tel: (925) 422-4148

Fax: (925) 423-4908

Email: [dlobell@llnl.gov](mailto:dlobell@llnl.gov)

## **Abstract**

Atmospheric general circulation models (GCMs) used to project climate responses to increased CO<sub>2</sub> generally omit irrigation of agricultural land. Using the NCAR CAM3 GCM coupled to a slab-ocean model, we find that inclusion of an extreme irrigation scenario has a small effect on the simulated temperature and precipitation response to doubled CO<sub>2</sub> in most regions, but reduced warming by as much as 1° C in some agricultural regions, such as Europe and India. This interaction between CO<sub>2</sub> and irrigation occurs in cases where agriculture is a major fraction of the land surface and where, in the absence of irrigation, soil moisture declines are projected to provide a positive feedback to temperature change. The reduction of warming is less than 25% of the temperature increase modeled for doubled CO<sub>2</sub> in most regions; thus greenhouse warming will still be dominant. However, the results indicate that land use interactions may be an important component of climate change uncertainty in some agricultural regions. While irrigated lands comprise only ~2% of the land surface, they contribute over 40% of global food production. Climate changes in these regions are therefore particularly important to society despite their relatively small contribution to average global climate.

## Introduction

The value of models that project societal impacts of climate change depend largely on the ability to make clear and accurate statements about the uncertainties associated with such projections. One impact of great interest is the effect of climate change on regional or global crop production, and the associated changes in food security. Uncertainties for these impacts depend largely on uncertainties associated with climate model projections over agricultural regions [Parry, *et al.*, 2005]. Therefore, a clear understanding of climate uncertainty over agricultural regions is important for evaluating climate change mitigation and adaptation strategies. Irrigated systems are particularly relevant, as they provide roughly 40% of global food production [FAO, 2002] despite covering just 2% of global land surface area [FAO, 2004].

Quantification of climate uncertainty is commonly achieved by comparing outputs from several different general circulation models (GCMs) [IPCC, 2001; Palmer, *et al.*, 2005]. However, this approach can only capture uncertainties due to processes that are represented differently in different models. As most GCMs were designed primarily to study global scale climate changes, they often treat very simply or entirely omit processes that are important in agricultural regions. For example, according to the documentation for the 22 models used for projection of future climate for the next Intergovernmental Panel on Climate Change (IPCC) report (available at [www.pcmdi-llnl.gov](http://www.pcmdi.llnl.gov)), only eight include some representation of land use, and none include irrigation.

Previous work has shown that changes in irrigation extent can have substantial effects on local climate [Adegoke, *et al.*, 2003; Lobell, *et al.*, 2006]. Thus, climate projections in areas with expansion or retraction of irrigation are subject to uncertainties

related to the magnitude of, and climatic response to, this land use change. However, if the effect of irrigation is independent of other climate forcings, then estimates of irrigation's impact on climate can simply be added to projections based on other forcings, such as greenhouse gas increases. This study investigates whether the effects of irrigation and CO<sub>2</sub> on climate are indeed independent or, instead, interacting. If the latter is true, then estimates of CO<sub>2</sub> induced warming over irrigated regions may be biased in the current suite of GCMs that ignore irrigation.

One reason to expect an interaction is that GCMs often simulate a reduction of soil moisture in future climate [*Manabe, et al.*, 1981; *Wetherald and Manabe*, 1995; *Manabe, et al.*, 2004; *Wang*, 2005], in particular for the summer months in middle latitudes where much of agricultural activity occurs. Moisture decreases are driven largely by the nonlinear increase in atmospheric saturation vapor pressure, and thus evaporation rates, with higher temperatures [*Wetherald and Manabe*, 1995]. These moisture decreases subsequently lead to an increase in the fraction of incident radiation partitioned to sensible heat flux, which provides a positive feedback to temperature change as more energy is used to heat the land surface rather than evaporate water. Irrigation eliminates the potential for this feedback because soil moisture is maintained via human activity (assuming that irrigation water availability is unaffected by climate change). At issue here is whether the presence of irrigation has the potential to significantly modify the response of climate to CO<sub>2</sub> over agricultural regions, and if so where these interactions are most likely to be important.

## **Model and Experiment Design**

We used the Community Atmosphere Model version 3.0 (CAM) [Collins, *et al.*, 2004] coupled to version 3.0 of the Community Land Model (CLM) [Oleson, *et al.*, 2004]. CAM has 26 levels in the vertical dimension and was run using 2.0° latitude x 2.5° longitude resolution using the Finite Volume (FV) dynamical core. We used a version of CAM that was coupled to a slab-ocean / thermodynamic sea-ice mode to allow interaction between atmosphere, ocean and sea ice. The horizontal ocean heat transport and mixed layer depth for the oceanic surface mixed layer were prescribed to ensure realistic sea surface temperatures and ice distributions for the present climate.

The CLM model, described in detail by Bonan *et al.* [2002], includes up to four different plant function types (PFT's) within each 2.0° x 2.5° grid cell, with a single PFT used to represent croplands. Multiple soil columns are simulated for grid cells with croplands, with cropland soils treated separately from natural vegetation. The percent of each grid cell with cropland (Figure 1a) is defined in CLM from satellite-based land cover maps [Loveland, *et al.*, 2000]. Monthly values of LAI for each PFT in CLM are prescribed based on satellite measurements from 1992-1993 [Bonan, *et al.*, 2002], and thus do not respond to CO<sub>2</sub> changes. However, the physiological effects of CO<sub>2</sub> on canopy photosynthesis and stomatal conductance are modeled within CLM [Oleson, *et al.*, 2004].

Four 50-year simulations were performed, with the first 20 years of each simulation treated as spin-up and the last 30 years used for analysis. The first two runs included the default land surface in CLM, which does not include irrigation, and atmospheric CO<sub>2</sub> levels of 355 ppm (CLM default) and 710 ppm, respectively. The difference between these two runs provided an estimate of the climate response to

doubled CO<sub>2</sub> in the absence of irrigation. These experiments were then repeated, but with soil moisture in all agricultural soil columns maintained at soil saturation percentage throughout the simulation. This extreme irrigation scenario was not intended to represent reality, where many regions are not irrigated and even irrigated soils are usually below saturation. Instead, the experiments provide a simple measure of the sensitivity of greenhouse warming to irrigation practices. Figure 1b shows a recent estimate of the percent of actual land surface that is irrigated [*Doll and Siebert, 2000*], for comparison with the modeled distribution (Figure 1a).

## **Results and Discussion**

Doubling of CO<sub>2</sub> in the absence of irrigation resulted in increases of global mean temperature by 2.2 °C and precipitation by 4 %. Consistent with nearly all GCMs, the simulated warming was greatest at high latitudes (Figure 2a) because of decreases in sea ice and snow extents [*Dai, et al., 2001; IPCC, 2001*]. Soil moisture exhibited significant changes in many regions (Figure 2b) with notable reductions in Europe, southern India, Southeast Asia, western United States, and northern South America. Increases in soil moisture were simulated in northeast Africa, Saudi Arabia, and western South America. Reductions in soil moisture corresponded to increases in the ratio of sensible to latent heat fluxes (not shown), which thus provided a positive feedback to temperature changes in these regions.

In the presence of irrigation, doubling of CO<sub>2</sub> resulted in global mean increases of temperature by 2.1 °C and precipitation by 4 %. Thus, even an extreme scenario of irrigating all agricultural soils to saturation had little effect on the global climate



sensitivity to CO<sub>2</sub>. In many agricultural regions, such as the corn belt of the U.S. or most of South America and China, irrigation had negligible effects on the climate response to CO<sub>2</sub> (Figure 2c). However, temperature changes differed significantly in several regions, in particular those with a positive soil moisture feedback in Figure 2b. Compared to the simulations without irrigation, the annual mean temperature response to doubled CO<sub>2</sub> was roughly 0.5°C lower in southern India and Southeast Asia in the presence of irrigation, and 1.0 °C lower throughout much of Europe. While significant, these temperature differences represented less than 25% of the simulated warming from doubled CO<sub>2</sub> in most regions (Figure 2d).

Another way irrigation could modify CO<sub>2</sub> response is through the addition of extra water vapor, a strong greenhouse gas, to the atmosphere in response to elevated CO<sub>2</sub>. However, if water vapor feedbacks were important one would expect to see interactions over all irrigated regions, which was not observed. This agrees with Boucher et al. [2004], who found that direct radiative effects of water vapor from irrigation were much smaller than temperature effects of increased surface latent heat flux.

The results presented here suggest that simulated climate changes in some agricultural regions may be biased if current models anticipate feedbacks from soil moisture that, in reality, will not occur because of irrigation. In the present model, irrigation tended to mitigate the simulated temperature response to doubled CO<sub>2</sub>. It is also possible, at least in principle, that irrigation could amplify the simulated response, if an agricultural region is located within an area simulated to have a strong negative soil moisture feedback.

A recent inter-comparison of 15 GCMs found that modeled soil moisture changes for an A1b emission scenario were consistently negative in the summertime for southern Europe, southwestern U.S., Australia, and northern and southern Africa [Wang, 2005]. Southeast Asia also exhibited significant declines in moisture content for most models, while changes in India were inconsistent among models. No regions were found to have a consistent increase in summer soil moisture. Thus, the results presented here for Europe appear more robust than those for South and Southeast Asia. In addition, it appears unlikely that including irrigation would ever increase projected warming, given a lack of agricultural regions with a negative soil moisture feedback.

The use of extreme irrigation scenarios in this study was useful for bracketing the potential impact of irrigation on GCM simulations of greenhouse gas responses. For example, it appears that irrigation is relatively unimportant for simulating temperature or precipitation responses at the global scale, given the limited intersection of agricultural land and regions of strong simulated soil moisture feedbacks. At the regional scale, the presence of irrigation may significantly mitigate greenhouse warming but is unlikely to aggravate it. Simulations of more realistic distributions of irrigation (e.g., Figure 1b) would be useful for assessing the actual impact of irrigation on future regional climates.

While this study focused on the interaction between irrigation and CO<sub>2</sub>, a comparison of temperature changes for doubled CO<sub>2</sub> with those for irrigation are also of interest and readily evaluated from the experiments. Table 1 shows that cooling from an extreme irrigation scenario was ~0.5° C larger in magnitude, on average, than warming from doubled CO<sub>2</sub> over cropland regions in summer months. Conversely, irrigation-induced cooling was lower than CO<sub>2</sub>-induced warming in winter. Table 1 also reveals

that the modification of CO<sub>2</sub> warming by irrigation, which was ~0.3° C in summer and ~0.2° C in winter, was roughly one-tenth the magnitude of the direct cooling influence of irrigation.

Thus, accounting for changes in irrigation are likely more important than the interaction between irrigation and CO<sub>2</sub> when projecting climate change in regions with substantial land use change. However, these results depend on how irrigation was represented in the model and should be compared with other models. The results are also potentially sensitive to the lack of LAI response in CLM to irrigation or CO<sub>2</sub>, and the modeled response of stomatal conductance to CO<sub>2</sub> [Betts, *et al.*, 1997].

## **Conclusions**

GCM's are nearly universal in their simulation of summertime soil drying in response to increased atmospheric CO<sub>2</sub> in many important agricultural regions, such as Europe and southern North America. Thus, modeled temperature changes over these regions due to historical or future CO<sub>2</sub> changes likely include, to some extent, a positive feedback between moisture change and near-surface air temperatures. The results presented here indicate that irrigation can significantly modify (i.e. interact with) greenhouse warming by eliminating this moisture feedback. Inclusion of irrigation in GCMs would thus likely reduce simulated warming in some agricultural regions, although the effect on global mean temperature is likely to be small relative to the consequences of the enhanced greenhouse effect by increased greenhouse gas concentrations. Given that a disproportionate amount of food production originates in irrigated regions, however, it is possible that the interaction of irrigation and CO<sub>2</sub> is an

important source of overall uncertainty for projecting societal impacts of climate change. Thus, future work to better understand and reduce this source of climate uncertainty in heavily irrigated regions appears warranted.

### **Acknowledgements**

We thank Richard Betts and an anonymous reviewer for helpful comments on the manuscript. This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48. CB was supported by the State of California through the Public Interest Energy Research Program.

### **References**

- Adegoke, J. O., et al. (2003), Impact of irrigation on midsummer surface fluxes and temperature under dry synoptic conditions: A regional atmospheric model study of the U.S. high plains, *Monthly Weather Review*, 131, 556-564.
- Betts, R. A., et al. (1997), Contrasting physiological and structural vegetation feedbacks in climate change simulations, *Nature*, 387, 796-799.
- Bonan, G. B., et al. (2002), Landscapes as patches of plant functional types: An integrating concept for climate and ecosystem models, *Global Biogeochemical Cycles*, 16, -.
- Boucher, O., et al. (2004), Direct human influence of irrigation on atmospheric water vapour and climate, *Climate Dynamics*, 22, 597-603.

- Collins, W. D., et al. (2004), Description of the NCAR Community Atmosphere Model (CAM 3.0), NCAR.
- Dai, A., et al. (2001), Climates of the twentieth and twenty-first centuries simulated by the NCAR climate system model, *Journal of Climate*, 14, 485-519.
- Doll, P., and S. Siebert (2000), A digital global map of irrigated areas, *Icid Journal*, 49, 55-66.
- FAO (2002), Crops and Drops: making the best use of water for agriculture, 28 pp, FAO, Rome.
- FAO (2004), Food and Agriculture Organization of the United Nations (FAO), FAO Statistical Databases, <http://apps.fao.org>
- IPCC (2001), Intergovernmental Panel on Climate Change Working Group 1, Climate Change 2001: The Scientific Basis, IPCC Working Group 1.
- Lobell, D. B., et al. (2006), Biogeophysical impacts of cropland management changes on climate, *Geophysical Research Letters*, 33, -.
- Loveland, T. R., et al. (2000), Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data, *International Journal of Remote Sensing*, 21, 1303-1330.
- Manabe, S., et al. (2004), Century-scale change in water availability: CO<sub>2</sub>-quadrupling experiment, *Climatic Change*, 64, 59-76.
- Manabe, S., et al. (1981), Summer Dryness Due to an Increase of Atmospheric Co-2 Concentration, *Climatic Change*, 3, 347-386.
- Oleson, K., et al. (2004), Technical Description of the Community Land Model (CLM), <http://www.cgd.ucar.edu/tss/clm/distribution/clm3.0>

- Palmer, T. N., et al. (2005), Representing model uncertainty in weather and climate prediction, *Annual Review of Earth and Planetary Sciences*, 33, 163-193.
- Parry, M., et al. (2005), Climate change, global food supply and risk of hunger, *Philosophical Transactions: Biological Sciences*, 360, 2125-2138.
- Wang, G. (2005), Agricultural drought in a future climate: results from 15 global climate models participating in the IPCC 4th assessment, *Climate Dynamics*, 25, 739-753.
- Wetherald, R. T., and S. Manabe (1995), The mechanisms of summer dryness induced by greenhouse warming, *Journal of Climate*, 8, 3096-3108.

Table 1. Average temperature change (°C) over land in response to extreme irrigation<sup>a</sup> (first 2 rows) or doubled CO<sub>2</sub> (last two rows).




| Model Experiments   | JJA      |               | DJF      |               |
|---|----------|---------------|----------|---------------|
|   | All land | Cropland only | All land | Cropland only |
| 1xCO <sub>2</sub> <sub>irr</sub> - 1xCO <sub>2</sub> <sub>no-irr</sub>    | -1.8     | -2.7          | -0.9     | -1.4          |
| 2xCO <sub>2</sub> <sub>irr</sub> - 2xCO <sub>2</sub> <sub>no-irr</sub>    | -2.1     | -3.0          | -1.1     | -1.6          |
| 2xCO <sub>2</sub> <sub>no-irr</sub> - 1xCO <sub>2</sub> <sub>no-irr</sub> | 2.7      | 2.2           | 2.6      | 2.1           |
| 2xCO <sub>2</sub> <sub>irr</sub> - 1xCO <sub>2</sub> <sub>irr</sub>       | 2.4      | 1.9           | 2.4      | 1.9           |

<sup>a</sup>Irrigation corresponds to all cropland grid cells being maintained at soil saturation. JJA = June-July-August; DJF = December-January-February.

#### Figure Legends:


1) (a) The percentage of each 2.0° x 2.5° grid cell defined as cropland within CAM3. (b) The estimated percentage of each 2.0° x 2.5° equipped for irrigation [*Doll and Siebert, 2000*]

2) Annual average surface (a) temperature (°C) and (b) soil moisture (%) changes for doubled CO<sub>2</sub> in the absence of irrigation, [2xCO<sub>2</sub> - 1xCO<sub>2</sub>]<sub>no-irr</sub>. (c) Difference in temperature response to doubled CO<sub>2</sub> with and without irrigation, [2xCO<sub>2</sub> - 1xCO<sub>2</sub>]<sub>irr</sub> - [2xCO<sub>2</sub> - 1xCO<sub>2</sub>]<sub>no-irr</sub>. (d) Ratio (%) of same responses, [2xCO<sub>2</sub> - 1xCO<sub>2</sub>]<sub>irr</sub> / [2xCO<sub>2</sub> - 1xCO<sub>2</sub>]<sub>no-irr</sub>. Gray cells are not significant at p = 0.05.







(a)



(b)



(c)



(d)

