

Evidence for charged B meson decays to $a_1^\pm(1260)\pi^0$ and $a_1^0(1260)\pi^\pm$

B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ X. Prudent,¹ V. Tisserand,¹
 A. Zghiche,¹ J. Garra Tico,² E. Grauges,² L. Lopez,³ A. Palano,³ G. Eigen,⁴ B. Stugu,⁴ L. Sun,⁴ G. S. Abrams,⁵
 M. Battaglia,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ J. A. Kadyk,⁵
 L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ D. Lopes Pegna,⁵ G. Lynch,⁵ L. M. Mir,⁵ T. J. Orimoto,⁵
 M. T. Ronan,^{5,*} K. Tackmann,⁵ W. A. Wenzel,⁵ P. del Amo Sanchez,⁶ C. M. Hawkes,⁶ A. T. Watson,⁶
 T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ T. Schroeder,⁷ M. Steinke,⁷ D. Walker,⁸ D. J. Asgeirsson,⁹
 T. Cuhadar-Donszelmann,⁹ B. G. Fulsom,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ A. Khan,¹⁰ M. Saleem,¹⁰
 L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ A. P. Onuchin,¹¹
 S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ K. Yu. Todyshev,¹¹ M. Bondioli,¹² S. Curry,¹² I. Eschrich,¹²
 D. Kirkby,¹² A. J. Lankford,¹² P. Lund,¹² M. Mandelkern,¹² E. C. Martin,¹² D. P. Stoker,¹² S. Abachi,¹³
 C. Buchanan,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ F. Liu,¹⁴ O. Long,¹⁴ B. C. Shen,¹⁴ L. Zhang,¹⁴ H. P. Paar,¹⁵
 S. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ A. Cunha,¹⁶ B. Dahmes,¹⁶ T. M. Hong,¹⁶
 D. Kovalskiy,¹⁶ J. D. Richman,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. J. Flacco,¹⁷ C. A. Heusch,¹⁷ J. Kroseberg,¹⁷
 W. S. Lockman,¹⁷ T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ M. G. Wilson,¹⁷ L. O. Winstrom,¹⁷ E. Chen,¹⁸
 C. H. Cheng,¹⁸ F. Fang,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ R. Andreassen,¹⁹
 G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ K. Mishra,¹⁹ M. D. Sokoloff,¹⁹ F. Blanc,²⁰ P. C. Bloom,²⁰ S. Chen,²⁰
 W. T. Ford,²⁰ J. F. Hirschauer,²⁰ A. Kreisel,²⁰ M. Nagel,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ J. G. Smith,²⁰
 K. A. Ulmer,²⁰ S. R. Wagner,²⁰ J. Zhang,²⁰ A. M. Gabareen,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹
 F. Winklmeier,²¹ D. D. Altenburg,²² E. Feltresi,²² A. Hauke,²² H. Jasper,²² J. Merkel,²² A. Petzold,²² B. Spaan,²²
 K. Wacker,²² V. Klose,²³ M. J. Kobel,²³ H. M. Lacker,²³ W. F. Mader,²³ R. Nogowski,²³ J. Schubert,²³
 K. R. Schubert,²³ R. Schwierz,²³ J. E. Sundermann,²³ A. Volk,²³ D. Bernard,²⁴ G. R. Bonneaud,²⁴ E. Latour,²⁴
 V. Lombardo,²⁴ Ch. Thiebaux,²⁴ M. Verderi,²⁴ P. J. Clark,²⁵ W. Gradl,²⁵ F. Muheim,²⁵ S. Playfer,²⁵
 A. I. Robertson,²⁵ Y. Xie,²⁵ M. Andreotti,²⁶ D. Bettoni,²⁶ C. Bozzi,²⁶ R. Calabrese,²⁶ A. Cecchi,²⁶ G. Cibinetto,²⁶
 P. Franchini,²⁶ E. Luppi,²⁶ M. Negrini,²⁶ A. Petrella,²⁶ L. Piemontese,²⁶ E. Prencipe,²⁶ V. Santoro,²⁶ F. Anulli,²⁷
 R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ S. Pacetti,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,†}
 M. Piccolo,²⁷ M. Rama,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Contri,²⁸ M. Lo Vetere,²⁸ M. M. Macri,²⁸ M. R. Monge,²⁸
 S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ K. S. Chaisanguanthum,²⁹
 M. Morii,²⁹ J. Wu,²⁹ R. S. Dubitzky,³⁰ J. Marks,³⁰ S. Schenk,³⁰ U. Uwer,³⁰ D. J. Bard,³¹ P. D. Dauncey,³¹
 R. L. Flack,³¹ J. A. Nash,³¹ W. Panduro Vazquez,³¹ M. Tibbetts,³¹ P. K. Behera,³² X. Chai,³² M. J. Charles,³²
 U. Mallik,³² V. Ziegler,³² J. Cochran,³³ H. B. Crawley,³³ L. Dong,³³ V. Eyges,³³ W. T. Meyer,³³ S. Prell,³³
 E. I. Rosenberg,³³ A. E. Rubin,³³ Y. Y. Gao,³⁴ A. V. Gritsan,³⁴ Z. J. Guo,³⁴ C. K. Lae,³⁴ A. G. Denig,³⁵
 M. Fritsch,³⁵ G. Schott,³⁵ N. Arnaud,³⁶ J. Béquilleux,³⁶ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ V. Lepeltier,³⁶
 F. Le Diberder,³⁶ A. M. Lutz,³⁶ S. Pruvot,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ J. Serrano,³⁶
 V. Sordini,³⁶ A. Stocchi,³⁶ W. F. Wang,³⁶ G. Wormser,³⁶ D. J. Lange,³⁷ D. M. Wright,³⁷ I. Bingham,³⁸
 C. A. Chavez,³⁸ I. J. Forster,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ D. E. Hutchcroft,³⁸ D. J. Payne,³⁸
 K. C. Schofield,³⁸ C. Touramanis,³⁸ A. J. Bevan,³⁹ K. A. George,³⁹ F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹
 G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ D. A. Hopkins,⁴⁰ S. Paramesvaran,⁴⁰ F. Salvatore,⁴⁰ A. C. Wren,⁴⁰
 D. N. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² Y. M. Chia,⁴² C. L. Edgar,⁴²
 G. D. Lafferty,⁴² T. J. West,⁴² J. I. Yi,⁴² J. Anderson,⁴³ C. Chen,⁴³ A. Jawahery,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³
 J. M. Tuggle,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ E. Salvati,⁴⁴
 S. Saremi,⁴⁴ R. Cowan,⁴⁵ D. Dujmic,⁴⁵ P. H. Fisher,⁴⁵ K. Koeneke,⁴⁵ G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵ M. Spitznagel,⁴⁵
 F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ M. Zhao,⁴⁵ Y. Zheng,⁴⁵ S. E. Mclachlin,^{46,*} P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶
 A. Lazzaro,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸
 D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ M. Simard,⁴⁹ P. Taras,⁴⁹
 F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ G. De Nardo,⁵¹ F. Fabozzi,^{51,‡} L. Lista,⁵¹ D. Monorchio,⁵¹ C. Sciacca,⁵¹
 M. A. Baak,⁵² G. Raven,⁵² H. L. Snoek,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ G. Benelli,⁵⁴ L. A. Corwin,⁵⁴
 K. Honscheid,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ J. P. Morris,⁵⁴ A. M. Rahimi,⁵⁴ J. J. Regensburger,⁵⁴ Q. K. Wong,⁵⁴
 N. L. Blount,⁵⁵ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ J. A. Kolb,⁵⁵ M. Lu,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵
 D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ N. Gagliardi,⁵⁶ A. Gaz,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ A. Pompili,⁵⁶

Published in the Physical Review Letters

Work supported in part by US Department of Energy contract DE-AC02-76SF00515

M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ E. Ben-Haim,⁵⁷ H. Briand,⁵⁷ G. Calderini,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ A. Perez,⁵⁷ L. Gladney,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ E. Manoni,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ A. Cervelli,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. A. Mazur,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ G. Rizzo,⁶⁰ J. J. Walsh,⁶⁰ M. Haire,⁶¹ J. Biesiada,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² E. Baracchini,⁶³ F. Bellini,⁶³ G. Cavoto,⁶³ A. D’Orazio,⁶³ D. del Re,⁶³ E. Di Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ P. D. Jackson,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Renga,⁶³ C. Voena,⁶³ M. Ebert,⁶⁴ T. Hartmann,⁶⁴ H. Schröder,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ G. Castelli,⁶⁵ B. Franek,⁶⁵ E. O. Olaiya,⁶⁵ S. Ricciardi,⁶⁵ W. Roethel,⁶⁵ F. F. Wilson,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ M. Escalier,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ X. R. Chen,⁶⁷ H. Liu,⁶⁷ W. Park,⁶⁷ M. V. Purohit,⁶⁷ J. R. Wilson,⁶⁷ M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ P. Bechtler,⁶⁸ N. Berger,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸ J. C. Dingfelder,⁶⁸ J. Dorfan,⁶⁸ G. P. Dubois-Felsmann,⁶⁸ W. Dunwoodie,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ M. T. Graham,⁶⁸ P. Grenier,⁶⁸ C. Hast,⁶⁸ T. Hryn’ova,⁶⁸ W. R. Innes,⁶⁸ J. Kaminski,⁶⁸ M. H. Kelsey,⁶⁸ H. Kim,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ S. Li,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ D. B. MacFarlane,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O’Grady,⁶⁸ I. Ofte,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ T. Pulliam,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va’vra,⁶⁸ N. van Bakel,⁶⁸ A. P. Wagner,⁶⁸ M. Weaver,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ L. Wilden,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ V. Jain,⁷⁰ B. Pan,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappler,⁷⁰ S. B. Zain,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. M. Ruland,⁷² C. J. Schilling,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Pelliccioni,⁷⁴ M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ V. Azzolini,⁷⁶ N. Lopez-March,⁷⁶ F. Martinez-Vidal,⁷⁶ § D. A. Milanes,⁷⁶ A. Oyanguren,⁷⁶ J. Albert,⁷⁷ Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ I. M. Nugent,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ P. F. Harrison,⁷⁸ J. Ilic,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ M. Pappagallo,⁷⁸ ¶ H. R. Band,⁷⁹ X. Chen,⁷⁹ S. Dasu,⁷⁹ K. T. Flood,⁷⁹ J. J. Hollar,⁷⁹ P. E. Kutter,⁷⁹ Y. Pan,⁷⁹ M. Pierini,⁷⁹ R. Prepost,⁷⁹ S. L. Wu,⁷⁹ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

²³Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²⁴Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

²⁵University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁶Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

- ²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ²⁹Harvard University, Cambridge, Massachusetts 02138, USA
- ³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³¹Imperial College London, London, SW7 2AZ, United Kingdom
- ³²University of Iowa, Iowa City, Iowa 52242, USA
- ³³Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁴Johns Hopkins University, Baltimore, Maryland 21218, USA
- ³⁵Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
- ³⁶Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France
- ³⁷Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁸University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³⁹Queen Mary, University of London, E1 4NS, United Kingdom
- ⁴⁰University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ⁴¹University of Louisville, Louisville, Kentucky 40292, USA
- ⁴²University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴³University of Maryland, College Park, Maryland 20742, USA
- ⁴⁴University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴⁵Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁶McGill University, Montréal, Québec, Canada H3A 2T8
- ⁴⁷Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁸University of Mississippi, University, Mississippi 38677, USA
- ⁴⁹Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
- ⁵⁰Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁵¹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵²NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵⁴Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁵University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁶Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁷Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France
- ⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁵⁹Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ⁶⁰Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶¹Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶²Princeton University, Princeton, New Jersey 08544, USA
- ⁶³Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁴Universität Rostock, D-18051 Rostock, Germany
- ⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁷University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁸Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁶⁹Stanford University, Stanford, California 94305-4060, USA
- ⁷⁰State University of New York, Albany, New York 12222, USA
- ⁷¹University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷²University of Texas at Austin, Austin, Texas 78712, USA
- ⁷³University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷⁴Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁶IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
- ⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁸Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
- ⁷⁹University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁸⁰Yale University, New Haven, Connecticut 06511, USA

We present measurements of the branching fractions for the decays $B^\pm \rightarrow a_1^\pm(1260)\pi^0$ and $B^\pm \rightarrow a_1^0(1260)\pi^\pm$ from a data sample of 232×10^6 $B\bar{B}$ pairs produced in e^+e^- annihilation through the $\Upsilon(4S)$ resonance. We measure the branching fraction $\mathcal{B}(B^\pm \rightarrow a_1^\pm(1260)\pi^0) \times \mathcal{B}(a_1^\pm(1260) \rightarrow \pi^-\pi^+\pi^\pm) = (13.2 \pm 2.7 \pm 2.1) \times 10^{-6}$ with a significance of 4.2σ , and the branching fraction $\mathcal{B}(B^\pm \rightarrow a_1^0(1260)\pi^\pm) \times \mathcal{B}(a_1^0(1260) \rightarrow \pi^-\pi^+\pi^0) = (20.4 \pm 4.7 \pm$

$3.4) \times 10^{-6}$ with a significance of 3.8σ , where the first error quoted is statistical and the second is systematic.

PACS numbers: 13.25.Hw, 12.39.St, 11.30.Er

The rare decays of B mesons to two-body final states with an $a_1(1260)$ and a π^\pm, π^0, K^\pm or K_s^0 are important processes for testing theoretical factorization model predictions for branching fractions, branching fraction ratios and CP -violation parameters. The measurements can be combined with assumptions about $SU(3)$ symmetries to form upper bounds on $\Delta\alpha = |\alpha - \alpha_{\text{eff}}|$, where α is the weak interaction phase $\alpha \equiv \arg[-V_{td}V_{tb}^*/V_{ud}V_{ub}^*]$ of the Unitarity Triangle [1] and α_{eff} is the measured phase. The difference $\Delta\alpha$ is a measurement of the poorly known strength of the penguin amplitudes in the decay and can be used to improve our understanding of the CP -violating mechanism.

The rare decays $B^\pm \rightarrow a_1^\pm(1260)\pi^0$ and $B^\pm \rightarrow a_1^0(1260)\pi^\pm$ are expected to be dominated by $b \rightarrow u\bar{u}d$ contributions. The branching fraction for $B^0 \rightarrow a_1^\pm\pi^\mp$ has been measured to be $(33.2 \pm 3.8 \pm 3.0) \times 10^{-6}$ [2] and this agrees well with the calculation of Bauer, Stech and Wirbel [3] within the framework of naive factorization and assuming $|V_{ub}/V_{cb}| = 0.08$. A more recent analysis using naive factorization and measured form factors predicts branching fractions in the range $(5 - 11) \times 10^{-6}$ and $(4 - 9) \times 10^{-6}$ for $B^\pm \rightarrow a_1^\pm\pi^0$ and $B^\pm \rightarrow a_1^0\pi^\pm$, respectively [4]. Previous measurements have placed 90% confidence level upper limits of 1.7×10^{-3} and 9×10^{-4} on the branching fractions for $B^\pm \rightarrow a_1^\pm\pi^0$ and $B^\pm \rightarrow a_1^0\pi^\pm$, respectively [5], and recently the *BABAR* collaboration reported the first measurements of the CP -violating asymmetries in the decay $B^0 \rightarrow a_1^\pm\pi^\mp$ [6].

We present measurements of the branching fractions for the two charmless B meson decays $B^\pm \rightarrow a_1^\pm\pi^0$ and $B^\pm \rightarrow a_1^0\pi^\pm$ where the final state contains one neutral and three charged pions. The $a_1 \rightarrow 3\pi$ decay proceeds mainly through the intermediate states $(\pi\pi)_\rho\pi$ and $(\pi\pi)_\sigma\pi$ [7]. We do not distinguish between the dominant P-wave $(\pi\pi)_\rho$ and the S-wave $(\pi\pi)_\sigma$ in the channel $\pi^+\pi^-$. Possible background contributions from $B \rightarrow a_2(1320)\pi$ are investigated. Charge conjugate modes are implied throughout this paper.

The data were collected with the *BABAR* detector [8] at the PEP-II asymmetric e^+e^- collider. An integrated luminosity of 211 fb^{-1} , corresponding to 232 million $B\bar{B}$ pairs, was recorded at the $\Upsilon(4S)$ resonance (“on-resonance”) at a center-of-mass (CM) energy $\sqrt{s} = 10.58 \text{ GeV}$. An additional 20 fb^{-1} were taken about 40 MeV below this energy (“off-resonance”) for the study of continuum background in which a charm or lighter quark pair is produced.

Charged particles are detected and their momenta measured by the combination of a silicon vertex tracker,

consisting of five layers of double-sided silicon detectors, and a 40-layer central drift chamber, both operating in the 1.5-T magnetic field of a solenoid. The tracking system covers 92% of the solid angle in the CM frame. Charged-particle identification (PID) is provided by the average energy loss (dE/dx) in the tracking devices and by an internally reflecting ring-imaging Cherenkov detector. A K/π separation of better than four standard deviations (σ) is achieved for momenta below 3 GeV/ c , decreasing to 2.5σ at the highest momenta in the B decay final states.

The off-resonance data together with the Monte Carlo (MC) simulations of the signal decay modes, continuum, $B\bar{B}$ backgrounds and detector response [9] are used to establish the event selection criteria and reconstruction efficiency. The MC signal events are simulated as B^+ decays to $a_1\pi$ with $a_1 \rightarrow \rho\pi$. The a_1 and a_2 line shapes are generated with EvtGen [10], where we use mass and width parameters from Refs. [2] and [7].

Two photons with a minimum energy of 30 MeV (100 MeV for $B^+ \rightarrow a_1^0\pi^+$) and an invariant mass of $120 < m_{\gamma\gamma} < 150 \text{ MeV}/c^2$ are used to reconstruct the π^0 . The intermediate dipion states $(\pi^+\pi^-)$ or $(\pi^+\pi^0)$ are required to have an invariant mass of $0.46 < m_{\pi\pi} < 1.1 \text{ GeV}/c^2$. We impose PID requirements to cleanly identify the charged pions and to suppress contamination from a_1K . We require the invariant mass reconstructed for candidate $a_1^+ \rightarrow \pi^-\pi^+\pi^+$ and $a_1^0 \rightarrow \pi^-\pi^+\pi^0$ decays to be $0.8 < m_{a_1} < 1.8 \text{ GeV}/c^2$.

A B meson candidate is characterized kinematically by the energy-substituted mass $m_{\text{ES}} = \sqrt{(s/2 + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - \mathbf{p}_B^2}$ and energy difference $\Delta E = E_B^* - \sqrt{s}/2$, where the subscripts 0 and B refer to the initial $\Upsilon(4S)$ and to the B candidate in the lab-frame, respectively, and the asterisk denotes the $\Upsilon(4S)$ frame. The resolutions in m_{ES} and in ΔE are about $3.0 \text{ MeV}/c^2$ and 20 MeV, respectively. Candidates are required to have $5.25 \leq m_{\text{ES}} \leq 5.29 \text{ GeV}/c^2$ and $|\Delta E| \leq 0.2 \text{ GeV}$. To reduce fake B meson candidates we require a B vertex χ^2 probability > 0.01 . The absolute value of the cosine of the angle between the direction of the π meson from $a_1 \rightarrow \rho\pi$ with respect to the flight direction of the B in the a_1 meson rest frame is required to be less than 0.85 to suppress misreconstructed candidates. The distribution of this variable is flat for signal and peaks near ± 1 for misreconstructed candidates.

To reject continuum background, we use the angle θ_T between the thrust axis of the B candidate’s decay products and that of the rest of the tracks and neutral clusters in the event, calculated in the CM frame. The distribution of $\cos\theta_T$ is sharply peaked near ± 1 for combinations

drawn from jetlike $q\bar{q}$ pairs and is nearly uniform for the isotropic B meson decays; we require $|\cos\theta_T| < 0.65$.

The decay mode $B \rightarrow a_2\pi$ can also give background contributions. It is suppressed by using the angular variable \mathcal{A} , defined as the cosine of the angle between the normal to the plane of the 3π resonance and the flight direction of the bachelor pion evaluated in the 3π resonance rest frame. Since the a_1 and a_2 have spins of 1 and 2, respectively, the distributions of \mathcal{A} for these two resonances differ. We require $|\mathcal{A}| < 0.6$, which reduces the a_2 background by more than a factor of two in both decay channels.

After all the above selections, we have on average 1.20 and 1.56 candidates per event in events where there is at least one candidate, for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$, respectively, and we select the B candidate with the $(\pi\pi)$ mass nearest to the nominal ρ mass [7]. From the simulation, we find that this algorithm selects the correct-combination candidate in $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$ in 65% and 55% of events containing multiple candidates, respectively.

We use an unbinned maximum-likelihood fit using five variables to extract the background and signal yields of $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$. We describe the B decay kinematics with the two variables ΔE and m_{ES} . We also include the invariant mass of the 3π system (m_{a_1}), the variable \mathcal{A} and a Fisher discriminant \mathcal{F} . This discriminant combines four variables: the angles with respect to the beam axis of the B momentum and B thrust axis in the CM frame, and the zeroth and second angular moments of the energy flow around the B thrust axis [2].

The extended likelihood function is

$$\mathcal{L} = \frac{1}{N!} \exp\left(-\sum_j n_j\right) \prod_{i=1}^N \left[\sum_j n_j \mathcal{P}_j(\vec{x}_i; \vec{\alpha}_j) \right], \quad (1)$$

where n_j is the yield of events for hypothesis j (signal, a_2 , $B\bar{B}$ charmless, $B\bar{B}$ charm or continuum) and N is the number of events in the sample. The probabilities \mathcal{P}_j are products of probability density functions (PDF) for each of the independent variables $\vec{x}_i = \{m_{\text{ES}}, \Delta E, m_{a_1}, \mathcal{F}, \mathcal{A}\}$ evaluated for each event i . The $\vec{\alpha}_j$ are the parameters of the distributions in \vec{x}_i . By minimizing the quantity $-\ln\mathcal{L}$ in two separate fits, we determine the yields for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$.

To take into account the relatively large number of misreconstructed signal events, the signal is separated into two components, representing the correctly reconstructed (true) and the self cross-feed (SCF) candidates, with proportions fixed in the fit for each mode. SCF occurs when a track from an $a_1^+\pi^0$ or $a_1^0\pi^+$ is exchanged with a track from the rest of the event. The fraction of SCF, determined from MC, is 35% and 44% for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$, respectively.

In addition to the a_2 , there are three main categories of backgrounds: $B\bar{B}$ charmless, $B\bar{B}$ charm and contin-

uum. $B\bar{B}$ backgrounds are studied using MC simulations of $B^0\bar{B}^0$ and B^+B^- decays, using a large sample equivalent to $\sim 0.8 \text{ ab}^{-1}$. There are 17 $B\bar{B}$ charmless decays for $B^+ \rightarrow a_1^+\pi^0$ and 20 for $B^+ \rightarrow a_1^0\pi^+$ that contribute as background. Those decays with similar distributions are grouped to form 13 and 10 hypotheses, respectively, and are included in the fit with a fixed yield as determined from MC. The total $B\bar{B}$ charmless yields are 368 ± 92 and 755 ± 164 for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$, respectively. These are dominated by $B \rightarrow \rho\rho$, $B \rightarrow a_1\rho$ and the other $B \rightarrow a_1\pi$ mode under study. The $B\bar{B}$ charm backgrounds are included as a single hypothesis, with the normalization of the $B\bar{B}$ charm yield as a free parameter. Continuum events come from light quark production. We establish the functional forms and parameter values of the PDFs for $B\bar{B}$ charm and $B\bar{B}$ charmless backgrounds from MC simulations. For continuum, we use off-resonance data for the Fisher, on-resonance data with $|\Delta E| > 0.1 \text{ GeV}$ for m_{ES} , and on-resonance data with $5.25 < m_{\text{ES}} < 5.27 \text{ GeV}/c^2$ for the other variables.

We model the distributions using appropriate functions. The \mathcal{A} distributions are modeled with polynomials. For the true signal component, the remaining distributions are fitted using modified Gaussians [11], and a relativistic Breit-Wigner line-shape with a mass-dependent width [12], as necessary. The SCF component and the a_2 have similar shapes to the true signal but have broader or more asymmetric distributions and shifted means. The $B\bar{B}$ backgrounds and continuum distributions are modeled with modified Gaussians, polynomials, nonparametric functions [13] and, for m_{ES} , a phase-space-motivated empirical function [14]. The PDF variables are assumed to be independent except for $B^+ \rightarrow a_1^0\pi^+$, where a two dimensional nonparametric PDF [13] in m_{a_1} and ΔE accounts for observed correlations in the MC for both true signal events and SCF.

In the fit there are six free parameters: four yields (signal, continuum, a_2 and $B\bar{B}$ charm background), and two continuum background parameters (ΔE polynomial coefficient and m_{ES} shape coefficient ξ [14]).

For $B^+ \rightarrow a_1^+\pi^0$, there are 24608 events in the data sample. We measure the raw signal yield to be 459 ± 78 events with a reconstruction efficiency of $12.5 \pm 0.1\%$, corrected for differences in tracking and neutral particle reconstruction between data and MC. The yield of the decay $B^+ \rightarrow a_2^+\pi^0$ is 28 ± 65 events. For $B^+ \rightarrow a_1^0\pi^+$, there are 33375 events in the data sample and we measure the raw signal yield to be 382 ± 79 events with a corrected reconstruction efficiency of $7.2 \pm 0.1\%$. The yield of the decay $B^+ \rightarrow a_2^0\pi^+$ is 107 ± 65 events.

We confirm our fitting procedure by generating and fitting MC samples containing signal and background populations using the yields as found from data. We identify a signal yield bias for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$ of $16.8 \pm 0.1\%$ and $10.9 \pm 0.1\%$, respectively. We fit for the branching fractions taking into account the fit-

ted signal yield, the yield bias, the corrected reconstruction efficiency, daughter branching fractions, and the number of produced B mesons, assuming equal production rates of $B^0\bar{B}^0$ and B^+B^- pairs. The statistical significance is taken as the square root of the difference between the value of $-2\ln\mathcal{L}$ for zero signal and the value at its minimum. We measure the branching fraction $\mathcal{B}(B^+ \rightarrow a_1^+\pi^0) \times \mathcal{B}(a_1^+ \rightarrow \pi^-\pi^+\pi^+) = (13.2 \pm 2.7) \times 10^{-6}$ with a statistical significance of 5.3σ and the branching fraction $\mathcal{B}(B^+ \rightarrow a_1^0\pi^+) \times \mathcal{B}(a_1^0 \rightarrow \pi^-\pi^+\pi^0) = (20.4 \pm 4.7) \times 10^{-6}$ with a statistical significance of 4.7σ , where the errors are statistical.




FIG. 1: (color online). Projections of a) ΔE , b) m_{ES} , c) m_{a_1} , and d) \mathcal{F} for $B^+ \rightarrow a_1^+\pi^0$. Points represent on-resonance data, dashed lines the signal, dotted lines the continuum, dashed-dotted lines the $B\bar{B}$ charm background, the filled region the a_2 background and solid lines the full fit function. These plots are made with a requirement on the signal likelihood to enhance the signal, and thus do not show all events in the data sample.

Figs. 1 and 2 show the ΔE , m_{ES} , m_{a_1} , and \mathcal{F} projections for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$ made by selecting events with a signal likelihood (computed without the variable shown in the figure) exceeding a threshold that optimizes the expected sensitivity.

The systematic errors are summarized in Table I. We determine the sensitivity to the parameters of the signal and background PDF components by varying these within their uncertainties. The effect of varying the mass and width of the a_1 by the errors as reported in Ref. [2] is included in the PDF parameters' variation systematic. The uncertainty in the fit bias correction is taken as half of the fit bias correction. The effect of possible interference between a_2 and a_1 is estimated by adding the




FIG. 2: (color online). Projections of a) ΔE , b) m_{ES} , c) m_{a_1} , and d) \mathcal{F} for $B^+ \rightarrow a_1^0\pi^+$, using the same criteria and line styles as Fig. 1.

TABLE I: Summary of systematic errors for the $a_1^+\pi^0$ and $a_1^0\pi^+$ branching fraction measurements.

Systematic	$a_1^+\pi^0$	$a_1^0\pi^+$
PDF Parameter Variation	8.6%	8.8%
Fit Bias	8.4%	5.5%
$a_1 - a_2$ Interference	6.6%	7.4%
SCF Variation	4.4%	8.2%
Tracking Efficiency	3.9%	3.9%
π^0 Efficiency	3.0%	3.0%
Flight Direction Criteria	2.0%	2.0%
P-wave and S-wave Reconstruction	1.6%	-
Charmless $B\bar{B}$ Background	1.4%	3.1%
Number of $B\bar{B}$ Pairs	1.1%	1.1%
$\cos\theta_T$ Selection Criteria	1.1%	1.8%
Track Multiplicity	1.0%	1.0%
$\rho\pi\pi$, 4π Cross-Feed	0.9%	0.5%
a_1K Cross-Feed	-	0.4%
Total	16%	16%

a_2 and a_1 amplitudes together with a varying phase difference and using half the maximum change in yield as an uncertainty. The uncertainty in SCF is investigated by varying the SCF fraction. We also perform a separate fit treating the SCF as an independent background component. The fitted branching fraction is compatible with the nominal fit within the increased statistical uncertainty, but the statistical significance is reduced to 3.5σ and 3.0σ for $B^+ \rightarrow a_1^+\pi^0$ and $B^+ \rightarrow a_1^0\pi^+$, respectively. A systematic uncertainty of 1.6% is estimated for the difference in reconstruction efficiency in the decay

modes through the dominant P-wave $(\pi\pi)_\rho$ and the S-wave $(\pi\pi)_\sigma$. An error is assigned for the uncertainty in the fixed charmless $B\bar{B}$ background yields and possible interference effects by varying the individual components by the reported error on the branching fractions [7]. The systematic errors for the flight direction criteria, number of $B\bar{B}$ pairs, $\cos\theta_T$ selection criteria, track multiplicity, potential backgrounds from $\rho\pi\pi$ and 4π , and a_1K cross-feed are small. The total systematic error for both modes is 16%. The significance of the branching fractions, combining both statistical and systematic errors, is 4.2σ for $B^+ \rightarrow a_1^+\pi^0$ and 3.8σ for $B^+ \rightarrow a_1^0\pi^+$.

In conclusion, we have measured the branching fractions $\mathcal{B}(B^\pm \rightarrow a_1^\pm(1260)\pi^0) \times \mathcal{B}(a_1^\pm(1260) \rightarrow \pi^-\pi^+\pi^\pm) = (13.2 \pm 2.7 \pm 2.1) \times 10^{-6}$ and $\mathcal{B}(B^\pm \rightarrow a_1^0(1260)\pi^\pm) \times \mathcal{B}(a_1^0(1260) \rightarrow \pi^-\pi^+\pi^0) = (20.4 \pm 4.7 \pm 3.4) \times 10^{-6}$. Neglecting isoscalar contributions to the two-pion state, we assume $\mathcal{B}(a_1^\pm(1260) \rightarrow \pi^-\pi^+\pi^\pm)$ is equal to $\mathcal{B}(a_1^\pm(1260) \rightarrow \pi^\pm\pi^0\pi^0)$ and $\mathcal{B}(a_1^\pm(1260) \rightarrow (3\pi)^\pm)$ is equal to 100% [7], resulting in $\mathcal{B}(B^\pm \rightarrow a_1^\pm(1260)\pi^0) = (26.4 \pm 5.4 \pm 4.1) \times 10^{-6}$. We measure $\mathcal{B}(B^\pm \rightarrow a_1^0(1260)\pi^\pm) = (20.4 \pm 4.7 \pm 3.4) \times 10^{-6}$, assuming $\mathcal{B}(a_1^0(1260) \rightarrow \pi^-\pi^+\pi^0)$ is equal to 100%. The first errors quoted are statistical and the second are systematic. The signals are seen with significances of 4.2σ and 3.8σ , respectively, and are in agreement with factorization model predictions [3].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and STFC (United Kingdom). Indi-

viduals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

* Deceased

† Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡ Also with Università della Basilicata, Potenza, Italy

§ Also with Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

¶ Also with IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom

- [1] B. Aubert *et al.* (*BABAR* Collaboration), hep-ex/0703008.
- [2] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **97**, 051802 (2006).
- [3] M. Bauer, B. Stech and M. Wirbel, Z. Phys. C **34**, 103 (1987).
- [4] V. Laporta, G. Nardulli and T. N. Pham, Phys. Rev. D **74**, 054035 (2006); hep-ph/0602243v4 (erratum submitted to Phys. Rev. D).
- [5] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **241**, 278 (1990).
- [6] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **98**, 181803 (2007).
- [7] S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [8] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [9] S. Agostinelli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [10] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).
- [11] J. E. Gaiser *et al.*, Phys. Rev. D **34**, 711 (1986).
- [12] T. A. Armstrong *et al.* (WA76 Collaboration), Z. Phys. C **48**, 213 (1990).
- [13] K. S. Cranmer, Comput. Phys. Commun. **136**, 198 (2001).
- [14] With $x \equiv m_{ES}/E_b$ and ξ a parameter to be fit, $f(x) \propto x\sqrt{1-x^2} \exp[-\xi(1-x^2)]$. See Ref. [5].