

The biogeochemistry of Pu mobilization and retention

B.D. Honeyman (PI)¹, A.J. Francis², C. J. Dodge²,
J.B. Gillow² and P.H. Santschi³

¹Environmental Science and Engineering Division, Colorado School of Mines; ²Environmental Sciences Department, Brookhaven National Laboratory; ³Texas A&M University at Galveston



Historically, expectations of minimal aqueous Pu transport of have been confounded by the apparent contradiction:


- an understanding of low Pu solubility (based on its inorganic speciation) and
- the observed transport of Pu sometimes at substantial distances from its presumed source.

Of the relatively limited number of papers on Pu environmental speciation, a majority have implicated 'organic matter' as a transport agent*.


This project focuses on the role of bacteria in the production of EPS: as Pu mobilization and immobilization agents.

*(e.g., Nelson *et al.*, 1990; Orlandini *et al.*, 1990; Loyland *et al.*, 2001; Honeyman, 1998; Santschi *et al.*, 2002)


Comparison of U and Pu solubility




Examples of environmental measurements




Examples of environmental measurements




Pu release from vegetated soils as a function of quality and quantity of DOC



PAGE (Gel electrophoresis) of $^{239,240}\text{Pu}$ from Soil Resuspension Experiment








Pourbaix Diagram (Eh/ pH) diagram for the system Pu-O₂-CO₂-H₂O at 25 °C and 1 bar total pressure for $\Sigma \text{Pu(aq)} = 10^{-8} \text{ M}$ and total carbonate (C_T) = $10^{-2.0} \text{ M}$



- A: Aerobic bacteria
- B: Denitrifiers
- C: Mn-reducers
- D: Fe reducers
- E: Fermenters
- F: Sulfate reducers

Pu mobility / immobility as a transformational process






Key: **A.** Pu carrier-phase dissolution; **B.** Trapping of colloidal Pu by EPS; **C.** Biodegradation of Pu-EPS; **D.** Enhanced Transport of Pu by EPS.


Characterization of EPS

TEM of <math><0.45 \mu\text{m}</math> EPS from
Shewanella putrefaciens



Bar = 200 nm

TEM image of *Clostridium* sp.



J.B.Gillow (unpublished)

TEM image of *Clostridium* sp. after
48 hours growth showing
polysaccharide surrounding cell (bar =
1000nm)



Soil Bacterial cultures

(C) ↓ Centrifugation (3600 g, 30 min)

Pellet

Dissolved in D.W. with 3% NaCl for 1 hr (C)

Capsular exopolymer

Supernatant

(C) ↓

Pellet

(C) ↓

Dissolved in D.W. with 3% NaCl, stirring 1 hr, add EtOH

Crude exopolymer

(C) ↓

Purification of EPS for three times with EtOH,

Mixture

(C) ↓


Removing Protein with proteinase or pronase for 12 hr at 37 C, then dialyzed, lyophilized





Pure EPS

EtOH precipitation (overnight, filtration)

Objectives

- Determination of amphiphilic character of EPS through evaluation of chemical composition of hydrophilic carbohydrates and uronic acids, and more hydrophobic proteins, through the use of Hydrophobic Interaction Chromatography (HIC).
- Pu(IV,V) partitioning to EPS from *Pseudomonas* with(w/) and without(w/o) proteins.








Key: **A.** Pu carrier-phase dissolution; **B.** Trapping of colloidal Pu by EPS; **C.** Biodegradation of Pu-EPS; **D.** Enhanced Transport of Pu by EPS.



GC-MS Spectra of EPS (C.-C. Hung): *Clostridium* sp. Exudates (> 6 kDa)

RT: 12.17 - 65.49





Hydrophobicity and MW determinations:




- Waters HPLC
- Amersham HiTrap, 2 butyl 1 ml columns in series

- Waters HPLC
- Tosoh Biosciences G4000 PWxl, 7.8 mm x 30 cm, particle size 10 μm , with guard column 6 mm x 4 cm

Results


HCA, Protein, Charged Groups



Results


Molecular Weight (MW) by SEC

Overlay SEC Chromatograms of *Pseudomonas fluorescens*




Results

Molecular Weight (MW) by SEC




Summary of characterization work:

- The **neutral monosaccharides** in this EPS consist of rhamnose, fucose, ribose, arabinose, xylose, mannose, galactose and glucose.
- The **acidic groups** in this EPS are mainly composed of **carboxylic acid and minor polyanionic groups**, e.g., sulphate and phosphate.
- **45 - 70 % of total carbohydrates are uronic acids**, and total carbohydrates made up 26-31% of organic carbon.
- Besides the neutral and acidic sugars in the EPS, **EPS also contained 9 % of proteins** (% of carbon), which makes the EPS amphiphatic (amphiphilic).




Pu / EPS complexes

- Evaluation of Pu binding sites ('empirical adequacy'): affinity distribution
- Determination of conditional Pu /EPS complexes: ligand competition method




Linear combination of independent ionizable sites?

'Smoothness Hypothesis'




'Discrete' peaks




} Affinity distributions

Comparison of EPS 'ligand' specific concentrations



Comparison of Pu / EPS binding



Microbially-enhanced Pu mobilization

- Soil analysis
- Batch studies
- 'Static' columns

Radioactive Properties of Key Plutonium Isotopes

Isotope	Half-Life (years)	Specific Activity (Ci/g)	Decay Mode	Alpha (MeV)	Beta (MeV)	Gamma (MeV)
Pu-238	88	17	α	5.5	0.011	0.0018
Pu-239	24,000	0.063	α	5.1	0.0067	<
Pu-240	6,500	0.23	α	5.2	0.011	0.0017
Pu-241	14	100	β	<	0.0052	<
Pu-242	380,000	0.0040	α	4.9	0.0087	0.0014


[US DOE, Plutonium Fact Sheet, ANL, 2001]

$^{239,240}\text{Pu}$: *in situ* ('aged')


^{241}Pu : tracer

^{242}Pu : tracer

μ -XRF of Rocky Flats Soil







[1 mm² map of Fe (red) and Sr (blue) of 'as-rec'd' RF soil; incident beam energy 17.5 KeV]



[Microbeam (10 x 20 μ m spot) X-ray fluorescence of RF soil elements characteristic of loam/sandy soil]

[Analyses performed at NSLS Envirosuite beamline X27A; note that 'as-rec'd' the soil contained 1.6 ng ^{239,240}Pu g⁻¹ dry wt., 5 mg Fe g⁻¹ dry wt. (0.5 wt. %), 1.97% TOC]




Key: **A.** Pu carrier-phase dissolution; **B.** Trapping of colloidal Pu by EPS; **C.** Biodegradation of Pu-EPS; **D.** Enhanced Transport of Pu by EPS.

Fe(II)

Pu_{aq}

Pu_s



Biotransformation of Pu in Soil from Rocky Flats



Incubation time = 10 days




Biostimulation of RF soil:

A. The pH dropped to 7 due to metabolism of lactate; and to 5 due to glucose fermentation. 25 days, E_H glucose = -180 mV, lact. = -123 mV.

B. Glucose fermentation released 50 wt. % of the total reducible iron oxide from the soil; lactate metabolism released 3.5 wt. %.




Formation of Pu Colloids in Soil due to Microbial Action



[^{242}Pu nitrate ($1.5 \times 10^{-6}\text{M}$) added to soil resulted in 1) inorganic colloid formation, 2) sorption to soil and 3) remobilization as a colloid ($<0.45 \mu\text{m}$) upon incubation with glucose and lactate. $<0.1\%$ of total ^{242}Pu was in this fraction of unamended samples.]


$^{239,240}\text{Pu}$ and ^{242}Pu Fate at 77 Days Incubation



[Indigenous $^{239,240}\text{Pu}$ and spiked ^{242}Pu were mobilized ($<0.45 \mu\text{m}$) from RF soil with both glucose and lactate; the majority of the $^{239,240}\text{Pu}$ in the 'as-rec'd' soil was associate with the organic and inert fraction.]


[mol % of total added]

Re-distribution of ^{242}Pu in Soil at 77 Days Incubation



[^{242}Pu was recovered with the reducible iron oxide fraction (CBD extraction) in unamended samples, however, it was redistributed to another phase after biostimulation with glucose or lactate]





^{242}Pu and Total Carbohydrates



[At 77 days, the colloidal ^{242}Pu was correlated with an increase in suspended carbohydrates ($<0.45 \mu\text{m}$); this indicates that microbial exudates may play a role in Pu mobilization in the incubation experiments]

Summary of Soil Biotransformation Studies

- Pu was below detection for microprobe XRF, however **discrete Fe phases** were observed.
- Biostimulation with **glucose released** a significant amount of **Fe** while Fe(II) was readsorbed in lactate amended samples.
- A **^{242}Pu spike** rapidly sorbed to soil, however it was **remobilized** due to microbial action; under highly reducing, fermentative conditions 0.7% of the total was detected in the $<0.45\mu\text{m}$, $>30\text{kDa}$ fraction.
- The majority of Pu was released with the reducible iron oxide fraction of unamended samples, however this was not the case after biostimulation.
- **The indigenous $^{239,240}\text{Pu}$ was remobilized, to a lesser extent than the spike**, but this may be due to different mineralogical association (majority resided with the organic and inert fraction).
- There was an increase in soluble carbohydrates in the biostimulated samples implicating **microbial exudates in stabilizing Pu** in the colloidal fraction.




Key: **A.** Pu carrier-phase dissolution; **B.** Trapping of colloidal Pu by EPS; **C.** Biodegradation of Pu-EPS; **D.** Enhanced Transport of Pu by EPS.

'Static column' experiments


Assess Pu mobility under solid / solution ratios appropriate to *in situ* conditions

Static Column Incubation Experiment: 10 vs. 47 Day Incubation




$^{239,240}\text{Pu} = \sim 70 \text{ pCi / g}; 0.5 \text{ w/v glucose}; 0.015\% \text{ w/v NH}_4\text{Cl}$

Pu transport enhancement by EPS




Sorbed Pu-241 to RF soil for 24hrs then 22 mg/L EPS (~ 10 mg/L OC) injected

EPS Facilitated Pu Mobilization




^{239,240}Pu mobilized: 0.018% v. 0.004% in the control

● $^{239,240}\text{Pu}$ vs. ^{241}Pu Tracer



Summary: Pu mobility as a transformational process



Acknowledgements

- Cetin Kantar (Ph.D. Student / post-doc)
- Ruth Tinnacher (Ph.D. student)
- Angelique Diaz (Ph.D. student)
- NABIR Program