


## THE KINETICS OF QUARTZ DISSOLUTION AND PRECIPITATION

J. A. Apps  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

Silica precipitation in geothermal power plants and in reservoir formations is considered to be a potential problem area in the successful development of geothermal power from liquid dominated resources. In order to gain insight into the significance of this problem and to estimate the rates of precipitation of silica under varying conditions, a literature review and evaluation was made.

Data on the kinetics of quartz dissolution and precipitation in water was fitted to an expression derived from absolute rate theory, assuming that the mechanism could be represented by the equation:



$k'_f$  and  $k'_b$ , the specific forward and back rate constants were obtained for various sets of literature data at temperatures ranging from 25°C to 625°C, primarily along the saturation curve for water. The rate constants were plotted as  $\log k'_f$  or  $\log k'_b$  against  $1/T$  (K<sup>-1</sup>) and a linear fit obtained.  $\log k'_f$  is found to vary between -9.67 (s<sup>-1</sup>) at 25°C and -0.07 at 350°C, and  $\log k'_b$  between -6.19 and +1.83 for the same temperature range.

Although there is sufficient literature data to observe trends in quartz dissolution and precipitation in water, the data are too variable in quality to estimate precipitation rates accurately. Kinetic data on other silica polymorphs or in other solutions is quite inadequate to draw any significant conclusions.

## **DISCLAIMER**

**This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

Figures 1, 2, 3.

Absolute rate theory to describe kinetics of silica solubility in the aqueous phase.

(See S. Glasstone, K. J. Laidler and H. Eyring, Theory of Rate Processes, McGraw Hill Book Co., New York, 1941).

A = surface area,  $\sigma$  = surface site density. For other symbols, see reference.





Figure 1

$$\begin{aligned}
 \frac{dn_f}{dt} &= \frac{\kappa k T}{h} [C^*] A\sigma \\
 &= \boxed{\frac{\kappa k T}{h} K_f^*} [\text{SiO}_2] [\text{H}_2\text{O}]^2 A\sigma \\
 &= k'_f [\text{SiO}_2] [\text{H}_2\text{O}]^2 A\sigma
 \end{aligned}$$

Figure 2


$$\begin{aligned}
 \frac{dn_f - dn_b}{dt} &= k'_f A\sigma [\text{SiO}_2] [\text{H}_2\text{O}]^2 - k'_b A\sigma [\text{Si}(\text{OH})_4] \\
 \frac{n}{m} &= \frac{n_0}{m} e^{\frac{-A\sigma k'_b t}{m}} + \frac{k'_f}{k'_b} \left( 1 - e^{\frac{-A\sigma k'_b t}{m}} \right)
 \end{aligned}$$

Figure 3




XBL 767-3228

Figure 4. The dissolution of quartz in water at 264°C. The data are by Kolvoord and Yorgason (Kennecott Exploration, Inc., Salt Lake City). The solid line represents the best fit to the data using equation in Figure 3.



XBL 767-3229


Figure 5. The dissolution and precipitation of quartz in water at 100°C. The data is by Van Lier et al. (J. Phys. Chem. 64 p. 1675 (1960)). The solid lines represent best fits to the data using equation in Figure 3.



XBL767-3230


Figure 6. The specific forward rate constants,  $k'_f$ , for the dissolution of quartz in water plotted against the reciprocal of the absolute temperature. Values of  $k'_f$  are calculated from literature data.





XBL767-3232

Figure 7. The specific back rate constant,  $k'_b$ , for the dissolution of quartz in water, plotted against the reciprocal of the absolute temperature. Values of  $k'_b$  are calculated from literature data.



XBL 767-3231

Figure 8. The equilibrium constant, K, for the dissolution of quartz in water, plotted against the reciprocal of the absolute temperature. Values of K are computed from literature data.