

UCRL-PROC-234670



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Detailed chemical kinetic oxidation mechanism for a biodiesel surrogate

Olivier Herbinet, William J. Pitz, Charles K.
Westbrook

September 18, 2007

2007 Fall Meeting of the Western States Section of the
Combustion Institute
Livermore, CA, United States
October 16, 2007 through October 17, 2007

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

2007 Fall Meeting of the Western States Section of the Combustion Institute
Sandia National Laboratories, Livermore, CA
October 16 & 17, 2007.

Detailed chemical kinetic oxidation mechanism for a biodiesel surrogate

O. Herbinet, W. J. Pitz and C. K. Westbrook

Lawrence Livermore National Laboratory, California 94550, USA

A detailed chemical kinetic mechanism has been developed and used to study the oxidation of methyl decanoate, a surrogate for biodiesel fuels. This model has been built by following the rules established by Curran et al. for the oxidation of n-heptane and it includes all the reactions known to be pertinent to both low and high temperatures. Computed results have been compared with methyl decanoate experiments in an engine and oxidation of rapeseed oil methyl esters in a jet stirred reactor. An important feature of this mechanism is its ability to reproduce the early formation of carbon dioxide that is unique to biofuels and due to the presence of the ester group in the reactant. The model also predicts ignition delay times and OH profiles very close to observed values in shock tube experiments fueled by n-decane. These model capabilities indicate that large n-alkanes can be good surrogates for large methyl esters and biodiesel fuels to predict overall reactivity, but some kinetic details, including early CO₂ production from biodiesel fuels, can be predicted only by a detailed kinetic mechanism for a true methyl ester fuel. The present methyl decanoate mechanism provides a realistic kinetic tool for simulation of biodiesel fuels.

1. Introduction

In recent years, biodiesel has become interesting as an additive to diesel fuel for two main reasons. This renewable alternative fuel can reduce dependence on imported petroleum and can also contribute to environment preservation by lowering net emissions of greenhouse gases. The use of biodiesel in diesel engine decreases emissions of pollutants such as carbon monoxide, unburned hydrocarbons and particulate matters [1], although a slight increase in emissions of nitrogen oxides is observed in some cases [2].

Biodiesel is a multiple component mixture of mono-alkyl esters of long-chain fatty acids derived from vegetable oils and animal fats. Most biodiesel fuels used in the world are made from soy oil and rapeseed oil by transesterification with an alcohol. The derived soy and rapeseed biodiesels are complex mixtures composed of mainly five saturated and unsaturated methyl esters (when methanol is used for the transesterification process): methyl palmitate (C₁₇H₃₄O₂), methyl stearate (C₁₉H₃₆O₂), methyl oleate (C₁₉H₃₄O₂), methyl linoleate (C₁₉H₃₂O₂) and methyl linolenate (C₁₉H₃₀O₂). Average compositions of soybean and rapeseed biodiesels [3] are given in

Table 1.

Table 1: Average compositions (%) of soybean and rapeseed biodiesels [3].

Esters	Soybean biodiesel	Rapeseed biodiesel
methyl palmitate	6-10%	4.3%
methyl stearate	2-5%	1.3%
methyl oleate	20-30%	59.9%
methyl linoleate	50-60%	21.1%
methyl linolenate	5-11%	13.2%

The structures of these components are shown in Figure 1, showing the considerable structural similarities in these chemical species, each with a methyl ester attached to a large hydrocarbon fragment.




Figure 1: Structure of the main components found in soybean and rapeseed oils methyl esters and of methyl decanoate.

Very few kinetic studies of biodiesel fuel combustion have been carried out, either experimentally or using computational modeling. There are several reasons for this lack of attention; biodiesel fuels have become important only quite recently, and they are generally very large fuel molecules that challenge the capabilities of kinetic modeling. As a result, past research in this area has followed two major paths. Experiments and kinetic modeling of much smaller methyl esters have addressed the special features of methyl ester oxidation, and combustion of large biofuels has been studied by assuming that large methyl esters can be approximated as being fundamentally the same as large n-alkanes. The largest methyl ester that has been studied kinetically is methyl butanoate, with a chain of only 4 carbon atoms connected to the methyl ester group. Kinetic modeling of methyl butanoate has concluded that this fuel reproduces kinetic features of the oxidation of the methyl ester but does a poor job of reproducing kinetic features of diesel fuels with their chains of 16 - 18 carbon atoms. Other

studies have used kinetic models for n-alkanes as large as n-hexadecane to simulate the combustion of the large methyl ester molecules in actual biodiesel fuels. The present work is intended to provide a reliable kinetic model for a methyl ester fuel that is much larger than the previous methyl butanoate. Instead of the 4 carbon atoms chain of methyl butanoate, the current work provides a kinetic mechanism for methyl decanoate, with a chain of 10 carbon atoms with a methyl ester group attached (Figure 1). Methyl decanoate reacts in a manner that is much closer to actual biodiesel fuel than methyl butanoate, including both early production of CO₂ from the methyl ester group and burning in a manner very similar to conventional diesel fuel.

We will review past work in the area of methyl ester combustion, leading to a description of the goals of the present work.

The oxidation of methyl butanoate (C₅H₁₀O₂) has been the subject of several papers. A detailed chemical kinetic mechanism for the combustion of methyl butanoate was developed by Fisher et al. [4], which was validated against the limited available data obtained at low temperature, subatmospheric conditions in closed vessels, using pressure measurements as the main diagnostic. More recently, Metcalfe et al. [5] studied the oxidation of methyl butanoate and ethyl propanoate in a shock tube. A revised detailed kinetic mechanism based on the work of Fisher et al. [4] for methyl butanoate and a new submechanism for ethyl propanoate were used to simulate measured ignition delay times with good agreement. Gail et al. performed a wide-ranging kinetic modeling study of the oxidation of methyl butanoate [6]. They obtained experimental ignition data and species profiles in a jet stirred reactor, a variable pressure flow reactor and an opposed-flow diffusion flame. A revised kinetic model based on the Fisher et al. mechanism was validated from the jet stirred reactor data. This model was shown to reproduce data obtained in a variable pressure flow reactor and in an opposed-flow diffusion flame. Sarathy et al. [7] performed an experimental study of methyl crotonate (C₅H₈O₂ unsaturated methyl ester) in a jet stirred reactor and an opposed-flow diffusion flame in order to compare with experimental data obtained for methyl butanoate [6] and understand the role of the double bond in the methyl ester.

Vaughn et al. [8] studied the combustion of bio-ester fuel droplets in microgravity. They measured ignition time of neat methyl esters (such as methyl butanoate, methyl decanoate, methyl dodecanoate, and methyl oleate) and commercial soy oil methyl esters. Ignition delay times obtained during this study showed that methyl decanoate and methyl dodecanoate are better surrogates for commercial soy oil methyl esters than methyl butanoate, in agreement with conclusions of Fisher et al. [4] and Gail et al. [6].

Dagaut et al. [9] performed an experimental study of the oxidation of rapeseed oil methyl ester in a jet stirred reactor at 1-10 atm over the temperature range 800-1400 K. Experimental species profiles were compared with computed mole fractions from a mechanism for oxidation of n-hexadecane, which had been validated against experiments in a JSR [10]. The agreement was shown to be satisfactory and n-hexadecane appeared to be a good surrogate for rapeseed oil methyl ester in the conditions of the study. However, the n-hexadecane mechanism was unable to predict the early production of CO₂ which was observed in the experiments.

A good knowledge of the kinetics of the reaction of biodiesel fuels at both high and low temperature is necessary to perform reliable simulations of ignition, combustion and emissions in homogeneous charge compression ignition (HCCI) and diesel engines. Modeling of the oxidation of methyl butanoate provided a better understanding of the chemistry of methyl esters combustion, but methyl butanoate is not a good surrogate for commercial biodiesel fuels because

its alkyl chain is too short. In this work, a detailed chemical kinetic mechanism has been developed and used to study the oxidation of methyl decanoate, which we feel is a much better surrogate for biodiesel fuel than methyl butanoate. This model is compared with the limited available experimental data obtained in a motored engine [11,12], and it is used to model rapeseed oil methyl ester experiments in a JSR [9] and shock tube ignition of n-decane [13].

2. Description of the chemical kinetic mechanism

The detailed chemical kinetic mechanism of the oxidation of methyl decanoate has been developed by using the same systematic rules which have been described by Curran et al. for n-heptane and iso-octane [14,15]. Some kinetic parameters and thermochemical properties used in this mechanism have been updated from more recent data from the literature. The entire mechanism will be available, including the kinetic parameters and thermochemistry, in Chemkin format on our web page at:

http://www-cmls.llnl.gov/?url=science_and_technology-chemistry-combustion

Structure of the methyl decanoate kinetic chemical mechanism

The model presented in this paper was developed from previous n-heptane and iso-octane [14,15] and methyl butanoate mechanisms [4] by combining them with the low and high temperatures chemistry specific to methyl decanoate. The overall primary oxidation reaction pathways in the methyl decanoate mechanism are shown in Figure 2. The same general pathways apply to all hydrocarbon fuels, although the details of each step depend on the size and structure of the specific fuel being studied. In general, the reaction classes from Curran et al. were used, but accommodations were required to take into account the fact that the methyl ester group in methyl decanoate changes some of the details of the mechanism.




Figure 2: Primary oxidation reactions taken into account for the development of the methyl decanoate mechanism.

High temperature part. At high temperatures, unimolecular decompositions of the fuel and H-atom abstractions from the fuel lead to the formation of alkyl and alkyl-ester radicals. Reactions













of these radicals, which are known to be pertinent at high temperature, are isomerizations, decompositions to olefins or unsaturated esters plus smaller radicals, and direct abstractions by O_2 to olefins or unsaturated esters plus HO_2 . Olefins and unsaturated esters formed through these primary routes react in turn through the same types of reactions as the fuel and through other reactions specifically due the presence of the double bond (additions of radicals to the double bond, decomposition by retro-ene reactions).

The unimolecular initiation reactions of methyl decanoate were written in the recombination direction and the kinetic parameters for the decomposition direction were then calculated using the thermodynamic properties. A rate constant of $1.0 \times 10^{14} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ was used for the recombination of an H atom and any C_{11} ester radical, $3.0 \times 10^{13} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ for the two reactions of recombination involving C-O bonds, $1.8 \times 10^{13} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ for the recombination leading to the formation of the C-CO bond, $3.0 \times 10^{13} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ for the formation of the C-C bond involving a methyl radical and an alkyl-ester radical, and $8.0 \times 10^{12} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ for the remaining C-C bonds formed through the reactions of recombination of alkyl and alkyl-ester radicals. Reactions of recombination involving C-C and C-O bonds in methyl decanoate are shown in Table 2.

H-atom abstractions from methyl decanoate by H, CH_3 , C_2H_3 , C_2H_5 , O, O_2 , OH, HO_2 , CH_3O , and CH_3O_2 have been included, using kinetic parameters recommended by Curran et al. [14]. Distinctions between three types of H atoms were made: primary H atoms in the two methyl groups at each end of the molecule, secondary H atoms bonded to the conventional secondary, internal carbon atoms, and the two H atoms bonded to the carbon atom adjacent to the carbonyl group. There is a lack of data concerning the rate constants of H-atom abstractions involving these two H atoms. These H atoms have C-H bond energies similar to those for tertiary C-H bonds, so we have followed [4] and used H atom abstraction rates from tertiary bonds in other molecules for these H atoms.

Alkyl and alkyl-ester radical decompositions were written in the reverse direction (addition of a radical to a double bond). Kinetic parameters are based on a recent review by Curran for the alkyl radicals [16] and from the methyl butanoate mechanism [4] for reactions involving atoms of the ester group. The kinetic parameters for addition of radicals to the oxygen of the C=O bond have been updated from the study of methyl radical addition to the C=O bond by Henry et al. [17]. Kinetic parameters used for isomerizations, or H-atom shifts, of radicals were taken from quantum calculations performed by Matheu et al. [18]. Some required rate constants not calculated by these authors are estimated using “structure-reactivity” relationships. The rate constant used for the direct abstraction from alkyl and alkyl-ester radicals by O_2 is $1.6 \times 10^{12} \times \exp[-5000(\text{cal} \cdot \text{mol}^{-1})/RT] \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ [19].

Table 2: Reactions of recombination involving C-O and C-C bonds in methyl decanoate.

Reaction of recombination	Rate constant ($\text{cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$)
	3.0×10^{13}
	3.0×10^{13}
	1.8×10^{13}
	8.0×10^{12}
	8.0×10^{12}
	8.0×10^{12}
	8.0×10^{12}
	8.0×10^{12}
	8.0×10^{12}
	8.0×10^{12}
	8.0×10^{12}
	3.0×10^{13}

As far as olefins and unsaturated esters are concerned, H-atom abstractions and molecular decompositions by retro-ene reactions were written in a systematic way. Rate constants for primary, secondary and tertiary H-atoms abstractions from olefins and unsaturated esters are the same as those described above for the methyl decanoate molecule. For allylic and vinylic H-atoms, kinetic parameters are those recommended by Curran et al. [14] for small species (propene, 1-butene). The rate constant for the molecular decomposition of olefins and unsaturated esters by retro-ene reaction is from King [20]: $3.98 \times 10^{12} \times \exp[-57630(\text{cal} \cdot \text{mol}^{-1})/\text{RT}] \text{ s}^{-1}$. Only unimolecular initiations involving C-C and C-H bonds in the beta position of the double bond have been taken in account. Other C-C, C-O and C-H bond breakings were not included because of their higher activation energies. Unimolecular decompositions of olefins and unsaturated esters by scission of the allylic C-C bond were written in the forward direction and the scission of the allylic C-H bonds in the reverse, recombination direction. Rate constants of $2.5 \times 10^{16} \times \exp[-71000(\text{cal} \cdot \text{mol}^{-1})/\text{RT}] \text{ s}^{-1}$ and $1.0 \times 10^{14} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ [14] were used, respectively. Additions of OH radicals to the double bond of olefins and unsaturated radicals were written (rate constants of $1.129 \times 10^5 \times T^{2.28} \times \exp[1241(\text{cal} \cdot \text{mol}^{-1})/\text{RT}] \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ from Zhu et al. [21]) and additions of H atoms and HO₂ radicals were considered in two other parts of the mechanism

(alkyl and alkyl-ester radicals C-H β -scission decompositions in the high temperature part and QOOH C-O β -scission decompositions in the low temperature part).

Rate constants for isomerizations of alkenyl, allylic and vinylic radicals are from [18]. Decompositions of these radicals were considered through the reverse additions and kinetic parameters are the same as those presented in the methyl decanoate section above.

Low temperature part. The low temperature part of the mechanism was built by adapting the kinetic scheme used in the well-validated n-heptane and iso-octane mechanisms. Again, some accommodations were required due to the presence of the methyl ester group in the fuel. Figure 3 displays a potential energy diagram (derived from the n-heptane and iso-octane mechanisms [14,15]) showing the major species and the main reaction pathways involved in the low temperature part of the methyl decanoate mechanism.




Figure 3: Potential energy diagram for the addition of R to O₂ and subsequent reactions (low temperature scheme used in the methyl decanoate mechanism).

The first step of the low temperature mechanism is the addition of alkyl and alkyl-ester radicals to O₂ (Figure 4). Rate constants of $4.52 \times 10^{12} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$, $7.54 \times 10^{12} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ and $1.41 \times 10^{13} \text{ cm}^3 \cdot \text{mol}^{-1} \cdot \text{s}^{-1}$ were used for additions of primary, secondary and tertiary radicals to O₂. The subsequent alkyl and alkyl-ester peroxy radicals (RO₂) react then by isomerizations to hydroperoxy alkyl and hydroperoxy alkyl-ester radicals (QOOH). Isomerizations through 5, 6, 7 and 8 member cyclic transition states have been included (Figure 5). Rate constants for isomerizations are from Curran et al. [15] and are presented in Table 3.




Figure 4: Addition of an alkyl-ester radical (R) to O₂.




Figure 5: Examples of isomerizations permitted for peroxy alkyl-ester radicals (RO₂).

Reactions of hydroperoxy alkyl and hydroperoxy alkyl-ester radicals (QOOH) are displayed in Figure 6, including the second addition of O₂ forming hydroperoxy peroxy radicals (O₂QOOH), the decomposition to cyclic ethers plus OH, and the C-O β-scission decomposition to HO₂ and alkyl or alkyl-ester radicals. Other reactions of C-C β-scissions have not been taken in account because of their higher activation energy [22]. Rate constants for QOOH decompositions to cyclic ethers plus OH and to olefin plus HO₂ are those recommended by Curran et al. [14] (Table 3). The rate expression of the last reaction type is written as the reverse addition of olefin + HO₂, with different activation energies depending on whether the HO₂ adds to a primary or secondary carbon atom. The direct eliminations from RO₂ (leading to olefins + HO₂) which intervenes in the more recent scheme proposed for the low temperature of alkyl radicals [22-24] were not included in the methyl decanoate mechanism and were also not included in the n-heptane and iso-octane mechanisms on which it is based. When developing these mechanisms it was believed that the channel of formation of olefins plus HO₂ occurred via C-C β-scissions following isomerizations through 5 member ring and kinetic parameters of these and related reactions were

calibrated to reproduce the formation of olefins. Reaction rate rules for direct elimination of HO₂ from RO₂ in concert with RO₂ isomerizations need to be derived and validated for lower molecular weight systems where much more experimental data is available before they can be successfully applied to high molecular weight systems like the present work. We anticipate that future revisions of the present mechanism may address this direct molecular elimination reaction pathway, but the present mechanism represents an internally consistent and predictive modeling tool in its present form.




Figure 6: Reactions of an hydroperoxy ester radical (QOOH) involved in the low temperature part of the mechanism.

For O₂QOOH species, only those specific isomerizations leading to ketohydroperoxide plus OH have been included (Figure 7). Other isomerizations described as “alternative paths” by Silke et al. [25], have not been included and are not expected to be significant for long, straight chain hydrocarbons such as n-alkanes or the straight-chain methyl esters such as methyl decanoate, both of which have many possible paths for ketohydroperoxide + OH production. Kinetic parameters for O₂QOOH isomerizations (producing ketohydroperoxide + OH) are derived from the rate constants for RO₂ isomerizations to QOOH, using the same A factors and activation energies 3 kcal.mol⁻¹ smaller than for the analogous RO₂ isomerizations [14,15].








Figure 7: Reaction of isomerization of O₂QOOH to ketohydroperoxide + OH.

Decomposition reactions of ketohydroperoxides lead to the formation of a new OH radical and another radical, providing chain branching, following reaction type 24 from Curran et al.. The rate of hydroperoxide decomposition (O-O scission) used in the mechanism is $1.05 \times 10^{16} \times \exp[-41600(\text{cal} \cdot \text{mol}^{-1})/RT] \text{ s}^{-1}$ [26]. The following reactions of disproportionation between radicals have been included with rate expressions suggested by Curran et al. [14,15]:



The above described mechanism involves 3012 species and includes 8820 reactions. The high numbers of reactions and species are caused by the numerous types of reactions taken in account but also to the fact that methyl decanoate is not a symmetric molecule like an n-alkane. Isomerizations of RO₂ species in the low temperature regime are also responsible for the large increase in the number of reactions because of the numerous permitted H-shifts.

Thermodynamic properties

Standard enthalpies of formation, entropies and specific heats of the molecules and radicals involved in the mechanism have been calculated using the THERM program developed by Ritter and Bozzelli [27]. This program is based on the group and bond additivity methods proposed by Benson [28].

The C-H bond dissociation energy of the carbon atom adjacent to the carbonyl group has been updated from the recent work of El-Nahas et al. [29] who studied the thermochemistry of methyl butanoate by performing quantum calculations. The value used in the mechanism for this specific bond is $94.1 \text{ kcal} \cdot \text{mol}^{-1}$. This compares closely to tertiary bond dissociation energies ($96.5 \text{ kcal} \cdot \text{mol}^{-1}$) as noted above for H atom abstractions from this site in methyl decanoate.

Table 3: Rate constants of the main reactions involving RO₂ and QOOH radicals (k=A×T^b×exp(Ea/RT), Units: cal, cm³, mol, s).

Reactions		A	b	Ea
R + O ₂ = RO ₂	R is a primary radical	4.52×10 ¹²	0.0	0.0
	R is a secondary radical	7.54×10 ¹²	0.0	0.0
	R is a tertiary radical	1.41×10 ¹³	0.0	0.0
RO ₂ = QOOH	5 atoms ring			
	primary H shifted	1.0×10 ¹¹	0.0	29400.0
	secondary H shifted	1.0×10 ¹¹	0.0	26850.0
	tertiary H shifted	1.0×10 ¹¹	0.0	24100.0
	6 atoms ring			
	primary H shifted	1.25×10 ¹⁰	0.0	24400.0
	secondary H shifted	1.25×10 ¹⁰	0.0	20850.0
	tertiary H shifted	1.25×10 ¹⁰	0.0	19100.0
	7 atoms ring			
	primary H shifted	1.563×10 ⁹	0.0	22350.0
	secondary H shifted	1.563×10 ⁹	0.0	19050.0
	tertiary H shifted	1.563×10 ⁹	0.0	17050.0
8 atoms ring				
primary H shifted	1.953×10 ⁸	0.0	25550.0	
secondary H shifted	1.953×10 ⁸	0.0	22050.0	
tertiary H shifted	1.953×10 ⁸	0.0	20050.0	
HO ₂ addition to C=C bonds	to a primary carbon atom	1.0×10 ¹¹	0.0	10750.0
	to a secondary carbon atom	1.0×10 ¹¹	0.0	11750.0
QOOH = OH + cyclic ether	oxirane	6.0×10 ¹¹	0.0	22000.0
	oxetane	7.5×10 ¹⁰	0.0	15250.0
	oxalane	9.375×10 ⁹	0.0	7000.0
	oxane	1.172×10 ⁹	0.0	1800.0

3. Results and discussion

A detailed kinetic mechanism is incomplete without a validation study comparing computed results from the mechanism with measurements from appropriate experiments. In many cases, laboratory experiments from shock tubes, laminar flames, stirred and flow reactors, and many other idealized systems are available, as well as experiments in engines or other practical systems. The laboratory experiments are especially valuable when they provide species-dependent and time-dependent information that provide a particularly demanding test of a mechanism, in contrast with experiments that provide only an integrated test, such as an ignition delay time or laminar burning velocity. In the present case of methyl decanoate, however, we could not identify any fundamental laboratory experiments using methyl decanoate, and only one engine experiment was found that specifically used methyl decanoate as a fuel.

As a result, we have been able only to compare computed results for methyl decanoate combustion with experimental results for two closely related fuels, n-decane, for which we have used experimental results from low and high temperatures shock tube experiments, and rapeseed oil methyl ester, for which we have used experimental results from a jet-stirred reactor. Finally, we compared computed results with experimental data obtained in a motored engine [11,12], which used methyl decanoate as a fuel, in addition to other cases using n-heptane and commercial diesel fuel.

Comparison with rapeseed oil methyl esters jet stirred reactor experiments

Dagaut et al. [9] studied oxidation of rapeseed oil methyl ester (RME) in a jet stirred reactor (JSR) at pressures of 1 and 10 atm, at temperatures from 800 to 1400 K and at several residence times (0.07, 0.1 and 1s) and equivalence ratios (0.25 – 1.5). Quantification of the species leaving the reactor was performed by gas chromatography (FID, TCD) and a GC/MS was used for their identification. Quantified species were 1-alkenes from ethylene to 1-heptene, methane, hydrogen, carbon dioxide, carbon monoxide and oxygen. Dagaut et al. did not report any data about the formation of unsaturated methyl esters.

Two sets of simulations have been performed to draw comparisons with data performed at 10 atm (approximating pressures met in engines): first by using neat methyl decanoate as surrogate and second by using a surrogate composed of methyl decanoate and n-heptane. The choice of this last surrogate is justified later in the paper.

JSR simulations with neat methyl decanoate. Methyl decanoate ($C_{11}H_{22}O_2$) is a smaller molecule than those found in rapeseed oil methyl esters (global formula of $C_{17.92}H_{33}O_2$ from [9]). Therefore, the experimental inlet mole fraction of RME (0.005) has been multiplied by a factor 18/11 in order to match the inlet flux of carbon atoms in rapeseed oil methyl esters. This leads to slightly higher numbers of H- and O-atoms in our C_{18} surrogate: 36 and 3.27 respectively, compared with 33 and 2 in RME. Inlet oxygen mole fractions were deduced from the values used by Dagaut et al. for their simulations with n-cetane. These values were reduced slightly in order to take in account the presence of oxygen atoms in methyl decanoate. Inlet compositions of the reacting mixture used for the present simulations are given in

Table 4.

Table 4 : Inlet compositions of reacting mixtures used for methyl decanoate simulations.

Experimental conditions	P = 10 atm, $\phi = 0.5$, $\tau = 1$ s	P = 10 atm, $\phi = 1$, $\tau = 1$ s
Methyl decanoate	0.000818	0.000818
Oxygen	0.024352	0.011767
Nitrogen	0.974830	0.987415

The comparison between rapeseed oil methyl ester experimental data and computed results is shown in Figs. 8 and 9 and is globally satisfactory. The model allows matching the concentration profiles of most products of the reaction. At $\phi = 0.5$, mole fractions of 1-alkenes are well reproduced by the model except for ethylene and 1-hexene, whose concentrations are slightly underpredicted. At $\phi = 1$, the ethylene concentration is still underpredicted whereas the mole

fractions of 1-butene is too high, especially when the temperature increases. The same trend was also observed by Dagaut et al. using their n-cetane mechanism [9]. The model predicts formation of hydrogen at temperatures somewhat lower than in the experiments.

At both equivalence ratios, CO and CO₂ mole fractions are slightly under predicted but their concentration profiles are much better than the concentrations profiles obtained using the n-cetane mechanism. The early production of CO₂ occurring in the range of temperatures 800 - 850 K is much better reproduced by the methyl decanoate mechanism than by the n-cetane mechanism. Dagaut et al. recognized this shortcoming of using an n-hexadecane mechanism for RME simulations, and they discussed how the structure of the methyl esters in RME lead to early CO and CO₂ that cannot be captured in an n-alkane reaction mechanism. Routes of formation of these species are detailed later in the paper. Dagaut et al. also commented that the use of n-hexadecane produced concentrations of large 1-olefins larger than they found in their RME experiments, which they attributed to the longer uninterrupted carbon chains in n-hexadecane than in the methyl ester fuels.




Figure 8: Comparison of the methyl decanoate model with rapeseed oil methyl ester experiments in a jet stirred reactor ($P = 10$ atm, $\phi = 0.5$, $\tau = 1$ s) [9].




Figure 9: Comparison of the methyl decanoate model with rapeseed oil methyl ester experiments in a jet stirred reactor ($P = 10$ atm, $\phi = 1$, $\tau = 1$ s) [9].

JSR simulations with a mixture of methyl decanoate and n-heptane. The methyl decanoate mechanism was modified slightly to run a mixture of methyl decanoate and n-heptane:

unimolecular decomposition reactions and H atom abstractions of radicals from n-heptane added to take into account for the presence of this additional species in the inlet flow. Reactions of the resulting n-heptyl radicals were already included in the methyl decanoate mechanism. N-heptane was chosen as a co-reactant because it has been used frequently as a surrogate for diesel fuels and because it did not require the addition of too many reactions. An equimolar blend of methyl decanoate ($C_{11}H_{22}O_2$) and n-heptane (C_7H_{16}) enabled us to match the number of carbon and oxygen atoms in rapeseed oil methyl ester ($C_{17.92}H_{33}O_2$) used in the experiments [9]. It is worth noticing that the number of H atoms in the surrogate (38) is still higher than the number of H-atoms in rapeseed oil methyl esters. Inlet mole fractions have also been deduced from the values used by Dagaut et al. for their simulations. Inlet compositions used for the simulations are summarized in Table 5.

Table 5 : Inlet compositions of the reacting mixture used for the methyl decanoate – n-heptane surrogate simulations.

Experimental conditions	P = 10 atm, $\phi = 0.5$, $\tau = 1$ s	P = 10 atm, $\phi = 1$, $\tau = 1$ s
Methyl decanoate	0.000500	0.000500
n-Heptane	0.000500	0.000500
Oxygen	0.024670	0.012085
Nitrogen	0.974330	0.986915

Computed results from the methyl decanoate/n-heptane mechanism are very close to those obtained with the methyl decanoate model, and both simulations lead to similar reactivity. The CO_2 concentration is slightly lower in the case of the blend surrogate than in the case of the neat methyl decanoate surrogate. This is due to the fact that the inlet mole fraction of methyl decanoate (and therefore of the ester group) was higher in the case of the simulations run with neat methyl decanoate than in the case of the simulations performed with the blend surrogate. However, the predicted CO_2 levels from both surrogates containing methyl decanoate were considerably larger than those calculated from the n-hexadecane mechanism of Dagaut et al. [9].

Mole fractions of the 1-alkenes are higher in the case of the blend surrogate than for the neat surrogate, due to the presence of n-heptane as co-reactant and to the fact that β -scission decomposition products of alkyl and ester-alkyl radicals are not quite identical. Decomposition of alkyl radicals leads to 1-alkenes and smaller alkyl radicals which then decompose in turn, while decomposition of ester-alkyl radicals leads to the formation of 1-alkenes and smaller alkyl-ester radicals or unsaturated ester and alkyl radicals.




Figure 10: Comparison of the methyl decanoate – n-heptane model with rapeseed oil methyl ester experiments in a jet stirred reactor ($P = 10$ atm, $\phi = 0.5$, $\tau = 1$ s) [9].




Figure 11: Comparison of the methyl decanoate – n-heptane model with rapeseed oil methyl ester experiments in a jet stirred reactor ($P = 10$ atm, $\phi = 1$, $\tau = 1$ s) [9].




Route of formation of CO_2 at low temperature. The two models (neat methyl decanoate and methyl decanoate/n-heptane) make it possible to reproduce the early formation of CO_2 observed in the experiments by Dagaut et al.. A rate of production analysis was performed for the simulation at $\phi = 0.5$, $T = 800$ K, $P = 10$ atm and $\tau = 1$ s with the methyl decanoate mechanism to highlight the route of production of CO_2 at low temperature.


At these conditions, CO_2 is formed through four elementary reactions: the decomposition of the radical $OCHO$ (+M) to $H + CO_2$ (+M) (38%), the decomposition of $HOCHO$ to $H_2 + CO_2$ (31%), the reaction of $OH + CO$ leading to $H + CO_2$ (17%), and the decomposition of the radical CH_3OCO to $CH_3 + CO_2$ (7%). With the exception of the reaction of CO with OH and the decomposition of $HOCHO$ to $H_2 + CO_2$, the other two reactions derive uniquely from the methyl ester group in methyl decanoate and would not occur in the oxidation of n-hexadecane or any other n-alkane fuel.

The radical $OCHO$ derives primarily from the decomposition of radicals involved in the low temperature part of the mechanism (Table 6). These radicals have a carbonyl group on the ester methyl group and a radical center on the carbon atom in the β position of the ester carbonyl group. They come from the decomposition of ester cyclic ethers formed in the low temperature part of the mechanism. The radical $HOCHO$ mainly comes from the reaction of addition of the OH radical to formaldehyde. The radical CH_3OCO mainly comes from the decomposition of the numerous radicals having a radical center on the carbon atom in the γ position of the ester carbonyl group (Figure 12).

According to the rate of production analysis performed with the blend surrogate, the CO_2 production occurs via the same four main routes. The rates of these reactions are very similar except for the reactions involving indirectly the ester group of methyl decanoate: the decomposition of the radicals $OCHO$ and CH_3OCO (**Error! Reference source not found.**). Rates of these reactions are lower in the case of the blend surrogate because the inlet mole fraction of methyl decanoate is less than in the case of the neat surrogate.

Table 6: Reactions leading to the formation of the radical OCHO.

Reaction	Normalized Rate of Production (%)
	68
	4
	22

**Figure 12: Formation of the radical CH₃OCO from a 3-alkyl ester radical.****Table 7: Rates of the main reactions leading to the formation of CO₂ ($\phi = 0.5$, $T = 800$ K, $P = 10$ atm and $\tau = 1$ s).**

Reactions	Rate of Production ($\times 10^8$, mol.cm ⁻³ .s ⁻¹)	
	Blend Surrogate	Neat Surrogate
OCHO (+M) = H + CO ₂ (+M)	1.54	2.43
HOCHO = H ₂ + CO ₂	2.20	2.03
OH + CO = H + CO ₂	0.94	0.90
CH ₃ OCO = CH ₃ + CO ₂	0.41	0.61

These results are also consistent with the rate of production analysis we performed in similar conditions with the methyl butanoate mechanism developed by Fisher et al. [4]. For methyl butanoate, the main source of CO₂ is the decomposition of the cyclic ether formed in the low temperature part of the mechanism via OCHO (38%). The second source of formation of CO₂ is not the decomposition of HOCHO, but the decomposition of the radical CH₃OCO (24%). This difference can be explained by the fact that the kinetic parameters of the two reactions of decomposition of this radical (CH₃OCO = CH₃ + CO₂ and CH₃OCO = CH₃O + CO) have been updated in the methyl decanoate mechanism from a recent work of Glaude et al. about dimethyl carbonate [30].

Early CO₂ production from the methyl ester group in methyl decanoate has important practical implications in diesel ignition and soot production. Recent kinetic modeling of ignition under diesel conditions [1] showed how the presence of oxygen atoms in the fuel can reduce soot production. However, if that oxygen immediately produces CO₂, as in methyl butanoate and

methyl decanoate (and, by implication, in all biodiesel fuels), that fuel-bound oxygen is less effective in reducing soot production.

Methyl decanoate ignition delay times comparison with n-decane and n-heptane.

In the previous section, we have examined the differences between the use of kinetic mechanisms for n-alkane and methyl ester fuels to simulate experimental results for combustion of methyl ester fuels. In this section, we address the reverse problem, the use of a kinetic mechanism for a methyl ester to simulate experimental results for combustion of an n-alkane fuel. The overall goal of this work is to compare how closely the combustion properties of large n-alkane fuels and related large methyl esters resemble each other, and how well their kinetic reaction mechanisms can reproduce the combustion properties of each other.

Ignition delay times calculated from the methyl decanoate mechanism were compared to experimental results for n-decane. N-decane was selected for the comparison because it has the same number of carbon atoms as the alkyl chain of methyl decanoate, and numerous experimental data of the oxidation of n-decane are available.

Davidson et al. [13] measured OH concentration/time histories behind reflected shocks in n-decane ignition. Experiments performed at a fuel concentration of 300 ppm, at temperatures ranging from 1479 to 1706 K, at pressures from 2.08 to 2.21 atm, and at an equivalence ratio of 1.0 (99.505 % Argon) have been compared to simulations from the methyl decanoate mechanism (simulations were performed with an average pressure of 2.15 atm). The agreement between the two sets of data is quite good in these conditions (Figure 13). Calculated OH concentration time histories for three temperatures are shown in Figure 14 (actual pressures were used for the simulations). These three profiles are very close to the experimental ones obtained by Davidson et al. for n-decane [13] except for the experiment at 1525 K, where the computed results are very similar to the measurements but delayed compared to the experimental results. The methyl decanoate mechanism reproduces the two stage ignition observed during the n-decane experiments: a first, very early and rapid increase of the OH concentration (with a small overshoot visible at the lowest temperatures), then an induction period which is more or less well defined according to the temperature, and a second rapid increase of the OH concentration to a level that is very well reproduced by the kinetic mechanism.




Figure 13: Comparison of fuel reactivity at 0.03% of fuel, $\phi=1$, 2.15 atm, 99.505 % Ar. Line – methyl decanoate prediction. Open squares – n-decane experimental data [13].




Figure 14: OH profiles (0.03% of fuel, $\phi=1$, 99.505 % Ar). Lines – methyl decanoate model predictions. Open symbols – n-decane experimental data [13].

Simulations performed with the methyl decanoate mechanism were also compared to shock tube experiments for n-decane/air mixtures performed by Pfahl et al. [31] over a broader range of temperatures covering both the high and low temperature regions. Experimental conditions of these measurements were 700 – 1300 K, 12 – 50 atm, equivalence ratio of 1 (in air). These experimental conditions are particularly relevant to conditions in internal combustion engines. Figure 15 displays a comparison of experimental ignition delay times for n-decane and calculated ones from the methyl decanoate mechanism with constant volume combustion behind the reflected shock wave assumed. At the highest temperatures (more than 800 K at 12 atm and more than 1000 K at 50 atm) experimental and calculated data are in a relatively good agreement. At the lowest temperatures, below the negative temperature coefficient region, the methyl decanoate mechanism leads to ignition delay times slightly longer than experimental ones for n-decane. This trend was also observed in the case of the modeling of n-heptane. It is likely due to uncertainties in kinetic parameters used in the mechanism (reaction of decomposition of hydroperoxy compounds involved in the low temperature part of the mechanism).




Figure 15: Comparison of fuel reactivity under shock tube conditions, in air, $\phi=1$, Lines – methyl decanoate model predictions. Open symbols – n-decane experimental data [31].

The molecular structure of methyl decanoate can be visualized as very similar to n-decane; both have the same chain of 10 C atoms, populated by a lot of secondary C-H bonds. This mechanism, and prior mechanisms for n-alkanes in general and n-heptane in particular, show how the

negative temperature coefficient (NTC) region is caused by the temperature dependence of the alkylperoxy radical isomerization reaction pathways. This NTC feature has been seen here for the combustion of methyl decanoate. In comparing methyl decanoate and n-decane, however, it is evident that the presence of the methyl ester group at one end of the carbon atom chain tends to reduce the number of RO_2 isomerization reactions that can occur in this NTC region. When the straight-chain is as long as 10 C atoms, elimination of a few RO_2 and O_2QOOH isomerization pathways due to the presence of the methyl ester group does not produce a significant reduction in the rate of low temperature chain branching, but it is a reduction large enough to be observed as seen in Fig. 15. This has important implications with respect to development of surrogate fuels and mechanisms for practical hydrocarbon fuels. As already observed earlier, the n-alkane mechanism cannot reproduce the early CO_2 production of the methyl ester fuels, but these calculations indicate that the methyl ester mechanism can reproduce the heat release and ignition delay of the corresponding n-alkane fuel very well.

Comparison with engine experiments.

In the next series of calculations, predictions using the chemical kinetic model were compared to experiments performed in a motored engine. The experiments were performed by Szybist et al. using premixed charges of fuel and air in a CFR engine with adjustable compression ratio [11,12]. Fuels used in this study were n-heptane, commercial diesel fuel and methyl decanoate. Exhaust analysis was performed with a FTIR spectrometer. Quantified compounds were carbon monoxide, carbon dioxide, formaldehyde, and acetaldehyde. In the case of methyl decanoate, condensable compounds in the exhaust gas, which were trapped before the FTIR analysis, were analyzed with a GC/MS system, allowing identification of many products of the reaction but not their quantification. Pressure measurements in the cylinder were performed through a piezoelectric pressure transducer.

Simulations have been performed for both n-heptane (with the mechanism developed by Curran et al. [14]) and methyl decanoate with the “Internal Combustion Engine” model (single zone) of the software Chemkin 3. Characteristics of the engine required for the simulations are the displaced volume (612 cm^3), the engine speed (900 rpm), and the ratio of the length of the engine connecting rod to the crank radius (4.5). Intake valve closure (IVC) was 24 degrees after dead bottom center which corresponds to a starting crank angle of 214 degrees in the “Internal Combustion Engine” model. The starting pressure used for the simulations was the measured pressure at IVC (1.05 atm). Very little information about the starting temperature and the heat losses was available, making simulations difficult and requiring some assumptions. Intake charge temperature was 383.15 K in the case of n-heptane and 503.15 K in the case of methyl decanoate (boiling point: 497 K at 1 atm) whereas the wall of the cylinder was cooled with water at 373.15 K. So for the simulations with n-heptane, the starting temperature was chosen equal to 383.15 K. In the case of methyl decanoate, the intake charge temperature was much higher than the cylinder wall temperature and the charge was likely cooled during the intake. We chose 423.15 K as starting temperature so that the calculated critical compression ratio matches the experimental one. The critical compression ratio is the minimum compression ratio at which autoignition occurs. No heat losses were considered in the calculations because they were not quantified and simulations have been performed with the lowest experimental equivalence ratio (0.25) in order to have the least heat release and minimizing heat transfer from the gas to the wall of the cylinder.

The residual gases remaining in the combustion chamber after the end of a cycle influence the reactivity of the next cycle. To account for the effect of residual gases, the gases remaining at the end of the expansion stroke of the first computed cycle were used to specify the composition and the temperature of the residual gases for the next computed cycle. At the start of the next cycle, the fraction of residual gases to total charge was assumed to be the inverse of the compression ratio. In this manner, consecutive cycles were performed until the steady state was reached [32].




Figure 16: Gases in the residual part of the cylinder were taken in account by considering consecutive cycles.

Motored Engine simulations with n-heptane. We first compared model predictions to the experiments for n-heptane because the n-heptane mechanism is more mature and well-validated than the methyl decanoate mechanism. If the comparison is successful, it can give confidence that the modeling approach used is valid. Simulations were performed by varying the compression ratio over the range of experimental investigation (4.5 to 9.5). The fresh charge was composed of n-heptane in air with an equivalence ratio of 0.25. Table 8 displays the calculated mass fractions (% fuel C) of CO, CO₂, CH₂O and CH₃CHO at the end of the expansion stroke for six consecutive cycles at a compression ratio of 6, where the 1st cycle starts with 100% fresh charge. The evolution in the mass fractions from cycle 1 to cycle 6 shows that the remaining gases in the cylinder have an influence on the kinetics of the reaction, for this compression ratio where ignition does not occur.

Table 8: Calculated mass fractions (% of fuel C) for six consecutive cycles at CR=6.

Cycle	CO	CO ₂	CH ₂ O	CH ₃ CHO
1	19.53	1.54	7.54	6.86
2	17.55	1.22	6.17	5.18
3	15.61	1.04	6.01	5.05
4	15.58	1.03	6.07	5.10
5	15.57	1.02	6.07	5.10
6	15.57	1.02	6.07	5.10

Figure 17 shows the comparison between the computed mass fractions at the end of the expansion stroke to those measured in the exhaust of the motored engine fueled by n-heptane. The agreement between calculated and experimental data [12] is globally satisfactory. The calculated ignition occurs at about the same compression ratio as in the experiments (slightly below CR=8). This shows that the overall reactivity of the n-heptane model is about right. The simulated ignition is rather sharp whereas the experimental one is much less abrupt. This can be explained by the fact that the “Internal Combustion Engine” model used for the simulation is a single zone model (concentrations are assumed to be homogeneous inside the cylinder) which is not the case in the real engine used for the experiments. CO and CO₂ concentration profiles are rather well reproduced by the model, with very low levels of CO₂ prior to ignition. Before the ignition occurs the model under-predicts the concentration of acetaldehyde by a factor of 1.3 and over-predicts that of formaldehyde by a factor of 2. These levels of precision are quite good for engine data analyses and illustrate the challenges of comparing single-zone simulations with real engine experiments.




Figure 17 : Mass fractions (% of fuel C) of CO (\diamond), CO₂ (\triangle), CH₂O (\circ) and CH₃CHO (\square) at an equivalence ratio of 0.25 in the case of n-heptane simulations in an engine. Open symbols correspond to experiments [12] and lines to simulations.

Motored engine simulations with methyl decanoate. Based on the positive results for n-heptane, simulations were performed with the methyl decanoate mechanism and compared to experimental measurements over the range of compression ratios 4.4 to 5.6. The fresh charge was composed of methyl decanoate in air with an equivalence ratio of 0.25. These calculations were very time consuming compared to the simulations with n-heptane because of the size of the mechanism (about 3 hours CPU time for one cycle on a 4 GHz Intel Pentium™ 4 processor PC, compared with 10 CPU minutes for n-heptane). With the added need to compute at least 6 cycles to achieve steady state, these calculations become very ambitious and resource intensive. One set of comparison curves represents about over 50 computed cycles and 160 hours of CPU time. The comparison between the computed and experimental results is shown in Figure 18. The starting temperature has been adjusted so that the computed critical compression ratio is very close to the experimental one. The computed critical compression ratio is sensitive to the charge temperature assumed at intake valve closing (423 K). This temperature is a reasonable value lying between the limiting temperatures of the wall (373 K) and the intake (503 K). If the chemical kinetic model was too reactive or too unreactive, the needed temperature at intake valve closing would have been outside these experimental temperature limits. The agreement

between computed and experimental mass fractions is again globally satisfactory. With methyl decanoate as a fuel, the model predicts a concentration of formaldehyde higher than that of acetaldehyde, in agreement with the experiments. The concentration of carbon dioxide is well reproduced and is much higher than in the n-heptane case, as shown both in the experimental measurements and in the model predictions for methyl decanoate. This is due to the additional formation paths for carbon dioxide from the ester chemistry present in methyl decanoate combustion. On the other hand, the concentration of carbon monoxide is over predicted. As the compression ratio increases, mass fractions of CO and CO₂ go up whereas mass fractions of acetaldehyde and formaldehyde go down. These trends and the high concentration of CO are due to overpredictions of the temperature. For example, at CR=4.8 the temperature at the top dead center reaches 889 K, which is likely too high because heat losses were not taken into account in these simulations. This elevated temperature is confirmed by an elevated pressure at the end of compression (the calculated pressure is about 1 bar higher than the experimental value). However, if heat losses are included, it is likely that the computed critical compression ratio of methyl decanoate would be too low. At this point, we thought we had gained about as much information as we could derive from these comparisons, given all the uncertainties in modeling these experiments (e.g. nonhomogeneities in the chamber) and the large computing and manpower resources required. Overall, the most significant result of these simulations is the kinetic explanation of the elevated CO₂ levels at low compression ratios that were observed in the experiments for methyl decanoate, which translates into early CO₂ production in actual engine cycles, a feature which makes biodiesel fuels different from conventional diesel fuels.




Figure 18: Mass fractions of CO (◇), CO₂ (△), CH₂O (○) and CH₃CHO (□) at an equivalence ratio of 0.25. Open symbols correspond to experiments [12] and lines to simulations.

Comparison with condensable compounds in the exhaust. Szybist et al. performed a qualitative analysis of the exhaust condensates from the engine by GC/MS. The species which were identified during this study were mainly methyl esters, methyl esters with a ketone group, and carboxylic acids. Other species like ketones and aldehydes were also observed.

In the range of temperatures corresponding to the region where compression ratios do not lead to ignition (below 900 K), the methyl decanoate mechanism predicts the formation of numerous methyl esters with one double bond (either at the extremity of the alkyl chain or conjugated with the double C=O bond of the ester group) as well as 1-alkenes. 1-alkenes and methyl esters with the double bond at the terminal position mainly come from the decomposition by C-C bond β-

scission of alkyl and alkyl-ester radicals. Methyl esters with the double bond at the conjugated position are obtained from the decomposition by C-H bond β -scission of 3-alkyl-ester radicals (Figure 19).




Figure 19 : decomposition by C-H bond β -scission of a 3-alkyl-ester radical.

The model also predicts the formation of aldehydes and methyl esters with a carbonyl group at the extremity of the alkyl chain. These species are obtained by decomposition by C-C bond β -scission of alkoxy and alkoxy-ester radicals. These last radicals are formed by addition of OH radicals to the double bond of alkenes and unsaturated esters followed by internal isomerizations involving the H-atom of the hydroxyl group. Organic acids are generated from the recombination of H-atoms and carboxylate radicals. Carboxylate radicals are derived from decomposition of some hydroperoxides (Figure 20).




Figure 20 : Pathway for the formation of carboxylate radicals.

The model also predicts the formation of hetero-cyclic species like cyclic ethers, but not 2-(3H)-furanone-5-ethylidihydro and 5-methoxycarbonylpentan-4-olide which were identified by GC/MS.

4. Conclusions

In this study, a detailed chemical kinetic oxidation mechanism for methyl decanoate, a surrogate for biodiesel fuel, has been developed by following the rules previously used for modeling n-heptane oxidation. Experimental data for the validation were very scarce and some of them have been performed in non ideal conditions that are difficult to model. The model was compared to rapeseed oil methyl ester experiments in a jet stirred reactor. This model (and the blend surrogate model including n-heptane chemistry) reproduced the overall reactivity as well as the concentrations of the products of the reaction. An important feature of this mechanism is its ability to predict the early formation of carbon monoxide and carbon dioxide. The kinetic

analysis of the model showed that early formation of these two species is linked to the presence of the ester group in the methyl decanoate molecule.

The methyl decanoate model was also compared with n-decane experimental results from shock tube experiments. Calculated ignition delay times and OH profiles were in very good agreement with n-decane experiments, showing that the reactivity of large methyl esters is very similar to the reactivity of n-alkanes of similar size. This mechanism was compared with methyl decanoate oxidation experiments in a motored engine. Although these experiments were not well characterized and the numerical model used for the computations was too simplified, the agreement between calculated and experimental concentrations was qualitatively satisfactory.

Further refinements are possible to improve capabilities for biodiesel simulations. Real biodiesel fuels are mixtures of several esters, some of them having one, two or three double bonds in their alkyl chains, as shown above in Fig. 1. We intend to develop additional submodels, based on methyl decanoate, to highlight the influences of the presence of double bonds in the alkyl chain and provide very realistic surrogate fuel mechanisms for real biodiesel fuels from various origins and having different compositions.

Overall, it appears that both n-hexadecane and methyl decanoate are acceptable surrogates for biodiesel fuel, based on the comparisons with the work of Dagaut et al. [9] and the present work. One strength of the present mechanism is its ability to reproduce the effects of the methyl ester group in all of the major components of soybean and rapeseed methyl esters. The n-hexadecane mechanism of Dagaut et al. predicts concentrations of large olefin species that are larger than those measured in rapeseed methyl ester fuel combustion, but the present methyl decanoate mechanism does not include olefins larger than C_{10} and cannot predict concentrations of any species larger than that. The present methyl decanoate mechanism is also unique by including low temperature reaction pathways that enable it to address such important practical problems as diesel ignition and sooting, as well as combustion in HCCI engines, all of which require a kinetic description of the low temperature kinetics of the fuel. Since the boiling point of methyl decanoate is 497 K, future experimental data in heated shocktubes and rapid compression machines may be acquired and used to further test this chemical kinetic mechanism. This is not the case for even higher molecular weight methyl esters whose high boiling points make acquisition of experimental data in these devices very difficult and limit the validation of their associated chemical kinetic mechanisms. Thus, methyl decanoate is a convenient test fuel for detailed chemical kinetic mechanisms of biodiesel surrogates. Ultimately, a full, detailed, high and low temperature mechanism for the C_{16} and C_{18} saturated and unsaturated species that are in real biodiesel fuels will be required. One additional accomplishment of the present mechanism development for methyl decanoate is a demonstration that such a biodiesel kinetic mechanism is already feasible and accessible by extending this mechanism to include a longer alkane chain.

Acknowledgment

The authors thank Dr. James Szybist and Prof. Andre Boehman for sending us experimental pressure histories and for valuable discussions about their experimental work. This work was supported by the U.S. Department of Energy, Office of Freedom CAR and Vehicle Technologies, and the authors thank program managers Kevin Stork and Gurpreet Singh for their support of this work. This work was also performed under the auspices of the U.S. Department

of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

References

1. C.K. Westbrook, W.J. Pitz, H.J. Curran, *J. Phys. Chem. A* (110) (2006) 6912-6922.
2. R.L. McCormick, J.D. Ross, M.S. Graboski, *Environ. Sci. Technol.* 21 (4) (1997) 1144-1150.
3. J. Van Gerpen, B. Shanks, R. Pruszko, D. Clements, G. Knothe, Biodiesel Production Technology, National Renewable Energy Laboratory subcontractor report NREL/SR-510-36244, 2004.
4. E.M. Fisher, W.J. Pitz, H.J. Curran, C.K. Westbrook, *Proc. Combust. Inst.* 28 (2000) 1579-1586.
5. W.K. Metcalfe, S. Dooley, H.J. Curran, J.M. Simmie, A.M. El-Nahas, M.V. Navarro, *J. Phys. Chem.* 111 (19) (2007) 4001-4014.
6. S. Gaïl, M.J. Thomson, S.M. Sarathy, S.A. Syed, P. Dagaut, P. Diévar, A.J. Marchese, F.L. Dryer, *Proc. Combust. Inst.* 31 (1) (2007) 305-311.
7. S.M. Sarathy, S. Gaïl, S.A. Syed, M.J. Thomson, P. Dagaut, *Proc. Combust. Inst.* 31 (1) (2007) 1015-1022.
8. T. Vaughn, M. Hammill, M. Harris, A.J. Marchese, Ignition Delay of Bio-Ester Fuel Droplets, SAE Technical Paper Series, 2006-01-3302.
9. P. Dagaut, S. Gaïl, M. Sahasrabudhe, *Proc. Combust. Inst.* 31 (2) (2007) 2955-2961.
10. A. Ristori, P. Dagaut, M. Cathonnet, *Combust. Flame* 125 (3) (2001) 1128-1137.
11. J.P. Szybist, J. Song, M. Alam and A.L. Boehman, *Fuel Process. Technol.* 88 (7) (2007) 679-691.
12. J.P. Szybist, A.L. Boehman, D.C. Haworth, and H. Koga, *Combust. Flame* 149 (1-2) (2007) 112-128.
13. D.F. Davidson, J.T. Herbon, D.C. Horning DC, R.K. Hanson, *Int. J. Chem. Kinet.* 33 (12) (2001) 775-783.
14. H.J. Curran, P. Gaffuri, W.J. Pitz, C.K. Westbrook, *Combust. Flame* 129 (2002) 253-280.
15. H.J. Curran, P. Gaffuri, W.J. Pitz, C.K. Westbrook, *Combust. Flame* 114 (1998) 149-177.
16. H.J. Curran, *Int. J. Chem. Kinet.* 38 (4) (2006) 250-275.
17. D.J. Henry, M.L. Coote, R. Gomez-Balderas, L. Radom, *J. Am. Chem. Soc.* 126 (6) (2004) 1732-1740.
18. D.M. Matheu, W.H. Green, J.M. Grenda, *Int. J. Chem. Kinet.* 35 (3) (2003) 95-119.
19. V. Warth, F. Battin-Leclerc, R. Fournet, P.A. Glaude, G.M. Come, G. Scacchi, *Comput. Chem.* 24 (5) (2000) 541-560.
20. K.D. King, *Int. J. Chem. Kinet.* 11 (1979), 1071-1080.
21. R.S. Zhu, J. Park, M.C. Lin, *Chem. Phys. Lett.* 408 (2005) 25-30.
22. H. Sun, Joseph W. Bozzelli, *J. Phys. Chem. A* 108 (10) (2004) 1694-1711.
23. C.Y. Sheng, J.W. Bozzelli, A.M. Dean, A.Y. Chang, *J. Phys. Chem. A* 106 (2002) 7276-7293.
24. J.D. DeSain, S.J. Klippenstein, J.A. Miller, C.A. Taatjes, *J. Phys. Chem. A* 107 (2003) 4415-4427.
25. E.J. Silke, W.J. Pitz, C.K. Westbrook, M. Ribaucour, *J. Phys. Chem. A* 111 (19) (2007) 3761-3775.
26. K.A. Sahetchian, R. Rigny, J. Tardieu De Maleyssie, L. Batt, M. Anwar Khan, S. Mathews, *Proc. Combust. Inst.* 24 (1992) 637-643.
27. E.R. Ritter, J. W. Bozzelli, *Int. J. Chem. Kinet.* 23 (1991) 767-778.
28. S.W. Benson, *Thermochemical Kinetics*, John Wiley and Sons Inc., New York, 1976.
29. A.M. El-Nahas, M.V. Navarro, J.M. Simmie, J.W. Bozzelli, H.J. Curran, S. Dooley, W. Metcalfe, *J. Phys. Chem. A* 111 (19) (2007) 3727-3739.
30. P.A. Glaude, W.J. Pitz, M.J. Thomson, *Proc. Combust. Inst.* 30 (2005) 1111-1118.
31. U. Pfahl, K. Fieweger, G. Adomeit, *Proc. Combust. Inst.* (1996) 781-789.
32. H.J. Curran, P. Gaffuri, W.J. Pitz, C.K. Westbrook, W.R. Leppard, Autoignition Chemistry of the Hexane Isomers: An Experimental and Kinetic Modeling Study, SAE Technical Paper Series 952406, 1995.