

Presented at the Particle Accelerator  
Conference, San Francisco, CA,  
March 5-7, 1973

LBL-1710  
e. d.

HIGH-INTENSITY EFFECTS IN THE LONGITUDINAL  
MOTION OF STORED PARTICLE BEAMS

Andrew M. Sessler

February 28, 1973

RECEIVED  
LIBRARY  
EXTENSION DIVISION

MAR 1973

UNIVERSITY OF CALIFORNIA  
DOCUMENTS SECTION

Prepared for the U.S. Atomic Energy  
Commission under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545



LBL-1710  
e. d.

Andrew M. Sessler

Lawrence Berkeley Laboratory  
University of California  
Berkeley, CA 94720

Summary. A brief review is given of the various self-field phenomena associated with the longitudinal motion of particles in storage rings.

### Introduction

Although there are some high-intensity phenomena for which the coupling of longitudinal and transverse motion is essential, such as, for example, the head-tail effect; the great majority of high-intensity phenomena primarily involve either longitudinal or transverse degrees of freedom. In this review, we restrict our attention to phenomena which are essentially longitudinal in nature.

It is convenient to consider separately the behavior of unbunched (coasting) and bunched (external RF system in operation) beams. Detailed experimental information on coasting beams has been obtained on the ISR, on the (old) CERN electron model CESAR, and on electron ring accelerators. All high-energy electron storage rings have bunched beams and, of course, so do synchrotrons, so that there are a large number of sources of experimental information about the longitudinal motion of bunched beams.

### Unbunched Beams

The primary phenomenon, due to self-field effects, in the longitudinal motion of a coasting beam is spontaneous formation of longitudinal density variations. For a storage ring operating above the transition energy (so that  $df/dE$  is negative) the effect is physically very simple (the "negative mass instability"), and both growth rates in the linear regime and thresholds were theoretically derived<sup>1</sup> prior to the phenomenon being observed on a variety of machines.<sup>2</sup>

Although the original negative mass theory considered bunching arising from self-forces associated with a beam in a smooth, perfectly conducting, and essentially straight vacuum chamber, it was soon realized that the beam-surroundings played an essential role in

the instability. In particular, unexcited RF cavities,<sup>3</sup> resistive walls,<sup>4</sup> and ceramic chambers<sup>5</sup> were shown to be able to greatly enhance the phenomenon (even to cause it below transition where the mass is positive), and hence the beam environment became of considerable concern to builders of storage rings.

It is convenient to introduce a longitudinal coupling impedance  $Z_n$  defined by<sup>6</sup>

$$Z_n = - \frac{2\pi R E_n}{I_n}, \quad (1)$$

where  $E_n$  is the  $n$ th harmonic of the azimuthal electric field at the beam,  $I_n$  is the  $n$ th harmonic of the beam current and  $R$  is the orbit radius. An approximate criterion for stability of a coasting beam against self-bunching at the frequency  $n\omega_{rev}$  is<sup>1,4,7</sup>

$$\frac{|Z_n|}{n} \leq \frac{\eta \gamma U_0}{I_0} \left( \frac{\Delta E}{E} \right)^2, \quad (2)$$

where  $\omega_{rev} = \beta c/R$  is the particle revolution frequency,

$$\eta = \left| \frac{1}{\gamma_t^2} - \frac{1}{\gamma^2} \right|,$$

with  $\gamma_t$  the transition energy in units of  $m_0 c^2$  and  $\gamma$  the particle energy in units of  $m_0 c^2$  so that  $E = \gamma m_0 c^2$ ,  $U_0 = m_0 c^2/e = 0.511$  MV for electrons and 938 MV for protons,  $I_0 = Ne f_{rev}$  is the circulating beam current, and  $\Delta E/E$  is the full width of the beam energy distribution at half maximum. This stability criterion arises from the balance between the instability-driving forces characterized by  $Z_n$  and the Landau damping associated with energy (and hence frequency) spread.

At the present time a considerable body of knowledge exists concerning the coupling impedance  $Z_n$ . Theoretical calculations have been performed by a large number of workers for a variety of structures (see references cited in Ref. 6.), and methods have even been devised for measuring  $Z_n$ .<sup>8</sup>

\* Work supported by the U. S. Atomic Energy Commission.

Recently, there have been detailed experimental studies of the onset of azimuthal instabilities.<sup>9,10</sup> The work on electron ring accelerators so-far provides only qualitative accord with the criterion (2), but the observations on the ISR are in remarkably good quantitative agreement with (2). Spontaneous beam bunching due to interaction of a coasting beam with an unexcited RF cavity has been observed on the AGS, and avoided either by increasing the beam energy spread or by de-tuning; i.e., reducing the cavity contribution to  $Z_n$ .<sup>11</sup>

The experimental observations are particularly interesting because the phenomenon of beam bunching can be observed in the nonlinear regime where the theory, presently, is far from complete.<sup>12</sup> Generally, it is observed<sup>2</sup> that the instability is self-stabilizing; i.e., a certain degree of bunching occurs, and, subsequently the beam again becomes longitudinally uniform (but with a larger energy spread than it had initially). Because this process leads to beam widening it is generally bad as it results in particle loss if there are aperture stops, and in the case of the electron ring accelerator to degraded ring quality.

Finally, it should be noted that recently observations have been made on electron rings of RF signals without noticeable beam degradation,<sup>9</sup> which, perhaps, is related to a prior theoretical calculation on mini-instabilities.<sup>13</sup>

#### Bunched Beams

In a stored bunched beam there are two classes of high intensity effects; namely coherent bunch motion, and alterations in bunch size and shape.

#### Coherent Bunch Motion

The simplest coherent mode of a single bunch is the dipole mode ( $m = 1$ ); i.e. rigid-bunch motion. Most of the analysis in the literature is confined to this mode, although some authors have considered higher order modes ( $m > 1$ ).<sup>14,15</sup> For the simple case in which the phase of the bunch center  $\phi_1$  for bunch 1, may be treated in linear approximation and in which there is no external feedback we have<sup>11,16,17,18</sup>

$$\ddot{\phi}_1 + \omega_s^2 \phi_1 = \sum_{j=1}^B A_{1j} \phi_j, \quad (3)$$

where  $B$  is the number of bunches,  $\omega_s$  is the frequency of small phase oscillations, and the coefficients  $A_{ij}$  are given by

$$A_{ij} = - \frac{\omega_s^2}{V \cos \phi_s} \frac{\partial V_{ij}}{\partial \phi_j}. \quad (4)$$

In Formula (4),  $V$  is the voltage gain per turn,  $\phi_s$  is the stable phase angle, and  $V_{ij}$  is the voltage on bunch  $i$  caused by bunch  $j$ . Clearly  $V_{ij}$  can be expressed in terms of a coupling impedance.<sup>17</sup>

If all the bunches are equal then the  $A_{ij}$  are only functions of  $i-j$  and the  $B$  normal modes are simply the  $B$ -roots of unity:<sup>11,17,19</sup>

$$\begin{aligned} \phi_1 &= e^{i\omega_n^{(1)}t} \\ \phi_j &= \phi_1 e^{in(j-1)2\pi/B}, \end{aligned} \quad (5)$$

where  $n = 1, \dots, B$  characterizes the mode and

$$\omega_n^{(1)} \approx \omega_s - \frac{1}{2\omega_s} \sum_{j=1}^B A_{j1} e^{in(j-1)2\pi/B}. \quad (6)$$

Clearly if  $\text{Im } \omega_n^{(1)} < 0$  the  $n$ th mode is unstable. If the nonlinear nature of the synchrotron motion is included, then a dispersion analysis shows<sup>14,15,18</sup> that the  $n$ th mode will be stable, even if  $\text{Im } \omega_n^{(1)} < 0$ , provided

$$|\omega_n^{(1)} - \omega_s| < S/4 \quad (7)$$

where  $S$  is the full-spread in the synchrotron oscillation frequency of particles in a bunch.

The analysis for unequal bunches has resulted in a condition for decoupling of the bunches; namely that the spread in individual bunch frequencies must be greater than the shift in frequency due to coupling. Stability requires, in addition, that a condition analogous to (7) be satisfied;<sup>15</sup> namely that for the  $m$ th order mode, having frequency  $\omega^{(m)}$ ,

$$|\omega^{(m)} - m\omega_s| < \frac{\sqrt{m}}{4} S. \quad (8)$$

Also, in the literature, are numerical studies of unequal bunches and analysis of the influence of beam control systems and active feed-back damping.<sup>17</sup>

Observations have been made on the CERN PS,<sup>17</sup> the AGS,<sup>11</sup> and the CEA,<sup>20</sup> as well as on the storage rings in Novosibirsk and Frascati. In general there is

good agreement between the observations and theory, although in some cases the actual values of the  $A_{ij}$  have been larger than was a priori expected.

#### Bunch Size and Shape

It was first observed, at Orsay and Frascati, that the length of bunches in the storage rings was a function of the stored current. No such effect was observed at Stanford or Novosibirsk.<sup>21</sup> Recent observations at CEA,<sup>20</sup> SLAC,<sup>22</sup> and Frascati<sup>19</sup> have indicated that both bunch length and width increases with increasing stored current. The parametric dependence of bunch length,  $\Delta$ , on beam current,  $I$ , beam energy,  $E$ , and RF voltage,  $V$ , is (approximately) for bunch length large compared to the natural (low-current) bunch length,  $\Delta_0$ , (as has been summarized in Ref. 21):

$$\Delta(\text{ns}) = \frac{0.46 I(\text{mA})^{1/3}}{E(\text{GeV})^{7/6}} \left[ \frac{30}{V(\text{keV})} \right]^{1/6} \Delta_0(\text{ns})^{2/3}. \quad (9)$$

Prior to the observation of bunch widening, a general equilibrium theory of bunch length was developed.<sup>23</sup> This theory included the effect of coherent synchrotron radiation as well as the electrical interaction of a beam with various resonant structures that might be present in a storage ring. The parametric dependence of bunch length was found to be in good agreement with (9), and numerical estimates--based upon reasonable guesses for the characteristics of resonant structures--were in substantial agreement with (9). This theory, however, predicted no bunch widening which isn't necessarily a defect of the theory since the two phenomena may be unrelated.

An alternative, and older, theory suggested that the instability of internal coherent synchrotron oscillations could explain bunch lengthening.<sup>24</sup> However, the parametric dependence deduced in this work is not in good accord with observations, although the theory does predict both bunch lengthening and widening.

In some work which is still in progress, the present writer has, in the spirit of Ref. 24, developed a theory of the behavior of a beam whose natural energy width--resulting from the balance between quantum fluctuations and classical radiation damping--is less than that required for the stability of collective modes. It is shown that in this circumstance there is a turbulent equilibrium state in which the wave diffusion (due to stable collective

modes with finite amplitudes) augments the quantum diffusion so as to yield a bunch of increased width and length. The theory, perhaps when combined with the equilibrium theory of Ref. 23, appears capable of explaining the observations, but it is too early to be sure that the proposed explanation is indeed correct.

#### References

1. C. E. Nielsen, A. M. Sessler, and K. R. Symon, International Conference on High-Energy Accelerators, CERN, 1959, p. 239; A. A. Kolomenskij and A. N. Lebedev, *ibid*, p. 115.
2. M. Q. Barton and C. E. Nielsen, International Conference on High-Energy Accelerators, Brookhaven National Laboratory, 1961, p. 163; H. Bruck, G. Gendreau, M. Gouttefangeas, J. Hamelin, R. Levy-Mandel, and R. Vienet, *ibid*, p. 175; D. Keefe, Proceedings of the 8th International Conference on High-Energy Accelerators, CERN, 1971, p. 397.
3. V. K. Neil and A. M. Sessler, *Rev. Sci. Instr.* 32, 256 (1961).
4. V. K. Neil and A. M. Sessler, *Rev. Sci. Instr.* 36, 429 (1965).
5. R. J. Briggs and V. K. Neil, *Plasma Physics* 8, 255 (1966).
6. A. M. Sessler, Proceedings of the 1971 Particle Accelerator Conference, IEEE Trans. Nucl. Sci. NS-18, #3, June 1971, p. 1039.
7. A. G. Ruggiero and V. G. Vaccaro, Solution of the Dispersion Relation for Longitudinal Stability of an Intense Coasting Beam in a Circular Accelerator, CERN Report ISR-TH/68-33, 1968.
8. A. Faltens, E. C. Hartwig, D. M6hl, and A. M. Sessler, Proceedings of the 8th International Conference on High-Energy Accelerators, CERN, 1971, p. 338; K. Hübner, E. Keil, and B. Zotter, *ibid*, p. 295.
9. G. R. Lambertson (Lawrence Berkeley Laboratory), private communication.
10. B. Zotter and P. Bramham, Proceedings of this Conference (Contribution K2).
11. M. Barton and E. C. Raka, Proceedings of the 1971 Particle Accelerator Conference, IEEE Trans. Nucl. Sci. NS-18 #3, June 1971, p. 1032.
12. H. G. Hereward, Longitudinal Space Charge Effects in Circular Accelerators, Lawrence Berkeley Laboratory Report BEV-524 (1960), (unpublished); E. A. Perelshtein, Proceedings of the 5th

- International Conference on High-Energy Accelerators, Frascati, 1965, p. 362; C. Pellegrini and A. M. Sessler, Theory of the Non-Linear Negative Mass Instability, Lawrence Berkeley Laboratory Report ERAN-203 (1973), (unpublished).
13. H. G. Hereward, A Mini-Instability, CERN Report MPS/Int. DL68-1 (1968), (unpublished).
  14. A. N. Lebedev, Proceedings of the 6th International Conference on High Energy Accelerators, Cambridge Electron Accelerator Report CEAL-2000, 1967, p. 284; I. Gumowski, Proceedings of the 8th International Conference on High-Energy Accelerators, CERN, 1971, p. 360.
  15. F. J. Sacherer, Proceedings of this Conference (Contribution K1).
  16. K. W. Robinson, Stability of Beam in Radio-frequency System, Cambridge Electron Accelerator Report CEAL-1010 (1964), (unpublished).
  17. D. Boussard and J. Gareyte, Proceedings of the 8th International Conference on High Energy Accelerators, CERN, 1971, p. 317.
  18. V. L. Auslander, M. M. Karliner, A. A. Naumov, S. G. Popov, A. N. Skrinsky, and I. A. Shekhtman, Proceedings of the 5th International Conference on High-Energy Accelerators, Frascati, 1965, p. 339.
  19. C. Pellegrini (Laboratori Nazionali di Frascati), private communication.
  20. R. Averill, A. Hofmann, R. Little, H. Mieras, J. Paterson, K. Strauch, G-A. Voss, and H. Winick, Proceedings of the 8th International Conference on High-Energy Accelerators, CERN, 1971, p. 301.
  21. F. Amman, Proceedings of the 1969 Particle Accelerator Conference, IEEE Trans. Nucl. Sci., NS-16 #3, 1073 (1969).
  22. M. Allen (Stanford Linear Accelerator Center), private communication.
  23. C. Pellegrini and A. M. Sessler, Il Nuovo Cimento 3A, 116 (1971).
  24. A. N. Lebedev, "On the Bunch-Lengthening Effect in Storage Rings," in Physics With Intersecting Storage Rings, edited by B. Touschek (Academic Press, N. Y., 1971), p. 184.