



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Physics of reshock and mixing in single-mode Richtmyer-Meshkov instability

O. Schilling, M. Latini, W. Don

December 15, 2006

Physical Review E

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Physics of reshock and mixing in single-mode Richtmyer-Meshkov instability

Oleg Schilling*

University of California, Lawrence Livermore National Laboratory, Livermore, California 94551

Marco Latini[†]

Applied and Computational Mathematics, California Institute of Technology, Pasadena, California 91125

Wai Sun Don[‡]

Division of Applied Mathematics, Brown University, Providence, Rhode Island 02912

(Dated: December 12, 2006)

The ninth-order weighted essentially non-oscillatory (WENO) shock-capturing method is used to investigate the physics of reshock and mixing in two-dimensional single-mode Richtmyer-Meshkov instability to late times. The initial conditions and computational domain were adapted from the Mach 1.21 air(acetone)/SF₆ shock tube experiment of Collins and Jacobs [J. Fluid Mech. **464**, 113 (2002)]: the growth of the bubble and spike perturbation amplitudes from fifth- and ninth-order WENO simulations of this experiment were compared to the predictions of amplitude growth models, and were shown to be in very good agreement with the experimental data prior to reshock [Latini, Schilling and Don, Phys. Fluids (2007), in press]. In the present investigation, the density, vorticity, baroclinic vorticity production, and simulated density Schlieren fields are first presented to qualitatively describe reshock. The baroclinic circulation deposition on the interface is shown to agree with the predictions of the Samtaney and Zabusky [J. Fluid Mech. **269**, 45 (1994)] model and linear instability theory. The time-evolution of the positive and negative circulation on the interface is considered before and after reshock: it is shown that the circulations are equal before, as well as after reshock, until the interaction of the reflected rarefaction with the layer leads to flow symmetry breaking and different evolutions of the positive and negative circulation. The post-reshock mixing layer growth is shown to be in very good agreement with three models predicting linear growth for a short time following reshock. Next, a comprehensive investigation of local and global mixing properties as a function of time is performed. The distribution and amount of mixed fluid along the shock propagation direction is characterized using averaged mole fraction profiles, a fast kinetic reaction model, and molecular mixing fractions. The modal distribution of energy in the mixing layer is quantified using the spectra of the fluctuating kinetic energy, fluctuating enstrophy, pressure variance, density variance, and baroclinic vorticity production variance. It is shown that a broad range of scales already exists prior to reshock, indicating that the single-mode Richtmyer-Meshkov instability develops non-trivial spectral content from its inception. At reshock, fluctuations in all fields (except for the density) are amplified across all scales. Reshock strongly amplifies the circulation, profiles and mixing fractions, as well as the energy spectra and statistics, leading to enhanced mixing, followed by a decay. The mole and mixing fraction profiles become nearly self-similar at late times following reshock; the mixing fraction approaches unity across the layer at the latest time, signifying nearly complete mixing. The comparison of the spectra to the predictions of classical inertial subrange scalings in two-dimensional turbulence shows that the post-reshock spectra are consistent with most of these scalings over short wave number ranges. To directly quantify the amplification of fluctuations by reshock, the previously considered quantities are compared immediately after and before reshock. Finally, to investigate the decay of fluctuations in the absence of additional waves interacting with the mixing layer following reshock, the boundary condition at the end of the computational domain is changed from reflecting to outflow to allow the reflected rarefaction wave to exit the domain. It is shown that the reflected rarefaction has an important role in breaking symmetry and achieving late-time statistical isotropy of the velocity field.

PACS numbers: 47.20.-k, 47.20.Lz, 47.20.Ma, 47.20.Cq, 47.51.+a, 47.40.Nm, 47.40.-x, 47.35.Lf, 47.11.Bc, 47.11.-j

I. INTRODUCTION

The Richtmyer-Meshkov instability is a fundamental fluid instability that occurs when perturbations on an interface separating different gases grow following the passage of a shock. In the weighted essentially non-oscillatory (WENO) simulations of a model of the Mach 1.21 air(acetone)/SF₆ experiments of Collins and Jacobs

*Author to whom correspondence should be addressed: schilling1@llnl.gov.

[†]Electronic address: mlatini@acm.caltech.edu

[‡]Electronic address: wson@cfm.brown.edu

[1] considered here (see Ref. 2 for details on the experiment and simulations), following the initial interaction at 0 ms, a transmitted shock enters the second gas. This shock reflects from the end wall of the shock tube test section and interacts with the evolving mixing layer at ≈ 6.4 ms in a process called *reshock*. Following reshock, a transmitted shock continues into the first gas and a reflected rarefaction wave returns into the second gas. The rarefaction wave reflects from the end wall and interacts again with the evolving interface at ≈ 11 ms, generating a compression wave. The compression wave reflects from the end wall and again interacts with the evolving interface at ≈ 15 ms. The growth and other properties of the mixing layer are affected by these wave-interface interactions, with each subsequent interaction further increasing turbulent fluctuations contributing to the mixing and formation of complex small-scale structures.

The dynamics and physics of the reshocked Richtmyer-Meshkov instability is under intensive investigation due to its relevance to both core-collapse supernovae dynamics [3] and to inertial confinement fusion (ICF) [4–6]. Very little previous investigation of reshocked Richtmyer-Meshkov instability-induced mixing has been performed in either two or three dimensions [7, 8]. The vast majority of simulations of capsule implosions in ICF involving the Rayleigh-Taylor and Richtmyer-Meshkov instabilities are still performed in one and two dimensions, as the resolutions and computational resources required to adequately resolve such flows in three dimensions are still beyond the reach of available computing facilities. Thus, ICF designers continue to rely on lower dimensional simulations, and it is therefore, valuable to better understand the dynamics of these instabilities in two dimensions.

Here, data from ninth-order WENO simulations using a grid resolution of 512 points per initial perturbation wavelength λ (the WENO9F simulation in Ref. 9) is used to investigate the physics of reshock and mixing. In particular, the effects of reshock, as well as the decay properties at late times, are quantified and assessed. The analysis considered here includes local and global quantities. Two types of local quantities are considered at different times—profiles across the mixing layer and energy spectra. In the present geometry, mixing profiles are quantities averaged across the spanwise (y) direction perpendicular to the shock propagation direction (so that they are only a function of the streamwise coordinate x and time) characterizing the distribution of mixing within the layer. Spectra of fluctuations within the mixing layer as a function of the wave number k characterize the modal energy distribution within the layer. Global quantities include mixing fractions and statistics as a function of time. Mixing fractions characterize the efficiency of mixing, including the effects of reshock and wave-interface interactions, and statistics characterize the energy evolution as the mixing progresses. To investigate the effects of the reflected rarefaction wave on the mixing, simulations using outflow and reflecting boundary conditions (the latter corresponding to the WENO9M simulation in

Ref. 9) at the end wall with resolutions of 256 points per initial perturbation wavelength were compared.

This paper is organized as follows. The equations solved and the suitability of the WENO method for investigating the physics of reshock and mixing are briefly discussed in Sec. II. The density, vorticity, baroclinic vorticity production, and simulated density Schlieren fields are used to visualize and qualitatively describe the reshock process in Sec. III. The circulation deposited on the interface by the shock is compared with the predictions of the Samtaney-Zabusky model and linear instability theory in Sec. IV. The post-reshock mixing layer width is compared to the predictions of three phenomenological reshock models in Sec. V. The analysis of the local and global properties of mixing, including profiles, mixing fractions, energy spectra, and statistics, is presented in Sec. VI. Also presented are a comparison of mixing quantities and spectra immediately before and after reshock, and a comparison of results from reflecting and outflow boundary conditions at the end wall. A summary of the results and conclusions are given in Sec. VII.

II. THE WENO METHOD FOR THE SIMULATION OF RESHOCK

The reshocked Richtmyer-Meshkov instability must be simulated with a shock-capturing method. The fundamental issue with a direct numerical simulation (DNS) to simulate this flow is that high-order (e.g., spectral or high-order finite-difference) schemes used for DNS are incapable of simulating shocked flows such as those arising from the Richtmyer-Meshkov instability due to Gibbs oscillations. In addition, it is not possible to fully resolve shocks in a continuum simulation in the sense required to consider it a DNS. Also, in simulations of the reshocked Richtmyer-Meshkov instability using explicit molecular dissipation terms in formally high-order shock-capturing schemes, the implicit dissipation may exceed the molecular dissipation upon shock passage (and especially upon reshock of a mixing layer). Typically, implicit numerical diffusion present in the scheme or some type of subgrid-scale model (e.g., an “artificial viscosity”) must be used.

In the present simulations, the two-dimensional Euler equations augmented by the mass fraction conservation equation for the denser gas

$$\frac{\partial \phi}{\partial t} + \frac{\partial \mathbf{F}}{\partial x} + \frac{\partial \mathbf{G}}{\partial y} = 0 \quad (1)$$

are solved, where the conservative variables, ϕ , and inviscid fluxes, \mathbf{F} and \mathbf{G} , are

$$\phi = (\rho, \rho u, \rho v, \rho e, \rho m)^T, \quad (2)$$

$$\mathbf{F} = (\rho u, \rho u^2 + p, \rho u v, (\rho e + p) u, \rho m u)^T, \quad (3)$$

$$\mathbf{G} = (\rho v, \rho u v, \rho v^2 + p, (\rho e + p) v, \rho m v)^T. \quad (4)$$

Here, ρ is the density, $\mathbf{u} = (u, v)$ is the velocity, p is the pressure, $e = (u^2 + v^2)/2 + p/(\gamma - 1)$ is the total

energy per unit mass, m is the mass fraction (here of the sulfur hexafluoride gas, SF_6) and $p = \rho RT$ is the ideal gas pressure (R is the gas constant).

The simulations were performed using the finite-difference WENO shock-capturing method [2, 9]. The conservative finite-difference discretization of the Euler equations with WENO flux reconstruction contains implicit truncation errors that can be regarded as a non-linear, adaptive numerical dissipation. Although the Euler equations are solved in the present work, it is possible to compare the kinematic viscosity of the mixture, ν_{mix} , in the Collins and Jacobs experiment to the implicit numerical viscosity, ν_{num} . The kinematic viscosities of air(acetone) and SF_6 are $0.156 \text{ cm}^2/\text{s}$ and $0.026 \text{ cm}^2/\text{s}$, respectively [10]. Estimating the mixture dynamic viscosity by [11]

$$\mu_{\text{mix}} = \frac{X_1 \mu_1 \sqrt{M_1} + X_2 \mu_2 \sqrt{M_2}}{X_1 \sqrt{M_1} + X_2 \sqrt{M_2}}, \quad (5)$$

where X_r are the mole fractions of the mixture components and M_r are the molecular weights [2], gives $\nu_{\text{mix}} = \mu_{\text{mix}}/\rho_{\text{mix}} = 0.047 \text{ cm}^2/\text{s}$, where $\rho_{\text{mix}} = (\rho_1 + \rho_2)/2$. The computed numerical viscosity $\nu_{\text{num}} \approx 0.1 \text{ cm}^2/\text{s}$ following reshock [9] is close to the estimated ν_{mix} . In this previous study [9], the effects of order of WENO flux reconstruction and grid resolution on the quantities considered here showed that prior to reshock, these quantities exhibited little sensitivity to order and resolution. However, following reshock, higher order methods preserved more small-scale structure and had much lower numerical dissipation than lower-order methods at the same grid resolution. Furthermore, in Ref. 2 the agreement between the WENO densities and the experimental PLIF images, as well as the agreement of the amplitude with the experimental data and with the predictions of amplitude growth models, provided an element of validation of this numerical method. All of these results support using the ninth-order WENO method in investigating the physics of shock-interface interactions, including the reshocked Richtmyer-Meshkov instability.

III. THE PHENOMENOLOGY OF RESHOCK

Here, a temporal sequence of the density, vorticity, baroclinic vorticity production, and simulated density Schlieren fields is used to qualitatively describe the phenomenology of reshock, including the effects of vorticity deposition on the interface, shock focusing, and the effects of reflected waves on the interface evolution. Images of the vortex roll-up are also considered to describe the interior structure of the core during reshock. As in Refs. 2 and 9, the simulation fields are rotated 90° counterclockwise.

A. The density fields

Reshock is first visualized through the time sequence of density fields shown in Fig. 1 from 6 to 7.18 ms. At 6 ms (prior to reshock), the densities exhibit small-scale structures on the roll-up of the interface, which form as a result of the secondary deposition of vorticity occurring from the interface roll-up [12]. Reshock is a refraction from a heavier gas (SF_6) into a lighter gas [air(acetone)], resulting in a transmitted shock that enters the air(acetone) and a reflected rarefaction that returns back into the SF_6 . Reshock occurs over a timescale of $\approx 0.2\text{--}0.3$ ms. The densities show the arrival of the reflected shock at 6.38 ms, the shock refraction with the transmitted shock in air(acetone), the reflected rarefaction returning into the SF_6 at 6.48 and 6.58 ms, the *inversion* process when the bubble transforms into a spike (and vice versa) from 6.58–7 ms, and the formation of additional small roll-ups and complex, disordered structures at 7 and 7.18 ms. Curved waves can be seen in the SF_6 during inversion. At late times (11, 15 and 18 ms) shown in Fig. 2, a large spike of SF_6 develops. A finely-mixed complex layer is observed, which is the remnant of the spike prior to reshock. The creation of small disordered structures breaks symmetry and forms a complex interface topology.

B. The vorticity and baroclinic vorticity production fields

The vorticity evolution equation (in two dimensions and in the absence of dissipation) is

$$\frac{d\omega}{dt} = \mathcal{P} + \mathcal{C}, \quad \mathcal{P} \equiv \frac{\nabla\rho \times \nabla p}{\rho^2}, \quad \mathcal{C} \equiv -\omega \nabla \cdot \mathbf{u} \quad (6)$$

where $d/dt = \partial/\partial t + \mathbf{u} \cdot \nabla$, \mathcal{P} is the baroclinic vorticity production term, and \mathcal{C} is the vortex compression term (which is small compared to \mathcal{P}); ω and \mathcal{P} are shown in Fig. 3 at the same times as the density fields in Fig. 1. Baroclinic vorticity production causes the deposition of vorticity with opposite sign on the interface during reshock: this term is approximately two orders of magnitude larger than prior to reshock. The vorticity rolls-up into strong cores prior to reshock, as seen at 6 and 6.38 ms. At the same times, the density and pressure fluctuations along the interface cause the deposition of additional vorticity, as seen in \mathcal{P} at the corresponding times. During reshock at 6.48 ms, the baroclinic vorticity production is strongly increased along the interface, particularly on the sides of the bubbles, where the misalignment of $\nabla\rho$ and ∇p is greatest. The vorticity deposited is also approximately four times larger than the vorticity in the cores of the spikes prior to reshock. The vorticity of opposite sign subsequently drives the inversion process, which transforms the bubbles into spikes and vice versa, and induces the formation of additional disordered complex structures at 7 and 7.18 ms. Follow-




FIG. 1: (Color). The density fields at 6, 6.38, 6.48, 6.58, 6.58, 6.68, 6.78, 7 and 7.18 ms. The gases are air(acetone) (blue) and SF_6 (red).




FIG. 2: (Color). Same as Fig. 1 but at 11, 15 and 18 ms.

ing shock passage, baroclinic production becomes much weaker.

The vorticity and baroclinic vorticity production fields are shown in Fig. 4 at late times following reshock (11, 15 and 18 ms). The vorticity fragments into small localized vortices at late times. Baroclinic production is active at the interface and contributes to the development of complex structures and to increased mixing. At late times, \mathcal{P} is approximately two orders of magnitude smaller than at reshock. Furthermore, it remains approximately constant at late times, indicating that the arrival of the reflected rarefaction at ≈ 11 ms and of the reflected compression at ≈ 15 ms does not significantly affect baroclinic vorticity production.

A close-up of the density, vorticity, and baroclinic vorticity production fields inside the roll-up is shown in Fig. 5 during and after reshock. The density shows the formation of complex structures, the vorticity shows the dynamical mechanisms driving the formation of these structures, and the baroclinic vorticity production shows the mechanisms driving the vorticity through the misalignment of $\nabla\rho$ and ∇p inside the vortex core as the roll-up progresses. The baroclinic vorticity production increases at 6.58 ms in the roll-up, corresponding to the passage of one of the transmitted shocks during reshock. This shock can also be seen in the Schlieren images discussed in the next section. The passage of this shock causes the deposition of vorticity of opposite sign on the interface, which drives the inversion process and the formation of additional complex small-scale structures.




FIG. 3: (Color). The vorticity, ω , and baroclinic vorticity production, \mathcal{P} , fields at 6, 6.38, 6.48, 6.58, 6.68, 6.78, 7 and 7.18 ms.

C. The simulated density Schlieren fields

The detailed wave structure can be investigated using simulated density Schlieren fields, which visualize density gradients associated with the waves and fine-scale mixing structures. The density Schlieren is that used in two-dimensional fifth-order WENO simulations of a shock-cylinder interaction [13],

$$\Phi(x, y, t) = \exp \left[-\alpha(m) \frac{|\nabla \rho|}{\max |\nabla \rho|} \right], \quad (7)$$

where $\alpha(m) = 20$ for $m > 0.25$ and $\alpha(m) = 100$ otherwise, and m is the mass fraction of SF_6 .

A temporal sequence of simulated density Schlieren fields during reshock and at late times is shown in Fig. 6. The fields sharply capture the diffuse interface, the complex wave structure during reshock, and the small-scale structures at late times. Note the focusing effect of the waves and the bubble transforming into the spike and vice versa. The field at 11 ms shows the arrival of the reflected rarefaction produced by the second shock after it has reflected from the end wall of the domain. The rich




FIG. 4: (Color). Same as Fig. 3 but at 11, 15 and 18 ms.




FIG. 5: (Color). Details of the roll-up in the density, ρ , vorticity, ω , and baroclinic vorticity production, \mathcal{P} , fields at 6.48, 6.58, 6.68 and 7 ms. The images show the complex structure in the roll-up, including the effects of reshock on this structure.

structure displayed by Φ shows weak, curved waves that intersect and cause focusing. As these waves are weak due to the small Mach number, it is unclear whether they have an important role in the flow dynamics. At late times, the waves have dissipated and are no longer present in the unmixed gases. Similar complex wave structures were also observed following a shock-interface interaction in the $Ma = 1.32$ experimental Schlieren images of Brouillette and Sturtevant [14].

IV. BAROCLINIC CIRCULATION DEPOSITION ON THE INTERFACE

The circulation deposited on the interface by the shock quantifies the principal mechanism driving the evolution of the Richtmyer-Meshkov instability. Reshock imparts additional circulation on the interface, contributing to the rapid development of complex flow structure through baroclinic production.

Consider a closed oriented curve C enclosing a surface S , so that the circulation is

$$\Gamma(t) = \oint_C \mathbf{u} \cdot d\mathbf{r} = \int_S \boldsymbol{\omega} \cdot d\mathbf{S},$$

where $d\mathbf{r}$ is tangent to C and $d\mathbf{S}$ is the outward oriented area element. Here, the vorticity vector and the area element are parallel; as a result, the latter definition for computing the circulation is adopted, with $d\mathbf{S}$ taken to be a rectangle. In the limit of a rectangle with infinite length in the streamwise (x) direction and negligible width in the spanwise (y) direction, the circulation on the interface is approximated as $\langle \omega \rangle(y, t) \equiv \int_{-\infty}^{\infty} \omega(x, y, t) dx$.

Samtaney and Zabusky [15] analytically derived

$$\langle \omega \rangle(y, 0^+) = \Gamma'_1 k a_0^+ \sin(ky) \quad (8)$$

for the circulation deposition on the sinusoidal interface by the initial incident shock in a fast/slow interaction, i.e., when the refraction is from a lighter gas into a heavier gas, where

$$\Gamma'_1 = \frac{c_1}{Ma_s} \left[\frac{1}{\gamma_2 - 1} \frac{1 - \psi(p_4/p_2)}{\eta \gamma_1 / \gamma_2} - \frac{1 - \psi(p_5/p_3) \psi(p_3/p_1)}{\gamma_1 - 1} \right], \quad (9)$$

$p_1 = p_2$ is the initial pressure ahead of the incident shock, p_3 is the pressure behind the incident shock, $p_4 = p_5$ is the pressure behind the reflected and transmitted shocks, η is the initial density ratio, c_1 is the initial sound speed, and the ratio of sound speed across the incident, reflected and transmitted shocks is given by $\sqrt{\psi(p_3/p_1)}$, $\sqrt{\psi(p_5/p_3)}$ and $\sqrt{\psi(p_4/p_2)}$, respectively, where $\psi(r) \equiv r(1 + \mu r)/(\mu + r)$ and $\mu \equiv (\gamma + 1)/(\gamma - 1)$. In the present simulation, $\Gamma'_1 = 8736.12$ cm/s, the perturbation wave number is $k = 2\pi/\lambda = 1.065$ cm⁻¹, and the post-shock

perturbation amplitude is $a_0^+ = 0.162$ cm. The initial interfacial circulation deposited by the incident shock, together with the prediction of Eq. (8), are shown in Fig. 7, indicating that the circulation deposition is sinusoidally distributed and closely follows the Samtaney-Zabusky model prediction. The model underpredicts the simulation data by $\approx 4\%$. Also shown is a comparison with linear instability theory $\langle \omega(y, 0^+) \rangle = d\mu/dy$, where $\mu(y) = 2v_0 \cos(ky)$ and $v_0 = 702$ cm/s is the initial instability growth rate, indicating that linear instability theory is in close agreement with the Samtaney-Zabusky model and also underpredicts the simulation data.

As the circulation deposition is sinusoidal, the regions $S^- = [0, L_x] \times [0, \lambda/2]$ and $S^+ = [0, L_x] \times [\lambda/2, \lambda]$ enclose the circulation, $\langle \omega \rangle(y, t)$, of the same sign. Thus, define the positive and negative circulations

$$\Gamma^\pm(t) \equiv \iint_{S^\pm} \omega(x, y, t) dx dy \quad (10)$$

(see [9] for details on their computation) which, as shown in Fig. 8, overlap prior to reshock, indicating symmetry as the instability evolves. At ≈ 6 ms, a small decrease resulting from the deposition of vorticity of opposite sign due to baroclinic vorticity deposition [12] is observed (this can also be seen in the roll-up in Fig. 5, where the vorticity forms positive and negative patches). Reshock at ≈ 6.4 ms deposits vorticity of opposite sign on the interface, as the refraction is from a heavier gas into a lighter gas. Consequently, both circulations increase sharply. The large values of Γ^\pm following reshock are due to the more complex topology of the interface, resulting in a larger amount of vorticity deposited by the shock. The secondary baroclinic circulation also explains the more pronounced increase in $|\Gamma^\pm|$ following reshock, but prior to the arrival of the reflected rarefaction. Following the arrival of the reflected rarefaction, Γ^+ and $|\Gamma^-|$ begin to differ as a result of symmetry breaking. Vortical structures become disordered and may cross from the positive region into the negative region, inducing the asymmetry. Baroclinic secondary instability and the rapid growth of positive and negative circulation was investigated in a two-dimensional stratified mixing layer [16–18].

V. THE POST-RESHOCK MIXING LAYER WIDTH

The post-reshock mixing layer width is compared here to the phenomenological Mikaelian [19], Brouillette and Sturtevant [20], and Charakhch'yan [21] reshock models, which predict linear growth for short times following reshock. These models apply up to the time of the interaction with the reflected rarefaction at ≈ 11 ms. The numerical determination of the mixing layer width is discussed in detail in Ref. 2.




FIG. 6: (Color). Simulated density Schlieren fields Φ showing the complex wave structure during reshock, shock focusing, and the fine-scale structure of the interface at late times.

A. The Mikaelian reshock model

A linear power-law model for the post-reshock mixing layer width was developed [19] based on the rocket-rig experimental results of Read [22] and Youngs [23] for the growth of the Rayleigh-Taylor unstable mixing layer. Assuming no dependence on the initial conditions and neglecting molecular dissipation (and other) effects [24]

$$\frac{dh}{dt} = 0.28 \Delta u_1 A_1^+, \quad (11)$$

where Δu_1 is the change in the interface velocity resulting from reshock, A_1^+ is the post-reshock Atwood number, and the empirical coefficient 0.28 is based on the mixing layer growth rate from the experiments [22].

B. The Brouillette-Sturtevant reshock model

Brouillette and Sturtevant [20] studied the effect of a diffuse interface on Richtmyer-Meshkov instability growth in shock tube experiments in which a thin metal

plate initially separated air and SF₆ or air and Freon-22 gas. Prior to launching the shock wave in the air, the plate was slowly withdrawn to generate a quasi-sinusoidal perturbation, leaving a thick diffuse interface separating the gases. The $Ma = 1.12$ – 1.66 shock refracted at the interface and was transmitted into the second gas. The transmitted shock reflected from the end wall of the shock tube test section and reshocked the interface.

The growth of the total width of a single-mode interface following $N + 1$ impulsive accelerations (or decelerations) was empirically modeled by generalizing the Richtmyer model:

$$\left(\frac{dh}{dt}\right)_N = 2k \sum_{i=0}^N \frac{\Delta u_i A_i^+ a_i^+}{\psi_i^+}, \quad (12)$$

where k is the initial perturbation wave number, Δu_i is the change in the velocity of the interface corresponding to the i th wave, A_i^+ and a_i^+ are the post-(re)shock Atwood number and amplitude, ψ_i^+ is the post-shock growth reduction factor [25] and the factor of two accounts for the width (twice the amplitude a). As the growth is based on the Richtmyer model for each impul-




FIG. 7: The initial circulation deposition on the interface $\langle \omega \rangle(y, 0^+)$ (solid line) by the incident shock at time $t = 0^+$, together with the predictions of the Samtaney-Zabusky model [Eq. (8)] (dashed line) and linear theory (dashed grey line).

sive acceleration phase, the Brouillette-Sturtevant model predicts piecewise-linear-in-time growth phases. This model also accounts for the multiple waves that successively interact with the interface. Brouillette and Sturtevant reported good agreement between their data and the prediction of Eq. (12).

C. The Charakhch'yan reshock model

Charakhch'yan [21] assumed that the reshock is much weaker than the initial incident wave and occurs during the nonlinear phase when $a(t) \sim \lambda$. As a result, the change in Atwood number caused by reshock can be neglected and

$$\frac{dh}{dt} = 2v_0 - 2.5 \Delta u_1 A_0^+, \quad (13)$$

where A_0^+ is the Atwood number following the passage of the *initial* shock, the empirical coefficient 2.5 is based on two-dimensional simulations, and da_r^-/dt in Ref. 21 was interpreted as the Richtmyer velocity v_0 . A wide range of numerical simulations was reported to support this model.

D. Comparison of the post-reshock mixing layer width to model predictions

The comparison of the post-reshock mixing layer width to the predictions of the models is shown in Fig. 9. Reshock in the simulation occurs at nearly the same time as in the experiment, as confirmed by both the $x-t$ diagram and the mixing layer amplitude in Ref. 2. For the Mikaelian model, the values $\Delta u_1 = 8000$ cm/s and




FIG. 8: The time-evolution of the positive and negative circulation $\Gamma^\pm(t)$ on the interface before (top) and after (bottom) reshock.

$A_1^+ = 0.6448$ from the simulation were used. For the Brouillette-Sturtevant model,

$$\left(\frac{dh}{dt} \right)_1 = 2k \left(\frac{\Delta u A_0^+ a_0^+}{\psi_0^+} + \frac{\Delta u_1 A_1^+ a_1^+}{\psi_1^+} \right) \quad (14)$$

was used, where the post-reshock amplitude is $a_1^+ = 1.5$ cm, $\Delta u_1 = 8000$ cm/s, and the value of ψ_1^+ was extrapolated as follows. First, recall that ψ is linear in δ/λ , where δ is the diffuse mixing layer width (linear in the post-shock perturbation amplitude a_0^+). As $a_1^+ \approx 10a_0^+$ and the value of ψ_0^+ is known, the new value of a_1^+ gives the new value of ψ_1^+ . The value $A_0^+ = 0.6045$ from Table I of Ref. 2 was used for the Charakhch'yan model.

Agreement between the mixing layer width and the predictions of the models can only be expected following reshock for $t > 6.5$ ms and prior to the arrival of the reflected rarefaction at ≈ 11 ms. Comparison of $h(t)$ with the model predictions confirms that the growth is approximately linear for very short times (< 2 ms) following reshock. The simulation width generally agrees well with the both the Mikaelian and Charakhch'yan model predictions at early times following reshock. Vetter and Sturtevant [24] also concluded that the widths from their reshocked air/SF₆ experiments agreed with the Mikaelian model within 1.5–23%, depending on the shock strength.




FIG. 9: (Color). The mixing layer width $h(t)$ showing reshock at ≈ 6.4 ms and the arrival of the reflected rarefaction at ≈ 11 ms. Also shown are the predictions of the models, which are shown starting at $t_r^+ = 6.58$ ms and at $h(t_r^+) = 3.1$ cm. Using the values in the text yields $(dh/dt)_1 = 1444.35$ cm/s, $(dh/dt)_1 = 1195.1$ cm/s and $(dh/dt)_1 = 1492$ cm/s for the Mikaelian, Brouillette-Sturtevant, and Charakhch'yan models, respectively. The variations in the growth rate at ≈ 11 and 15 ms are due to further wave interactions with the evolving interface, and are not accounted for by these models.

For later times, the Mikaelian and Charakhch'yan models overestimate the growth, while the Brouillette-Sturtevant model underestimates the growth at all times following reshock. These reshocked amplitude growth predictions only depend on Δu_1 and A_1^+ . The Mikaelian and Charakhch'yan models do not depend on k (in contrast to the Brouillette-Sturtevant model) and contain empirical parameters. The Mikaelian model is based on an estimate of the growth rate of a *three-dimensional* multi-mode Rayleigh-Taylor experiment, while the Brouillette-Sturtevant and Charakhch'yan models are fundamentally *two-dimensional* single-mode (and thus more consistent with the present simulations). The comparison shown here cannot distinguish which model best agrees with the simulation data.

VI. ANALYSIS OF MIXING AND THE PHYSICS OF RESHOCK

Presented here is an investigation of quantities characterizing the evolution of mixing induced by the reshocked Richtmyer-Meshkov instability. The quantities are adapted from similar investigations of the Rayleigh-Taylor instability [26] and, more generally, of turbulent flows. The variety of quantities considered—mixing profiles and parameters, production and mixing fractions, energy spectra, and statistics—allow a quantitative estimate of the effects of reshock for the first time. The

effects of additional waves on the evolution of the layer are also considered.

The initial condition is deterministic in the present study. Furthermore, only one realization of the flow is simulated. Thus, the profiles, spectra and statistics exhibit variations that would otherwise be reduced if an ensemble average of realizations with stochastic initial conditions was used. The observed *trends*, however, are not expected to differ significantly between a single realization and an ensemble average.

A. Definitions of averages, fluctuations, spectra, and statistics

Several averages are introduced to define streamwise profiles across the mixing layer, as well as fluctuations required to define energy spectra. Let an overbar denote an instantaneous average of a field $\phi(x, y, t)$ over the spanwise direction y with length L_y :

$$\bar{\phi}(x, t) = \frac{1}{L_y} \int_0^{L_y} \phi(x, y, t) dy \quad , \quad \tilde{\phi}(x, t) = \frac{\rho \phi}{\bar{\rho}} \quad , \quad (15)$$

which can be interpreted as an instantaneous Reynolds average (see Ref. 7 for the three-dimensional analog) and an instantaneous Favre average, respectively; the latter is used for the statistical analysis of variable-density and compressible flows [27, 28]. Then the corresponding Reynolds and Favre fluctuating fields are

$$\phi(x, y, t)' = \phi(x, y, t) - \bar{\phi}(x, t) \quad (16)$$

$$\phi(x, y, t)'' = \phi(x, y, t) - \tilde{\phi}(x, t) \quad . \quad (17)$$

Thus, in the present analysis, the Reynolds- and Favre-averaged fields are functions only of the streamwise coordinate x and of time, while the fluctuating fields are functions of both the streamwise and spanwise coordinates x and y , and of time.

The time-evolution of the fluctuating kinetic energy, fluctuating enstrophy, density variance, pressure variance, and baroclinic vorticity production variance spectra quantify the growth of different scales of the flow. The spectral analysis used here adapts the formulations previously applied to inhomogeneous flows with one direction of statistical-homogeneity [8, 12, 29, 30]. The general procedure used to compute the spectrum of a quantity $\phi(x, y, t)$ within the mixing layer $x \in [\ell_s, \ell_b]$ is described in Ref. 9 (where ℓ_b and ℓ_s are the positions of the bubble and spike, respectively). The energy associated with each Fourier mode k is obtained by averaging over the mixing layer to obtain the one-dimensional energy spectrum

$$E_{\phi\phi}(k, t) = \frac{1}{h(t)} \int_{\ell_s(t)}^{\ell_b(t)} \hat{\phi}(k, x, t)^* \hat{\phi}(k, x, t) dx \quad , \quad (18)$$

where the superscript $*$ denotes complex conjugation. In the results presented below, all modes above the Nyquist

wave number $\pi/\Delta x$ are neglected [31]. The spectrum $E_{\phi\phi}(k, t)$ provides information on the energy content of all of the scales in the autocorrelation of ϕ as a function of time. The characteristic scale of a structure with wave number k is $\sim 1/k$. The time-evolution of statistics

$$E_{\phi\phi}(t) = \int_0^\infty E_{\phi\phi}(k, t) dk \quad (19)$$

is also considered.

B. Local and global mixing properties

Quantities previously used in the analysis of Rayleigh-Taylor mixing [7, 26] are adapted here to the two-dimensional Richtmyer-Meshkov instability, including mixing profiles, production and mixing fractions, energy spectra, and statistics [9].

1. Mole and mixing fraction profiles

To simplify the comparison of the profiles across the mixing layer at different times, the streamwise coordinate is recentered by the midpoint between the bubble and spike location $x_{\text{mid}}(t)$, and divided by the mixing layer width $h(t)$. Thus the mixing layer extent is mapped to $[x - x_{\text{mid}}(t)]/h(t) \in [-0.5, 0.5]$, where $[-0.5, 0]$ is the air(acetone) side and $[0, 0.5]$ is the SF₆ side, which facilitates the analysis of self-similarity.

a. Mole fraction profile Consider the mole fraction

$$X(x, y, t) = \frac{m(x, y, t) M_1}{[1 - m(x, y, t)] M_2 + m(x, y, t) M_1} \quad (20)$$

averaged over the spanwise direction y , $\bar{X}(x, t)$. The gases have densities ρ_1 and $\rho_2 > \rho_1$; M_1 and M_2 are the molecular weights of the air(acetone) and SF₆, respectively, and m is the mass fraction of SF₆. The mole fraction profile $\bar{X}(x, t)$ gives the relative mass distribution of the two gases within the mixing layer. If the gases are uniformly-distributed over the entire domain, as expected in complete homogeneous mixing, then $\bar{X} = 0.5$.

The mole fraction profiles shown in Fig. 10 at 1 ms time intervals increase from $\bar{X} = 0$ in the air(acetone) to $\bar{X} = 1$ in the SF₆. The profiles show the varying mass distribution inside the layer prior to reshock. At the initial time, \bar{X} increases monotonically as the initial interface is slightly diffused. After the initial shock passage, \bar{X} widens and becomes non-monotonic, with a peak developing within the air(acetone) that moves left and increases in magnitude with time. This is due to spikes of SF₆ penetrating into the air(acetone), forming the characteristic roll-ups resulting from the entrainment of additional SF₆ within the air(acetone). The bubbles of air(acetone) “rise” in the SF₆, causing an overall decrease in \bar{X} in the SF₆. Reshock compresses \bar{X} between 6 and 7 ms, and also generates additional structure reflected

in the distribution of \bar{X} (now with localized peaks). At later times, the layer growth slows, the peaks decrease in magnitude, and \bar{X} begins to approach 0.5 on average, indicating a well-mixed distribution of mass within the layer.

b. Product mole fraction profile A quantitative measure of mixing can be defined using $X(x, y, t)$ as follows. Suppose that the two fluids undergo a fast kinetic reaction, so that the amount of ‘product’ produced is [26, 32]

$$X_p(x, y, t) = \begin{cases} \frac{X(x, y, t)}{X_s} & \text{for } X \leq X_s \\ \frac{1 - X(x, y, t)}{1 - X_s} & \text{for } X > X_s \end{cases}, \quad (21)$$

where $X_s = 1/2$ (indicating that the product is composed of one mole of each ‘reactant’), and is limited by the amount of reactant (either heavy or light fluid). The product mole fraction profile $\bar{X}_p(x, t) \in [0, 1]$ is a measure of how well mixed the ‘reactants’ are. Here, numerically induced mixing is a surrogate for a chemical reaction.

The time-evolution of the product mole fraction profile \bar{X}_p is shown in Fig. 10. The initial diffusion layer is well-mixed, with a rapid decrease in \bar{X}_p between 0 and 1 ms, as expected; \bar{X}_p broadens and rapidly develops sharp cusps near the mixing layer edges $x/h(t) \approx \pm 0.5$ as the heavy gas spikes penetrate into the lighter gas and the light gas bubbles “rise” in the lighter gas (and the gases become less mixed). At these early times in the instability evolution, \bar{X}_p is nearly-symmetric, with values ≈ 0.04 – 0.06 over most of the layer. A pronounced asymmetry develops for $t > 2$ ms: as the roll-ups form, additional peaks within the air(acetone) develop, corresponding to well-mixed fluid in the cores. Reshock significantly increases \bar{X}_p , as additional fine-scale structures form; \bar{X}_p oscillates following reshock, which persists to late times. At late times the gases are more mixed and larger peaks develop in the SF₆, corresponding to the arrival of multiple reflected waves that increase mixing at this end of the layer. The gases are less mixed in the region closer to the air(acetone) than in the region closer to the SF₆.

c. Mixing fraction profile Averaging the volume fractions of gases $r = 1$ and 2 ,

$$f_r(x, y, t) \equiv \frac{\frac{m_r(x, y, t)}{\rho_1}}{\frac{m_1(x, y, t)}{\rho_1} + \frac{m_2(x, y, t)}{\rho_2}}, \quad (22)$$

over the spanwise direction gives the volume fraction profiles $\bar{f}_r(x, t)$. The molecular mixing fraction $\theta(x, t)$ is then obtained from the volume fraction profiles and the average of their product [7, 33],

$$\theta(x, t) = \frac{\bar{f}_1 \bar{f}_2}{\bar{f}_1 \bar{f}_2}, \quad (23)$$

where $f_2 = 1 - f_1$. When $\theta = 0$ (corresponding to $\bar{f}_1 \bar{f}_2 = 0$), the two fluids are completely segregated, and when $\theta = 1$ (corresponding to $\bar{f}_1 \bar{f}_2 = \bar{f}_1 \bar{f}_2$), they are




FIG. 10: (Color). The time-evolution of the mole fraction profile \bar{X} (top row), product mole fraction profile \bar{X}_p (middle row) and mixing fraction profile θ (bottom row) across the mixing layer at 1 ms intervals. The profiles before reshock (0–5 ms) are shown in the first column, and the profiles following reshock (6–17 ms) are shown in the second and third columns.

completely molecularly mixed. Prior to reshock, the evolution of θ in Fig. 10 indicates increased mixing toward the air(acetone) side of the layer, corresponding to the formation of the roll-up with a well-mixed core. Additional mixed structures within the roll-up are indicated by the peaks at 5 and 6 ms. At reshock (≈ 6.4 ms), the layer is compressed by the passage of the shock [which continues into the air(acetone)] and is “pulled back” on the SF_6 side by the reflected rarefaction. This effect is captured by θ , which is larger on the air(acetone) side and smaller on the SF_6 side. Following reshock, the layer contains complex, disordered, well-mixed structures, as measured by a large value of θ across the layer (varying from approximately 0.6 to 1). On average, the largest values of θ across the entire layer are attained at the latest time, with the degree of mixedness increasing, as indicated by θ approaching unity. *This ‘molecular mixing’ is induced by stirring and by the numerical diffusion in the algorithm, rather than by molecular processes.* Comparison of the profiles at 6–11 ms to those at 12–17 ms shows

that \bar{X} , \bar{X}_p and θ become approximately self-similar at late times, with intermittent variations across the layer.

2. Production and mixing fractions

The total chemical product $P_t(t)$ and maximum chemical product $P_m(t)$ are [26, 32]

$$P_t(t) = \frac{1}{h(t)} \int_{\ell_s(t)}^{\ell_b(t)} \bar{X}_p dx \quad (24)$$

$$P_m(t) = \frac{1}{h(t)} \int_{\ell_s(t)}^{\ell_b(t)} X_p(\bar{X}) dx .$$

In general $P_m(t) > P_t(t)$, and is the amount of product obtained if both ‘reactants’ were homogeneously mixed.

The mixing fractions $\Xi(t)$ and $\Theta(t)$ are [7, 26, 33]

$$\Xi(t) = \frac{P_t(t)}{P_m(t)}, \quad \Theta(t) = \frac{\int_{\ell_s(t)}^{\ell_b(t)} \overline{f_1 f_2} dx}{\int_{\ell_s(t)}^{\ell_b(t)} \overline{f_1} \overline{f_2} dx}. \quad (25)$$

Note that Ξ is a measure of the total product formed relative to the product that would be formed if all entrained fluid were completely mixed. Larger Ξ and Θ correspond to more complete mixing.

The time-evolution of the global production quantities P_t and P_m is shown in Fig. 11. Before reshock, P_t increases, indicating an increase in mixing, while P_m decreases. The decrease in P_m indicates that the maximum molar product decreases due to the bubble (which creates large, unmixed structures within the mixing layer). During reshock, the mixing layer is compressed, inducing additional mixing as measured by P_t . Following reshock, P_t increases rapidly, indicating significantly increased mixing. As the mixing progresses, large unmixed structures form, decreasing P_m .

The time-evolution of the mixing fractions Ξ and Θ , also shown in Fig. 11, provide qualitatively similar information, with $\Theta(t) > \Xi(t)$ for all time. As the gases mix following the initial shock passage, the fractions decrease and then increase. Immediately after reshock, the layer is compressed, causing a well-mixed region to form, as shown by the sharply-peaked values of Ξ and Θ . This well-mixed region is characterized by the rapid production of small-scale flow structures by reshock, as shown in Sec. III. Following reshock, Ξ and Θ increase faster than before reshock, indicating that *reshock increases the mixing rate*, as expected physically. Subsequently, the mixing layer undergoes a rapid growth as the bubbles and spikes invert, causing the overall mixing rate to decrease rapidly over a time interval of ≈ 2 ms. Following this decrease, the formation of complex flow features with roll-ups again increases the mixing.

C. Energy spectra and statistics

The time-evolutions of fluctuating energy spectra provide a measure of the growth and decay of the energy at different scales in the flow at a given time. The post-reshock energy spectra are compared here to the scalings suggested by two-dimensional turbulence phenomenology. The time-evolution of statistics (integrated energy spectra) are also considered.

1. Energy spectra

The time-evolution of the total, streamwise, and spanwise fluctuating kinetic energy spectra per unit volume $E(k, t) = E_{u''u''}(k, t) + E_{v''v''}(k, t)$, $E_{u''u''}(k, t)$, and $E_{v''v''}(k, t)$, respectively, is shown in Fig. 12. The oscillations for small and intermediate wave numbers k at




FIG. 11: The time-evolution of the production fractions $P_t(t)$ and $P_m(t)$ (top), and mixing fractions $\Xi(t)$ and $\Theta(t)$ (bottom).

early times and following reshock are damped out at late times, as indicated by the more smoothly decaying spectra at large t . Reshock sharply increases the spectra between 6 and 7 ms by exciting a wide range of scales, thereby imparting additional energy into the mixing layer. For $t > 8$ ms, the spectra decay very slowly in time. The streamwise fluctuating kinetic energy spectrum $E_{u''u''}(k, t)$ is peaked at $k \approx 1.5 \text{ cm}^{-1}$, while the peak of the spanwise fluctuating kinetic energy spectrum $E_{v''v''}(k, t)$ corresponds to the largest scale of the flow. As expected, there is more energy content in the streamwise than spanwise velocity fluctuations, i.e., the energy spectra are anisotropic. Thus, the evolution of $E(k, t)$ is dominated by that of $E_{u''u''}(k, t)$.

The time-evolution of the ratio of the streamwise and spanwise fluctuating kinetic energy spectra $E_{u''u''}(k, t)/E_{v''v''}(k, t)$ is shown in Fig. 13. The energy content is similar at early times, approaching a constant for all scales at 1 ms. At later times prior to reshock, the ratio increases at intermediate and large k , with similar increases at reshock. At late times ($t \geq 13$ ms) the ratio changes very little with time, with a value ≈ 2 at the smallest k increasing nearly linearly to ≈ 6 at $k \approx 170 \text{ cm}^{-1}$, and then decreasing to ≈ 4.5 at $k \approx 270 \text{ cm}^{-1}$. Reshock, the arrival of the reflected rarefaction wave at

≈ 11 ms, and the compression wave at ≈ 15 ms decrease the ratio, indicating that these waves excite more spanwise fluctuations relative to streamwise fluctuations. At late times, the ratio indicates that *spectral isotropy* (i.e., a ratio of unity) is not achieved, as the streamwise fluctuating kinetic energy spectrum contains more small-scale energy than the spanwise spectrum by a factor ≈ 2 –6.

The fluctuating enstrophy density per unit mass is $\Omega(x, y, t)'' = [\omega(x, y, t)'']^2/2$ and the time-evolution of the fluctuating enstrophy spectrum $E_{\omega''\omega''}(k, t)$ is shown in Fig. 14. In a homogeneous flow, the fluctuating enstrophy and kinetic energy spectra are related by $E_{\omega''\omega''}(k, t) = k^2 E(k, t)$, so that the peak of $E_{\omega''\omega''}(k, t)$ is weighted toward smaller scales than that of $E(k, t)$. The fluctuating enstrophy spectrum is less steep than the fluctuating kinetic energy spectrum. As in the case of $E(k, t)$, reshock primarily amplifies $E_{\omega''\omega''}(k, t)$ but does not change its shape. The spectrum decays very slowly at late times.

The time-evolution of the density variance spectrum $E_{\rho'\rho'}(k, t)$ and pressure variance spectrum $E_{p'p'}(k, t)$ is also shown in Fig. 14. The density variance spectrum varies very little with time, indicating that density fluctuations are not as strongly affected by reshock as other quantities. Moreover, the magnitude of $E_{\rho'\rho'}(k, t)$ is very small for all times, indicating that the density variance is very small and the flow is nearly-incompressible: additional numerical evidence exists to support this conclusion in Richtmyer-Meshkov unstable flows [34, 35]. The spectrum also shows the interaction of reflected waves with the mixing layer, which have a profound effect on $E_{p'p'}(k, t)$. The pressure variance spectrum increases very rapidly upon reshock at ≈ 6.4 ms, and also exhibits a rapid increase at ≈ 11 ms when the reflected rarefaction interacts with the mixing layer. Both $E_{\rho'\rho'}(k, t)$ and $E_{p'p'}(k, t)$ decay very slowly at late times.

To determine the scales at which the baroclinic vorticity production [Eq. (6)] is large, consider the baroclinic vorticity production variance spectrum $E_{\mathcal{P}\mathcal{P}}(k, t)$ shown in Fig. 14. This spectrum is large and nearly constant over all scales for $t > 2$ ms and decreases rapidly at the smallest scales, where numerical dissipation dominates. The spectrum increases rapidly after 2 ms (prior to reshock), consistent with additional baroclinic production at the interface where the roll-up occurs. Prior to reshock, $E_{\mathcal{P}\mathcal{P}}(k, t)$ is highly oscillatory. A large increase occurs at reshock, followed by a rapid decay. A further increase occurs between 11 and 12 ms, corresponding to the arrival of the reflected rarefaction. At late times, $E_{\mathcal{P}\mathcal{P}}(k, t)$ decays at a significantly faster rate than the other spectra.

2. Comparison to classical two-dimensional turbulence phenomenology

The fluctuating kinetic energy, enstrophy, and density and pressure variance spectra are compared here to the

inertial subrange scalings predicted by two-dimensional turbulence phenomenology [36–41] at 7 and 18 ms (late time). It is of interest to explore whether these scalings apply in the reshocked two-dimensional Richtmyer-Meshkov unstable flow considered here. This phenomenology applies to incompressible and isotropic turbulence; here it is assumed that the flow is nearly-incompressible and approximately isotropic at late times after reshock.

For sufficiently large Reynolds numbers, the inertial subrange fluctuating (turbulent) kinetic energy spectrum scales as

$$E(k, t) = Ko' \eta(t)^{2/3} k^{-3}, \quad k_f \ll k \ll k_\eta(t), \quad (26)$$

which was originally proposed for turbulence forced at a wave number k_f , with kinetic energy and enstrophy injection rates ϵ and η , respectively. Here, Ko' is the two-dimensional analog of the Kolmogorov constant and $k_\eta(t) = [\eta(t)/\nu^3]^{1/6}$ is the enstrophy dissipation wave number (with ν the kinematic viscosity). A subsequent logarithmic modification to the k^{-3} law explicitly depending on k_f was found to be required for self-consistency [42]. The inertial subrange enstrophy spectrum corresponding to (26) is

$$E_{\omega'\omega'}(k, t) = Ko' \eta(t)^{2/3} k^{-1}. \quad (27)$$

These are the fluctuating kinetic energy and enstrophy spectra in the *enstrophy cascade subrange* in which the turbulent kinetic energy and enstrophy flux are $\Pi(k, t) \approx 0$ and $\Pi_\Omega(k, t) \approx \eta(t)$, respectively.

In the *inverse kinetic energy cascade subrange* in which $\Pi(k, t) \approx \epsilon(t)$ and $\Pi_\Omega(k, t) \approx 0$, the inertial subrange kinetic energy spectrum for small wave numbers is

$$E(k, t) = Ko'' \epsilon(t)^{2/3} k^{-5/3}, \quad k \ll k_f \quad (28)$$

with a different Kolmogorov constant Ko'' . The corresponding inertial subrange enstrophy spectrum is

$$E_{\omega'\omega'}(k, t) = Ko'' \epsilon(t)^{2/3} k^{1/3}, \quad k \ll k_f. \quad (29)$$

The above scaling laws are a consequence of the conservation of kinetic energy *and* enstrophy in two-dimensional, nearly-inviscid, high Reynolds number turbulent flows.

The fluctuating kinetic energy spectrum $E(k)$ shortly after reshock at 7 ms is shown in Fig. 15, together with the expected power-laws: $k^{-5/3}$ for the small, energy-containing wave numbers and k^{-3} for larger wave numbers. Also shown are the compensated spectra used to verify the approximate scalings and the wave number extent of these scalings (less than one decade). Peng, Zabusky and Zhang [12] also found that the density-weighted kinetic energy spectrum was consistent with a k^{-3} power-law in their two-dimensional piecewise-parabolic method simulation of a model of the Jacobs and Krivets [43] $Ma = 1.3$ single-mode, air/SF₆ Richtmyer-Meshkov instability experiment.

The fluctuating enstrophy spectrum $E_{\omega''\omega''}(k)$ is also shown in Fig. 15. As the vorticity is the spatial derivative of the velocity field, the fluctuating enstrophy and fluctuating kinetic energy spectra are related by $E_{\omega''\omega''}(k, t) = k^2 E(k, t)$. Consequently, the expected inertial subrange scalings for the intermediate and large wave numbers are $k^{1/3}$ and k^{-1} , respectively. The compensated spectra show that a possible k^{-1} scaling is observed over less than one decade; however, there is an insufficient range of wave numbers in the largest scales to ascertain whether a $k^{1/3}$ scaling exists.

The fluctuating kinetic energy spectrum $E(k)$ and fluctuating enstrophy spectrum $E_{\omega''\omega''}(k)$ are shown at late time (18 ms) in Fig. 16. The fluctuating kinetic energy spectrum exhibits a scaling similar to the one immediately following reshock, viz. $k^{-5/3}$ for intermediate scales and k^{-3} for small scales: this is further verified by the compensated energy spectrum. However, $E_{\omega''\omega''}(k)$ exhibits a different scaling than expected in classical two-dimensional turbulence: a decay close to $k^{-2/3}$ and k^{-2} is observed for intermediate and small scales, respectively, as shown by the compensated spectra. *The fluctuating enstrophy scalings $k^{1/3}$ and k^{-1} can only be expected in the presence of forcing that sustains the turbulence.* In the absence of such forcing, the spectra may satisfy different scalings. In the case of the Richtmyer-Meshkov instability, a *transient forcing* in the momentum equation, $F \sim \rho \Delta v_1 \delta(t)$, is induced by the interaction of the reflected shock with the evolving mixing layer (reshock), where Δv_1 is the velocity jump. Following the passage of the shock, no additional forcing is provided, and the turbulence enters a *quasi-decay phase*, in which only interactions with smaller amplitude waves occur: this results in *different* scaling laws for the late-time enstrophy spectrum than expected classically.

In a nearly-incompressible flow, density fluctuations are expected to be weakly coupled to the velocity field, so that the density fluctuations may be passively advected by the velocity field. If the Reynolds number is sufficiently large to support an inertial subrange, and the Schmidt number is small enough so that molecular diffusivity is negligible over a large wave number range, then there is an inertial-convective subrange in which density fluctuations are advected and the density variance spectrum scales as

$$E_{\rho'\rho'}(k, t) = Ba \frac{\epsilon_{\rho'\rho'}(t)}{\epsilon(t)^{1/3}} k^{-5/3}, \quad (30)$$

where Ba is the Batchelor (or Corrsin-Obukhov) constant and $\epsilon_{\rho'\rho'}(t)$ is the density variance dissipation rate [44–48]. Dimensionally, the inertial subrange pressure variance spectrum in two-dimensional large Reynolds number turbulence is proportional to $kE(k, t)^2$, so that [49]

$$E_{p'p'}(k, t) = Ko_p' \bar{\rho}^2 \eta(t)^{4/3} k^{-5}, \quad k_f \ll k \ll k_\eta(t), \quad (31)$$

$$E_{p'p'}(k, t) = Ko_p'' \bar{\rho}^2 \epsilon(t)^{4/3} k^{-7/3}, \quad k \ll k_f, \quad (32)$$

in the enstrophy cascade and inverse kinetic energy cascade subranges, respectively, where Ko_p' and Ko_p'' are pressure Kolmogorov constants.

The density variance spectrum $E_{\rho'\rho'}(k)$ is shown following reshock at 7 ms and at late time (18 ms) in Figs. 15 and 16, respectively. The density variance spectrum shows an approximate $k^{-5/3}$ scaling for intermediate-to-large k , which is further verified by the compensated spectrum. The pressure variance spectrum $E_{p'p'}(k)$ in Figs. 17 and 18 shows a scaling of $k^{-7/3}$ over a slightly broader range of scales than the density variance spectrum at late time, which is also verified by the compensated spectrum. A k^{-5} decay of $E_{p'p'}(k)$ at large k is inconclusive. The $k^{-5/3}$ scaling [Eq. (30)] for the density variance spectrum and the $k^{-7/3}$ scaling [Eq. (31)] for the pressure variance spectrum *may* be approximately valid in the two-dimensional case considered here. Note that the apparent scalings of $E_{p'p'}(k, t)$ are shifted to larger k than the short apparent inverse kinetic energy and enstrophy cascade subranges exhibited by $E(k, t)$.

3. Statistics

Statistics within the mixing layer were obtained by integrating the spectra over all wave numbers. The time-evolution of the streamwise, spanwise, and total fluctuating kinetic energy is shown in Fig. 19. Initially, $E_{v''v''}(t) < E_{u''u''}(t)$ as the initial shock primarily excites modes in the streamwise direction. The kinetic energy decreases following the initial shock passage, while reshock deposits additional vorticity on the interface and amplifies the kinetic energy. Reshock and the subsequent reflected wave interactions also excite spanwise velocity fluctuations, as can also be seen by considering the time-evolution of the ratio $E_{u''u''}(t)/E_{v''v''}(t)$, also shown in Fig. 19. On average, this ratio is $\gtrsim 3$, with a peak value ≈ 20 at reshock. At 18 ms this ratio decreases to ≈ 1.3 , showing that the contributions from the spanwise and streamwise fluctuations become comparable at late times, and indicating an approach of the flow to *statistical (but not spectral) isotropy at late times*.

The fluctuating enstrophy $E_{\omega''\omega''}(t)$ in Fig. 20 increases significantly during reshock. Following reshock, $E_{\omega''\omega''}(t)$ decays, but the arrival of the reflected waves compensates for the decrease, and a nearly constant value is observed for 8–11 ms. Finally, after the interaction of the reflected rarefaction with the mixing layer at ≈ 11 ms, $E_{\omega''\omega''}(t)$ begins to decay once again.

The density variance $E_{\rho'\rho'}(t)$ in Fig. 20 is not as significantly affected by reshock or by other reflected waves as the other statistics, and remains nearly constant in time. The pressure variance $E_{p'p'}(t)$ is much more sensitive, exhibiting peaks as waves interact with the mixing layer. The first peak at ≈ 6.4 ms corresponds to the arrival of the reflected shock; the second peak at ≈ 11 ms corresponds to the arrival of the reflected rarefaction, and; the third peak at ≈ 15 ms corresponds to the ar-




FIG. 12: (Color). The time-evolution of the total, streamwise, and spanwise fluctuating kinetic energy spectra $E(k, t)$ (top row), $E_{u''u''}(k, t)$ (middle row), and $E_{v''v''}(k, t)$ (bottom row), respectively, at 1 ms intervals (in units of cm^3/s^2). The spectra before reshock are shown in the first column and the spectra after reshock are shown in the second and third columns.




FIG. 13: (Color). The time-evolution of the ratio of the streamwise and spanwise fluctuating kinetic energy spectra $E_{u''u''}(k, t)/E_{v''v''}(k, t)$ at 1 ms intervals. The spectra before reshock are shown in the first column and the spectra after reshock are shown in the second and third columns.




FIG. 14: (Color). The time-evolution of the fluctuating enstrophy spectrum $E_{\omega''\omega''}(k, t)$ (in units of cm^2/s^2), density variance spectrum $E_{\rho'\rho'}(k, t)$ (in units of g^2/cm^5), pressure variance spectrum $E_{p'p'}(k, t)$ (in units of bar^2cm), and baroclinic vorticity production variance spectrum $E_{\mathcal{P}\mathcal{P}}(k, t)$ (in units of cm/s^4) at 1 ms intervals. The spectra before reshock are shown in the first column and the spectra after reshock are shown in the second and third columns.




FIG. 15: The fluctuating kinetic energy spectrum $E(k)$ and compensated spectrum $k^n E(k)$ shortly after reshock at 7 ms showing a $k^{-5/3}$ decay for small k with $Ko''\epsilon^{2/3} = 1.0987 \times 10^6 \text{ cm}^2/\text{s}^3$ and an approximate k^{-3} decay for large k with $Ko'\eta^{2/3} = 7.6912 \times 10^7 \text{ s}^{-2}$ (left column). The fluctuating entrophy spectrum $E_{\omega''\omega''}(k)$ showing an approximate $k^{1/3}$ growth for small k with $Ko''\epsilon^{2/3} = 2.7469 \times 10^6 \text{ cm}^2/\text{s}^3$ and an approximate k^{-1} decay for large k with $Ko'\eta^{2/3} = 2.4722 \times 10^7 \text{ s}^{-2}$ (middle column). The density variance spectrum $E_{\rho'\rho'}(k)$ showing an approximate $k^{-5/3}$ decay with $Ba\epsilon_{\rho'\rho'}/\epsilon^{1/3} = 1.011 \times 10^{-6} \text{ g}^2/\text{cm}^{20/3}$ (right column).

rival of the reflected compression wave generated by the refraction of the rarefaction. *The pressure variance is the only statistic considered that captures the effect of this third interaction.* With the exception of $E_{\rho'\rho'}(t)$, all of the other statistics decrease sharply following reshock. With the reflecting boundary condition at the end wall, the simulation does not achieve a purely-decaying state at late times: later in Sec. VI E, the reflecting boundary condition is changed to outflow at the end wall in order to remove the effects of reflected waves following reshock.

The baroclinic vorticity production variance $E_{\mathcal{P}\mathcal{P}}(t)$ is also shown in Fig. 20. Prior to reshock, $E_{\mathcal{P}\mathcal{P}}(t)$ increases between 3.5 and 4 ms, corresponding to the vortex roll-up. As the roll-up develops, $E_{\mathcal{P}\mathcal{P}}(t)$ is large as a result of the density gradient between the two gases, and later decreases as the gases mix. Reshock sharply increases the production, followed by a rapid decay as the shock passes through the interface. At ≈ 11 ms, another increase corresponding to the arrival of the reflected rarefaction is observed, followed by a decay; $E_{\mathcal{P}\mathcal{P}}(t)$ does not increase upon the arrival of the reflected compression wave.

D. Comparison of mixing quantities and spectra immediately after and before reshock

As in the case of the interaction of a shock with a turbulent flow [50–52], reshock of the evolving Richtmyer-Meshkov mixing layer amplifies quantities characterizing turbulence and mixing. This amplification is investigated here by comparing quantities at $t = t_r^- = 6.38$ and $t = t_r^+ = 6.58$ ms, immediately prior to and following reshock, respectively. The mole fraction profile \bar{X} , product mole fraction profile \bar{X}_p , and the mixing fraction profile θ prior to and following reshock are shown in Fig. 21 as a function of $[x - x_{\text{mid}}(t)]/h(t)$. The mole fraction profile shows that reshock compresses the interface, shifting mass from the SF_6 side to the air(acetone) side. The product mole fraction profile shows that reshock accelerates mixing at the diffuse interfaces, including in the cores of the vortex roll-ups. This results in increased overall mixing, as shown by the increase in θ .

The fluctuating kinetic energy spectrum $E(k, t)$ and its streamwise and spanwise components, $E_{u''u''}(k, t)$ and $E_{v''v''}(k, t)$, are shown in Fig. 22. The ratio of the spectra after and before reshock quantifies the amplification in energy as a function of wave number k . The spectra are all amplified most at the intermediate and small scales. Reshock sharply increases the modal energy content: the increase in the total and streamwise spectra is nearly




FIG. 16: The fluctuating kinetic energy spectrum $E(k)$ and compensated spectrum $k^n E(k)$ at 18 ms showing a $k^{-5/3}$ decay for small k with $Ko''\epsilon^{2/3} = 2.4546 \times 10^5 \text{ cm}^2/\text{s}^3$ and an approximate k^{-3} decay for large k with $Ko'\eta^{2/3} = 9.9937 \times 10^6 \text{ s}^{-2}$ (left column). The fluctuating enstrophy spectrum $E_{\omega''\omega''}(k)$ showing an approximate $k^{-2/3}$ decay for small k with $Ko''\epsilon^{2/3} = 1.2376 \times 10^6 \text{ cm}^2/\text{s}^3$ and an approximate k^{-2} decay for large k with $Ko'\eta^{2/3} = 1.7533 \times 10^8 \text{ s}^{-2}$ (middle column). The density variance spectrum $E_{\rho'\rho'}(k)$ showing an approximate $k^{-5/3}$ decay with $Ba\epsilon_{\rho'\rho'}/\epsilon^{1/3} = 9.3085 \times 10^{-7} \text{ g}^2/\text{cm}^{20/3}$ (right column).

uniform for small k , with an increase for intermediate-to-large k , as indicated by the ratio after and before reshock (also shown in Fig. 22). The spanwise component $E_{v''v''}(k, t)$ increases nearly uniformly from small to large k , as indicated by the increasing ratio. Reshock primarily excites fluctuations in the spanwise direction, as the ratio is considerably larger for $E_{v''v''}(k, t)$ than for $E_{u''u''}(k, t)$. The ratio of the streamwise and spanwise fluctuating kinetic energy spectra, $E_{u''u''}(k, t)/E_{v''v''}(k, t)$, is shown in Fig. 23 after and before reshock: this ratio decreases with k following reshock. The ratio is of $O(1)$ at intermediate k ($k > 5 \text{ cm}^{-1}$) and approaches ≈ 0.3 for large k : reshock significantly increases the spanwise fluctuations to a level comparable to the streamwise fluctuations at the largest scales. As mentioned in Sec. VIC 1, this ‘equilibration’ is only transient, as the spanwise component dominates shortly after reshock once inversion occurs.

The fluctuating enstrophy, pressure variance and baroclinic vorticity production variance spectra in Figs. 24 and 25 also sharply increase following reshock. The largest amplification of $E_{\omega''\omega''}(k, t)$ occurs at intermediate to small k . The density variance spectrum $E_{\rho'\rho'}(k, t)$ is not significantly amplified compared to the other spectra, and shows an average amplification of ≈ 3 . By contrast, the pressure variance spectrum undergoes the most amplification, having a peak of $\approx 1.4 \times 10^7$ for $k \approx 110 \text{ cm}^{-1}$. The baroclinic vorticity production variance spec-

trum also shows an amplification of $\approx 8 \times 10^5$ for small k , with the amplification decreasing at larger k .

E. The effects of reflected waves on the mixing dynamics

Following reshock, the mixing layer interacts with the rarefaction formed during reshock at $\approx 11 \text{ ms}$. This interaction causes the formation of a compression wave, which subsequently interacts with the layer at $\approx 15 \text{ ms}$. Here, the boundary condition at the right end of the computational domain (corresponding to the end wall of the test section) is changed from reflecting to outflow immediately following reshock to allow the reflected rarefaction to exit the computational domain, so that no further interactions of waves with the layer occur. The purpose of this is to investigate the properties of mixing in the *decay regime*, as distinguished from those in the *quasi-decay regime* occurring when reflected waves interact with the evolving interface following reshock. Another objective is to understand the effects of a reflected rarefaction on the evolution of a mixing layer in the same spirit that the study of reshock investigates the effects of a shock wave on an evolving mixing layer. The results presented are from simulations with resolution of 256 points per initial perturbation wavelength (the WENO9M simulation




FIG. 17: The pressure variance spectrum $E_{p'p'}(k)$ (top) and the compensated spectrum $k^n E_{p'p'}(k)$ (bottom) at 7 ms showing an approximate $k^{-7/3}$ decay for small k with $Ko_p'' \bar{\rho}^2 \epsilon^{4/3} = 5.6737 \times 10^9 \text{ g}^2/(\text{cm}^{10/3} \text{ s}^4)$ and a very short k^{-5} decay for large k with $Ko_p' \bar{\rho}^2 \eta^{4/3} = 2.8368 \times 10^{15} \text{ g}^2/(\text{cm}^{10/3} \text{ s}^4)$.




FIG. 18: The pressure variance spectrum $E_{p'p'}(k)$ (top) and the compensated spectrum $k^n E_{p'p'}(k)$ (bottom) at 18 ms showing an approximate $k^{-7/3}$ decay for small k with $Ko_p'' \bar{\rho}^2 \epsilon^{4/3} = 2.4885 \times 10^8 \text{ g}^2/(\text{cm}^{10/3} \text{ s}^4)$ and a very short k^{-5} decay for large k with $Ko_p' \bar{\rho}^2 \eta^{4/3} = 3.9598 \times 10^{13} \text{ g}^2/(\text{cm}^{10/3} \text{ s}^4)$.




FIG. 19: The time-evolution of the fluctuating kinetic energy $E(t)$ (solid line) and its streamwise and spanwise components $E_{u''u''}(t)$ and $E_{v''v''}(t)$ (dashed and dot-dashed lines, respectively) (in units of cm^2/s^2) (top), and the time-evolution of the ratio of the streamwise and spanwise component of the fluctuating kinetic energy $E_{u''u''}(t)/E_{v''v''}(t)$ (bottom).

in Ref. 9).

A comparison of the mixing layer width, production fractions and mixing fractions is shown in Fig. 26 for reflecting and outflow boundary conditions. As expected, the quantities overlap from the time of reshock ≈ 6.4 ms until ≈ 11 ms when the reflected rarefaction interacts with the mixing layer. Generally, quantities computed with the outflow boundary condition are smoother compared to those computed with the reflecting boundary condition, as a consequence of the absence of wave interactions with the layer.

A comparison of the spectra obtained with the outflow and reflecting boundary conditions is shown in Figs. 27–30 at 12 ms, following the arrival of the reflected rarefaction wave and at 18 ms. When ratios are shown, the quantity computed from the reflecting case is divided by the corresponding quantity from the outflow case. The reflected rarefaction has a modest effect on the fluctuating kinetic energy (and its streamwise and spanwise components), fluctuating enstrophy, and pressure and density variance spectra. The reflected rarefaction contributes to approximately a factor of two difference in the energy content, which is significantly less than the corresponding contribution by energy deposition during reshock. However, the effects on the ratio of the spanwise and streamwise components in Fig. 28 are significant, indicating




FIG. 20: The time-evolution of the fluctuating enstrophy $E_{\omega''\omega''}(t)$ (in units of s^{-2}), density variance $E_{\rho'\rho'}(t)$ (in units of g^2/cm^6), pressure variance $E_{p'p'}(t)$ (in units of bar^2), and baroclinic vorticity production variance $E_{PP}(t)$ (in units of s^{-4}).




FIG. 21: Comparison of the mole fraction and production fraction profiles \bar{X} and \bar{X}_p , and of the mixing fraction θ before reshock at $t_r^+ = 6.38$ ms (solid line) and after reshock at $t_r^+ = 6.58$ ms (dashed line).

that the reflected rarefaction excites spanwise velocity fluctuations and contributes to the late-time isotropization of the velocity field. The reflected rarefaction increases $E_{PP}(k, t)$ in Fig. 30 by an order of magnitude.

A comparison of the evolution of the statistics obtained with reflecting and outflow boundary conditions is shown in Figs. 31 and 32. The fluctuating kinetic energy per unit mass $E(t)$ and its components $E_{u''u''}(t)$ and $E_{v''v''}(t)$ are shown in Fig. 31. In the outflow case, the energy is not increased by the reflected waves, and therefore decays. The reflected rarefaction increases the spanwise energy, contributing to the statistical isotropization of the flow. In the outflow case, both components retain

their separation, as shown by the ratio of the components in Fig. 31. In particular, statistical isotropy is approached faster at late time in the reflecting than in the outflow case.

The evolution of the fluctuating enstrophy $E_{\omega''\omega''}(t)$ in Fig. 32 is very similar for the reflecting and outflow cases, indicating that the reflected waves do not increase $E_{\omega''\omega''}(t)$ significantly. However, $E_{\rho'\rho'}(t)$ corresponding to the outflow case is *greater* than $E_{\rho'\rho'}(t)$ corresponding to the reflecting case, indicating that the reflected rarefaction decreases the density fluctuations. The pressure variance $E_{p'p'}(t)$ indicates that the pressure fluctuations are nearly constant in the absence of reflected waves. By




FIG. 22: Comparison of the fluctuating kinetic energy spectrum $E(k, t)$ and its streamwise and spanwise components, $E_{u''u''}(k, t)$ and $E_{v''v''}(k, t)$, respectively (in units of cm^3/s^2) before reshock at $t_r^- = 6.38$ ms and after reshock at $t_r^+ = 6.58$ ms (top row). The ratios of these spectra after and before reshock are also shown (bottom row).




FIG. 23: Comparison of the ratio of the streamwise and spanwise fluctuating kinetic energy spectrum $E_{u''u''}(k, t)/E_{v''v''}(k, t)$ before reshock at $t_r^- = 6.38$ ms and after reshock at $t_r^+ = 6.58$ ms (top). The ratio $[E_{u''u''}(k, t_r^+)/E_{v''v''}(k, t_r^+)]/[E_{u''u''}(k, t_r^-)/E_{v''v''}(k, t_r^-)]$ after and before reshock is also shown (bottom).

contrast, the fluctuations in the reflecting case undergo a sudden transient increase following the interaction with reflected waves, and approach the nearly constant value corresponding to the outflow case shortly thereafter. The baroclinic vorticity production variance, $E_{\mathcal{P}\mathcal{P}}(t)$, decays in the absence of the reflected rarefaction.

VII. DISCUSSION AND CONCLUSIONS

Presented in this study is a comprehensive investigation of the physics and mixing properties before and after reshock, obtained from two-dimensional ninth-order WENO simulations using initial and boundary conditions and flow geometry modeling the test section in the Mach 1.21 Collins and Jacobs air(acetone)/SF₆ shock tube experiment. The details of the simulations are discussed in Ref. 2, which considered the growth of the perturbation amplitude into the nonlinear regime prior to reshock. The dependence of the quantities considered here on the order of WENO flux reconstruction and grid resolution was considered in Ref. 9.

A. Phenomenology of reshock

Density fields were used to visualize reshock, including the inversion process and the formation of complex disordered structures. The vorticity and baroclinic vorticity production fields were also used to investigate vorticity




FIG. 24: Comparison of the fluctuating entrophy spectrum $E_{\omega''\omega''}(k, t)$ (in units of cm^2/s^2), density variance spectrum $E_{\rho'\rho'}(k, t)$ (in units of g^2/cm^5), and pressure variance spectrum $E_{p'p'}(k, t)$ (in units of bar^2cm) before reshock at $t_r^- = 6.38$ ms and after reshock at $t_r^+ = 6.58$ ms (top row). The ratios of the spectra after and before reshock are also shown (bottom row).




FIG. 25: Comparison of the baroclinic vorticity production variance spectrum $E_{ppp}(k, t)$ (in units of cm^4/s^4) before reshock at $t_r^- = 6.38$ ms and after reshock at $t_r^+ = 6.58$ ms (top). The ratio of the spectra after and before reshock is also shown (bottom).

production during reshock, and the deposition of vorticity of opposite sign on the interface, which drives the inversion process. Simulated density Schlieren fields were used to visualize the complex wave interactions occurring during reshock, i.e., the reflected and transmitted waves, and the focusing effects. The following qualitative picture emerges for the roll-up dynamics: the misalignment of the density and vorticity gradient during the roll-up causes vorticity deposition of opposite sign on the roll-up. This vorticity is responsible for the formation of large-scale structures as the roll-up develops, resulting in the formation of complex structures.

B. Baroclinic interfacial circulation deposition

To quantify the baroclinic mechanism driving the evolution of the Richtmyer-Meshkov instability, the circulation deposited on the interface by the initial shock was shown to be in agreement with the predictions of the Samtaney-Zabusky model and linear instability theory. The positive and negative circulations increase following the shock interaction, consistent with the physical picture of ‘vortex-accelerated secondary baroclinic vorticity deposition’ [12]. This secondary instability is responsible for the complex structures in the roll-ups and for additional vorticity deposition of opposite sign on the interface, decreasing the circulation. Reshock causes the deposition of additional vorticity of opposite sign, which




FIG. 26: (Color). Comparison of the time-evolution of the mixing layer width $h(t)$, production fractions $P_t(t)$ and $P_m(t)$, and mixing fractions $\Xi(t)$ and $\Theta(t)$ for reflecting and outflow boundary conditions.




FIG. 27: Comparison of the fluctuating kinetic energy spectrum $E(k)$ and its streamwise and spanwise components, $E_{u''u''}(k)$ and $E_{v''v''}(k)$, respectively (in units of cm^3/s^2) at 12 ms (black) and 18 ms (grey) for reflecting and outflow boundary conditions (top row). The ratios of the spectra for the reflecting and outflow cases are also shown (bottom row).

is further amplified by the secondary baroclinic vorticity deposition. Following the interaction with the reflected rarefaction wave, complex structures form and the flow breaks symmetry, resulting in large asymmetry between the positive and negative circulations and a non-zero total circulation.

C. The post-reshock mixing layer width

The mixing layer width immediately following reshock was shown to be consistent with the linear growth predicted by the Mikaelian [19] and Charakhch'yan [21]

reshock models. The Brouillette-Sturtevant model underestimated the growth at all times following reshock. At later times following reshock, the amplitude is affected by waves interacting with the layer.

D. Analysis of mixing and the physics of reshock

Mixing was investigated by considering the time-evolution of the: (1) mole fraction, production fraction, and mixing fraction profiles to quantitatively assess mixing across the layer; (2) spatially-averaged mixing fractions to quantify the efficiency and rate of mixing; (3)




FIG. 28: Comparison of the ratio of the streamwise and spanwise fluctuating kinetic energy spectrum $E_{u''u''}(k)/E_{v''v''}(k)$ at 12 ms (black) and 18 ms (grey) for reflecting and outflow boundary conditions (top). The ratio of the spectra from the reflecting and outflow cases is also shown (bottom).

fluctuating kinetic energy, fluctuating enstrophy, density variance, pressure variance, and baroclinic vorticity production variance spectra to determine the modal distribution of the fluctuations, and; (4) statistics to characterize the time-evolution of energy.

1. Mole and mixing fraction profiles

The mole and mixing fraction profiles quantify mixing along the shock propagation direction. The mole fraction profile measures the evolving redistribution of mass within the mixing layer caused by the creation of bubble- and spike-like structures, and the mixing fraction profile characterizes the efficiency of mixing. Prior to reshock, the main mechanism affecting these profiles is the formation of the single bubble and spike, which roll up at late times creating a region of well-mixed fluid within. Single localized peaks are observed in the mole and mixing fraction profiles prior to reshock. Prior to reshock, the evolution of the mixing fraction profile indicates increased mixing toward the air(acetone) side of the layer, corresponding to the formation of the roll-up with a well-mixed core.

During reshock, the layer is compressed by the passage of the shock [which continues into the air(acetone)] and is “pulled back” on the SF_6 side by the reflected rarefaction. This effect is captured by the mixing fraction profile, which is larger on the air(acetone) side and

smaller on the SF_6 side. Reshock significantly increases the mixing fractions as the mixing layer is compressed.

Following reshock, the inversion process, followed by the late-time formation of small bubbles and spikes, contributes to the formation of highly irregular, topologically-complex structure on the interface. These additional bubbles and spikes elongate the mixing layer, resulting in a nearly uniformly increasing distribution of mass across the layer from the air(acetone) to the SF_6 side. Several smaller roll-ups also form, indicating large regions of locally well-mixed air and SF_6 . The mixing fractions also show that these regions grow larger at late times, corresponding to more complete mixing. This is consistent with the inverse cascade of small-scale velocity fluctuations to larger scales in two-dimensional turbulence. *The profiles indicate a late-time collapse towards self-similarity within the mixing layer.*

2. Production and mixing fractions

The total chemical product, maximum chemical product, and global mixing fractions provide a physical interpretation of the enhanced mixing caused by reshock. Before reshock, the total chemical product increases, indicating an increase in mixing. During reshock, the mixing layer is compressed, inducing additional mixing as measured by the total chemical product. Following reshock, the total chemical product increases rapidly, indicating significantly increased mixing. The mixing fractions are sharply-peaked at reshock: these fractions increase faster than before reshock, indicating that reshock increases the mixing rate.

3. Energy spectra

The spectral evolution prior to and following reshock provides a modal quantification of the complex features in the density, vorticity, and baroclinic vorticity production fields. It was shown that a broad range of scales already exists prior to reshock, indicating that the Richtmyer-Meshkov instability develops non-trivial spectral content from its inception. At reshock, the fluctuations in all fields (except for the density) are amplified across all scales. Reshock does not appear to broaden the range of scales, but primarily imparts energy into structures of all sizes. At late times, numerical dissipation smooths the spectra. When the shock is far away from the layer and the reflected waves become weaker, the spectra decay very slowly.

The decomposition of the fluctuating kinetic energy spectrum into its streamwise and spanwise components indicates that the former is dominant, and that *significant spectral anisotropy in the velocity field exists even at late times*. As the flow is nearly-incompressible at small Mach numbers, density-weighted spectra [53, 54] are not expected to provide different conclusions. Compared to




FIG. 29: Comparison of the fluctuating entrophy spectrum $E_{\omega''\omega''}(k)$ (in units of cm^2/s^2), density variance spectrum $E_{\rho'\rho'}(k)$ (in units of g^2/cm^5), and pressure variance spectrum $E_{p'p'}(k)$ (in units of bar^2cm) at 12 ms (black) and 18 ms (grey) for reflecting and outflow boundary conditions (top row). The ratios of the spectra from the reflecting and outflow cases are also shown (bottom row).




FIG. 30: Comparison of the baroclinic vorticity production variance spectrum $E_{\mathcal{P}\mathcal{P}}(k,t)$ (in units of cm^4/s^4) at 12 ms (black) and 18 ms (grey) for reflecting and outflow boundary conditions (top). The ratio of the spectra from the reflecting and outflow cases is also shown (bottom).

the fluctuating kinetic energy spectrum, the fluctuating entrophy spectrum exhibits a more pronounced cascade toward smaller scales, as expected from two-dimensional turbulence phenomenology. The pressure variance spectrum increases upon reshock and during the arrival of the reflected rarefaction. The density variance spectrum shows that density fluctuations remain small throughout the flow evolution and are not strongly affected by reshock. To our knowledge, *this is the first examination of the fluctuating entrophy, density variance, pressure variance, and baroclinic vorticity production variance spectra in a reshocked Richtmyer-Meshkov mixing layer.*

Possible inertial subrange scalings of the fluctuating kinetic energy, fluctuating entrophy, density variance and pressure variance spectra were also investigated by comparing to the scaling predictions for two-dimensional turbulence. Immediately following reshock, the compensated spectra showed that the turbulence induced by the reshocked Richtmyer-Meshkov instability has a transient behavior in which the energy is transferred both to smaller scales (forward cascade) and to larger scales (inverse cascade), and approximate inertial subranges *may* exist for short wave number extents. However, at late time the fluctuating entrophy spectra exhibit large and intermediate wave number scalings different from those expected from classical two-dimensional isotropic turbulence phenomenology, as a consequence of the transient forcing mechanism in the reshocked Richtmyer-Meshkov




FIG. 31: (Color). Comparison of the time-evolution of the fluctuating kinetic energy $E(t)$ (solid line) and its streamwise and spanwise components $E_{u''u''}(t)$ and $E_{v''v''}(t)$ (dashed and dot-dashed lines, respectively) (in units of cm^2/s^2) for reflecting and outflow boundary conditions (red and blue, respectively) (top). The time-evolution of the ratio of the streamwise and spanwise component of the fluctuating kinetic energy $E_{u''u''}(t)/E_{v''v''}(t)$ is also shown (bottom).

instability. The density variance spectrum may exhibit an inertial-convective subrange scaling expected for a passive scalar, and the pressure variance spectrum may scale as expected in two-dimensional turbulence. However, *an even larger range of scales must be simulated to definitively examine the putative scalings examined here. Reshock is also necessary to attain the intensity of turbulence and to develop the broad range of structures necessary to achieve such scalings.*

4. Statistics

The time-evolution of statistics was also investigated. The fluctuating kinetic energy and enstrophy were sharply peaked at reshock, with a rapid increase followed by a rapid decrease, then followed by a decay. The decay becomes more established following the interaction with the reflected rarefaction. The streamwise and spanwise fluctuating kinetic energies indicate that statistical isotropy is approached at very late times. The spanwise component increases following the interaction with reflected waves and then decreases due to dissipation. The pressure variance showed narrow sharp peaks corre-

sponding to reshock and to the arrival of the reflected rarefaction, and further showed the arrival of the reflected compression wave at late times. These waves excite fluctuations that are apparent in the increased values of the statistics. The fluctuations are damped and the variances decrease in magnitude as time evolves. The density variance is very small in magnitude and remains nearly constant, although signatures of the reflected shock and of the reflected rarefaction are evident.

E. Comparison of mixing quantities and spectra immediately after and before reshock

The comparison of mixing quantities and spectra immediately after and before reshock quantified the effects of reshock on the mixing layer and the energy deposition into the layer by baroclinic production mechanisms. Reshock compresses the layer and shifts mass from the SF_6 side toward the air(acetone) side, consistent with the direction of reshock from the heavier gas into the lighter gas. Reshock further generates structures within the mixing layer, creating a well-mixed region close to the SF_6 and peaks closer to the air(acetone). This is also reflected by the mixing fraction θ .

The energy deposition by reshock was quantified by examining the amplification of the energy spectra. Reshock amplifies the fluctuating kinetic energy spectra most at larger wave numbers (smaller scales), consistent with the formation of small-scale, vortical structures following reshock. The pressure variance is amplified most by reshock, while the density variance is only slightly amplified, indicating that it does not have a significant role in the post-reshock dynamics for the flow considered here. Thus, the fluctuating kinetic energy, fluctuating enstrophy, and pressure variance spectra are useful for the characterization of the modal distribution of energy within the mixing layer following reshock. *The amplifications above are dependent upon the shock Mach number and on other details of the flow configuration.*

F. The effects of reflected waves on mixing

The effects of the reflected rarefaction wave on the mixing layer were also investigated through a comparison of simulations with reflecting and outflow boundary conditions at the end wall of the domain. The reflected rarefaction does not significantly increase the energy content in the layer, but contributes to flow symmetry breaking and driving the flow towards statistical isotropy at late times by exciting fluctuations in the spanwise direction. The reflected rarefaction also has a small effect on the fluctuating enstrophy and pressure variance evolution. By contrast, reshock compresses the layer and causes a significant deposition of energy that induces rapid growth of the layer, concomitant with the formation of complex structures. The reflected rarefaction instead “stretches”




FIG. 32: Comparison of the time-evolution of the fluctuating enstrophy $E_{\omega'\omega''}(t)$ (in units of s^{-2}), density variance $E_{\rho'\rho'}(t)$ (in units of g^2/cm^6), pressure variance $E_{p'p'}(t)$ (in units of bar^2), and baroclinic vorticity production variance $E_{\mathcal{P}\mathcal{P}}(t)$ (in units of s^{-4}) for reflecting and outflow boundary conditions (solid and dashed lines, respectively).

the mixing layer (causing it to grow), but provides little additional energy, so that the overall mixing decreases, as quantified by the mixing fractions. The additional mixing induced by the reflected waves is responsible for a faster approach to late-time statistical isotropy. This comparison further supports the important role that reflected waves have in the post-reshock flow dynamics. The interactions of the waves with the evolving layer can be interpreted as small, transient forcings at all scales. To our knowledge *this is the first study that removed the effects of reflected waves on the post-reshock Richtmyer-Meshkov mixing layer evolution, and showed their important role on the dynamics of the reshocked flow.*

G. Final remarks

The modern shock-capturing WENO method was used here to investigate the physics of reshock and mixing by computing and physically interpreting quantities not presently available from experiments. Specifically, a wide range of quantities that have been used to characterize other mixing layers, including profiles across the layer, mixing fractions, energy spectra, and statistics, were considered here. A comparison of such quantities just after and before reshock quantified the effects of reshock, including the mechanisms of energy deposition. A com-

parison of these quantities for outflow and reflecting boundary conditions quantified the important role that successive wave-interface interactions have on mixing. Thus, the quantities presented here complement those currently available from Richtmyer-Meshkov instability experiments. Additional experimental data pertaining to the post-reshock phase, including quantities depending on spatial derivatives (vorticity or enstrophy) or quantities sensitive to small-scale mixing (mixing fractions), is needed to further validate numerical simulations and constrain models.

Acknowledgments

Marco Latini acknowledges support from the Air Force Office of Scientific Research through the National Defense Science and Engineering Graduate Fellowship and the Caltech Accelerated Strategic Computing (ASC) Center. Wai Sun Don acknowledges support of this work by the DOE under contract number DE-FG02-98ER25346 and the AFOSR under contract number FA9550-05-1-0123. This work was performed under the auspices of the U.S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

- [1] B. D. Collins and J. W. Jacobs, *J. Fluid Mech.* **464**, 113 (2002).
- [2] M. Latini, O. Schilling, and W.-S. Don, *Phys. Fluids* (2007), in press.
- [3] W. D. Arnett, *Astrophys. J.* **319**, 136 (1987).
- [4] J. D. Lindl, *Inertial Confinement Fusion: the Quest for Ignition and Energy Gain Using Indirect Drive* (Springer-Verlag, New York, 1997).
- [5] J. M. Taccetti, S. H. Batha, J. R. Fincke, N. D. Delamater, N. E. Lanier, G. R. Magelssen, R. M. Hueckstaedt, S. D. Rothman, C. J. Horsfield, and K. W. Parker, *Astrophys. Space Sci.* **298**, 327 (2005).
- [6] J. R. Rygg, Ph.D. thesis, Massachusetts Institute of Technology (2006).
- [7] D. L. Youngs, *Laser Part. Beams* **12**, 725 (1994).
- [8] C. Mügler and S. Gauthier, *Phys. Rev. E* **58**, 4548 (1998).
- [9] M. Latini, O. Schilling, and W. S. Don, *J. Comput. Phys.* (2007), in press.
- [10] P. J. Linstrom and W. G. Mallard, *NIST Chemistry Web-Book*, no. 69 in NIST Standard Reference Database (National Institute of Standards and Technology, Gaithersburg, MD, 2005), <http://webbook.nist.gov>.
- [11] N. Themelis, *Transport and Chemical Rate Phenomena* (Taylor and Francis, New York, 1995).
- [12] G. Peng, N. J. Zabusky, and S. Zhang, *Phys. Fluids* **15**, 3730 (2003).
- [13] A. Marquina and P. Mulet, *J. Comput. Phys* **185**, 120 (2003).
- [14] M. Brouillette and B. Sturtevant, *Phys. Fluids A* **8**, 916 (1993).
- [15] R. Samtaney and N. J. Zabusky, *J. Fluid Mech.* **269**, 45 (1994).
- [16] M. C. Soteriou and A. F. Ghoniem, *Phys. Fluids* **7**, 2036 (1995).
- [17] C. Staquet, *J. Fluid Mech.* **296**, 73 (1995).
- [18] J. Reinaud, L. Joly, and P. Chassaing, *Phys. Fluids* **12**, 2489 (2000).
- [19] K. O. Mikaelian, *Physica D* **36**, 343 (1989).
- [20] M. Brouillette and B. Sturtevant, *Physica D* **37**, 248 (1989).
- [21] A. A. Charakhch'yan, *Plasma Phys. Control. Fusion* **43**, 1169 (2001).
- [22] K. I. Read, *Physica D* **12**, 45 (1984).
- [23] D. L. Youngs, *Physica D* **12**, 32 (1984).
- [24] M. Vetter and B. Sturtevant, *Shock Waves* **4**, 247 (1995).
- [25] R. E. Duff, F. H. Harlow, and C. W. Hirt, *Phys. Fluids* **5**, 417 (1962).
- [26] A. W. Cook and P. E. Dimotakis, *J. Fluid Mech.* **443**, 69 (2001), corrigendum, *J. Fluid Mech.* **457**, 410 (2002).
- [27] A. Favre, in *Studies in Turbulence*, edited by T. B. Gatski, S. Sarkar, and C. G. Speziale (Springer-Verlag, New York, 1992), pp. 324–341.
- [28] P. Chassaing, R. A. Antonia, F. Anselmet, L. Joly, and S. Sarkar, *Variable Density Fluid Turbulence*, vol. 69 of *Fluid Mechanics and its Applications* (Kluwer Academic, New York, 2002).
- [29] M. Lesieur, C. Staquet, P. Le Roy, and P. Comte, *J. Fluid Mech.* **192**, 511 (1988).
- [30] C. Mügler and S. Gauthier, *Phys. Fluids* **12**, 1783 (2000).
- [31] S. B. Pope, *Turbulent Flows* (Cambridge University Press, Cambridge, 2000).
- [32] M. M. Koochesfahani and P. E. Dimotakis, *J. Fluid Mech.* **170**, 83 (1986).
- [33] D. L. Youngs, *Phys. Fluids A* **3**, 1312 (1991).
- [34] D. J. Hill, C. Pantano, and D. I. Pullin, *J. Fluid Mech.* **557**, 29 (2006).
- [35] M. R. Meloon, Ph.D. thesis, California Institute of Technology (1997).
- [36] R. H. Kraichnan, *Phys. Fluids* **10**, 1417 (1967).
- [37] G. K. Batchelor, *Phys. Fluids Suppl.* **12**, 233 (1969).
- [38] R. H. Kraichnan and D. Montgomery, *Rep. Prog. Phys.* **43**, 547 (1980).
- [39] M. Lesieur, *Turbulence in Fluids*, vol. 40 of *Fluid Mechanics and its Applications* (Kluwer Academic, New York, 1997), 3rd ed.
- [40] J. Sommeria, in *New Trends in Turbulence (Les Houches Summer School on Theoretical Physics LXXIV, 2000)*, edited by M. Lesieur, A. Yaglom, and F. David (Springer-Verlag, 2001), pp. 385–447.
- [41] P. Tabeling, *Phys. Rep.* **362**, 1 (2002).
- [42] R. H. Kraichnan, *J. Fluid Mech.* **47**, 525 (1971).
- [43] J. W. Jacobs and V. V. Krivets, *Phys. Fluids* **17**, 034104 (2005).
- [44] A. M. Obukhov, *Izv. Akad. Nauk SSSR* **13**, 58 (1949).
- [45] S. Corrsin, *J. Appl. Phys.* **22**, 469 (1951).
- [46] S. Corrsin, *J. Aeronaut. Sci.* **18**, 417 (1951).
- [47] G. K. Batchelor, *J. Fluid Mech.* **5**, 113 (1959).
- [48] A. S. Monin and A. M. Yaglom, *Statistical Fluid Mechanics*, vol. 1 (MIT Press, Cambridge, 1971).
- [49] M. Lesieur, S. Ossia, and O. Métais, *Phys. Fluids* **11**, 1535 (1999).
- [50] Y. Andreopoulos, J. H. Agui, and G. Briassulis, *Ann. Rev. Fluid Mech.* **32**, 309 (2000).
- [51] G. Briassulis, J. H. Agui, and Y. Andreopoulos, *J. Fluid Mech.* **432**, 219 (2001).
- [52] J. H. Agui, G. Briassulis, and Y. Andreopoulos, *J. Fluid Mech.* **524**, 143 (2005).
- [53] S. Kida and S. A. Orszag, *J. Sci. Comput.* **5**, 1 (1990).
- [54] S. Kida and S. A. Orszag, *J. Sci. Comput.* **7**, 1 (1992).