

UCB-PTH-05/37
LBNL-59061
BUHEP-05-16

Tensor Mesons in AdS/QCD

Emanuel Katz^{a1}, Adam Lewandowski^{a,b2}, and Matthew D. Schwartz^{c3}

^a *Department of Physics,
Boston University, Boston, MA 02215, USA*

^b *Department of Physics, United States Naval Academy,
Annapolis, Maryland 21403, USA*

^c *Department of Physics, University of California, Berkeley,
and
Theoretical Physics Group, Lawrence Berkeley National Laboratory,
Berkeley, CA 94720, USA*

Abstract

We explore tensor mesons in AdS/QCD focusing on $f_2(1270)$, the lightest spin-two resonance in QCD. We find that the f_2 mass and the partial width $\Gamma(f_2 \rightarrow \gamma\gamma)$ are in very good agreement with data. In fact, the dimensionless ratio of these two quantities comes out within the current experimental bound. The result for this ratio depends only on N_c and N_f , and the quark and glueball content of the operator responsible for the f_2 ; more importantly, it does not depend on chiral symmetry breaking and so is both independent of much of the arbitrariness of AdS/QCD and completely out of reach of chiral perturbation theory. For comparison, we also explore $f_2 \rightarrow \pi\pi$, which because of its sensitivity to the UV corrections has much more uncertainty. We also calculate the masses of the higher spin resonances on the Regge trajectory of the f_2 , and find they compare favorably with experiment.

¹amikatz@buphy.bu.edu

²lewandow@usna.edu

³mdschwartz@lbl.gov

1 Introduction

It is well known that QCD, of the real world, cannot be studied through the AdS/CFT correspondence. After all, QCD is not a conformal field theory, N_c is not large, the string dual of QCD is a complete mystery, and if there is such a dual, the string scale must be low. One can even make more practical objections, such as that any low energy predictions which do come out cannot be original. They must be the same as the predictions of chiral perturbation theory, since the symmetries are the same. Or at least, since AdS/CFT employs the operator product expansion, it should not be more powerful than QCD sum rules, which should be able to extract as much information as possible out of the OPE. Therefore, why bother with AdS/QCD at all? We are motivated by several reasons. First, with QCD we have experimental results which can help determine the elements of AdS/CFT essential for establishing a predictive duality. Second, after we know what works and what does not, there is the possibility that we may learn something about QCD itself, and perhaps about strongly coupled gauge theories in more generality. Finally, as our framework depends on the holographic map, we are also indirectly exploring this map experimentally.

To elaborate on these motivations, we turn to the topic of the current work: tensor mesons. We will focus on the f_2 , a spin-two isospin singlet meson with mass 1275 MeV, although higher spin mesons will be discussed as well. As we will shortly see (Section 2), not much can be said about tensor mesons in perturbation theory. Even if we couple the f_2 universally, assuming general coordinate invariance (GC), there is an unknown dimensional coupling constant, the analog of the Planck scale for the spin-2 graviton. The decay rates to pions and to kaons can be related by SU(3), but the decay rate to photons has a free NDA factor of order 1. The f_2 mass is another free parameter. What's worse is that the cutoff on the chiral Lagrangian is $\Lambda = 4\pi f_\pi \approx 1200$ MeV, which is below the f_2 mass. So even if chiral symmetry did make predictions, we would not be able to trust them.

In contrast, on the AdS side, the f_2 will be treated as a spin-2 gauge field in the bulk (Section 3). Equivalently, we can think of it as a KK excitation of the graviton. Thus, we secure its interactions by GC. But now the coupling constant can be calculated by matching to the perturbative OPE. The f_2 mass is set by the eigenvalue equation for the KK mode. We define our units by fixing the IR cutoff with the ρ mass. This leads to the simple formula that the ratio of the f_2 mass to the ρ mass is simply given by the ration of the zeros of Bessel functions \mathcal{J}_1 and \mathcal{J}_0 . Thus

$$\frac{m_{f_2}}{m_\rho} = \frac{\text{zero of } \mathcal{J}_1}{\text{zero of } \mathcal{J}_0} = 1.59 \quad (\text{AdS/QCD}) \quad (1)$$

The experimental value for this ratio is 1.64, a difference of 3% . Remarkably, mass predictions can be extended up the Regge trajectory for higher spin mesons, without a noticeable loss of accuracy (Section 5).

The coupling to photons is also fixed in this theory – it is determined by the

overlap of the flat photon wavefunction with that of the f_2 . Thus we can compute

$$\Gamma(f_2 \rightarrow \gamma\gamma) = 2.71 \text{ keV} \quad (\text{AdS/QCD}) \quad (2)$$

The observed width is 2.60 ± 0.24 keV, so we get this rate correct to within experimental error. These two calculations indicate that, for some reason, AdS/QCD is more accurate than we have any right to expect. Because the decay to photons is sensitive to the admixture of the tensor glueball operator, we actually obtain some non-trivial information about the f_2 from this calculation. This is discussed in section 3.3.

Not everything in the tensor sector works so well. We also calculate the rate for f_2 decaying to pions, still without introducing any new free parameters, and find it too small by a factor of 4. This is more in line with our naive expectations. It indicates that either some of our assumptions, such as the those about boundary conditions, should be reconsidered, or that higher dimension operators are relevant. In fact, we know roughly where the string scale is on the AdS side. Some of the tensor modes, such as $a_2(1320)$, must be string states, because they carry isospin. So we are at least very near the regime where string corrections become important. Gravity corrections are also important because we know the effective Planck scale for the f_2 , and it is also low. So it is not surprising that our calculations can receive stringy corrections. In Section 4 we show that higher dimension bulk operators are very relevant to the f_2 to pion decays. More importantly, we also show that higher dimension bulk operators are *not* relevant for the decay to photons, so we have more reason to trust our $f_2 \rightarrow \gamma\gamma$ calculation.

2 Introducing the f_2

Before we describe the AdS/QCD construction, we review and elaborate on what is known about the f_2 from other methods. In particular, we discuss chiral perturbation theory, which allows us to couple the f_2 to pions. We will also establish notation and present some formulae which will be used later on.

2.1 Chiral Perturbation Theory

In chiral perturbation theory, the pions are Goldstone bosons for a spontaneously broken $SU(N_f)_L \times SU(N_f)_R$ global symmetry [1]. They are embedded in a matrix π which is exponentiated to get

$$U = e^{2i\pi/f_\pi} \quad (3)$$

This matrix U transforms under $SU(N_f)_L \times SU(N_f)_R$

$$U \rightarrow g_L U g_R^\dagger \quad (4)$$

and the non-linear transformations of the pions follow. The lowest order chiral Lagrangian (with massless quarks) is

$$\mathcal{L}^{(2)} = \frac{f_\pi^2}{4} \text{Tr}[D_\mu U^\dagger D^\mu U] = \frac{1}{2} (\partial_\mu \pi^a) (\partial_\mu \pi^a) + \dots \quad (5)$$

To next order, there are additional terms [37]

$$\begin{aligned} \mathcal{L}^{(4)} = & L_1 \text{Tr}[D_\mu U^\dagger D^\mu U]^2 + L_2 \text{Tr}[D_\mu U^\dagger D_\nu U] \text{Tr}[D^\mu U^\dagger D^\nu U] + L_3 \text{Tr}[D_\mu U^\dagger D^\mu U D_\nu U^\dagger D^\nu U] \\ & + iL_9 \text{Tr}[F_R^{\mu\nu} D_\mu U D_\nu U^\dagger + F_L^{\mu\nu} D_\mu U^\dagger D_\nu U] + L_{10} \text{Tr}[U^\dagger F_R^{\mu\nu} U F_{\mu\nu}^L] \end{aligned} \quad (6)$$

where $F_{R/L}^{\mu\nu}$ are the field strengths for external right- and left-handed gauge fields. At this point, all the L_i 's and f_π are unknown parameters to be fit to data. Often, assumptions additional to chiral symmetry can predict some of these L_i 's. For example, vector meson dominance assumes that the bare L_i are all zero, and the observed L_i come from integrating out heavy mesons, in particular the ρ meson [26–28]. This predicts some relations between the L_i . Another example is AdS/QCD which predicts similar, but different, relations [3, 4, 10]. The L_i are fairly well known, and are therefore handy for distinguishing different theories.

To add the f_2 , we introduce, by hand, a massive spin-2 meson. Its kinetic term should be of the Fierz-Pauli form [24]

$$\mathcal{L}_{\text{kin}}^{(f)} = \frac{1}{2} h_{\mu\nu} \square h_{\mu\nu} + h_{\mu\alpha, \alpha}^2 - h_{\mu\nu, \mu} h_{, \mu} + \frac{1}{2} h_{, \mu}^2 + \frac{1}{2} m_f^2 (h_{\mu\nu}^2 - h^2) \quad (7)$$

where $h = h_{\mu\mu}$. Then the most general set of interactions with the chiral Lagrangian begins

$$\mathcal{L}_{\text{int}} = c_1 h_{\mu\nu} \text{Tr}[D_\mu U D_\nu U^\dagger] + c_2 h \text{Tr}[D_\mu U D_\mu U^\dagger] + c_3 h_{\mu\nu} \text{Tr}[U^\dagger F_{\mu\alpha}^R U F_{\alpha\nu}^L] + \dots \quad (8)$$

By chiral symmetry alone, the c_i , like the L_i , are completely undetermined.

To progress, we can assume that the f_2 couples universally to the energy momentum tensor for matter, like in general relativity. And, like in general relativity, this would let us bootstrap the f_2 's self-couplings as well, at least in the absence of an f_2 mass. But because of the mass, the final Lagrangian will have no exact symmetry lacking in the general expression (8). Nevertheless, the assumption that the f_2 couples to the energy momentum tensor of matter is consistent within effective field theory, as it parametrically raises the scale at which the f_2 interactions become strong [35, 36].

Adding an f_2 coupling strength G_f (if f_2 were a graviton, $G_f \approx M_P^{-1}$) to the Lagrangian, we have

$$\mathcal{L} = \mathcal{L}^{(2)} + \mathcal{L}_{\text{kin}}^{(f)} + \frac{1}{2} G_f h_{\mu\nu} \Theta_{\mu\nu}^{(2)} + \dots \quad (9)$$

where $\Theta_{\mu\nu}^{(2)}$ is the energy momentum tensor following from $\mathcal{L}^{(2)}$

$$\Theta_{\mu\nu}^{(2)} = \frac{f_\pi^2}{4} [\eta_{\mu\nu}(D_\kappa U)(D^\kappa U^\dagger) - 2(D_\mu U)(D_\nu U^\dagger)] \quad (10)$$

This is of the form $\mathcal{L}_{\text{int}}^{(f)}$ with $c_1 = -2c_2$. We can integrate out the f_2 now, recalling that because of the Fierz-Pauli mass its propagator is [25, 29]

$$\langle h_{\mu\nu} h_{\rho\sigma} \rangle = \frac{P_{\mu\nu\rho\sigma}}{p^2 - m_f^2} \quad (11)$$

where the projector, in the particle's rest frame, has the form

$$P_{\mu\nu\rho\sigma} = \frac{1}{2}(\eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho} - \frac{2}{3}\eta_{\mu\nu}\eta_{\rho\sigma}) \quad (12)$$

(In the massless case, or if the mass had just the $h_{\mu\nu}^2$ term, the residue would have a 1 instead of a 2/3). Then integrating out the f_2 gives an $\mathcal{L}^{(4)}$ term with coefficients

$$L_1^f = -\frac{G_f^2 f_\pi^4}{96m_f^2} \quad \text{and} \quad L_2 = -3L_1 \quad (13)$$

Of course, these are not predictions, as the L_i can have contributions from integrating out other fields as well, or simply from adding a bare term.

We will find in the next paragraph, that from the f_2 decay rate into pions we can set $G_f = g_{f\pi\pi} = 0.019 \text{ MeV}^{-1}$. This leads to, $L_1 = -0.17 \times 10^{-3}$ and $L_2 = 0.50 \times 10^{-3}$. These are fairly substantial contributions to the experimental values of $L_1 = 0.4 \pm 0.3 \times 10^{-3}$ and $L_2 = 1.4 \pm 0.4 \times 10^{-3}$. Of course, this tells us nothing about the f_2 , as we must make assumptions about what else contributes to the L_i , which are just as likely to be wrong as our universality hypothesis for the f_2 couplings.

Now let us turn to this the f_2 decays. The f_2 decays predominantly into pions. The minimal couplings can be written as

$$\mathcal{L}_{f\pi\pi} = \frac{1}{2}g_{f\pi\pi}h_{\mu\nu}\Theta_{\mu\nu}^\pi, \quad \Theta_{\mu\nu}^\pi = \frac{1}{2}\eta_{\mu\nu}(\partial_\alpha\pi)^2 - (\partial_\mu\pi)(\partial_\nu\pi) \quad (14)$$

then the decay rate is [9, 12, 18]

$$\Gamma(f \rightarrow \pi^+\pi^-, \pi^0\pi^0) = g_{f\pi\pi}^2 \frac{m_f^3}{1280\pi} \left(1 - 4\frac{m_\pi^2}{m_f^2}\right)^{5/2} = g_{f\pi\pi}^2 (4.56 \times 10^5 \text{ MeV}^3) \quad (15)$$

where we have used $m_f = 1275.4 \text{ MeV}$ and $m_\pi = 139.5 \text{ MeV}$. We are also interested in the decay into photons, which is well measured also. If we define the coupling as

$$\mathcal{L}_{f\gamma\gamma} = \frac{1}{2}g_{f\gamma\gamma}h_{\mu\nu}\Theta_{\mu\nu}^\gamma, \quad \Theta_{\mu\nu}^\gamma = \frac{1}{4}\eta_{\mu\nu}F_{\alpha\beta}^2 - F_{\nu\alpha}F_{\alpha\mu} \quad (16)$$

then the formula for this decay rate is [9, 12]

$$\Gamma(f_2 \rightarrow \gamma\gamma) = g_{f\gamma\gamma}^2 \frac{m_f^3}{320\pi} = g_{f\gamma\gamma}^2 (2.04 \times 10^6 \text{MeV}^3) \quad (17)$$

The observed decay rates are $\Gamma(f \rightarrow \pi\pi) = 156 \text{ MeV}$ and $\Gamma(f_2 \rightarrow \gamma\gamma) = 2.6 \text{ keV}$ which imply $g_{f\pi\pi} = 0.0185 \text{ MeV}^{-1}$ and $g_{f\gamma\gamma} = 3.55 \times 10^{-5} \text{ MeV}^{-1}$. A helpful way to compare theory and experiment is through the ratio of decay rates with the phase space factored out

$$\frac{g_{f\pi\pi}^2}{g_{f\gamma\gamma}^2} = 2.71 \times 10^5 \quad (\text{Exp}) \quad (18)$$

Assuming strict universality of the f_2 couplings would lead to $g_{f\pi\pi} = g_{f\gamma\gamma} = G_f$, and get this ratio wrong by six orders of magnitude.

To get a better prediction, we must use the fact that the photon is really external to QCD, and so its coupling must be suppressed by powers of the weak coupling constant $e/4\pi$. Naive dimensional analysis [38] instructs us to take $g_{f\gamma\gamma} = c_{\text{NDA}} e^2 / 16\pi^2 G_f$, for some constant c_{NDA} of order one. Then

$$\frac{g_{f\pi\pi}^2}{g_{f\gamma\gamma}^2} = \frac{1}{c_{\text{NDA}}^2} \left(\frac{16\pi^2}{e^2} \right)^2 = \frac{1}{c_{\text{NDA}}^2} 2.96 \times 10^6 \quad (\text{NDA}) \quad (19)$$

This is a fairly good estimate. We see that $c_{\text{NDA}} = 3.31$. But we cannot get c_{NDA} by NDA or by assuming universality of the f_2 couplings (which would say $c_{\text{NDA}} = 16\pi^2/e^2$).

One way to estimate c_{NDA} has been proposed in [12]. Similar to the assumption in vector meson dominance, it declares that the decay to photons is mediated purely by ρ exchange. That is, f decays into two ρ 's which decay into photons. So the bare coupling $g_{f\gamma\gamma}$ is zero. This assumption can be called tensor meson dominance (TMD) [12, 18, 23]. Taking $g_{f\pi\pi}$ from experiment, it gives

$$\frac{g_{f\pi\pi}^2}{g_{f\gamma\gamma}^2} = 3.6 \times 10^6 \quad (\text{TMD}) \quad (20)$$

This is not better than naive dimensional analysis with $c_{\text{NDA}} = 1$.

Tensor meson dominance also relates the f_2 decay constant f_f to $g_{f\pi\pi}$. The f_2 decay constant is defined by

$$\langle \Theta_{\mu\nu} | f_2 \rangle = f_f m_f^3 \epsilon_{\mu\nu} \quad (21)$$

where $\epsilon_{\mu\nu}$ is the polarization tensor for the f_2 and $\Theta_{\mu\nu}$ is the energy momentum tensor. We have chosen f_f to be dimensionless, and so it is off by factors of m_f from what is often called g_f in the literature. If we assume, via TMD, that the pion tensor

form factors are determined by f_2 exchange, even as $p \rightarrow 0$, then the normalization of the pion's energy determines

$$f_f = \frac{1}{m_f g_{f\pi\pi}} = 0.042 \quad (\text{TMD}) \quad (22)$$

We cannot compare f_f to experiment, but we will compare this value to predictions from AdS/QCD below.

In summary, we wrote down the most general Lagrangian involving a spin-2 field, the f_2 , which respects chiral symmetry. We assumed the f_2 couples to the energy momentum tensor of the various matter fields, like the graviton, even though the f_2 is massive. Its mass, its coupling constant, and an order one factor in its coupling to photons are all free parameters. We can fit these to the observed mass, decay rate to pions, and decay rate to photons, but then there are no predictions. Instead, we can assume tensor meson dominance *ad hoc*, which eliminates one parameter, and makes a prediction which turns out not to much better than naive dimensional analysis.

3 AdS/QCD

Now we turn to the predictions from AdS/QCD. We will mostly follow the notation of [3], although [2] presents an equivalent formulation. Our review will be quick, and we refer the reader to these two papers, or to the original AdS/CFT literature [5–8] for more details.

The basic premise of AdS/CFT is that there is a duality between a 4D conformal field theory and a 5D gravity theory on a curved AdS background. Position in the extra dimension corresponds to energy in the 4D theory. We will use conformal coordinates, with the curvature scale normalized to 1, so the metric is

$$ds^2 = \frac{1}{z^2}(dx_\mu^2 - dz^2) \quad (23)$$

Energy independence in the CFT corresponds to an AdS isometry which shifts z . Small z is the high energy UV region, and large z is the IR. Though approximately conformal in the UV (in the sense that $\beta \sim g^3 \rightarrow 0$), QCD becomes strongly coupled, and breaks conformality in IR. We avoid this awkward region by explicitly cutting off the space at $z = z_m$, where z_m (which will be of order $1/\Lambda_{QCD}$) is to be fit to data. We thus implicitly assume that asymptotic freedom sets in very quickly as we go the UV in QCD, motivating our choice of the AdS metric.

3.1 The f_2 mass

Operators in the CFT, or in this case QCD, correspond to fields in the AdS bulk. For example, the ρ , which is a isospin triplet spin-1 meson in QCD, corresponds to

a bulk gauge field V_M in bulk. The action for V_M is just

$$\mathcal{L} = \int d^5x \sqrt{g} \frac{1}{4} F_{MN} F^{MN} = \int d^5z \frac{1}{z} \left(\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2} F_{\mu 5}^2 \right) \quad (24)$$

More precisely, ρ is the first KK excitation of the 4-vector part $V_\mu(x, z)$, while the other KK modes, ρ_n , correspond to heavier resonances with the same quantum numbers. Thus the ρ_n are 4D fields with 5D profiles determined by the vector KK equation in AdS

$$z \partial_z \frac{1}{z} \partial_z V_n(z) + (m_n^V)^2 V_n(z) = 0. \quad (25)$$

The generic solution is

$$V_n(z) = N_n z \left[\mathcal{J}_1(m_n^V z) + \beta_n \mathcal{Y}_1(m_n^V z) \right] \quad (26)$$

We then impose boundary conditions $\rho'(z_m) = \rho(0) = 0$, which sets $\beta_n = 0$ and quantizes the masses to be solutions of $\mathcal{J}_0(m_n^V z_m) = 0$. The normalization is set by $\int dz \rho_n(z)^2 / z = 1$ and so the ρ wavefunction is

$$V(z) = 2.72 \frac{z}{z_m} \mathcal{J}_1\left(2.40 \frac{z}{z_m}\right) \quad (27)$$

We can then use the observed ρ mass of 775 MeV to fix $z_m = (323 \text{ MeV})^{-1}$.

For the f_2 , we need a spin-2 field in the bulk, h_{MN} . The first KK mode of h_{MN} will be the f_2 . Of course, we already have a spin-2 field in the bulk, the graviton. Expanding linearized gravity excitations about the AdS background

$$ds^2 = \frac{1}{z^2} (\eta_{\mu\nu} + h_{\mu\nu}) dx^\mu dx^\nu - \frac{1}{z^2} dz^2 \quad (28)$$

produces a Lagrangian

$$\int d^5x \sqrt{g} R_5 = \int d^5x \frac{1}{2} \frac{1}{z^3} [\partial_z h_{\mu\nu} \partial_z h_{\mu\nu} - h_{\mu\nu} \square h_{\mu\nu} + \dots] \quad (29)$$

But since we are not demanding a fully consistent theory, we could just as well have introduced a different spin-2 field. In fact, we expect from QCD that there are a number of isospin singlet spin-2 particles, including glueballs, and so a complete formulation should include a number of bulk spin-2 fields.

Each field will have a tower of KK modes, whose profiles are determined by the tensor equation of motion

$$z^3 \partial_z \frac{1}{z^3} \partial_z h_n(z) + (m_n^h)^2 h_n(z) = 0. \quad (30)$$

The generic solution is

$$h_n(z) = N_n z^2 \left[\mathcal{J}_2(m_n^h z) + \beta_n \mathcal{Y}_2(m_n^h z) \right] \quad (31)$$

Boundary conditions $h'(z_m) = h(0) = 0$ again put $\beta_n = 0$ and now quantize the masses according to $\mathcal{J}_1(m_n^h z_m) = 0$. For the spin-2 case, the normalization is set by $\int dz h_n^2 / z^3 = 1$. Thus the f_2 profile is

$$h(z) = 3.51 \frac{z^2}{z_m} \mathcal{J}_2\left(3.83 \frac{z}{z_m}\right) \quad (32)$$

Thus we get our first prediction. The f_2 mass is predicted to be $3.83 z_m^{-1} = 1236$ MeV, which, as noted in the introduction is, only 3% off of the observed mass of 1275 MeV. In other words, we have made the z_m independent prediction $m_{f_2}/m_\rho = 1.59$ of equation (1). The experimental central values give

$$\frac{m_{f_2}}{m_\rho} = 1.64 \quad (\text{Exp}) \quad (33)$$

To be fair, we have been working in the free field approximation, so that, at this point, the f_2 and ρ are infinitely narrow resonances. In fact, the widths of the f_2 and ρ are 185 MeV and 146 MeV respectively, and thus there is an uncertainty in what we should expect the resonance mass to be. So the AdS/CFT prediction is entirely satisfactory. It is remarkable how little we had to introduce to arrive at this prediction – no chiral symmetry breaking, no mention of the number of flavors or colors, in fact, no matching to QCD at all. We simply assumed that the lightest spin-2 mode is captured by a 5D massless spin-2 field (i.e. the dual of a spin-2 operator of the lowest dimension).

3.2 Matching to QCD

Next, we will calculate the coupling constant of the f_2 by matching to the operator product expansion for the tensor-tensor two point function in QCD. We prefer to express all the dimensionful scales in terms of z_m , so we write the coupling constant as $G_f = z_m g_f$, leaving g_f as the dimensionless coupling constant associated with the f_2 , directly analogous to g_5 for the ρ .

For each quark, we can construct a tensor bilinear,

$$\Theta_q^{\mu\nu} = \frac{1}{4} i \bar{q} (\gamma^\mu \overleftrightarrow{\partial}_\nu + \gamma^\nu \overleftrightarrow{\partial}_\mu) q \quad (34)$$

where $\overleftrightarrow{\partial} = \overleftarrow{\partial} - \overrightarrow{\partial}$. This is just the energy momentum tensor for quarks in QCD [9]. The OPE we need is the $\Theta\Theta$ two-point function, which can be written as

$$\langle \Theta_q^{\mu\nu} \Theta_q^{\rho\sigma} \rangle = P^{\mu\nu\rho\sigma} \Pi_{\theta\theta}^q(p^2) \quad (35)$$

where $P^{\mu\nu\rho\sigma}$ is the transverse projector (12). The constant term in the $\Theta\Theta$ OPE comes from a 1-loop quark diagram [18, 20]

$$\Pi_{\theta\theta}^q(p^2) = -\frac{N_c}{160\pi^2} p^4 \ln\left(\frac{p^2}{\Lambda^2}\right) + \dots \quad (36)$$

Since we have the correlator as a function of momentum p for fixed cutoff Λ , on the 5D side, we should think of the current as a source on the UV boundary at $z = \varepsilon$,

$$\mathcal{L} \supset \frac{1}{2} g_f \delta(z - \varepsilon) h_{\mu\nu} \Theta^{\mu\nu} \quad (37)$$

Then the $\Theta\Theta$ correlator can be derived from an effective 4D action at $z = \varepsilon$, with the fifth dimension integrated out:

$$\langle \Theta^{\mu\nu} \Theta^{\rho\sigma} \rangle = \frac{4}{g_f^2} \left(\frac{\delta}{\delta h_{\mu\nu}} \frac{\delta}{\delta h_{\rho\sigma}} e^{iS_{\text{eff}}} \right) \Big|_{h=0} \quad (38)$$

The easiest way to compute this is through the bulk-to-boundary propagators $K(z)$, which propagates a source via $h_{\mu\nu}(z) = h_{\mu\nu}(\varepsilon)K(z)$. Then the action (29) reduces to a boundary term on the equations of motion, and we have simply

$$\langle \Theta^{\mu\nu} \Theta^{\rho\sigma} \rangle = P^{\mu\nu\rho\sigma} \frac{4}{g_f^2} \frac{K'(\varepsilon)}{\varepsilon^3} \quad (39)$$

For the purpose of evaluating g_f , the IR boundary conditions are irrelevant and taking $z_m \rightarrow \infty$ we solve for $K(z)$ directly. The solution involves the second order Bessel function \mathcal{K}_2 [11]

$$K(p, z) = \frac{z^2 \mathcal{K}_2(pz)}{\varepsilon^2 \mathcal{K}_2(p\varepsilon)} \quad (40)$$

It is not hard to check that this satisfies $(z^3 \partial_z \frac{1}{z^3} \partial_z + p^2)K(p, z) = 0$ and $K(\varepsilon) = 1$. Thus the AdS prediction for the leading log in the tensor-tensor correlator is

$$\Pi_{\theta\theta}(p^2) = \frac{4}{g_f^2} \frac{K'(\varepsilon)}{\varepsilon^3} = -\frac{1}{2g_f^2} p^4 \log p^2 + \dots \quad (41)$$

The right hand side of this equation presents the leading log in the small z expansion of $\mathcal{K}_2(z)$. Once we identify the quark content of the f_2 , we can match the logs in (41) and (36) to fix g_f .

To identify the f_2 note that there is more than one isospin singlet state we can construct in QCD. First, there are the quark states. But in addition, in QCD there is a tensor glueball state. Its energy momentum tensor is

$$\Theta_{\mu\nu}^G = \frac{1}{4} \eta_{\mu\nu} (G_{\alpha\beta}^a)^2 - G_{\nu\alpha}^a G_{\alpha\mu}^a \quad (42)$$

Conveniently, the two-point function for the glueball in QCD has also been studied. Its OPE begins [14]

$$\langle \Theta_{\mu\nu}^G \Theta_{\rho\sigma}^G \rangle = -P^{\mu\nu\rho\sigma} \frac{1}{10\pi^2} p^4 \ln p^2 + \dots \quad (43)$$

Now, in general, there will be kinetic mixing among the glueball and the quark states, proportional to α_s . Although, the explicit mixing is known, we will be content to observe that one of the eigenstates should couple to the conserved current, the energy momentum tensor. This will be the lightest state, as the α_s corrections can only lift the mass, so we identify it as the f_2 .

Thus we are led to take the f_2 coupling to be

$$\mathcal{L} = \frac{1}{2}g_f h_{\mu\nu} \Theta^{\mu\nu}, \quad \Theta_{\mu\nu} = \Theta_{\mu\nu}^u + \Theta_{\mu\nu}^d + \Theta_{\mu\nu}^s + \Theta_{\mu\nu}^G \quad (44)$$

Hence, from QCD,

$$\langle \Theta_{\mu\nu} \Theta_{\rho\sigma} \rangle = -P^{\mu\nu\rho\sigma} \left(\frac{N_c N_f}{160\pi^2} + \frac{N_c^2 - 1}{80\pi^2} \right) p^4 \ln p^2 \quad (45)$$

Taking $N_c = N_f = 3$ and matching (41) produces to (45)

$$g_f = \frac{4\pi}{\sqrt{5}}. \quad (46)$$

And using $z_m^{-1} = 322$ MeV, determined from the ρ mass, the f_2 coupling constant is

$$G_f = g_f z_m = 0.0174 \text{ MeV}^{-1} \quad (47)$$

This coupling cannot be directly measured, but it lets us calculate the f_2 decay rates.

It is instructive to calculate the f_2 decay constant f_f as well, defined in (21). f_f also appears in residue of the decomposition of the two point function $\Theta\Theta$ into meson resonances.

$$\langle \Theta\Theta \rangle = \sum_{\text{mesons}} \frac{f_n^2 m_f^4}{p^2 - m_n^2} \quad (48)$$

On the AdS side, the easiest way to compute the residue is by observing that the Dirichlet Green's function has a similar decomposition over KK modes

$$G(z, z') = \sum_{\text{KK modes}} \frac{h_n(z) h_n(z')}{p^2 - m_n^2} \quad (49)$$

This Green's function is related to the bulk-to-boundary propagator by $K(z') = (z_m^2/z^3) \partial_z G(z, z')|_{z=\varepsilon}$ and so using (39) we can relate the decay constant directly to the wavefunction. For the f_2 , with wavefunction (32), we get

$$f_f = \frac{2}{g_f} \frac{1}{m_f^3 z_m^3} \frac{h'(\varepsilon)}{\varepsilon^3} = \frac{8.47}{m_f^3 z_m^3 g_f} = 0.024 \quad (50)$$

This also cannot be measured, but we can compare it to the prediction from TMD ($f_f = 0.042$). We will discuss the significance of this discrepancy after calculating the decay rates.

3.3 $f_2 \rightarrow \gamma\gamma$

We now discuss how to introduce the photon and calculate $\Gamma(f_2 \rightarrow \gamma\gamma)$. The photon is a vector field which is external to QCD. Nevertheless, we can consider it as a bulk gauge field, whose zero mode, the photon, is massless. Thus the photon will have a flat profile in the bulk

$$\gamma(z) = c_\gamma \quad (51)$$

Generally, a flat mode in AdS is not normalizable. In fact, regulating with a UV cutoff at $z = \varepsilon$

$$\int_\varepsilon^{z_m} \frac{1}{z} \gamma(z)^2 dz = c_\gamma^2 \log \frac{z_m}{\varepsilon}. \quad (52)$$

Normally, this would be a serious problem, as this divergent normalization appears as the coefficient of the photon kinetic term in the 4D action when the extra dimension is integrated out

$$\mathcal{L} = c_\gamma^2 \log \frac{z_m}{\varepsilon} \left(\frac{1}{4} F_{\mu\nu}^2 \right) + \dots \quad (53)$$

However, the coefficient of the term is supposed to be $1/e^2$, and the electric charge is divergent – it has a Landau pole. Explicitly, including only the quark contributions

$$\frac{1}{e^2(\mu)} = \left(\left(\frac{2}{3}\right)^2 + \left(\frac{1}{3}\right)^2 + \left(\frac{1}{3}\right)^2 + \dots \right) \frac{N_c}{6\pi^2} \ln\left(\frac{\mu}{\Lambda}\right) \quad (54)$$

But this means that if we set $c_\gamma^2 = e^2/(3\pi^2)$ (for $N_f = 3$) then the photon kinetic term becomes canonically normalized, $-\frac{1}{4}F_{\mu\nu}^2$. We can use this divergent c_γ in calculations by just taking the renormalized value for the electric charge at the relevant energy scale. It is satisfying that in contrast to chiral perturbation theory, where the electric charge must be added by hand, including an order one uncertainty, here the electric charge appears naturally from matching to QED.

Another way to model the photon is as a linear combination of the diagonal generators of $SU(3)_{\text{flavor}}$. Its coupling can then be fit from the AdS/CFT matching to the two-point function of vector currents in QCD. Since the relevant diagram is the same as the one-loop contribution as in the QED β -function, it leads to the same value for c_γ . Although the two ways of modeling the photon are equivalent, they have complimentary advantages. The former nicely demonstrates that the z -dependence of the photon profile corresponds to its running coupling, while the latter shows that it is really only the quark contributions which we must include. For example, we do not include the electron contribution to c_γ because the electron is not part of our AdS/QCD model.

The coupling of the f_2 to the photon is determined by general coordinate invariance. Using the parametrization (28), we get

$$\mathcal{L} = \int d^5x \sqrt{g} F_{MN} F^{MN} = \frac{1}{2} g_{f\gamma\gamma} h_{\mu\nu} \Theta_{\mu\nu}^\gamma + \dots \quad (55)$$

where $\Theta_{\mu\nu}^\gamma$ is given in (16) and the coupling constant is determined by the overlap integral

$$g_{f\gamma\gamma} = g_f \int_0^{z_m} \frac{dz}{z} h(z) \gamma(z)^2 = 0.671 c_\gamma^2 z_m g_f = 0.671 \frac{e^2}{3\pi^2} z_m g_f = 2.08 \times 10^{-3} G_f \quad (56)$$

If we use the value of G_f , calculated in the previous subsection by matching to QCD (including the 3 quark and gluon contributions), this leads to $g_{f\gamma\gamma} = 3.62 \times 10^{-5} \text{ MeV}^{-1}$ and a decay rate of

$$\Gamma(f_2 \rightarrow \gamma\gamma) = 2.70 \text{ keV} \quad (\text{AdS/QCD}) \quad (57)$$

Compared to the experimental rate of 2.60 ± 0.24 this is quite remarkable. All we needed to get this rate was N_c and N_f , and the ρ mass to define MeV. In fact, we do not even need the ρ at all. If we take z_m from the measured f_2 mass, $z_m = 333 \text{ MeV}$, then $\Gamma(f_2 \rightarrow \gamma\gamma) = 2.54 \text{ keV}$, which is also within experimental error.

At this point, since we have achieved a very favorable comparison to experiment, it is useful to return to some of our assumptions. Recall that we identified the f_2 with the energy-momentum tensor, because it is lightest spin-2 field in the theory. Thus, in our model, the f_2 is part glueball, and equal parts up, down, and strange. In fact, some quark models suggest that the f_2 and its nonet partner, the $f_2'(1525)$, are close to ideally mixed. That is, the f_2 is mostly up and down, and the f_2' is almost pure strange. We have not attempted to include quark masses at this point, so we cannot split the f_2 from the f_2' , but we can nevertheless compare our result to the results of making other assumptions about the f_2 quark content. For example, if we decouple the strange, so the f_2 is up, down and glue, we would get $\Gamma(f_2 \rightarrow \gamma\gamma) = 2.13 \text{ keV}$. It is reassuring that the 3 quark value (2.70 keV) and the 2 quark value (2.13 keV) straddle the experimental decay rate ($2.60 \pm 0.24 \text{ keV}$). If we had not included the glueball component, by ignoring the last term in Eq. (45), the decay rates would be 7.51 keV and 7.82 keV for the 3 quark and 2 quark cases respectively. This lets us conclude that the f_2 must have a large glueball component, consistent with a universal coupling to the energy momentum tensor.

3.4 $f_2 \rightarrow \pi\pi$

Thus far, we have derived two predictions – the f_2 mass and the partial width $\Gamma(f_2 \rightarrow \gamma\gamma)$ – and the only input from experiment (besides N_c and N_f) was the ρ mass, which just set the scale. To work out the rate $f_2 \rightarrow \pi\pi$ we need more information. We need to model chiral symmetry breaking, including both the quark condensate and quark masses. Needless to say, the construction we will use is not unique, and ambiguities make these predictions less accurate, and less convincing. Since this is not the main focus of this paper, we will quickly review, and then simply use, the formalism suggested in [3].

The global symmetries $SU(3)_L \times SU(3)_R$ of QCD map to gauge symmetries in the AdS. So we introduce bulk gauge bosons A_M^L and A_M^R for the two subgroups. In 4D, pions appear as the Goldstone bosons when the global chiral symmetry is spontaneously broken. Since in 5D the global symmetry is gauged, the Goldstone bosons are eaten and show up in unitary gauge as a mass term $m_A^2 A_M^2$ for the axial vector boson $A_M = A_M^R - A_M^L$. In the 4D description of the 5D Higgs mechanism, the vectors A_μ eat the scalar A_5 components. So in order to have massless 4D pions, we must arrange for A_5 to have a massless excitation.

The mass m_A corresponds to the formation of vacuum condensates, such as $\langle \bar{q}q \rangle$ in QCD. Thus we should think of m_A as being spontaneously generated by the expectation value of some bulk scalar field X . Since this breaking occurs in the IR, the mass should be localized near the IR brane. However, it should have some z dependence corresponding to the energy dependence of the operator $\langle \bar{q}q \rangle$. AdS/CFT tells us to match the dimension of the 4D operator to the bulk mass of a 5D field. This is somewhat intuitive – the energy dependence of an operator is determined by its scaling dimension, and the z dependence of a bulk field by its mass. So we assume $\langle \bar{q}q \rangle$ is represented by the expectation value of a bulk scalar field X . The scalar equations of motion in AdS lead to

$$\langle X \rangle = mz + \xi z^3 \equiv v(z) \quad (58)$$

Spontaneous symmetry breaking occurs in the IR, at large z , so ξ represents the strength of this effect. Similarly, explicit symmetry breaking due to quark masses, which is apparent even in the UV, corresponds to the m term, which is more relevant at small z . We will work in the massless quark limit for simplicity, and thus set $m = 0$. Then the free parameter ξ corresponds to the chiral symmetry breaking scale $\Lambda_{\text{QCD}}^3 \sim \langle \bar{q}q \rangle$.

So, the bulk action for the gauge fields in the AdS background is

$$\mathcal{S} = \int d^4x dz - \frac{1}{4z} (F_{MN}^L)^2 - \frac{1}{4z} (F_{MN}^R)^2 + \frac{v(z)^2}{2z^3} (A_M)^2 \quad (59)$$

Note that the vector combination $V_M = A_M^L + A_M^R$, does not have a bulk mass. So its first KK mode will be the same ρ we introduced previously. For the axial section, we want to decouple A_5 and A_μ . To do this, we introduce Goldstone bosons and gauge-fix, following [3]. This separates the pions from the axial vectors.

After this gauge fixing, the KK equation for the axial vector is

$$\left[m^2 - 2\frac{v(z)^2}{z^2} + z\partial_z \frac{1}{z} \partial_z \right] a_n(z) = 0 \quad (60)$$

We can use this and the experimental a_1 mass to fix ξ . With boundary conditions $a'(z_m) = a'(0) = 0$, and using $z_m^{-1} = 322\text{MeV}$ from the ρ mass, we find, numerically, an eigenvalue at the a_1 mass of 1230 MeV for $\xi = 3.77z_m^{-3}$.

The zero mode of A_5 represents the physical pions. The KK profile for this mode satisfies

$$\partial_z \left(\frac{z^3}{2v(z)^2} \partial_z \left(\frac{1}{z} A_5(z) \right) \right) = A_5(z) \quad (61)$$

The solution is

$$A_5(z) = N \xi z^3 \left[\mathcal{I}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z^3 \right) - \beta_\pi \mathcal{K}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z^3 \right) \right] \quad (62)$$

Although we have not imposed boundary conditions, $A_5(0) = 0$ automatically. This is essentially set by the fact that we insist on having a massless mode. The boundary condition $A_5(z_m) = 0$ then sets β_π . The normalization is determined by a z integral

$$- \int dz \left[\frac{1}{z} A_5(z)^2 + \frac{1}{z} \frac{z^3}{2v(z)^2} \left(\partial_z \frac{1}{z} A_5(z) \right)^2 \right] = 1 \quad (63)$$

On the solution for $A_5(z)$, the expression in brackets is a total derivative, which lets us calculate the normalization analytically

$$N^{-1/2} = \frac{\mathcal{I}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)}{2\mathcal{K}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)^2} \left[\sqrt{3} \pi \mathcal{I}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right) + 3\mathcal{K}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right) \right] \quad (64)$$

Thus, identifying this mode with the pion wavefunction, we have

$$\pi(z) = N \xi z^3 \left[\mathcal{I}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z^3 \right) - \frac{\mathcal{I}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)}{\mathcal{K}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z_m^3 \right)} \mathcal{K}_{2/3} \left(\frac{\sqrt{2}}{3} \xi z^3 \right) \right] \quad (65)$$

Now we can calculate $g_{f\pi\pi}$.

The coupling of f_2 to the pions is determined by general coordinate invariance. Since everything is canonically normalized, we simply evaluate the overlap of the f_2 with the combination that appears in the kinetic term for π .

$$g_{f\pi\pi} = g_f \int dz h(z) \left[\frac{1}{z} \pi(z)^2 + \frac{z^3}{2v(z)^2} \left(\partial_z \frac{1}{z} \pi(z) \right)^2 \right] = 0.519 z_m g_f = 0.519 G_f \quad (66)$$

So we predict for the G_f -independent ratio

$$\frac{g_{f\pi\pi}^2}{g_{f\gamma\gamma}^2} = 6.24 \times 10^4 \quad (\text{AdS/QCD}) \quad (67)$$

Compared to the experimental value of 2.72×10^5 this is off by a factor of 4. Using our previous result, $G_f = 0.0174 \text{ MeV}^{-1}$, we get $g_{f\pi\pi} = 9.05 \times 10^{-3} \text{ MeV}^{-1}$, and predict

$$\Gamma(f_2 \rightarrow \pi^+ \pi^-, \pi^0 \pi^0) = 37.4 \text{ MeV} \quad (\text{AdS/QCD}) \quad (68)$$

This also off from the experimental value of 156.9 ± 3.8 MeV by a factor of 4.

This factor of 4 is from the same origin as the factor of 2 discrepancy between f_f from AdS/QCD ($f_f \sim 0.02$) and f_f from TMD. After all, in TMD, f_f is extracted from the $f_2 \rightarrow \pi\pi$ rate. In AdS/QCD, there is a conflict between the decay rates to photons and to pions, because they depend on the same parameters; but we will now see that it is justifiable, and it is good that the photon rate is the one which comes out right.

4 Higher order operators

We can understand why the decay to photons was so accurate, but the decay to pions was so far off by looking at higher dimension operators. Because the f_2 is heavy, 1275 MeV, compared to the naive cutoff of the chiral Lagrangian, $4\pi f_\pi = 1168$ MeV, we expect chiral perturbation theory not to be accurate in the f_2 sector. We can also see the breakdown of perturbation theory on the AdS side, by looking at where higher dimension derivative operators become relevant. Let us now consider some of these operators, relevant to the $f_2 \rightarrow \pi\pi$ process.

Since we are interested in $f_2 \rightarrow \pi\pi$ we want operators linear in h and quadratic in A_M . The term we have been using to calculate $g_{f\pi\pi}$ was determined by general coordinate invariance and the π kinetic term. It is, roughly,

$$\int d^5x \frac{1}{z} g_f h_{\mu\nu} (\partial_\mu A_5) (\partial_\nu A_5) = (0.209) G_f \int d^4x h_{\mu\nu} (\partial_\mu A_5) (\partial_\nu A_5) \quad (69)$$

The number 0.209 comes from the z -integral, with the f_2 and π wavefunctions. The difference from (66) is because we have simplified the tensor structure, but it is the same order of magnitude.

A possible higher order term, leading to exactly the same 4D structure is

$$\int d^5x \frac{1}{z} g_f h_{\mu\nu} (\partial_\mu A_5) (z^2 \partial_z^2) (\partial_\nu A_5) = (-1.57) G_f \int d^4x h_{\mu\nu} (\partial_\mu A_5) (\partial_\nu A_5) \quad (70)$$

This could come from the 5D general coordinate invariant term

$$\int d^5x \sqrt{g} F_{MN} D^N D^P F_{PM} \quad (71)$$

Or

$$\int d^5x \frac{1}{z} g_f (z^2 \partial_z^2 h_{\mu\nu}) (\partial_\mu A_5) (z^2 \partial_z^2) (\partial_\nu A_5) = (5.05) G_f \int d^4x h_{\mu\nu} (\partial_\mu A_5) (\partial_\nu A_5) \quad (72)$$

which might come from

$$\int d^5x \sqrt{g} R_{MNO P} (h) F^{MN} F^{OP}, \quad (73)$$

where $R_{\text{MNOP}}(h)$ is invariant under $h_{\mu\nu} \rightarrow h_{\mu\nu} + \partial_\mu \xi_\nu(x) + \partial_\nu \xi_\mu(x)$. Note that linearized $\tilde{R}_{\mu\nu}(h) = R_{\mu\nu}(h) - 4/z^2 h_{\mu\nu}$ (the combination invariant under the above shift) vanishes when the f_2 is on shell.

We could assume that these terms are suppressed compared to the leading piece by some factors of the weak coupling constant g_f . At best, we could use the 5D loop factor, and suppress by $g_f^2/24\pi^3 = 0.04$. Thus the correction (70) contributes 30% relative to the original term (69), which is not a small effect. With the 4D loop factor, $16\pi^2$, the new term is 1.5 times as important as the original. Either way, there is no excuse to ignore the higher order corrections and we cannot reasonably expect our rate $f_2 \rightarrow \pi\pi$ to be at all accurate. In fact, it is surprising that we were even able to get the coupling constant to a factor of 2.

In contrast, consider the terms contributing to corrections to $f_2 \rightarrow \gamma\gamma$. At leading order, we had

$$\int d^5x \frac{1}{z} g_f h_{\mu\nu} F_{\mu\alpha} F_{\nu\alpha} = (0.671) G_f \int d^4x h_{\mu\nu} F_{\mu\alpha} F_{\nu\alpha} \quad (74)$$

Any correction to the $f_2 \rightarrow \gamma\gamma$ rate has to come from a vertex with one h and two F 's. But the photon profile is flat, so any term with ∂_z 's acting on F will vanish. Terms with ∂_μ acting on F (like the Riemann term above) may appear, but they cannot contribute to the decay rate because the photon is massless and transverse – there is nothing to which we may contract the new momentum factor.

One might worry that term such as

$$\int d^5x \sqrt{g} F_{\text{MN}} D^2 F^{\text{MN}} \quad (75)$$

could contribute if the derivatives are ∂_z 's and they act on the background metric. But the effect is then only to shift the coupling constant g_5 , which is unobservable because the renormalized coupling has been matched to QCD. Another way to see this is that on shell $D^2 F_{\text{MN}} \sim 1/z^2 F_{\text{MN}}$ which is the standard gauge kinetic term. Thus, the contribution of (75) has already been accounted for in the definition of g_5^2 . So no operators, analogous to the ones relevant for the pion decay, affect the photon decay mode at all.

Throughout, we have been ignoring boundary operators on the IR brane. But these can affect the photon decay rate, and should be included in a consistent effective field theory. The natural size of a term like $\delta(z - z_m) h F^2$ is of order z_m times $g_f^2/16\pi^2 = 0.20$, or about $.04 G_f$. Comparing to (74) we expect corrections of order 10% to the f_2 photon coupling. In addition, some other assumption could be wrong. For example, α_s corrections have not been included – we have assumed that QCD is conformal. So this can change the result as well. Really, we could not have known ahead of time that we would get $f_2 \rightarrow \gamma\gamma$ so precisely. Nevertheless, the rate came out right, and it is suggestive that the contribution of boundary operators and violations of conformality are small.

5 Higher Spin States

In addition to the f_2 , which we have studied here, QCD has a whole tower of high spin resonances (for example, the f_6 with spin 6 sits at 2510 MeV). In this section, we derive the predictions of AdS/QCD for their masses, and compare to experiment.

A spin s state in QCD corresponds to a field $\phi_{M_1 M_2 \dots M_s}$ in AdS. Ideally, we would like to write down an action for ϕ , but higher spin Lagrangians are in general very complicated. For example, even at the free field level in flat space, one needs to introduce on the order of s auxiliary fields to eliminate the propagation of unphysical modes. Instead, we will take a more direct approach, and simply use symmetry arguments to guess the equations of motion for the transverse spin- s modes.

It is natural to expect the AdS theory for ϕ to respect a linearized gauge invariance, under which

$$\phi_{M_1 M_2 \dots M_s} \rightarrow \phi_{M_1 M_2 \dots M_s} + D_{\{M_1} \xi_{M_2 \dots M_s\}}, \quad (76)$$

Such a symmetry allows us to go to an axial gauge in which $\phi_{5 M_2 \dots M_s} = 0$. In AdS, it is not hard to show that this gauge is preserved under residual transformations with $\xi_{5 M_3 \dots M_s} = 0$ and $\xi_{\mu_2 \dots \mu_s}(x, z) = z^{2-2s} \xi_{\mu_2 \dots \mu_s}(x)$. This means that a field $\phi(x, z) = z^{2-2s} \tilde{\phi}(x)$ simply shifts under the 4D gauge transformation, and therefore represents a zero mode. In terms of $\tilde{\phi}$, in this axial gauge, the Lagrangian contains a piece $\sqrt{g}(\partial_\mu \phi)^2 = z^{1-2s}(\partial_\mu \tilde{\phi})^2$ and a similar piece with ∂_z 's acting directly on $\tilde{\phi}$. Thus the KK modes for a field of spin s will satisfy

$$\partial_z \frac{1}{z^{2s-1}} \partial_z \tilde{\phi} + \frac{m^2}{z^{2s-1}} \tilde{\phi} = 0 \quad (77)$$

Note that this equation is just a naive generalization of (25) and (30) and matches previous results for spins 1 and 2.

We can confirm this result from AdS/CFT reasoning as well. On the QCD side, the lowest dimension operator of spin s , may contain terms like

$$\begin{aligned} & \bar{q} \gamma_{\{\mu_1} \partial_{\mu_2} \dots \partial_{\mu_s\}} q \\ & G_{\{\mu_1}^\rho \partial_{\mu_2} \dots \partial_{\mu_{s-1}} G_{\mu_s\} \rho}. \end{aligned} \quad (78)$$

These operators are of dimension $\Delta = s + 2$. Under conformal transformations, operators scale based on their twist $\Delta - s$. Thus a constant bulk field, representing the vev of such an operator, should vary as $\phi \sim z^2$. The source for the operator \mathcal{J}_Δ has dimension $4 - \Delta = 2 - s$ and so it sources a field which varies like $\phi \sim z^{2-2s}$. This is the same AdS zero mode solution we derived by symmetry arguments above.

The normalizable solutions to (77) are $z^s \mathcal{J}_s(mz)$. Requiring Neumann boundary conditions in the IR, we find that the mass of the lightest spin- s particle is given by the first zero of \mathcal{J}_{s-1} . This leads to the following predictions for spin- s resonances

in QCD

Particle	ρ	f_2	ω_3	f_4	ρ_5	f_6
Experiment(MeV)	775.8	1275 ± 1	1667 ± 4	2025 ± 8	2330 ± 35	2465 ± 50
AdS/QCD(MeV)	--	1236	1656	2058	2448	2829

Note that we have neglected the effects of quark masses and so we are not sensitive to isospin. In Figure 5, we plot the squared masses from this table as a function of spin. We also display the linear fit to the experimental values, which is the Regge trajectory, and the quadratic curve through the AdS/QCD points. Note that the AdS/QCD results take only m_ρ from data, but provide a good estimate of both the intercept and slope of the Regge trajectory. Although the linear fit is a good match to data, the quadratic fit from AdS/QCD seems to produce deviations at low energy in the right direction. It is expected that at higher energy, stringy effects are dominant, and we should merge smoothly back into the linear regime.




Figure 1: Squared masses from AdS/QCD (red) versus experiment (blue), as a function of spin. Linear (Regge) fit to experimental values and quadratic fit to AdS/QCD values are shown. The errors shown are due to the widths of the resonances.

We might try to calculate the decay rates of these particles as well. There are two main impediments. First, none of the decay modes are well measured. Second, we do not know how to write down interactions for higher spin fields. For the spin-2 case, we used general coordinate invariance to guess the f_2 couplings. But for higher spin fields, it is impossible to find a consistent set of interactions (there can be no higher

spin conserved current), at least in flat space⁴. Yet we know massive higher spin fields exist in QCD, and we can try to guess these interactions, or perhaps compute them using the holographic map.

6 Conclusions

In this paper, we have shown that despite obvious challenges, the AdS/CFT correspondence produces some remarkably accurate predictions about QCD. We saw that the mass for the f_2 meson and the rate for $f_2 \rightarrow \gamma\gamma$ are in fantastic agreement with experiment. The $f_2 \rightarrow \gamma\gamma$ prediction is particularly satisfying, because it is difficult to approach this decay through more traditional methods. Moreover, we have demonstrated that higher-dimension bulk operators do not affect this decay rate, and so our results are trustworthy. In contrast, we also calculated the rate for $f_2 \rightarrow \pi\pi$, which was off by a factor of 4. This rate is sensitive to higher order corrections, and to our representation of chiral symmetry breaking in AdS/CFT. Thus our results are predictions of a self-consistent effective field theory which matches remarkably well to QCD at leading order.

We have also shown that the naive expectation from AdS/CFT, that the lightest state of a given spin is captured by the lowest dimension operator with that spin, seems to be favored by data. This indicates that not only is AdS/QCD capable of quantitative success, but also that it may reveal interesting connections between QCD and its dual. For example, we have seen that the f_2 makes an awful 4D graviton. In contrast, on an AdS₅ background, general coordinate invariance allowed us to predict the f_2 coupling to photons correctly, making it a natural 5D graviton KK mode.

Acknowledgements

We would like to thank T. Bhattacharya, M. Chanowitz, and M. Suzuki for illuminating conversations about meson physics. EK and MS thank the Aspen Center of Physics for its hospitality during the completion of part of this work.

References

- [1] For a review, see e.g. A. Pich, Rept. Prog. Phys. **58**, 563 (1995) [arXiv:hep-ph/9502366].
- [2] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, arXiv:hep-ph/0501128.

⁴See [39] for the intriguing possibility of constructing consistent interacting theories of higher spin fields in AdS.

- [3] L. Da Rold and A. Pomarol, Nucl. Phys. B **721**, 79 (2005) [arXiv:hep-ph/0501218].
- [4] L. Da Rold and A. Pomarol, arXiv:hep-ph/0510268.
- [5] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [6] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
- [7] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].
- [8] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111].
- [9] T. Han, J. D. Lykken and R. J. Zhang, Phys. Rev. D **59**, 105006 (1999) [arXiv:hep-ph/9811350].
- [10] J. Hirn and V. Sanz, arXiv:hep-ph/0507049.
- [11] L. Randall and M. D. Schwartz, JHEP **0111**, 003 (2001) [arXiv:hep-th/0108114].
- [12] M. Suzuki, Phys. Rev. D **47**, 1043 (1993).
- [13] M. E. Peskin and D. V. Schroeder,
- [14] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **191**, 301 (1981).
- [15] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 448 (1979).
- [16] M. A. Shifman, Prog. Theor. Phys. Suppl. **131**, 1 (1998) [arXiv:hep-ph/9802214].
- [17] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979).
- [18] T. M. Aliev and M. A. Shifman, Phys. Lett. B **112**, 401 (1982).
- [19] P. Colangelo and A. Khodjamirian, arXiv:hep-ph/0010175.
- [20] L. J. Reinders, S. Yazaki and H. R. Rubinstein, Nucl. Phys. B **196**, 125 (1982).
- [21] E. Bagan and S. Narison, Phys. Lett. B **214**, 451 (1988).

- [22] E. Bagan, A. Bramon and S. Narison, Phys. Lett. B **196**, 203 (1987).
- [23] K. Ishikawa, I. Tanaka, K. F. Liu and B. A. Li, Phys. Rev. D **37**, 3216 (1988).
- [24] M. Fierz and W. Pauli, Proc. Roy. Soc. Lond. A **173**, 211 (1939).
- [25] P. Van Nieuwenhuizen, Nucl. Phys. B **60**, 478 (1973).
- [26] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B **321**, 311 (1989).
- [27] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, Phys. Lett. B **223**, 425 (1989).
- [28] M. Harada and K. Yamawaki, Phys. Rept. **381**, 1 (2003) [arXiv:hep-ph/0302103].
- [29] H. van Dam and M. J. G. Veltman, Nucl. Phys. B **22**, 397 (1970).
- [30] D. T. Son and M. A. Stephanov, Phys. Rev. D **69**, 065020 (2004) [arXiv:hep-ph/0304182].
- [31] G. F. de Teramond and S. J. Brodsky, Phys. Rev. Lett. **94**, 201601 (2005) [arXiv:hep-th/0501022].
- [32] C. Csaki, J. Erlich, T. J. Hollowood and Y. Shirman, Nucl. Phys. B **581**, 309 (2000) [arXiv:hep-th/0001033].
- [33] T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000) [arXiv:hep-ph/0003129].
- [34] C. R. Hagen, Phys. Rev. D **6**, 984 (1972).
- [35] N. Arkani-Hamed, H. Georgi and M. D. Schwartz, Annals Phys. **305**, 96 (2003) [arXiv:hep-th/0210184].
- [36] M. D. Schwartz, Phys. Rev. D **68**, 024029 (2003) [arXiv:hep-th/0303114].
- [37] J. Gasser and H. Leutwyler, Nucl. Phys. B **250**, 465 (1985).
- [38] A. Manohar and H. Georgi, Nucl. Phys. B **234**, 189 (1984).
- [39] M. A. Vasiliev, Phys. Lett. B **567**, 139 (2003) [arXiv:hep-th/0304049].