

LLNL-JRNL-400008



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Response to #A Madden-Julian Oscillation Event Realistically Simulated by a Global Cloud-Resolving Model#

K. R. Sperber

December 19, 2007

Science

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Response to “A Madden-Julian Oscillation Event Realistically Simulated by a Global  
Cloud-Resolving Model” (1)

Kenneth R. Sperber

Lawrence Livermore National Laboratory

Program for Climate Model Diagnosis and Intercomparison

P. O. Box 808, L-103

Livermore, CA 94551

I agree with the authors that forecasting the Madden-Julian Oscillation (MJO) in a high resolution global model is important for numerous reasons, including improved weather forecast skill beyond 10 days, and resolving small scale features embedded in the MJO that coarse resolution (~100-300km horizontal grid spacing) climate models do not (e.g., tropical cyclones). Unfortunately, the authors promote the (incorrect) overall impression that coarse resolution climate models cannot simulate the MJO by (a) only discussing aspects of works that indicate the poor ability of coarse resolution climate models to simulate the MJO (2, 3), and (b) by promoting the use of higher resolution models (1), and the use of embedded two-dimensional cloud resolving models embedded in coarse resolution climate models (4, 5) as the principal methods for realistically representing the MJO because of the difficulty of coarse resolution models “to estimate the vertical redistribution of heat and moisture by unresolved convective clouds.”

Regarding items (a) and (b), I have co-authored two of the works cited by Miura *et al.* that bemoan the poor ability of coarse resolution climate models to simulate the MJO (2, 3), and indeed simulating the MJO in coarse resolution climate models is a grand challenge. However, I would like to draw to their attention to work that has demonstrated that two different coarse resolution climate models, using conventional parameterizations of convection and clouds, can represent the MJO with high fidelity (6, 7). In the later study, where more complete model diagnostics were available, important aspects of the MJO that were realistically represented included the relationship between convection and low-level moisture convergence, surface fluxes, the vertical structure of

winds and divergence, and important air-sea interactions. Additionally, regarding item (b), convection is certainly of central importance in representing the MJO, but it is the interaction of convection (parameterized or otherwise) with other aspects of the model physics that is important. Thus, it is the implementation of the full physics package that is relevant, not simply the manner in which convection is represented.

The issue at hand is to determine what aspects of the physics present in the high resolution (and embedded 2-dimensional cloud resolving model) simulations are essential for incorporation into coarse resolution climate model parameterizations in order for them to adequately represent the MJO. In the foreseeable future, models used for climate and climate change studies will continue to require convective parameterization in order to produce simulations that span decadal to centennial time scales to estimate the impact of anthropogenic influences on the statistics of weather variability and extremes. As such, intercomparison across different classes of modeling is essential to better understand the physics relevant to the climate system.

Acknowledgment. K. R. Sperber was supported under the auspices of the U.S. Department of Energy Office of Science, Climate Change Prediction Program by Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344. I thank P. J. Gleckler (PCMDI) for helpful comments on a earlier version of this comment.

## References

- (1) H. Miura *et al.*, *Science* **318**, 1763 (2007)
- (2) J. M. Slingo *et al.*, *Clim. Dyn.* **12**, 325 (1996)
- (3) J.-L. Lin *et al.* *J. Clim.* **19** 265 (2006)
- (4) W. W. Grabowski, *J. Atmos. Sci.* **60**, 847 (2003)
- (5) M. F. Khairoutdinov, D. A. Randall, C. DeMotte, *J. Atmos. Sci.* **62**, 2136 (2005)
- (6) K. R. Sperber *et al.* *Clim. Dyn.* **13**, 769 (1997)
- (7) K. R. Sperber *et al.* *Clim. Dyn.* **25**, 117 (2005)