

UNIVERSITY OF
CALIFORNIA


*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-191
02



CLASSIFICATION CANCELLED BY AUTHORITY
OF THE DISTRICT ENGINEER
BY THE DECLASSIFICATION COMMITTEE
UNIVERSITY OF CALIFORNIA

Radiation Laboratory

THE ION-EXCHANGE SEPARATION OF ZIRCONIUM AND HAFNIUM

Kenneth Street, Jr. and Glenn T. Seaborg

October 11, 1943

Special Review of Declassified Reports
Authorized by USDOE JK Bretton
Unclassified TWX F182206Z May 79

REPORT PROPERLY DECLASSIFIED

| | |
|----------------------------------|----------------|
| <u>J N Green</u> | <u>8/16/79</u> |
| Authorized Derivative Classifier | Date |
| <u>R G Hunt</u> | <u>8/17/79</u> |
| By | Date |

Berkeley, California

STANDARD DISTRIBUTION: Series A

| | |
|--|-------|
| Argonne National Laboratory, | 9 |
| Armed Forces Special Weapons Project | 10-11 |
| Atomic Energy Commission, Washington | 12 |
| Battelle Memorial Institute | 13-22 |
| Brookhaven National Laboratory | 23-26 |
| Carbide & Carbon Chemicals Corporation (K-25 Area) | 27-30 |
| Carbide & Carbon Chemicals Corporation (Y-12 Area) | 31 |
| Columbia University (Failla) | 32-35 |
| General Electric Company | 36-42 |
| Hanford Directed Operations | 43 |
| Iowa State College | 44-45 |
| Kellex Corporation | 46-48 |
| Los Alamos | 49 |
| Massachusetts Institute of Technology | 50-51 |
| Monsanto Chemical Company, Dayton | 52-53 |
| National Bureau of Standards | 54 |
| Naval Radiological Defense Laboratory | 55 |
| NEPA | 56-57 |
| New York Directed Operations | 58-69 |
| Oak Ridge National Laboratory | 70 |
| Patent Advisor | 71-85 |
| Technical Information Division | 86 |
| UCLA Medical Research Laboratory (Warren) | 87-89 |
| University of California, Radiation Laboratory | 90 |
| Information Division | 91-92 |
| Chemistry Dept. | 93 |
| University of Rochester | 94 |
| Western Reserve University (Friedell) | |
| Office of Chicago Directed Operations | |

DECLASSIFICATION PROCEDURE:

| | |
|--------------------------|---------|
| Declassification Officer | 95-98 |
| Publication Officer | 99 |
| Patent Department | 100-101 |
| Area Manager | 102 |
| E. O. Lawrence | 103 |
| Information Div. | 104 |

CLASSIFICATION COPY NUMBERS
 OF THIS DISTRICT ENGINEER
 BY THE DECLASSIFICATION COMMITTEE

The Ion-Exchange Separation of Zirconium and Hafnium

Kenneth Street, Jr. and Glenn T. Seaborg
Department of Chemistry and Radiation Laboratory
University of California, Berkeley, California

October 11, 1948

In the course of a rather cursory examination of the elution of tetra-positive ions from the cation exchange resin Dowex 50 with hydrochloric acid solutions, we have discovered a very effective method of separating zirconium from hafnium. In view of the great labor involved in preparing even reasonably pure hafnium compounds by existing methods, we feel that this procedure will prove very valuable to those interested in obtaining hafnium compounds free of zirconium.

Although the conditions which give satisfactory separation were first worked out using microgram amounts of material and the radioactive tracer technique, the run described below, involving milligrams of material, illustrates the applicability of the method to the production of significant amounts of pure hafnium and zirconium.

35 mg. of zirconium oxide 15 mg. of hafnium oxide were dissolved in sulfuric and hydrofluoric acids, hafnium and zirconium tracer added, and the mixture fumed to dryness. The residue was taken up in concentrated hydrochloric acid and the hydroxides precipitated with ammonium hydroxide and washed. The hydroxides were again dissolved in hydrochloric acid and the oxychlorides crystallized by evaporation. 1 cc. of 250 to 500 mesh Dowex 50 spheres, in the ammonium form, were suspended in 30 cc. of 2 molal perchloric acid and the oxychlorides added a few mg. at a time over a period of 15 minutes, the mixture being continually agitated by bubbling

air through it. Under these conditions, i.e. $\sim 0.01M$ zirconium and hafnium in 2M perchloric acid, the zirconium and hafnium are not appreciably polymerized¹ and about 80 percent of each goes on the resin. The slurry of resin was placed on the top of an ion-exchange column 1 sq. cm. in area and 30 cm. long which had been packed with the same resin and washed with 6 molal hydrochloric acid to convert it to the acid form. On elution with 6 molal hydrochloric acid the curve shown in Fig. 1 was obtained. The outline of the curve was obtained by counting the tracers (Zr^{95} and Hf^{181}) and the dotted portions by optical spectrographic analysis.

Thus it can be seen that ~ 66 percent of the starting hafnium oxide, i.e. 10 mg., is obtained containing ~ 0.1 percent zirconium oxide by weight. The column used in these experiments was relatively small and thus gram amounts of material should be easily handled on columns of only moderate size.

This work was done under the auspices of the Atomic Energy Commission at the Radiation Laboratory, University of California, Berkeley, California.

1. Connick, R. E., and W. H. McVey, Private Communication.




FIG. 1
ELUTION OF Zr AND Hf WITH 60MHCl