

## The Highest Redshift Relativistic Jets

C. C. Cheung<sup>1</sup>, Ł. Stawarz

*Kavli Institute for Particle Astrophysics and Cosmology,  
Stanford University, Stanford, CA 94305*

A. Siemiginowska, D. E Harris, D. A. Schwartz

*Harvard-Smithsonian Center for Astrophysics,  
60 Garden St., Cambridge, MA 02138*

J. F. C. Wardle, D. Gobeille

*Physics Department, Brandeis University, Waltham, MA 02454*

N. P. Lee

*Institute for Astrophysical Research, Boston University,  
725 Commonwealth Ave., Boston, MA 02215*

**Abstract.** We describe our efforts to understand large-scale (10's–100's kpc) relativistic jet systems through observations of the highest-redshift quasars. Results from a VLA survey search for radio jets in  $\sim 30$   $z > 3.4$  quasars are described along with new *Chandra* observations of 4 selected targets.

### 1. Why High-redshift Jets?

It is now well established that X-ray emission is a common feature of kiloparsec-scale radio jets (see Harris & Krawczynski 2006, for a recent review and the associated website, <http://hea-www.harvard.edu/XJET/>). The spectral energy distributions (SEDs) of the powerful quasar jets are predominantly characterized as “optically faint”, with the spectra rising between the optical and X-ray bands. Current models for this ‘excess’ X-ray emission posit either inverse Compton (IC) scattering off CMB photons in a (still) relativistic kpc-scale jet or an additional high-energy synchrotron emitting component.

In the simplest scenario, such models have diverging predictions at high redshift. Specifically, we expect a strong redshift dependence in the monochromatic flux ratio,  $f_X/f_r \propto U_{\text{CMB}} \propto (1+z)^4$  for IC/CMB, whereas in synchrotron models, we expect no such dependence,  $f_X/f_r \propto (1+z)^0$ . As a first order test of this simple idea, our approach is to study the highest-redshift relativistic jets. Such jets probe the physics of the earliest (first  $\sim 1$  Gyr of the Universe in the quasars studied) actively accreting supermassive black hole systems and

---

<sup>1</sup>Jansky Postdoctoral Fellow of the National Radio Astronomy Observatory

are interesting for other reasons. For instance, the ambient medium in these high-redshift galaxies is probably different (e.g., De Young 2006) and this may manifest in jets with different morphologies, increased dissipation, and slower than their lower-redshift counterparts.




Figure 1. Examples of newly discovered arcsecond-scale radio jets from our VLA observations (§ 2.1). Clockwise from upper left, the sources are J0624+3856 ( $z=3.469$ ; Xu et al. 1995), J2042-2223 ( $z=3.630$ ; Hook et al. 2002), J2220-3336 ( $z=3.691$ ; Hook et al. 2002), and J2219-2719 ( $z=3.634$ ; Hook et al. 2002). The J2219-2719 image is at 1.4 GHz while the rest are at 5 GHz. The beam-sizes are  $0.41'' \times 0.41''$ ,  $0.73'' \times 0.38''$  at  $PA=-8.2^\circ$ ,  $1.13'' \times 0.39''$  at  $PA=10.2^\circ$ , and  $0.75'' \times 0.75''$  (super-resolved), respectively. The lowest contour levels begin at 0.125 mJy/bm for all images except for J2219-2719 where it is 0.2 mJy/bm, and increase by factors of  $\sqrt{2}$ .

Most *Chandra* studies of quasar jets have so far targeted known arcsecond-scale radio jets (e.g., Sambruna et al. 2004; Marshall et al. 2005), as most known examples are at  $z \lesssim 2$  (Liu & Zhang 2002). There are currently only two high- $z$  quasars with well-established kpc-scale X-ray jet detections: GB 1508+5714 at  $z=4.3$  (Siemiginowska et al. 2003; Yuan et al. 2003; Cheung 2004) and 1745+624 at  $z=3.9$  (Cheung et al. 2006). They are observed to have large  $f_X/f_r$  values as expected in the IC/CMB model (Schwartz 2002; Cheung 2004), although the

small number of high- $z$  detections preclude any definitive statements (Kataoka & Stawarz 2005; Cheung et al. 2006).

We have therefore carried out a VLA survey in search of new radio jets in a sample of high- $z$  quasars (§ 2.1.) and new *Chandra* observations of a small subset (§ 2.2.). This contribution presents some results from these observations. For the redshifts considered,  $z=3.4$  to 4.7,  $1''$  corresponds to 7.4 to 6.5 kpc ( $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_M = 0.3$  and  $\Omega_\Lambda = 0.7$ ).

## 2. Observations of a High-Redshift Quasar Sample

### 2.1. VLA Imaging Survey

Using NED, we assembled a sample of  $z>3.4$  flat-spectrum radio quasars for imaging with the VLA. We did not aim for our sample to be a complete one as current samples of lower- $z$  X-ray jets are inhomogenous also. With archival (Lee 2005) and new VLA observations, we find that radio jets in this redshift range are common with a  $\sim 50\%$  detection rate (Cheung et al. 2005, and in preparation). Examples of new radio jets detected from our observations are shown in Figure 1.

### 2.2. *Chandra* Observations

A small percentage of the radio jets from our radio study (§ 2.1.) are extended enough ( $>2.5''$  long) to study with *Chandra*. We observed four of them with short snapshot *Chandra* observations (Figure 2). We detected bright X-ray counterparts to the jets in the quasars J1421–0643 ( $z=3.689$ ; Ellison et al. 2001) and GB 1428+4217 ( $z=4.72$ ; Hook & McMahon 1998); the latter detection is currently the highest-redshift kpc-scale radio and X-ray jet known. We did not detect the X-ray counterparts to the radio jets in 1239+376 ( $z=3.819$ ; Vermeulen et al. 1996) and J1754+6737 ( $z=3.6$ ; Villani & di Serego Alighieri 1999). The 2/4 X-ray jet detection rate of our high- $z$  sample is comparable to that of lower- $z$  samples (Sambruna et al. 2004; Marshall et al. 2005).

## 3. Discussion and Summary

Previous *Chandra* imaging studies of a number of  $z>4$  radio loud quasars do not reveal significant extended X-ray emission (Bassett et al. 2004; Lopez et al. 2006). However, in these studies, there were no pre-existing information on possible radio structures in the target objects and any definitive statements regarding the nature of the X-ray emission mechanism in jets at high-redshifts may be premature. In fact, in one case where there was evidence of an extended X-ray structure (J2219–2719; Lopez et al. 2006), our VLA observation revealed a radio counterpart (Figure 1).

In our approach, we began with a VLA survey of a sample of  $z>3.4$  quasars and found radio jets to be relatively common ( $\sim 50\%$  detection rate). These jets are quite luminous; with a confident detection of a 1 mJy knot at 1.4 GHz, this corresponds to luminosities of  $1.5 \times 10^{42} \text{ erg s}^{-1}$  ( $z=3.4$ ) to  $3.1 \times 10^{42} \text{ erg s}^{-1}$  ( $z=4.7$ ).




Figure 2. *Chandra* X-ray images (colorscale) with VLA contours overlaid of the four high- $z$  radio jets observed. There are X-ray detections of the top two objects but not of the bottom two (§ 2.2).

With the radio survey results, we found only a few radio jets to have sufficient angular extent to be imaged with *Chandra*. The detection rate of X-ray counterparts of the high- $z$  radio jets (2/4) is similar to that of lower- $z$  radio jet samples (Sambruna et al. 2004; Marshall et al. 2005). The implications of these observations for models of X-ray emission from large-scale jets will be described in forthcoming publications.

**Acknowledgments.** The National Radio Astronomy Observatory is operated by Associated Universities, Inc. under a cooperative agreement with the National Science Foundation. This research was funded in part by NASA through contract NAS8-39073 (A. S., D. E. H., D. A. S.) and *Chandra* Award Numbers GO7-8114 (C. C. C., L. S., J. F. C. W., D. G.) issued by the *Chandra* X-Ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS8-39073. Radio astronomy at Brandeis University is supported by the NSF and NASA. L. S. is supported by MEiN through research project 1-P03D-003-29 from 2005-2008.

**References**

- Bassett, L. C., et al. 2004, *AJ*, 128, 523
- Cheung, C.C. 2004, *ApJ*, 600, L23
- Cheung, C.C., Wardle, J.F.C., & Lee, N.P. 2005, in 22nd Texas Symposium on Relativistic Astrophysics, Eds. P. Chen et al., (Palo Alto: SLAC) 1613
- Cheung, C.C., Stawarz, L., & Siemiginowska, A. 2006, *ApJ*, 650, 679
- De Young, D. 2006, *Astron. Nachr.*, 327, 231
- Ellison, S. L., Yan, L., Hook, I. M., Pettini, M., Wall, J. V., & Shaver, P. 2001, *A&A*, 379, 393
- Harris, D.E., Krawczynski, H. 2006, *ARAA*, 44, 463
- Hook, I.M., & McMahon, R.G. 1998, *MNRAS*, 294, L7
- Hook, I.M., McMahon, R.G., Shaver, P.A., & Snellen, I.A.G. 2002, *A&A*, 391, 509
- Kataoka, J., & Stawarz, L. 2005, *ApJ*, 622, 797
- Lee, N. P. 2005, Undergraduate Honors Thesis, Brandeis University
- Liu, F. K., & Zhang, Y. H. 2002, *A&A*, 381, 757
- Lopez, L. A., Brandt, W. N., Vignali, C., Schneider, D. P., Chartas, G., & Garmire, G. P. 2006, *AJ*, 131, 1914
- Marshall, H. L., et al. 2005, *ApJS*, 156, 13
- Sambruna, R. M., Gambill, J. K., Maraschi, L., Tavecchio, F., Cerutti, R., Cheung, C. C., Urry, C. M., & Chartas, G. 2004, *ApJ*, 608, 698
- Schwartz, D.A. 2002, *ApJ*, 569, L23
- Siemiginowska, A., Smith, R.K., Aldcroft, T.L., Schwartz, D.A., Paerels, F., & Petric, A.O. 2003, *ApJL*, 598, L15
- Vermeulen, R. C., Taylor, G. B., Readhead, A. C. S., & Browne, I. W. A. 1996, *AJ*, 111, 1013
- Villani, D., & di Serego Alighieri, S. 1999, *A&AS*, 135, 299
- Xu, W., Readhead, A.C.S., Pearson, T.J., Polatidis, A.G., & Wilkinson, P.N. 1995, *ApJS*, 99, 297
- Yuan, W., Fabian, A.C., Celotti, A., & Jonker, P.G. 2003, *MNRAS*, 346, L7