

LA-UR- 02-7611

Approved for public release;  
distribution is unlimited.

Title: Electronic Structure of Pu and Pu Compounds

Author(s): John J. Joyce, 108919, MST-10

Submitted to: APS March Meeting  
March 3-7, 2003  
Austin, TX



## Los Alamos

NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# Electronic Structure of Pu and Pu Compounds

*J.J. Joyce, J.M. Wills, T. Durakiewicz, M.T. Butterfield, E. Guzewicz, J.L. Sarrao, L.A. Morales, D.P. Moore, A.J. Arko (Los Alamos National Laboratory)*

We report the electronic structure of Pu metal and several Pu compounds including materials with magnetic (PuSb<sub>2</sub>, Pu<sub>2</sub>RhGa<sub>8</sub>) and superconducting (PuCoGa<sub>5</sub>) transitions. The materials are experimentally determined by photoelectron spectroscopy and computationally by the mixed level model. The photoemission measurements are generally conducted at low temperatures 10-80 K and experimental resolutions between 35 to 75 meV. The electronic structure calculations are a mixed level model which is an extension of the generalized gradient approximation (GGA) within density functional theory. As determined both experimentally and computationally, the 5f electrons of plutonium exhibit two different configurations, one localized and the other hybridized. The dual nature of the Pu 5f electrons clearly indicates the boundary between localized and itinerant in the actinide series. Interestingly, the Pu 5f electrons exhibit this dual behavior over a wide ligand and crystal structure range. Limiting cases are observed on the localized side of 5f characteristics and some attribute the characteristics of alpha phase Pu metal to nearly itinerant 5f electrons. The successes and limitations of this Pu research will be discussed.

**Presentation type:** Oral

**Sorting category:** 9. Phase Transitions and Strongly Correlated Systems

**Submitter:** John Joyce

**Submitting Member ID:** JO973590

**submitting member email address:** jjoyce@LANL.gov

**Submitting Member affiliation:** Los Alamos National Lab

**Email address[1]:** jjoyce@LANL.gov

**CategoryType:** E

**Subtitle:** Work supported by the Department of Energy