

SRNL-STI-2009-00821
Revision 0

DETERMINATION OF REPORTABLE RADIONUCLIDES FOR DWPF SLUDGE BATCH 5 (MACROBATCH 6)

C. J. Bannochie
N. E. Bibler
D. P. DiPrete

February 2010

Savannah River National Laboratory
Savannah River Nuclear Solutions
Aiken, SC 29808

Prepared for the U.S. Department of Energy Under
Contract Number DE-AC09-08SR22470



DISCLAIMER

This work was prepared under an agreement with and funded by the U.S. Government. Neither the U. S. Government or its employees, nor any of its contractors, subcontractors or their employees, makes any express or implied:

- 1. warranty or assumes any legal liability for the accuracy, completeness, or for the use or results of such use of any information, product, or process disclosed; or**
- 2. representation that such use or results of such use would not infringe privately owned rights; or**
- 3. endorsement or recommendation of any specifically identified commercial product, process, or service.**

Any views and opinions of authors expressed in this work do not necessarily state or reflect those of the United States Government, or its contractors, or subcontractors.

Printed in the United States of America

**Prepared For
U.S. Department of Energy**

Key Words: *DWPF, Sludge Batch 5,
WAPS, Radionuclides*

Retention: PERMANENT

**DETERMINATION OF REPORTABLE
RADIONUCLIDES FOR DWPF SLUDGE BATCH 5
(MACROBATCH 6)**

C. J. Bannochie
N. E. Bibler
D. P. DiPrete

February 2010

Savannah River National Laboratory
Savannah River Nuclear Solutions
Aiken, SC 29808

Prepared for the U.S. Department of Energy Under
Contract Number DE-AC09-08SR22470



REVIEWS AND APPROVALS

AUTHORS:

C. J. Bannochie, Process Technology Programs Date

N. E. Bibler, Process Technology Programs Date

D. P. DiPrete, Analytical Development Date

TECHNICAL REVIEWER:

R. S. Lee, Analytical Development Date

APPROVERS

C. C. Herman, Manager, Process Technology Programs Date

S. L. Marra, Manager, E&CPT Research Programs Date

J. E. Occhipinti, Manager, Waste Solidification Engineering Date

TABLE OF CONTENTS

LIST OF TABLES	v
LIST OF ACRONYMS AND ABBREVIATIONS	vi
1.0 EXECUTIVE SUMMARY	1
2.0 INTRODUCTION	3
2.1 Background	3
2.2 Radionuclides Identified For Consideration as Reportable	3
3.0 EXPERIMENTAL	5
3.1 Sample Adjustments	5
3.2 Direct Methods	5
3.2.1 ICP-MS	5
3.2.2 Gamma Counting	5
3.2.3 Liquid Scintillation Counting	5
3.3 Separation Methods	6
3.3.1 Cl-36 Method	6
3.3.2 Ni-59/-63 Method	6
3.3.3 Se-79 Method	6
3.3.4 Sr-90 Method	6
3.3.5 Gamma Counting Following Cs-137 Removal	7
3.3.6 I-129 Method	7
3.3.7 Pu-238/-241 Method	8
3.3.8 Am/Cm Method	8
3.3.9 Sm-151/Pm-147 Method	8
3.4 Calculated Activities of the Remaining Radionuclides	9
3.4.1 Nb-93m	9
3.4.2 Pd-107	9
3.4.3 Cd-113m	9
3.4.4 Cs-135	11
4.0 RESULTS AND DISCUSSION	12
4.1 Sample Solids and Densities	12
4.2 Summary of the Activities and Radionuclides for Input	12
4.3 Identification of Reportable Radionuclides	14
4.4 The Ratio by Weight of U and Pu Isotopes	16
5.0 CONCLUSIONS	18
6.0 RECOMMENDATIONS	19
7.0 REFERENCES	20
8.0 ACKNOWLEDGEMENTS	22
APPENDIX A. Activities of Dried Sludge in Year 2015 ($\mu\text{Ci/g}$)	23
APPENDIX B. Activities of Dried Sludge in Year 2115 ($\mu\text{Ci/g}$)	25
APPENDIX C. Activities of Dried Sludge in Year 2215 ($\mu\text{Ci/g}$)	27
APPENDIX D. Activities of Dried Sludge in Year 2315 ($\mu\text{Ci/g}$)	29
APPENDIX E. Activities of Dried Sludge in Year 2415 ($\mu\text{Ci/g}$)	31
APPENDIX F. Activities of Dried Sludge in Year 2515 ($\mu\text{Ci/g}$)	33
APPENDIX G. Activities of Dried Sludge in Year 2615 ($\mu\text{Ci/g}$)	35

APPENDIX H. Activities of Dried Sludge in Year 2715 ($\mu\text{Ci/g}$)	37
APPENDIX I. Activities of Dried Sludge in Year 2815 ($\mu\text{Ci/g}$)	39
APPENDIX J. Activities of Dried Sludge in Year 2915 ($\mu\text{Ci/g}$)	41
APPENDIX K. Activities of Dried Sludge in Year 3015 ($\mu\text{Ci/g}$)	43
APPENDIX L. Activities of Dried Sludge in Year 3115 ($\mu\text{Ci/g}$)	45

LIST OF TABLES

Table 2-1. List of Radioisotopes Considered for Sludge Batch 5.....	4
Table 2-2. Radioisotopes Excluded for Determination of Reportable Radioisotopes for Sludge Batch 5.....	4
Table 3-1. Calculated Values of the Fission Yield Scaling Factor (FYSF) for Fourteen U-235 Fission Products in SB5.....	10
Table 4-1. Weight Percent Solids and Densities for Tank 40 SB5 Sample Before and After Simulated Tank Farm Adjustments and H-Canyon Np Additions [Number of Samples Included in Average].....	12
Table 4-2. List of Radionuclides and Activities Used as Input to the <i>RadDecay</i> [®] Program.	13
Table 4-3. Reportable Radionuclides in DWPF Sludge Batch 5.....	15
Table 4-4. Reportable Radionuclides in DWPF Sludge Batch 1B through 4.....	16
Table 4-5. Uranium Isotope Distribution in DWPF Sludge Batch 5.....	17
Table 4-6. Plutonium Isotope Distribution in DWPF Sludge Batch 5.....	17

LIST OF ACRONYMS AND ABBREVIATIONS

AD	Analytical Development
ASP	Analytical Study Plan
DI	Deionized
DOE	Department of Energy
dpm	disintegrations per minute
DWPF	Defense Waste Processing Facility
FYSF	Fission Yield Scaling Factor
g	gram
HLW	High Level Waste
ICP-AES	Inductively Coupled Plasma – Atomic Emission Spectroscopy
ICP-MS	Inductively Coupled Plasma – Mass Spectrometry
MB2	Macrobatch 2
MB3	Macrobatch 3
MB4	Macrobatch 4
MB5	Macrobatch 5
MB6	Macrobatch 6
μCi	micro-Curies
PHA	Pulse Height Analysis
QA	Quality Assurance
SB1B	Sludge Batch 1B
SB2	Sludge Batch 2
SB3	Sludge Batch 3
SB4	Sludge Batch 4
SB5	Sludge Batch 5
SCO	SRNL Shielded Cells Operations
SpA	Specific Activity (Ci/g)
SRNL	Savannah River National Laboratory
SRS	Savannah River Site
$t_{1/2}$	half-life
TTQAP	Task Technical and Quality Assurance Plan
TTR	Technical Task Request
WAPS	Waste Acceptance Product Specifications
WCP	Waste Form Compliance Plan
wt%	Weight Percent
WQR	Waste Form Qualification Report

1.0 EXECUTIVE SUMMARY

The Waste Acceptance Product Specifications (WAPS)¹ 1.2 require that “The Producer shall report the inventory of radionuclides (in Curies) that have half-lives longer than 10 years and that are, or will be, present in concentrations greater than 0.05 percent of the total inventory for each waste type indexed to the years 2015 and 3115”. As part of the strategy to comply with WAPS 1.2, the Defense Waste Processing Facility (DWPF) will report for each waste type, all radionuclides (with half-lives greater than 10 years) that have concentrations greater than 0.01 percent of the total inventory from time of production through the 1100 year period from 2015 through 3115². The initial listing of radionuclides to be included is based on the design-basis glass as identified in the Waste Form Compliance Plan (WCP)² and Waste Form Qualification Report (WQR)³. However, it is required that this list be expanded if other radionuclides with half-lives greater than 10 years are identified that may meet the greater than 0.01% criterion for Curie content.

Specification 1.6 of the WAPS, International Atomic Energy Agency (IAEA) Safeguards Reporting for High Level Waste (HLW), requires that the ratio by weights of the following uranium and plutonium isotopes be reported: U-233, U-234, U-235, U-236, U-238, Pu-238, Pu-239, Pu-240, Pu-241, and Pu-242. Therefore, the complete set of reportable radionuclides must also include this set of U and Pu isotopes.

The DWPF is receiving radioactive sludge slurry from HLW Tank 40. The radioactive sludge slurry in Tank 40 is a blend of the heel from Tank 40 (Sludge Batch 4 (SB4)), Sludge Batch 5 (SB5) that was transferred to Tank 40 from Tank 51, and H-Canyon Np transfers completed after the start of processing. The blend of sludge in Tank 40 is also referred to as Macrobatches 6 (MB6). This report develops the list of reportable radionuclides and associated activities and determines the radionuclide activities as a function of time. The DWPF will use this list and the activities as one of the inputs for the development of the Production Records that relate to the radionuclide inventory.

This work was initiated through Technical Task Request (TTR) HLW-DWPF-TTR-2008-0010; Rev. 2 entitled *Sludge Batch 5 SRNL Shielded Cells Testing*⁴. Specifically, this report details results from performing Subtask II, 5 of the TTR and, in part, meets Deliverable 7 of the TTR. The work was performed following the Task Technical and Quality Assurance Plan (TTQAP), WSRC-RP-2008-00137, Rev. 2⁵ and Analytical Study Plan (ASP), WSRC-RP-2008-00138, Rev. 2⁶.

In order to determine the reportable radionuclides for SB5 (MB6), a list of radioisotopes that may meet the criteria as specified by the Department of Energy’s (DOE) WAPS was developed. All radioactive U-235 fission products and all radioactive activation products that could be in the SRS HLW were considered. In addition, all U and Pu isotopes identified in WAPS 1.6 were included in the list. This list was then evaluated and some isotopes excluded from the projection calculations.

Based on measurements and analytical detection limits, twenty-six radionuclides have been identified as reportable for DWPF SB5 as specified by WAPS 1.2. The 26 reportable radionuclides are:

Cl-36	Ni-59	Ni-63	Sr-90	Zr-93	Nb-93m
Tc-99	Sn-126	Cs-137	Sm-151	U-233	U-234
Np-237	U-238	Pu-238	Pu-239	Pu-240	Am-241
Pu-241	Pu-242	Am-242m	Am-243	Cm-244	Cm-245
Cm-246	Cf-251				

Chlorine-36 is reported for the first time based on the upper bounding activity determined from the aqua regia digested sludge slurry.

The WCP and WQR require that all of radionuclides present in the Design Basis glass be considered as the initial set of reportable radionuclides. For SB5 (MB6), all of the radionuclides in the Design Basis glass are reportable except for four radionuclides: Se-79, Pd-107, Cs-135, and Th-230. At no time through the year 3115 did any of these three radionuclides contribute to more than 0.01% of the radioactivity on a Curie basis.

Two additional uranium isotopes (U-235 and -236) must be added to the list of reportable radionuclides in order to meet WAPS 1.6. All of the Pu isotopes and other U isotopes (U-233, -234, and -238) identified in WAPS 1.6 were already determined to be reportable according to WAPS 1.2 This brings the total number of reportable radionuclides for SB5 to 28.

The radionuclide measurements made for SB5 are the most extensive conducted to date. Some method development/refinement occurred during the conduct of these measurements, leading to lower detection limits and more accurate measurement of some isotopes than was previously possible. Continuous improvement in the analytical measurements will likely continue, and this in turn should lead to improved detection limit values for some radionuclides and actual measurements for still others.

2.0 INTRODUCTION

2.1 Background

The Waste Acceptance Product Specifications (WAPS)¹ 1.2 require that “The Producer shall report the inventory of radionuclides (in Curies) that have half-lives longer than 10 years and that are, or will be, present in concentrations greater than 0.05 percent of the total inventory for each waste type indexed to the years 2015 and 3115.” As part of the strategy to comply with WAPS 1.2, the Defense Waste Processing Facility (DWPF) will report for each waste type, all radionuclides (with half-lives greater than 10 years) that have concentrations greater than 0.01 percent of the total inventory from time of production through the 1100 year period from 2015 through 3115². The initial listing of radionuclides to be included is based on the design-basis glass as identified in the Waste Form Compliance Plan (WCP)² and Waste Form Qualification Reports (WQR)³. However, it is required that this list be expanded if other radionuclides with half-lives greater than 10 years are identified that may meet the greater than 0.01% criterion for Curie content.

Specification 1.6 of the WAPS, International Atomic Energy Agency (IAEA) Safeguards Reporting for High Level Waste (HLW), requires that the ratio by weights of the following uranium and plutonium isotopes be reported: U-233, U-234, U-235, U-236, U-238, Pu-238, Pu-239, Pu-240, Pu-241, and Pu-242. Therefore, the complete set of reportable radionuclides must also include this set of U and Pu isotopes.

The DWPF is receiving radioactive sludge slurry from HLW Tank 40. The radioactive sludge slurry in Tank 40 is a blend of the heel from Sludge Batch 4 (SB4), Sludge Batch 5 (SB5) that was transferred to Tank 40 from Tank 51, and H-Canyon Np transfers completed after the start of processing. The blend of sludge in Tank 40 is also referred to as Macrobatches 6 (MB6). This report develops the list of reportable radionuclides and associated activities and determines the radionuclide activities as a function of time. The DWPF will use this list and the activities as one of the inputs for the development of the Production Records that relate to radionuclide inventory.

This work was initiated through Technical Task Request (TTR) HLW-DWPF-TTR-2008-0010, Rev. 2 entitled *Sludge Batch 5 SRNL Shielded Cells Testing*⁴. Specifically, this report details results from performing Subtask II, 5 of the TTR and, in part, meets Deliverable 7 of the TTR. The work was performed following the Task Technical and Quality Assurance Plan (TTQAP), WSRC-RP-2008-00137, Rev. 2⁵ and Analytical Study Plan (ASP), WSRC-RP-2008-00138, Rev. 2⁶.

2.2 Radionuclides Identified For Consideration as Reportable

In order to determine the reportable radionuclides for Sludge Batch 5, a list of radioisotopes that may meet the criteria as specified by the Department of Energy’s (DOE) WAPS was developed. All radioactive U-235 fission products and all radioactive activation products that could be in the SRS HLW were considered. New to the list in Table 2-1 is one isotope recently identified: Cl-36. In addition, all U and Pu isotopes identified in WAPS 1.6 have been included in this list.

Table 2-1 presents the list of radioisotopes identified for consideration as reportable. The radioisotopes that were deleted from the list and the arguments that support that decision are presented in Table 2-2.

Table 2-1. List of Radioisotopes Considered for Sludge Batch 5

Radioisotope	Radioisotope	Radioisotope	Radioisotope
C-14 ^a	Sn-121m ^b	Pb-210 ^d	Pu-240 ^a
Cl-36 ^a	Sn-126 ^b	Ra-226 ^d	Pu-241 ^a
Ni-59 ^a	I-129 ^b	Ac-227 ^d	Am-241 ^a
Co-60 ^a	Ba-133 ^b	Th-229 ^d	Pu-242 ^a
Ni-63 ^a	Cs-135 ^b	Th-230 ^d	Am-242m ^a
Se-79 ^b	Cs-137 ^b	Pa-231 ^d	Am-243 ^a
Rb-87 ^b	La-138 ^b	Th-232 ^c	Cm-243 ^a
Sr-90 ^b	Ce-142 ^{b,c}	U-232 ^a	Cm-244 ^a
Zr-93 ^b	Nd-144 ^b	U-233 ^a	Cm-245 ^a
Nb-93m ^b	Pm-147 ^b	U-234 ^d	Cm-246 ^a
Nb-94 ^b	Sm-147 ^b	U-235 ^c	Cm-247 ^a
Zr-96 ^b	Sm-149 ^b	U-236 ^a	Bk-247 ^a
Tc-99 ^b	Nd-150 ^b	Np-236 ^a	Cm-248 ^a
Cd-113 ^{b,c}	Sm-151 ^b	Np-237 ^a	Cf-249 ^a
Pd-107 ^b	Eu-152 ^b	U-238 ^c	Cf-250 ^a
Cd-113m ^b	Eu-154 ^b	Pu-238 ^a	Cf-251 ^a
In-115 ^b	Bi-210m ^d	Pu-239 ^a	

^a Activation product

^b Fission product

^c Naturally occurring radionuclide that resulted in the waste from processing at SRS

^d Decay product of an actinide isotope in SRS waste

Table 2-2. Radioisotopes Excluded for Determination of Reportable Radioisotopes for Sludge Batch 5

Radioisotope	Radioisotope
C-14 ⁱ	Nd-144 ⁱⁱⁱ
Nb-94 ⁱⁱ	Nd-150 ⁱⁱⁱ
Zr-96 ⁱⁱⁱ	Sm-147 ⁱⁱⁱ
Cd-113 ⁱⁱⁱ	Sm-149 ⁱⁱⁱ
In-115 ⁱⁱⁱ	Eu-152 ⁱⁱ
La-138 ⁱⁱⁱ	Np-236 ^{iv}
Ce-142 ⁱⁱⁱ	U-232 ^v

ⁱ C-14 is volatilized during DWPF Chemical and Melt Cell processing and is not immobilized in the glass⁷.

ⁱⁱ “Nb-94 and Eu-152 are shielded isotopes: the isobaric fission product decay chain for these stops at a stable isotope before reaching these. They are therefore produced predominately by secondary processes and are present only in very small amounts. They have not been observed in the sludge slurry”⁸.

ⁱⁱⁱ Zr-96, Cd-113, In-115, La-138, Ce-142, Nd-144, Nd-150, Sm-147 and Sm-149 were deleted because their long half-lives (> 1.05E11 years) make their activities negligible at all times⁸.

^{iv} “No data was available for Np-236 but it is known to be made in only very small amounts in reactor irradiation. Np-236 is a minor product of fast neutron spallation. It was neglected”⁸.

^v “U-232 is present only in very small amounts and decays rapidly compared to other actinide isotopes that are much more abundant (it is primarily obtained as a contaminant at a few ppm from the reactor irradiation of Th-232)”⁸.

3.0 EXPERIMENTAL

The details for sample acquisition from Tank 40 – SB5, preparation of the digestions, and measurement of the elemental composition have been published separately⁹. At the time the Tank 40 sample was pulled, the final supernate decant, NaNO₂ corrosion inhibitor addition, and the H-Canyon Np transfers had not yet been conducted. Those activities were reproduced on the Tank 40 sample at SRNL after they were completed in the Tank Farm so that they could be simulated accurately¹⁰. The adjusted sample of Tank 40 was then digested again as described previously⁹. The new solids and density measurements are reported here. The results for those radionuclides that required additional separation techniques, that were not included in the referenced report, have been included in this report. Details of the separation and detection methods are provided. All measurements and counting analyses were done by SRNL Analytical Development (AD).

3.1 Sample Adjustments

A 998 g subsample of Tank 40 material was adjusted to simulate the Tank Farm operations conducted since the acquisition of the original sample for SRNL. A decant of 187 g of settled sludge supernate was removed from the sample. 22.95 g of 40 wt% NaNO₂ (Tank Farm corrosion inhibitor) was added to the decanted sample. In a separate container 6.56 g of H-Canyon Np and 10.94 g of deionized water (DI H₂O) were “neutralized” with 4.76 g of 50 wt% NaOH. The adjusted Np sample was added to the Tank 40 subsample with 21.97 g of DI H₂O to rinse the contents of the Np adjustment container. The final adjusted sludge was mixed thoroughly¹⁰.

3.2 Direct Methods

3.2.1 ICP-MS

Inductively coupled plasma – mass spectrometry (ICP-MS) was employed to analyze separate sub-samples of the aqua regia digestions of the adjusted Tank 40 – SB5 dried solids described in a previous report⁹. Measurements were first converted to weight percents on a dried solids basis and then converted to activities using the specific activity of each isotope taken from References 11 and 18. The isotopes obtained from direct ICP-MS measurements included: Zr-93, Tc-99, Sm-151, Th-232, U-233, U-234, U-235, U-236, Np-237, U-238, Pu-239, Pu-240, and Pu-242.

3.2.2 Gamma Counting

Gamma Pulse Height Analysis (PHA) was performed on separate sub-samples of the alkali fusion (also known as “peroxide fusion”) digestions of the adjusted Tank 40 – SB5 dried solids described in a previous report⁹. Detectors were carefully calibrated with known standards. Since detection efficiencies for gamma-rays vary with energy, they were determined for these specific radionuclide energies during the calibration process. The counting geometry was established during calibration and carefully duplicated for these measurements. Samples were diluted as necessary to achieve accurate counting. The isotopes obtained from Gamma PHA counting included: Co-60, Cs-137, Eu-154, and Eu-155.

3.2.3 Liquid Scintillation Counting

Liquid scintillation counting was performed on separate sub-samples of the alkali fusion digestions of the adjusted Tank 40 – SB5 dried solids described in a previous report⁹. The scintillation cocktail used for the analysis was Ultima Gold AB since it is specifically formulated for alpha-beta discrimination and is the best choice for samples dissolved in mineral acids. Measurements were performed on one of three Packard Instruments which automatically correct for quenching and many other interference problems commonly associated with liquid scintillation counting. This method was used to determine total Beta activity. Diluted aliquots of digested slurry were analyzed for 10 minutes. Diluted aliquots of digested

slurry were also spiked with an alpha standard and analyzed for 10 minutes to measure and correct for any alpha/beta discrimination cross-talk issues.

3.3 Separation Methods

These analytical methods involved separation techniques that enabled radionuclides that were at low concentrations to be measured more accurately and to determine more reliable and lower detection limits. These techniques are now SRNL AD procedures and will only be summarized here. Aliquots of the alkali fusions or the aqua regia dissolutions were analyzed along with blanks. In all cases, the activity in the blanks did not contribute any significant activity to the radionuclides being analyzed.

3.3.1 Cl-36 Method

Aliquots from the pressurized aqua-regia dissolution of Sludge Batch 5 slurry were diluted and treated with a variety of cation exchange and chromatographic resins to decontaminate interfering beta-emitting isotopes from Cl-36 in the sample. Aliquots from the decontaminated solutions were then analyzed by liquid scintillation analysis to determine a gross beta activity of the treated solution. This residual beta activity was used as an upper bounding activity for the Cl-36 in the Sludge Batch 5 slurry.

3.3.2 Ni-59/-63 Method

This separation is based on isolation of Ni from the dissolved sludge using a column containing dimethylglyoxime as an extractant that is specific for Ni. Each of the solutions resulting from the alkali fusions of the four samples of dried sludge slurry was spiked with a stable Ni carrier to trace the Ni separation and was then passed through a column containing the above extractant. The absorbed Ni was then eluted from each column. The Ni-59 was measured in the eluted solutions by its characteristic X-rays and Ni-63 by its beta particles. Total Ni in each eluted solution was measured by inductively coupled plasma atomic emission spectroscopy (ICP-AES). The radiochemical Ni analyses were corrected for the Ni carrier recoveries as measured by ICP-AES.

3.3.3 Se-79 Method

Four aliquots of wet sludge slurry were spiked with a known amount of stable Se as a carrier. The samples were digested with concentrated nitric acid. The Fe in the dissolutions was reduced to Fe (II) using ascorbic acid to ensure it would not interfere with subsequent decontamination steps designed to extract Y-90, the lanthanides and the actinides from the Se traced dissolutions. The dissolutions were then treated with resins (Bio-Rad AMP-1, Eichrom Sr, RE, and Actinide resins) to reduce levels of Sr-90, Cs-137, Y-90, the lanthanides and the actinides to levels low enough to allow for their removal from the Shielded Cells. The Se traced dissolutions were then further decontaminated with a Bio-Rad AMP-1, Eichrom Sr and RE resin treatment. The total Se was reduced to Se metal using titanium (III) chloride, hydroxylamine hydrochloride, and ascorbic acid. The precipitated Se metal was then washed repeatedly with deionized water and dilute nitric acid. The Se metal was then dissolved with concentrated HBr, and the resulting SeBr_4 was extracted by solvent-solvent extraction using a tri-butyl phosphate/n-paraffin solvent extraction system. The Se was back extracted from the solvent. Aliquots of the purified Se fraction were then analyzed. A portion was neutron activated in a Cf-252 neutron source at SRNL to determine the total amount of Se present in order to calculate the recovery of Se from the radiochemical separation. A second portion was counted by liquid scintillation to determine the Se-79 beta activity. The yields of the stable Se carrier were applied to the Se-79 beta activity result to determine Se-79 activities in the sample aliquots initially treated.

3.3.4 Sr-90 Method

Aliquots of each sample from the alkali fusions⁹ were spiked with a stable Sr carrier, and a stable Ce carrier. The Sr carrier was used for separation yielding purposes and the Ce carrier was used to enhance the separation rates of undesirable isotopes such as Y-90, the lanthanides or the actinides. The spiked sample aliquots were initially oxidized using nitric acid. The Sr in the samples was extracted using

commercially available Sr extraction resin. This resin also extracts some of the Pu under the conditions used to extract the Sr. The Pu was washed from the resin using an oxalic acid/nitric acid mixture. The Sr was eluted from the resin, and the resulting solution concentrated. A portion of the purified Sr solution was neutron activated in a Cf-252 neutron activation facility at SRNL to determine the total Sr and in order to calculate the fraction of Sr isolated by the procedure. A second portion of each of the Sr fractions was stored for five to seven days to allow Y-90 to grow in. Each fraction was then counted by liquid scintillation analysis to determine the Y-90 activity. The Sr-90 beta activity in each case was calculated from the Y-90 activity. The yields of the stable Sr carriers were applied to the Sr-90 beta activity results to determine Sr-90 activities in the original aliquots of the solutions resulting from the dissolution of the dried sludge slurry samples.

3.3.5 Gamma Counting Following Cs-137 Removal

This method was used to determine Ru-106, Sn-121m, Sb-125, Sn-126, Ba-133, Ce-144, Eu-154, Eu-155, and Am-241. These gamma emitters could not be determined directly because of the high Cs-137 activity in the samples. Consequently the Cs-137 was removed. Aliquots of each of the four alkali fusions⁹ of the dried sludge slurry samples were treated with two batch additions of an ammonium phosphomolybdate resin to selectively remove the Cs-137 from the aliquots. This allowed gammas for isotopes at low concentrations to be detected or allowed lower detection limits to be determined for those isotopes that were not detected. The Cs-137 decontaminated aliquots were then gamma counted in two different detectors. The first was a high purity coaxial germanium detector to detect the gamma rays from Ru-106, Sb-125, Sn-126, Ba-133, Ce-144, Eu-154, Eu-155, and Am-241. Only Eu-154, Eu-155, and Am-241 were detected. Because of their low concentrations the other isotopes were not detected. To obtain reliable and lower detection limits for these radionuclides, each of the solutions was counted for four hours. The detection limits were used to calculate the maximum activity of each for input to the projection calculations. Of this group of radionuclides, only Sn-126 and Ba-133 have half-lives greater than 10 years. Even though the others have half-lives less than 10 years, their activities were included to calculate the total Curies present in SB5 at the selected decay times. The second detector was a Be thin window, semi-planar, high purity germanium detector. This detector has a high counting efficiency for low energy gamma rays (37.2 keV) that are used to measure Sn-121m. The Sn-121m concentration was so low that it was not detected. Again to obtain a reliable lower detection limit, each of the four aliquots was counted for 2.2 –13.9 hours.

3.3.6 I-129 Method

The radionuclide I-129 is a long-lived beta emitting fission product ($t_{1/2} = 1.6E+07$ years) that is in SRS wastes. Four aliquots of wet sludge slurry were spiked with a known amount of stable KI to act as an iodine tracer/carrier. The samples were digested with 8M nitric acid. The iodate/iodine in the samples was reduced with sodium sulfite to minimize losses of iodine in the Shielded Cells I-129 procedure. The Fe in the dissolutions was reduced to Fe (II) using ascorbic acid to ensure it would not interfere with subsequent decontamination steps designed to extract Y-90, the lanthanides and the actinides from the KI traced dissolutions. The dissolutions were then treated with resins (Bio-Rad AMP-1, Eichrom Sr, RE, and Actinide resins) to reduce levels of Sr-90, Cs-137, Y-90, the lanthanides and the actinides. The traced samples were then rendered caustic, treated a second time with a sodium sulfite reduction, and filtered to ensure Sr-90 and Y-90 levels were reduced low enough to allow for sample removal from the Shielded Cells. The samples were decontaminated a final time with a resin treatment to remove Cs-137 and the actinide elements. The solution was then treated with $AgNO_3$ in order to precipitate the iodide ion as AgI. The precipitate was analyzed by low energy gamma spectroscopy to determine the amount of I-129 present. I-129 is detected by its characteristic gamma and x-ray emissions. The precipitate was then neutron activated in a Cf-252 neutron source at SRNL to determine the total amount of iodine present in order to calculate the recovery of I-129 in the radiochemical separation.

3.3.7 Pu-238/-241 Method

Pu-241 is a beta-emitting Pu isotope that cannot be measured directly in the dissolved dried sludge slurry solutions because of its low concentration. Pu-241 has a relatively short half-life (15 years). Its concentration, along with that for Pu-238, was determined by isolating the Pu from each solution by a thenoyltrifluoroacetone extraction procedure. The extracted Pu was then analyzed by beta and alpha counting to determine the ratio of beta activity from Pu-241 to the alpha activity from the other isotopes of Pu (Pu-238, Pu-239, Pu-240, and Pu-242). In the original dissolution solutions, the total alpha activity from the Pu isotopes was determined by alpha counting and ICP-MS. Knowing the total alpha activity from Pu in the solutions resulting from the extraction allows the concentration of Pu-241 in the original dissolution solutions to be calculated using the beta/alpha ratio determined in the extracted solution. In the extracted solution, the alpha counting technique also gives the alpha counts due specifically to Pu-238 so that the total amount of Pu-238 can be determined. The activities of these two radionuclides were then used in the calculations to determine the reportable radionuclides.

3.3.8 Am/Cm Method

This method was used for Am-241, Am-242m, Cm-242, Am-243, Cm-243, Cm-244, Cm-245, Cm-246, Cm-247, Bk-247, Cm-248, Cf-249, Cf-250, and Cf-251. These radionuclides are neutron activation products produced in the SRS reactors. These isotopes are difficult to measure because of their low concentrations in the sludge slurry and the dilutions necessary to get the dissolved slurry samples out of the Shielded Cells. Of these isotopes, the Am-241 can be easily and accurately analyzed directly by long term gamma counting of the dissolved sludge (see Section 3.2.2). For the other radionuclides listed above, a separation method has been developed by AD for isolating Am, Cm, Bk and Cf from a wet sludge slurry solution. The slurry is digested in the Shielded Cells with concentrated nitric acid. The actinides are then extracted from the dissolution using a commercially available ion exchange resin (Eichrom RE). As Y-90 co-extracts with the trivalent actinides on RE resin, the treated samples were held in the Shielded Cells for nine days to allow the Y-90 to decay. The solutions were purified further with a second RE resin extraction followed by an Eichrom Ln resin extraction. The Am, Cm, Bk, and Cf extracts were then analyzed by alpha and low energy gamma counting techniques as well as by ICP-MS. The radionuclides Cm-242, Am-242m, and Cm-244 were measured by alpha spectroscopy, Am-241, Am-243, Cm-243, Cm-245, Cf-249, and Cf-251 were measured by low energy gamma spectroscopy, and Cm-246, Cm-247, Bk-247, Cm-248 and Cf-250 was measured by ICP-MS. The fraction of each actinide element isolated by this ion exchange technique was determined by comparing the measured concentrations of Am-241 in the eluted solutions with their respective concentration in the original dissolved slurry that was measured by direct gamma counting of Cs-137 removed aliquots of the dissolved slurry.

By using this technique, the radionuclides Am-243, Cm-244, Cm-246, and Cm-248 were detected and measured along with the Am-241. All the other radionuclides had concentrations below the detection limit of the analytical methods. These radionuclides were Am-242m, Cm-242, Cm-243, Cm-245, Cm-247, Bk-247, Cf-249, Cf-250, and Cf-251. For these, the detection limits were then used as the maximum concentrations or activities that could be present.

3.3.9 Sm-151/Pm-147 Method

Aliquots of each sample from the alkali fusions⁹ were spiked with a stable Sm carrier. The Sm carrier was used for separation yielding purposes. The spiked sample aliquots were initially oxidized using nitric acid. The Sm and Pm along with other trivalent species in the samples were extracted using Eichrom RE resin. The Sm and Pm were then extracted from the other radionuclides present using Eichrom Ln resin. A portion of the purified Pm/Sm solution was neutron activated in a Cf-252 neutron activation facility at SRNL to determine the total Sm and in order to calculate the fraction of Sm isolated by the procedure. A second portion of each of the Pm/Sm fractions was then counted by liquid scintillation analysis to determine the Pm-147 and Sm-151 activity. The Pm-147 measurement was

conducted using a higher energy beta window which was free of any interference from the low energy Sm-151 beta. The Sm-151 beta result is corrected for any Pm-147 events occurring in its beta counting window when necessary. The yields of the stable Sm carriers were applied to the Sm-151 and the Pm-147 beta activity results to determine Sm-151 and Pm-147 activities in the original aliquots of the solutions resulting from the dissolution of the dried sludge slurry samples. A Pm-147 spiked sample was run through the process to monitor and correct for any slight differences in the chemical recoveries of Sm and Pm.

3.4 Calculated Activities of the Remaining Radionuclides

3.4.1 Nb-93m

The radionuclide Nb-93m ($t_{1/2} = 16.1$ years)¹³ is in SRS HLW as the decay product of the radioactive fission product Zr-93 ($t_{1/2} = 1.53E+06$ years). For previous sludge batches both the Zr-93 and Nb-93m became reportable after the waste was ~100 years old^{21,22,23,24}. The concentration of Nb-93m during SB5 vitrification can be calculated if the age of the sludge and the concentration of Zr-93 are known.

The age of the waste can be calculated from the measured concentration at mass 90 that is composed of Sr-90 ($t_{1/2} = 28.5$ years) and its daughter Y-90 ($t_{1/2} = 2.67$ days) that decays to stable Zr-90. The radionuclide Sr-90 is the initial long lived radioactive product of the isobaric decay chain at mass 90. In the ICP-MS analysis of the aqua regia dissolutions of the SB5 total solids, the concentration at mass 90 which is composed of Sr-90, Y-90, and Zr-90 was 0.0231 wt%, but this value predicts a waste age of 14 years which is clearly unrealistic for SRS Tank Waste. Thus the concentration at mass 90 predicted by the fission yield scaling factor (FYSF) (see Section 3.4.3), 0.0334 wt%, was used in the calculation. The concentration of Sr-90 determined by radioactive counting was 0.0165 wt% (see Section 3.3.4). Thus the fraction of Sr-90 that remained in this sludge is 0.494. With the half-life of Sr-90 and standard equation for radioactive decay¹², the age of the sludge can be calculated. The result is 29 years.

The concentration of Zr-93 in SB5 measured by ICP-MS is 0.0208 wt%. Because of the large difference in the half-lives of Zr-93 and its radioactive daughter Nb-93m, these radionuclides are in secular equilibrium. Consequently, the initial concentration (wt%) of Nb-93m as input to the decay calculations can be calculated from the age of the waste and radioactive decay equation for two radionuclides that are in secular equilibrium as shown in Equation 5-4 on page 130 in Reference 12.

3.4.2 Pd-107

The noble metal Pd-107 is a pure beta emitter with a very long half-life ($6.5E+06$ years). This radionuclide could not be detected in the SB5 dissolved dried slurry samples by ICP-MS due to the presence of natural silver. Natural Ag contains the isotope Ag-107, which interferes with the measurement of Pd-107. The concentration of Pd-105 could be measured in the solutions thus the concentration of Pd-107 was calculated from the concentration of Pd-105. This was done by multiplying the ratio of the product of the fission yields and masses for Pd-107 and Pd-105 by the measured wt% for Pd-105 as determined by ICP-MS.

3.4.3 Cd-113m

With a half life of 13.7 years and specific activity of 217 Ci/g¹¹, Cd-113m may also qualify as a WAPS reportable radionuclide. However, Cd-113m primarily decays by β emission and thus would require a careful separation technique to measure in sludge slurry. Also, the determination of Cd-113m by ICP-MS of the dissolved dried solids is essentially impossible because of the presence of natural Cd in the sludge. Natural Cd is 12.3 % Cd-113 with a half-life greater than $1.1E11$ years¹³. This makes its activity negligible at all times⁸. Finally, the fission yield of Cd-113m is very small, $1.66E-04\%$ ¹⁴, and thus its concentration is expected to be very small in the HLW sludge.

An upper limit of the mass concentration of Cd-113m can be estimated by using the FYSF¹⁵. The FYSF relates the concentration of a fission product in the total sludge solids to the fission yield and the atomic mass of that fission product. The atomic mass of that isotope has to be included in the equation because fission yields are given in terms of atoms produced per 100 fissions of U-235 and not in terms of mass percent of the isotope produced. The equation for the concentration in weight percent is then:

$$\text{Concentration (wt\%)} = \text{FYSF (fission yield} \times \text{atomic mass)} \quad (1)$$

Thus the FYSF for each measured isotope can be calculated from the Equation 2.

$$\text{FYSF}_i = \text{wt\%}_i / (\text{fy}_i \times \text{am}_i) \quad (2)$$

Where FYSF_i \equiv the fission yield scaling factor based on isotope i
 wt\%_i \equiv the weight per percent of isotope i in the HLW total dried solids
 fy_i \equiv the fission yield of isotope i
 am_i \equiv the atomic mass of isotope i.

Several of the U-235 fission products have the six critical chemical and nuclear properties that allow calculation of a constant FYSF for a particular sludge. These properties are discussed in Reference 15. In SB5, there are 14 isotopes that have these six properties. The FYSF's calculated for these 14 isotopes are presented in the Table 3-1. The measured concentrations in the Table 3-1 were determined by ICP-MS analysis of four samples of dissolved SB5 total solids by the aqua regia method. The average of the FYSF's calculated for the 14 fission products is 7.13E-05 with a relative standard deviation of 12%.

Table 3-1. Calculated Values of the Fission Yield Scaling Factor (FYSF) for Fourteen U-235 Fission Products in SB5

Isotope	Measured		FYSF
	Wt% of Total Solids	Fission Yield	
Ru-101	3.48E-02	5.20	6.62E-05
Ru-102	3.02E-02	4.30	6.89E-05
Rh-103	1.78E-02	3.03	5.69E-05
Ru-104	1.75E-02	1.88	8.96E-05
La-139	6.36E-02	6.41	7.14E-05
Ce-140	7.01E-02	6.22	8.05E-05
Pr-141	5.57E-02	5.80	6.81E-05
Ce-142	6.03E-02	5.85	7.26E-05
Nd-143	5.39E-02	5.96	6.33E-05
Nd-144	6.04E-02	5.50	7.62E-05
Nd-145	3.87E-02	3.93	6.80E-05
Nd-146	3.25E-02	3.00	7.41E-05
Sm-147	2.03E-02	2.25	6.15E-05
Sm-148	1.99E-02	1.67	8.04E-05
Average	-	-	7.13E-05

With the FYSF for SB5, the maximum possible concentration in terms of weight percent of dried solids can be estimated for the other U-235 fission products. These estimations are a maximum for the other fission products because they do not have the necessary six properties as the fission products given in Table 3-1. For example, the radionuclide Cd-113m has a very large neutron absorption cross section

(20,000 barns¹³), and thus it was transmuted in the reactors at SRS to stable Cd-114 while the reactors were in operation. Consequently, the concentration calculated for Cd-113m with Equation 1 and a fission yield of 1.66E-04% can be considered the maximum concentration and is estimated below:

$$\text{Concentration Cd-113m (wt\%)} = 7.13\text{E-}05 \times 1.66\text{E-}04 \times 113 = 1.34\text{E-}06 \text{ wt\%}.$$

3.4.4 Cs-135

The radionuclide Cs-135 cannot be detected by ICP-MS in the sludge slurry due to its low concentration and the large amount of natural Ba-135 (same mass as the Cs-135). Cs-135 cannot be detected by counting techniques either because of its long half-life ($t_{1/2} = 2.3\text{E}+06$ years). The detection by ICP-MS of Cs-135 in the supernate is possible because Ba-135 is insoluble in caustic supernate. The same philosophy applies to Ba-137 and Cs-137. Thus, Ba does not interfere with the analyses of Cs-135 or Cs-137 in the supernate. By using the ratio of Cs-135 to Cs-137 in the supernate, and the amount of Cs-137 in the sludge slurry, the activity for Cs-135 in the sludge slurry can be calculated. The weight percent of Cs-135 was then converted to $\mu\text{Ci/g}$ and used in the calculations to determine if Cs-135 was reportable.

4.0 RESULTS AND DISCUSSION

4.1 Sample Solids and Densities

Table 4-1 presents the measured SB5 density and weight percent solids data¹⁶ measured for the SB5 WAPS sample after it was collected in December 2008 as well as the data for this sample after adjustment for Tank Farm operations (decant and corrosion inhibitor addition) and H-Canyon Np transfers.

Table 4-1. Weight Percent Solids and Densities for Tank 40 SB5 Sample Before and After Simulated Tank Farm Adjustments and H-Canyon Np Additions
[Number of Samples Included in Average]

Property	Tank 40 – WAPS w/o Np (% RSD) ⁹	Tank 40 – WAPS w/ Np (% RSD)
Slurry Density	1.094 (0.6) [4]	1.138 (1.8) [5]
Supernate Density	1.043 (1.0) [4]	1.041 (0.4) [4]
Wt % Total Solids	13.36 (0.6) [4]	14.28 (0.6) [4]
Wt % Calcined Solids	10.52 (0.4) [4]	10.15 (1.8) [4]
Wt % Dissolved Solids ^a	4.29 (0.9) [4]	5.14 (0.8) [4]
Wt % Insoluble Solids	9.48	9.63
Wt % Soluble Solids ^b	3.88	4.65

^a Also known as Uncorrected Soluble Solids

^b Also known as Corrected Soluble Solids

4.2 Summary of the Activities and Radionuclides for Input

The complete list of radionuclides and their activities that were considered in the determination of reportable radionuclides are provided in Table 4-2. For those radionuclides with measured concentrations, the initial activities were calculated by using the weight percent reported for each radioisotope and its specific activity with the following equation: $A_o = M_o * SpA$, where A_o = Initial Activity, M_o = mass in weight percent and SpA = specific activity of the isotope.

For each radionuclide listed in Table 4-2 there is an associated specific activity in units of Ci/g, wt% of total solids, activity in $\mu\text{Ci/g}$, activity in Ci/gal and the method used to determine or estimate the value.

The total measured alpha activity of the digested samples was $<423 \mu\text{Ci/g}$ of total dried solids. Total measured beta activity was measured at $3.29\text{E}+04 \mu\text{Ci/g}$ of total dried solids.

Table 4-2. List of Radionuclides and Activities Used as Input to the RadDecay® Program

Radionuclide	Specific Activity (Ci/g)	Wt% of Total Solids	Activity (μ Ci/g)‡	Activity (Ci/gal)	Method
Cl-36	3.30E-02	<2.95E-03	<9.75E-01	<6.00E-04	Cl-36
Ni-59	8.08E-02	1.61E-03	1.30E+00	8.02E-04	Ni-59/-63
Co-60	1.13E+03	2.12E-07	2.40E+00	1.47E-03	Direct Gamma Counting
Ni-63	6.17E+01	1.80E-04	1.11E+02	6.82E-02	Ni-59/-63
Se-79	6.97E-02	<3.99E-06	<2.78E-03	<1.71E-06	Se-79
Sr-90	1.36E+02	1.65E-02	2.25E+04	1.38E+01	Sr-90
Y-90*	5.44E+05	4.14E-06	2.25E+04	1.38E+01	Secular equilibrium w/ Sr-90
Zr-93	2.51E-03	2.08E-02	5.23E-01	3.22E-04	ICP-MS
Nb-93m	2.83E+02	1.39E-07	3.93E-01	2.42E-04	Calculated from Zr-93 and Waste Age
Tc-99	1.70E-02	9.04E-04	1.53E-01	9.42E-05	ICP-MS
Pd-107	5.14E-04	2.50E-04	1.29E-03	7.91E-07	Calculated from Pd-105
Cd-113m	2.17E+02	1.34E-06	2.91E+00	1.79E-03	Based on Fission Yield
Sn-121m	5.91E+01	<5.16E-07	<3.05E-01	<1.88E-04	Cs-Removed Gamma Counting
Sb-125	1.03E+03	<9.60E-09	<9.90E-02	<6.09E-05	Cs-Removed Gamma Counting
Te-125m*	1.80E+04	<5.50E-10	<9.90E-02	<6.09E-05	Secular equilibrium w/ Sb-125
Sn-126	2.84E-02	<1.27E-03	<3.60E-01	<2.21E-04	Cs-Removed Gamma Counting
I-129	1.77E-04	<2.14E-04	<3.77E-04	<2.32E-07	I-129
Ba-133	2.50E+02	<3.37E-08	<8.42E-02	<5.18E-05	Direct Gamma Counting
Cs-135	1.15E-03	1.40E-04	1.61E-03	9.93E-07	Calculated from Cs-135/-137 ratio
Cs-137	8.70E+01	3.71E-04	3.23E+02	1.98E-01	Direct Gamma Counting
Ba-137m*	5.38E+08	5.76E-11	3.10E+02	1.90E-01	Secular equilibrium w/ Cs-137
Pm-147	9.27E+02	<1.72E-05	<1.60E+02	<9.82E-02	Pm-147/Sm-151
Sm-151	2.63E+01	6.91E-04	1.82E+02	1.12E-01	Pm-147/Sm-151
Eu-154	2.70E+02	8.62E-06	2.33E+01	1.43E-02	Direct Gamma Counting
Eu-155	4.65E+02	5.27E-07	2.45E+00	1.51E-03	Direct Gamma Counting
Th-232	1.10E-07	6.63E-02	7.28E-05	4.48E-08	ICP-MS
U-233	9.68E-03	2.29E-04	2.22E-02	1.37E-05	ICP-MS
U-234	6.25E-03	9.85E-04	6.15E-02	3.79E-05	ICP-MS
U-235	2.16E-06	3.24E-02	7.00E-04	4.30E-07	ICP-MS
U-236	6.47E-05	2.08E-03	1.34E-03	8.26E-07	ICP-MS
Np-237	7.05E-04	9.82E-03	6.92E-02	4.26E-05	ICP-MS
U-238	3.36E-07	4.97E+00	1.67E-02	1.03E-05	ICP-MS
Pu-238	1.71E+01	1.61E-03	2.76E+02	1.70E-01	Pu-238/-241
Pu-239	6.22E-02	3.28E-02	2.04E+01	1.26E-02	ICP-MS
Pu-240	2.28E-01	2.89E-03	6.59E+00	4.05E-03	ICP-MS
Pu-241	1.03E+02	9.45E-05	9.74E+01	5.99E-02	Pu-238/-241
Pu-242	3.82E-03	1.79E-04	6.82E-03	4.19E-06	ICP-MS
Am-241	3.43E+00	9.47E-04	3.25E+01	2.00E-02	Direct Gamma Counting
Am-242m	9.72E+00	<4.67E-06	<4.54E-01	<2.79E-04	Am/Cm
Am-243	1.99E-01	4.83E-04	9.64E-01	5.93E-04	Am/Cm
Cm-242	3.31E+03	<1.14E-08	<3.75E-01	<2.31E-04	Am/Cm
Cm-243	5.16E+01	<4.69E-07	<2.42E-01	<1.49E-04	Am/Cm
Cm-244	8.09E+01	5.69E-05	4.61E+01	2.83E-02	Am/Cm
Cm-245	1.72E-01	<2.50E-05	<4.29E-02	<2.64E-05	Am/Cm
Cm-246	3.07E-01	4.19E-06	1.29E-02	7.93E-06	Am/Cm – ICP-MS
Cm-247	9.28E-05	<2.62E-07	<2.43E-07	<1.49E-10	Am/Cm – ICP-MS
Bk-247	1.03E+00	<2.63E-07	<2.71E-03	<1.66E-06	Am/Cm – ICP-MS
Cm-248	4.25E-03	8.87E-07	3.77E-05	2.32E-08	Am/Cm – ICP-MS
Cf-249	4.38E+00	<2.05E-07	<8.96E-03	<5.51E-06	Am/Cm
Cf-250	1.09E+02	<1.24E-10	<1.36E-04	<8.34E-08	Am/Cm – ICP-MS
Cf-251	1.86E+00	<1.24E-06	<2.31E-02	<1.42E-05	Am/Cm
Total			4.66E+04		

* Included because this isotope is in secular equilibrium with a parent for which a measured value was available.

‡ Less than values represent the minimum detection limit value and hence are an upper bound for that isotope's activity.

4.3 Identification of Reportable Radionuclides

Based on radionuclides and activities provided in Table 4-2, a commercially available computer program was used to identify which radionuclides were reportable through calendar year 3115. The program is called *RadDecay*[®] version 3.0¹⁷.

The initial activities for 51 isotopes were entered into *RadDecay*[®] and the results of two calculations with the index years 2015 and 3115 (1100 years) are presented in Appendix A and Appendix L. Those radionuclides that are reportable are designated in these tables by a “yes”. Additional calculations were performed for every 100 years up to 1100 years. These results are presented in Appendix B through K. Microsoft Excel spreadsheets were used to calculate the total activity in $\mu\text{Ci/g}$ of dried sludge solids at each time and the percent of the activity that each of the radionuclides contributed.

The calculations performed by *RadDecay*[®] v. 3.0 were verified against a separate program called *Radioactive Decay Calculator*¹⁸. A comparison between the outputs of the two programs for a series of calculations is given in the Software QA Plan¹⁹. This comparison demonstrates the accuracy of the *RadDecay*[®] v. 3.0 program. Both software packages have a software QA classification of “C”²⁰. In addition to this check, a separate independent evaluation utilizing the input radionuclides and *RadDecay*[®] v. 1.13 was conducted. The reportable radionuclides determined by this evaluation were the same as those reported here within a few tenths of a percent.

The total Curie content of the dried sludge in the year 2015 is $4.03\text{E}+04 \mu\text{Ci/g}$. This value is greater than the $2.05\text{E}+04 \mu\text{Ci/g}$ total represented by the reportable radionuclides in Appendix A. The difference is due to the significant contribution to the activity from radionuclides having half-lives shorter than ten years. The activity of Y-90 accounts for nearly the entire difference, with the balance attributable primarily to Ba-137m.

Appendix L presents the reportable radionuclides indexed to the year 3115. The total Curie content of the dried sludge in 3115 is $3.85\text{E}+01 \mu\text{Ci/g}$. This value is slightly greater than the $3.71\text{E}+01 \mu\text{Ci/g}$ total represented by the radionuclides identified as reportable. The difference is due to the minor contribution to the total activity from radionuclides having half-lives shorter than 10 years. These radionuclides include (in decreasing order of activity contribution): Np-239, Sb-126m, Pa-233, Sb-126, Th-234, and Pa-234m.

Twenty-six radionuclides have been identified as reportable for DWPF SB5 as specified by WAPS 1.2. Consistent with the strategy detailed in the WCP and WQR, each of these radionuclides has a half-life greater than 10 years and contributes more than 0.01 % of the radioactivity on a Curie basis at some point from production through the 1100-year period between 2015 and 3115. The 26 reportable radionuclides are given in Table 4-3. The calculations at every one hundred years out to 1100 years demonstrated that only one radionuclide became reportable during this time period, Am-242m, which was reportable for the years between 2215 and 3015, but no longer reportable in 3115. The data for these intermediate year calculations can be found in Appendices B-K.

Table 4-3. Reportable Radionuclides in DWPF Sludge Batch 5

Cl-36*	Ni-59	Ni-63	Sr-90	Zr-93	Nb-93m
Tc-99	Sn-126*	Cs-137	Sm-151	U-233	U-234
Np-237	U-238	Pu-238	Pu-239	Pu-240	Am-241
Pu-241	Pu-242	Am-242m*	Am-243	Cm-244	Cm-245*
Cm-246	Cf-251*				

* Based upon an analytical detection limit.

The WCP and WQR require that all of the radionuclides present in the Design Basis glass be considered as the initial set of reportable radionuclides. All of the radionuclides in the Design Basis glass are reportable except for four radionuclides: Se-79, Pd-107, Cs-135, and Th-230. At no time during the 1100-year period between 2015 and 3115 did any of these three radionuclides contribute to more than 0.01% of the radioactivity on a Curie basis.

Five of the 26 reportable radionuclides for SB5 are not part of either the design-basis list of radionuclides² or the list of Pu and U isotopes identified in WAPS 1.6. These include Cl-36, Am-242m, Cm-245, Cm-246, and Cf-251. One of these radionuclides (Cm-246) was also reported for SB1B (MB2)²¹, three of these radionuclides (Cm-245, Cm-246, and Cf-251) were also reported for SB2 (MB3)²², and four of these radionuclides (Am-242m, Cm-245, Cm-246, and Cf-251) plus Cm-247 and Cm-248 were reported for SB3 (MB4)²³, and four of these radionuclides (Am-242m, Cm-245, Cm-246, and Cf-251) plus Cm-247, Bk-247, and Cm-248 were reportable for SB4 (MB5)²⁴.

The list of reportable radionuclides that were determined for SB1B (MB2)²¹ contained two radionuclides that were not reportable for SB2 (MB3), SB3 (MB4) or SB4 (MB5). These radionuclides were I-129 and Th-229. Sn-121m was reportable for SB1B (MB2), SB2 (MB3), and SB3 (MB4), but due to an improved detection limit was not reportable for SB4 (MB5), nor is it reportable for SB5. Similarly, Cf-249 was reportable for SB3, but due to an improved detection limit for the input value, it was not reportable for SB4 (MB5), nor is it reportable for SB5. Cm-247, Cm-248, and Bk-247 were reportable for SB4 (MB5), but improved detection limits for these input values make all of them not reportable for the current SB5. For earlier sludge batches, the Se-79 input value was not a detection limit value as it is for SB5. The detection limit was low enough that the radionuclide was not reportable. C-14, reported in the first two batch analyses, has been excluded from consideration for future sludge batches⁷. For easier comparison, the reportable nuclides for SB1B, SB2, SB3, and SB4 have been reproduced in Table 4-4.

Table 4-4. Reportable Radionuclides in DWPF Sludge Batch 1B through 4

Isotope	SB1B (MB2) ²¹	SB2 (MB3) ²²	SB3 (MB4) ²³	SB4 (MB5) ²⁴
C-14	X	X		
Ni-59	X	X	X	X
Ni-63	X	X	X	X
Se-79	X	X	X	X
Sr-90	X	X	X	X
Zr-93	X	X	X	X
Nb-93m	X	X	X	X
Tc-99	X	X	X	X
Sn-121m	X	X	X	
Sn-126	X	X	X	X
I-129	X			
Cs-137	X	X	X	X
Sm-151	X	X	X	X
Th-229	X			
U-233	X	X	X	X
U-234	X	X	X	X
Np-237	X	X	X	X
U-238	X	X	X	X
Pu-238	X	X	X	X
Pu-239	X	X	X	X
Pu-240	X	X	X	X
Am-241	X	X	X	X
Pu-241	X	X	X	X
Pu-242	X	X	X	X
Am-242m			X	X
Am-243	X	X	X	X
Cm-244	X	X	X	X
Cm-245		X	X	X
Cm-246	X	X	X	X
Bk-247				X
Cm-247			X	X
Cm-248			X	X
Cf-249			X	
Cf-251		X	X	X

²¹ Fellingner, T. L, Bibler, N. E., and Harbour, J. R., *Characterization of and Waste Acceptance Radionuclides to be Reported for DWPF Macrobatches 2 (ESP 215 – ESP 221)*, WSRC-RP-99-00436, Revision 1, Savannah River Site, March 2004.

²² Bibler, N. E., DiPrete, D. P., and Harbour, J. R., *Determination of Reportable Radionuclides for DWPF Sludge Batch 2 (Macrobatches 3)*, WSRC-TR-2002-00255, Revision 0, Savannah River Site, September 2002.

²³ Bannoche, C. J. and Bibler, N. E., *Determination of Reportable Radionuclides for DWPF Sludge Batch 3 (Macrobatches 4)*, WSRC-TR-2005-00157, Rev. 0, Savannah River Site, May 2005.

²⁴ Bannoche, C. J., Bibler, N. E., and DiPrete, D. P., *Determination of Reportable Radionuclides for DWPF Sludge Batch 4 (Macrobatches 5)*, WSRC-STI-2008-00142, Rev. 0, Savannah River Site, May 2008.

4.4 The Ratio by Weight of U and Pu Isotopes

The WQR requires that the relative concentrations of the uranium and plutonium isotopes be provided from the analysis of each Macrobatches (in this case MB6 – Sludge Batch 5) in order to meet the WAPS

IAEA Safeguards Reporting for HLW Specification (WAPS 1.6). The data for uranium isotopes are given in Table 4-5.

Table 4-5. Uranium Isotope Distribution in DWPF Sludge Batch 5

Isotope	Wt% Total Solids	Percent Distribution
U-233	2.29E-04	0.00458
U-234	9.85E-04	0.0197
U-235	3.24E-02	0.647
U-236	2.08E-03	0.0414
U-238	4.97E+00	99.3
Total	5.01E+00	100

The data for the plutonium isotopes is given in Table 4-6.

Table 4-6. Plutonium Isotope Distribution in DWPF Sludge Batch 5

Isotope	Wt% Total Solids	Percent Distribution
Pu-238	1.61E-03	4.28
Pu-239	3.28E-02	87.3
Pu-240	2.89E-03	7.68
Pu-241	9.45E-05	0.251
Pu-242	1.79E-04	0.475
Total	3.76E-02	100

All of the Pu isotopes and U-233, U-234, and U-238 are already reportable since they meet the requirement of having half-lives greater than 10 years and a contribution to the overall activity of greater than 0.01% on a Curie basis through the year 3115. In order to be compliant with WAPS 1.6, U-235 and U-236 also become reportable even though they contribute less than 0.01% to the total activity (U-235 at 0.002% and U-236 at 0.004% in 3115).

5.0 CONCLUSIONS

Twenty-six radionuclides have been identified as reportable for DWPF SB5 as specified by WAPS 1.2. Consistent with the strategy detailed in the WCP and WQR, each of these radionuclides has a half-life greater than ten years and contributes more than 0.01% of the radioactivity on a Curie basis at some point from production through the 1100 year period between 2015 and 3115. The 26 reportable radionuclides are:

Cl-36	Ni-59	Ni-63	Sr-90	Zr-93	Nb-93m
Tc-99	Sn-126	Cs-137	Sm-151	U-233	U-234
Np-237	U-238	Pu-238	Pu-239	Pu-240	Am-241
Pu-241	Pu-242	Am-242m	Am-243	Cm-244	Cm-245
Cm-246	Cf-251				

Chlorine-36 is reported for the first time based on the upper limit determined from the aqua regia digested sludge slurry.

The WCP and WQR require that all of radionuclides present in the Design Basis glass be considered as the initial set of reportable radionuclides. For SB5, all of the radionuclides in the Design Basis glass are reportable except for four radionuclides: Se-79, Pd-107, Cs-135, and Th-230. At no time through the year 3115 did any of these three radionuclides contribute to more than 0.01% of the radioactivity on a Curie basis.

Two additional uranium isotopes (U-235 and -236) must be added to the list of reportable radionuclides in order to meet WAPS 1.6. All of the Pu isotopes and other U isotopes (U-233, -234, and -238) identified in WAPS 1.6 were already determined to be reportable according to WAPS 1.2. This brings the total number of reportable radionuclides for SB5 to 28.

6.0 RECOMMENDATIONS

The radionuclide measurements made for SB5 are the most extensive conducted to date. Some method development/refinement occurred during the conduct of these measurements, leading to lower detection limits and more accurate measurement of some isotopes than was previously possible. Continuous improvement in the analytical measurements will likely continue, and this in turn should lead to improved detection limit values for some radionuclides and actual measurements for still others.

7.0 REFERENCES

- ¹ Office of Environmental Management, *Waste Acceptance Product Specifications for Vitrified High-Level Waste Forms*, USDOE Document DOE/EM-0093, Rev. 2, December 1996.
- ² Washington Savannah River Company, *DWPF Waste Form Compliance Plan*, WSRC-IM-91-116-0, Rev. 8, Savannah River Site, March 2006.
- ³ Washington Savannah River Company, *DWPF Waste Form Qualification Report, Volume 4: Reporting the Radionuclide Inventory of the DWPF Product*, WSRC-IM-91-116-4, Rev. 3, Savannah River Site, 2006.
- ⁴ Fellingner, T. L., *Technical Task Request: Sludge Batch 5 SRNL Shielded Cells Testing*, HLW-DWPF-TTR-2008-0010, Rev. 2, Savannah River Site, November 2008.
- ⁵ Bannochie, C. J. and Pareizs, J. M., *Qualification of DWPF Sludge Batch 5 in the SRNL Shielded Cells: Task Technical and Quality Assurance Plan*, WSRC-RP-2008-00137, Rev. 2, Savannah River Site, December 2008.
- ⁶ Bannochie, C. J. and Pareizs, J. M., *Qualification of DWPF Sludge Batch 5 in the SRNL Shielded Cells: Analytical Study Plan*, WSRC-RP-2008-00138, Rev. 2, Savannah River Site, December 2008.
- ⁷ Bibler, N. E. and Fowler, J. R., *Technical Basis for Eliminating Carbon-14 as a Reportable Radionuclide In DWPF Glass*, WSRC-TR-2004-00629, Revision 0, Savannah River Site, December 2004.
- ⁸ Hyder, M. L., *Waste Acceptance Radionuclides to be Reported in Tank 51 Sludge Only Glass*, WSRC-TR-95-0485, Rev. 0, Savannah River Site, December 1995.
- ⁹ Bannochie, C. J., *Tank 40 Final SB5 Chemical Characterization Results Prior to Np Addition*, SRNL-STI-2009-00060, Rev. 2, Savannah River Site, January 2010.
- ¹⁰ Elder, H. H., *Neptunium Addition to the Sludge Batch 5 WAPS Sample*, LWO-WSE-2009-00170, Savannah River Site, June 2009.
- ¹¹ Oak Ridge National Laboratory, *Integrated Data Base Report – 1994: U. S. Spent Nuclear Fuel and Radioactive Waste Inventories, Projections, and Characteristics*, USDOE Document DOE/RW-0006, Rev. 11, Oak Ridge National Laboratory, September 1995.
- ¹² Friedlander, G. and Kennedy, J. W., *Nuclear and Radiochemistry*; John Wiley & Sons, Inc: New York, 1957; p7.
- ¹³ Baum, E. M., Knox, H. D., and Miller, T. R., *Nuclides and Isotopes (Chart of the Nuclides)*, 16th Edition, KAPL, Inc. and Lockheed Martin, 2002.
- ¹⁴ England, T. R. and Rider, B. F., *ENDF-349, Evaluation and Compilation of Fission Product Yields*, LA-UR-3106, Los Alamos National Laboratory, 1994.

- ¹⁵ Bibler, N. E., Fellingner, T. L., and Hobbs, D. T., *Technetium-99 Behavior in Savannah River Site HLW Sludges During Waste Processing*, WSRC-MS-2004-00614, Savannah River Site, Published as Paper No. 4 in Session 43 of WM'06 Proceedings, WM Symposium, Tucson, AZ, February 2006.
- ¹⁶ Marek, J. C., *Correction Factor for Soluble and Insoluble Solids*, SRTC-PTD-92-0040, Savannah River Site, Aiken, SC 29808 (1992).
- ¹⁷ *RadDecay*, Version 3.0, Grove Software, Inc., Lynchburg, VA 24502.
- ¹⁸ *Radioactive Decay Calculator*, D. W. James & Associates, North Oaks, MN 55127.
- ¹⁹ Pareizs, J. M. and Bannochie, C. J., *Software Quality Assurance Plan and Acceptance Testing for Commercial Radioactive Decay Calculation Packages Utilized by the Process Technology Programs*, G-SQP-A-00006, Rev. 2, Savannah River Site, November 2009.
- ²⁰ Bannochie, C. J., *Radioactive Decay Modeling Using RadDecay, Radioactive Decay Calculator, and Radcalc - Software Classification*, WSRC-RP-2005-01585, Rev. 2, Savannah River Site, November 2009.
- ²¹ Fellingner, T. L., Bibler, N. E., and Harbour, J. R., *Characterization of and Waste Acceptance Radionuclides to be Reported for DWPF Macro Batch 2 (ESP 215 – ESP 221)*, WSRC-RP-99-00436, Rev. 1, Savannah River Site, March 2004.
- ²² Bibler, N. E., DiPrete, D. P., and Harbour, J. R., *Determination of Reportable Radionuclides for DWPF Sludge Batch 2 (Macro Batch 3)*, WSRC-TR-2002-00255, Rev. 0, Savannah River Site, September 2002.
- ²³ Bannochie, C. J. and Bibler, N. E., *Determination of Reportable Radionuclides for DWPF Sludge Batch 3 (Macrobatch 4)*, WSRC-TR-2005-00157, Rev. 0, Savannah River Site, May 2005.
- ²⁴ Bannochie, C. J., Bibler, N. E., and DiPrete, D. P., *Determination of Reportable Radionuclides for DWPF Sludge Batch 4 (Macrobatch 5)*, WSRC-STI-2008-00142, Rev. 0, Savannah River Site, May 2008.

8.0 ACKNOWLEDGEMENTS

The authors would like to acknowledge the support of the SRNL Shielded Cells technicians and supervision. Additionally, we would like to thank Ceci DiPrete for assistance with the vast array of counting analyses.

APPENDIX A. ACTIVITIES OF DRIED SLUDGE IN YEAR 2015 ($\mu\text{Ci/g}$)

Nuclide	Y 2015 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2015 μCi/g	Fraction of Activity	Reportable
Ac-225	1.24E-05	3.07E-10		Po-213	1.21E-05	3.00E-10	
Ac-227	7.96E-09	1.97E-13		Po-214	4.40E-09	1.09E-13	
Ac-228	3.73E-05	9.25E-10		Po-215	7.66E-09	1.90E-13	
Am-241	3.30E+01	8.18E-04	Yes	Po-216	2.38E-05	5.90E-10	
Am-242	4.40E-01	1.09E-05		Po-218	4.40E-09	1.09E-13	
Am-242m	4.42E-01	1.10E-05		Pu-238	2.63E+02	6.53E-03	Yes
Am-243	9.63E-01	2.39E-05		Pu-239	2.04E+01	5.06E-04	Yes
At-217	1.24E-05	3.07E-10		Pu-240	6.61E+00	1.64E-04	Yes
At-218	8.79E-13	2.18E-17		Pu-241	7.30E+01	1.81E-03	Yes
Ba-133	5.72E-02	1.42E-06		Pu-242	6.82E-03	1.69E-07	
Ba-137m	2.66E+02	6.59E-03		Pu-243	2.49E-07	6.18E-12	
Bi-210	2.56E-10	6.35E-15		Pu-244	1.74E-12	4.32E-17	
Bi-211	7.66E-09	1.90E-13		Ra-223	7.66E-09	1.90E-13	
Bi-212	2.38E-05	5.90E-10		Ra-224	2.38E-05	5.90E-10	
Bi-213	1.24E-05	3.07E-10		Ra-225	1.25E-05	3.09E-10	
Bi-214	4.40E-09	1.09E-13		Ra-226	4.42E-09	1.10E-13	
Bk-247	2.70E-03	6.70E-08		Ra-228	3.73E-05	9.25E-10	
Cd-113m	2.14E+00	5.31E-05		Rn-219	7.66E-09	1.90E-13	
Cf-249	8.85E-03	2.20E-07		Rn-220	2.38E-05	5.90E-10	
Cf-250	9.90E-05	2.45E-09		Rn-222	4.40E-09	1.09E-13	
Cf-251	2.30E-02	5.70E-07		Sb-125	2.21E-02	5.47E-07	
Cl-36	9.75E-01	2.42E-05		Sb-126	5.04E-02	1.25E-06	
Cm-242	3.65E-01	9.04E-06		Sb-126m	3.60E-01	8.92E-06	
Cm-243	2.09E-01	5.19E-06		Se-79	2.78E-03	6.89E-08	
Cm-244	3.66E+01	9.08E-04	Yes	Sm-147	3.15E-09	7.80E-14	
Cm-245	4.29E-02	1.06E-06		Sm-151	1.74E+02	4.31E-03	Yes
Cm-246	1.29E-02	3.20E-07		Sn-121	2.19E-01	5.44E-06	
Cm-247	2.49E-07	6.18E-12		Sn-121m	2.83E-01	7.01E-06	
Cm-248	3.77E-05	9.35E-10		Sn-126	3.60E-01	8.92E-06	
Co-60	1.09E+00	2.70E-05		Sr-90	1.95E+04	4.84E-01	Yes
Cs-135	1.61E-03	3.99E-08		Tc-99	1.53E-01	3.79E-06	
Cs-137	2.81E+02	6.97E-03	Yes	Te-125m	5.34E-03	1.32E-07	
Eu-154	1.45E+01	3.60E-04		Th-227	7.66E-09	1.90E-13	
Eu-155	1.06E+00	2.63E-05		Th-228	2.39E-05	5.92E-10	
Fr-221	1.24E-05	3.07E-10		Th-229	1.26E-05	3.12E-10	
Fr-223	1.10E-10	2.72E-15		Th-230	3.45E-06	8.54E-11	
I-129	3.77E-04	9.35E-09		Th-231	7.00E-04	1.74E-08	
Nb-93m	4.27E-01	1.06E-05		Th-232	7.25E-05	1.80E-09	
Ni-59	1.30E+00	3.22E-05		Th-234	1.67E-02	4.14E-07	
Ni-63	1.06E+02	2.64E-03	Yes	Tl-207	7.64E-09	1.89E-13	
Np-237	6.93E-02	1.72E-06		Tl-208	8.55E-06	2.12E-10	
Np-238	2.10E-03	5.21E-08		Tl-209	2.67E-07	6.62E-12	
Np-239	9.63E-01	2.39E-05		U-233	2.22E-02	5.50E-07	
Np-240m	1.74E-12	4.31E-17		U-234	6.61E-02	1.64E-06	
Pa-231	8.88E-08	2.20E-12		U-235	7.00E-04	1.74E-08	
Pa-233	6.93E-02	1.72E-06		U-236	1.34E-03	3.32E-08	
Pa-234	5.51E-05	1.37E-09		U-237	1.79E-03	4.44E-08	
Pa-234m	1.67E-02	4.13E-07		U-238	1.67E-02	4.14E-07	
Pb-209	1.24E-05	3.07E-10		U-240	1.74E-12	4.31E-17	
Pb-210	2.59E-10	6.42E-15		Y-90	1.95E+04	4.84E-01	
Pb-211	7.66E-09	1.90E-13		Zr-93	5.23E-01	1.30E-05	
Pb-212	2.38E-05	5.90E-10		TOTAL	4.03E+04	1.00E+00	
Pb-214	4.40E-09	1.09E-13					
Pd-107	1.29E-03	3.20E-08					
Pm-147	3.28E+01	8.13E-04					
Po-210	1.98E-10	4.91E-15					
Po-211	2.14E-11	5.32E-16					
Po-212	1.53E-05	3.78E-10					

APPENDIX B. ACTIVITIES OF DRIED SLUDGE IN YEAR 2115 ($\mu\text{Ci/g}$)

Nuclide	Y 2115 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2115 μCi/g	Fraction of Activity	Reportable
Ac-225	2.21E-04	5.55E-08		Po-213	2.16E-04	5.43E-08	
Ac-227	1.12E-06	2.81E-10		Po-214	1.82E-06	4.56E-10	
Ac-228	7.25E-05	1.82E-08		Po-215	1.12E-06	2.81E-10	
Am-241	3.02E+01	7.59E-03	Yes	Po-216	7.25E-05	1.82E-08	
Am-242	2.79E-01	7.00E-05		Po-218	1.82E-06	4.56E-10	
Am-242m	2.80E-01	7.03E-05		Pu-238	1.20E+02	3.00E-02	Yes
Am-243	9.54E-01	2.40E-04	Yes	Pu-239	2.03E+01	5.11E-03	Yes
At-217	2.21E-04	5.55E-08		Pu-240	6.64E+00	1.67E-03	Yes
At-218	3.63E-10	9.13E-14		Pu-241	6.35E-01	1.59E-04	Yes
Ba-133	9.00E-05	2.26E-08		Pu-242	6.83E-03	1.72E-06	
Ba-137m	2.64E+01	6.63E-03		Pu-243	3.47E-07	8.72E-11	
Bi-210	1.00E-06	2.51E-10		Pu-244	3.08E-11	7.72E-15	
Bi-211	1.12E-06	2.81E-10		Ra-223	1.12E-06	2.81E-10	
Bi-212	7.25E-05	1.82E-08		Ra-224	7.25E-05	1.82E-08	
Bi-213	2.21E-04	5.55E-08		Ra-225	2.21E-04	5.55E-08	
Bi-214	1.82E-06	4.56E-10		Ra-226	1.82E-06	4.56E-10	
Bk-247	2.57E-03	6.45E-07		Ra-228	7.25E-05	1.82E-08	
Cd-113m	1.31E-02	3.29E-06		Rn-219	1.12E-06	2.81E-10	
Cf-249	7.27E-03	1.82E-06		Rn-220	7.25E-05	1.82E-08	
Cf-250	4.94E-07	1.24E-10		Rn-222	1.82E-06	4.56E-10	
Cf-251	2.13E-02	5.34E-06		Sb-125	3.00E-13	7.52E-17	
Cl-36	9.75E-01	2.45E-04	Yes	Sb-126	5.04E-02	1.26E-05	
Cm-242	2.31E-01	5.80E-05		Sb-126m	3.60E-01	9.03E-05	
Cm-243	1.84E-02	4.61E-06		Se-79	2.78E-03	6.97E-07	
Cm-244	7.98E-01	2.00E-04	Yes	Sm-147	3.96E-09	9.94E-13	
Cm-245	4.26E-02	1.07E-05		Sm-151	8.05E+01	2.02E-02	Yes
Cm-246	1.27E-02	3.19E-06		Sn-121	6.22E-02	1.56E-05	
Cm-247	3.47E-07	8.72E-11		Sn-121m	8.02E-02	2.01E-05	
Cm-248	3.77E-05	9.46E-09		Sn-126	3.60E-01	9.03E-05	
Co-60	2.12E-06	5.33E-10		Sr-90	1.80E+03	4.53E-01	Yes
Cs-135	1.61E-03	4.04E-07		Tc-99	1.53E-01	3.84E-05	
Cs-137	2.79E+01	7.01E-03	Yes	Te-125m	7.25E-14	1.82E-17	
Eu-154	5.51E-03	1.38E-06		Th-227	1.10E-06	2.77E-10	
Eu-155	9.04E-07	2.27E-10		Th-228	7.25E-05	1.82E-08	
Fr-221	2.21E-04	5.55E-08		Th-229	2.21E-04	5.55E-08	
Fr-223	1.55E-08	3.88E-12		Th-230	8.91E-05	2.24E-08	
I-129	3.77E-04	9.47E-08		Th-231	7.02E-04	1.76E-07	
Nb-93m	5.22E-01	1.31E-04	Yes	Th-232	7.25E-05	1.82E-08	
Ni-59	1.30E+00	3.26E-04	Yes	Th-234	1.67E-02	4.19E-06	
Ni-63	5.16E+01	1.30E-02	Yes	Tl-207	1.12E-06	2.80E-10	
Np-237	7.03E-02	1.77E-05		Tl-208	2.60E-05	6.54E-09	
Np-238	1.33E-03	3.35E-07		Tl-209	4.77E-06	1.20E-09	
Np-239	9.54E-01	2.40E-04		U-233	2.22E-02	5.58E-06	
Np-240m	3.07E-11	7.71E-15		U-234	1.18E-01	2.95E-05	
Pa-231	1.57E-06	3.94E-10		U-235	7.02E-04	1.76E-07	
Pa-233	7.03E-02	1.77E-05		U-236	1.36E-03	3.42E-07	
Pa-234	5.51E-05	1.38E-08		U-237	1.56E-05	3.91E-09	
Pa-234m	1.67E-02	4.19E-06		U-238	1.67E-02	4.19E-06	
Pb-209	2.21E-04	5.55E-08		U-240	3.07E-11	7.71E-15	
Pb-210	1.00E-06	2.51E-10		Y-90	1.81E+03	4.53E-01	
Pb-211	1.12E-06	2.81E-10		Zr-93	5.23E-01	1.31E-04	Yes
Pb-212	7.25E-05	1.82E-08		TOTAL	3.98E+03	1.00E+00	
Pb-214	1.82E-06	4.56E-10					
Pd-107	1.29E-03	3.24E-07					
Pm-147	1.10E-10	2.76E-14					
Po-210	9.86E-07	2.48E-10					
Po-211	3.13E-09	7.87E-13					
Po-212	4.65E-05	1.17E-08					

APPENDIX C. ACTIVITIES OF DRIED SLUDGE IN YEAR 2215 ($\mu\text{Ci/g}$)

Nuclide	Y 2215 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2215 μCi/g	Fraction of Activity	Reportable
Ac-225	4.28E-04	8.29E-07		Po-213	4.19E-04	8.11E-07	
Ac-227	2.59E-06	5.02E-09		Po-214	7.96E-06	1.54E-08	
Ac-228	7.25E-05	1.41E-07		Po-215	2.59E-06	5.01E-09	
Am-241	2.58E+01	5.00E-02	Yes	Po-216	7.25E-05	1.41E-07	
Am-242	1.77E-01	3.42E-04		Po-218	7.96E-06	1.54E-08	
Am-242m	1.77E-01	3.44E-04	Yes	Pu-238	5.44E+01	1.05E-01	Yes
Am-243	9.46E-01	1.83E-03	Yes	Pu-239	2.03E+01	3.93E-02	Yes
At-217	4.28E-04	8.29E-07		Pu-240	6.57E+00	1.27E-02	Yes
At-218	1.59E-09	3.08E-12		Pu-241	4.72E-02	9.14E-05	
Ba-133	1.42E-07	2.75E-10		Pu-242	6.84E-03	1.33E-05	
Ba-137m	2.62E+00	5.08E-03		Pu-243	4.38E-07	8.50E-10	
Bi-210	5.69E-06	1.10E-08		Pu-244	5.98E-11	1.16E-13	
Bi-211	2.59E-06	5.01E-09		Ra-223	2.59E-06	5.01E-09	
Bi-212	7.25E-05	1.41E-07		Ra-224	7.25E-05	1.41E-07	
Bi-213	4.28E-04	8.29E-07		Ra-225	4.28E-04	8.29E-07	
Bi-214	7.96E-06	1.54E-08		Ra-226	7.96E-06	1.54E-08	
Bk-247	2.44E-03	4.74E-06		Ra-228	7.25E-05	1.41E-07	
Cd-113m	8.02E-05	1.55E-07		Rn-219	2.59E-06	5.01E-09	
Cf-249	5.96E-03	1.16E-05		Rn-220	7.25E-05	1.41E-07	
Cf-250	2.47E-09	4.79E-12		Rn-222	7.96E-06	1.54E-08	
Cf-251	1.97E-02	3.82E-05		Sb-125	4.07E-24	7.88E-27	
Cl-36	9.75E-01	1.89E-03	Yes	Sb-126	5.03E-02	9.75E-05	
Cm-242	1.46E-01	2.84E-04		Sb-126m	3.59E-01	6.97E-04	
Cm-243	1.61E-03	3.13E-06		Se-79	2.77E-03	5.38E-06	
Cm-244	1.74E-02	3.37E-05		Sm-147	3.96E-09	7.67E-12	
Cm-245	4.23E-02	8.20E-05		Sm-151	3.72E+01	7.22E-02	Yes
Cm-246	1.25E-02	2.43E-05		Sn-121	1.76E-02	3.42E-05	
Cm-247	4.38E-07	8.50E-10		Sn-121m	2.27E-02	4.41E-05	
Cm-248	3.77E-05	7.30E-08		Sn-126	3.59E-01	6.97E-04	Yes
Co-60	4.13E-12	8.00E-15		Sr-90	1.67E+02	3.24E-01	Yes
Cs-135	1.61E-03	3.12E-06		Tc-99	1.53E-01	2.96E-04	Yes
Cs-137	2.77E+00	5.36E-03	Yes	Te-125m	9.84E-25	1.91E-27	
Eu-154	2.09E-06	4.06E-09		Th-227	2.55E-06	4.94E-09	
Eu-155	7.71E-13	1.49E-15		Th-228	7.25E-05	1.41E-07	
Fr-221	4.28E-04	8.29E-07		Th-229	4.28E-04	8.30E-07	
Fr-223	3.57E-08	6.92E-11		Th-230	2.07E-04	4.01E-07	
I-129	3.77E-04	7.31E-07		Th-231	7.04E-04	1.36E-06	
Nb-93m	5.23E-01	1.01E-03	Yes	Th-232	7.25E-05	1.41E-07	
Ni-59	1.30E+00	2.51E-03	Yes	Th-234	1.67E-02	3.24E-05	
Ni-63	2.51E+01	4.86E-02	Yes	Tl-207	2.58E-06	5.00E-09	
Np-237	7.12E-02	1.38E-04	Yes	Tl-208	2.60E-05	5.05E-08	
Np-238	8.45E-04	1.64E-06		Tl-209	9.24E-06	1.79E-08	
Np-239	9.46E-01	1.83E-03		U-233	2.22E-02	4.31E-05	
Np-240m	5.97E-11	1.16E-13		U-234	1.41E-01	2.73E-04	Yes
Pa-231	3.05E-06	5.92E-09		U-235	7.04E-04	1.36E-06	
Pa-233	7.12E-02	1.38E-04		U-236	1.38E-03	2.68E-06	
Pa-234	5.51E-05	1.07E-07		U-237	1.16E-06	2.24E-09	
Pa-234m	1.67E-02	3.23E-05		U-238	1.67E-02	3.24E-05	
Pb-209	4.28E-04	8.29E-07		U-240	5.97E-11	1.16E-13	
Pb-210	5.70E-06	1.10E-08		Y-90	1.67E+02	3.24E-01	
Pb-211	2.59E-06	5.01E-09		Zr-93	5.23E-01	1.01E-03	Yes
Pb-212	7.25E-05	1.41E-07		TOTAL	5.16E+02	1.00E+00	
Pb-214	7.96E-06	1.54E-08					
Pd-107	1.29E-03	2.50E-06					
Pm-147	3.69E-22	7.15E-25					
Po-210	5.66E-06	1.10E-08					
Po-211	7.24E-09	1.40E-11					
Po-212	4.65E-05	9.00E-08					

APPENDIX D. ACTIVITIES OF DRIED SLUDGE IN YEAR 2315 ($\mu\text{Ci/g}$)

Nuclide	Y 2315 μCi/g	Fraction of Activity	Reportable
Ac-225	6.33E-04	4.48E-06	
Ac-227	4.07E-06	2.88E-08	
Ac-228	7.25E-05	5.13E-07	
Am-241	2.20E+01	1.55E-01	Yes
Am-242	1.12E-01	7.92E-04	
Am-242m	1.12E-01	7.96E-04	Yes
Am-243	9.37E-01	6.63E-03	Yes
At-217	6.33E-04	4.48E-06	
At-218	3.84E-09	2.71E-11	
Ba-133	2.23E-10	1.58E-12	
Ba-137m	2.60E-01	1.84E-03	
Bi-210	1.53E-05	1.08E-07	
Bi-211	4.07E-06	2.88E-08	
Bi-212	7.25E-05	5.13E-07	
Bi-213	6.33E-04	4.48E-06	
Bi-214	1.92E-05	1.36E-07	
Bk-247	2.32E-03	1.64E-05	
Cd-113m	4.91E-07	3.48E-09	
Cf-249	4.89E-03	3.46E-05	
Cf-250	1.23E-11	8.74E-14	
Cf-251	1.82E-02	1.29E-04	Yes
Cl-36	9.74E-01	6.90E-03	Yes
Cm-242	9.28E-02	6.57E-04	
Cm-243	1.42E-04	1.00E-06	
Cm-244	3.78E-04	2.68E-06	
Cm-245	4.20E-02	2.97E-04	Yes
Cm-246	1.23E-02	8.73E-05	
Cm-247	5.23E-07	3.70E-09	
Cm-248	3.77E-05	2.67E-07	
Co-60	8.04E-18	5.69E-20	
Cs-135	1.61E-03	1.14E-05	
Cs-137	2.75E-01	1.94E-03	Yes
Eu-154	7.95E-10	5.63E-12	
Eu-155	6.59E-19	4.66E-21	
Fr-221	6.33E-04	4.48E-06	
Fr-223	5.62E-08	3.98E-10	
I-129	3.77E-04	2.67E-06	
Nb-93m	5.23E-01	3.70E-03	Yes
Ni-59	1.30E+00	9.18E-03	Yes
Ni-63	1.22E+01	8.63E-02	Yes
Np-237	7.20E-02	5.09E-04	Yes
Np-238	5.35E-04	3.79E-06	
Np-239	9.37E-01	6.63E-03	
Np-240m	8.87E-11	6.28E-13	
Pa-231	4.54E-06	3.21E-08	
Pa-233	7.20E-02	5.09E-04	
Pa-234	5.51E-05	3.90E-07	
Pa-234m	1.67E-02	1.18E-04	
Pb-209	6.33E-04	4.48E-06	
Pb-210	1.53E-05	1.08E-07	
Pb-211	4.07E-06	2.88E-08	
Pb-212	7.25E-05	5.13E-07	
Pb-214	1.92E-05	1.36E-07	
Pd-107	1.29E-03	9.13E-06	
Pm-147	1.24E-33	8.77E-36	
Po-210	1.52E-05	1.08E-07	
Po-211	1.14E-08	8.06E-11	
Po-212	4.65E-05	3.29E-07	

Nuclide	Y 2315 μCi/g	Fraction of Activity	Reportable
Po-213	6.19E-04	4.38E-06	
Po-214	1.92E-05	1.36E-07	
Po-215	4.07E-06	2.88E-08	
Po-216	7.25E-05	5.13E-07	
Po-218	1.92E-05	1.36E-07	
Pu-238	2.48E+01	1.75E-01	Yes
Pu-239	2.02E+01	1.43E-01	Yes
Pu-240	6.50E+00	4.60E-02	Yes
Pu-241	4.21E-02	2.98E-04	Yes
Pu-242	6.85E-03	4.85E-05	
Pu-243	5.23E-07	3.70E-09	
Pu-244	8.88E-11	6.28E-13	
Ra-223	4.07E-06	2.88E-08	
Ra-224	7.25E-05	5.13E-07	
Ra-225	6.33E-04	4.48E-06	
Ra-226	1.92E-05	1.36E-07	
Ra-228	7.25E-05	5.13E-07	
Rn-219	4.07E-06	2.88E-08	
Rn-220	7.25E-05	5.13E-07	
Rn-222	1.92E-05	1.36E-07	
Sb-125	5.53E-35	3.92E-37	
Sb-126	5.03E-02	3.56E-04	
Sb-126m	3.59E-01	2.54E-03	
Se-79	2.77E-03	1.96E-05	
Sm-147	3.96E-09	2.80E-11	
Sm-151	1.72E+01	1.22E-01	Yes
Sn-121	5.01E-03	3.54E-05	
Sn-121m	6.45E-03	4.57E-05	
Sn-126	3.59E-01	2.54E-03	Yes
Sr-90	1.55E+01	1.09E-01	Yes
Tc-99	1.53E-01	1.08E-03	Yes
Te-125m	1.34E-35	9.47E-38	
Th-227	4.01E-06	2.84E-08	
Th-228	7.25E-05	5.13E-07	
Th-229	6.33E-04	4.48E-06	
Th-230	3.39E-04	2.40E-06	
Th-231	7.06E-04	5.00E-06	
Th-232	7.25E-05	5.13E-07	
Th-234	1.67E-02	1.18E-04	
Tl-207	4.06E-06	2.87E-08	
Tl-208	2.60E-05	1.84E-07	
Tl-209	1.37E-05	9.68E-08	
U-233	2.23E-02	1.58E-04	Yes
U-234	1.52E-01	1.07E-03	Yes
U-235	7.06E-04	5.00E-06	
U-236	1.40E-03	9.91E-06	
U-237	1.03E-06	7.30E-09	
U-238	1.67E-02	1.18E-04	Yes
U-240	8.87E-11	6.28E-13	
Y-90	1.55E+01	1.09E-01	
Zr-93	5.23E-01	3.70E-03	Yes
TOTAL	1.41E+02	1.00E+00	

APPENDIX E. ACTIVITIES OF DRIED SLUDGE IN YEAR 2415 ($\mu\text{Ci/g}$)

Nuclide	Y 2415 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2415 μCi/g	Fraction of Activity	Reportable
Ac-225	8.36E-04	1.04E-05		Po-213	8.18E-04	1.02E-05	
Ac-227	5.56E-06	6.93E-08		Po-214	3.57E-05	4.45E-07	
Ac-228	7.25E-05	9.04E-07		Po-215	5.55E-06	6.92E-08	
Am-241	1.87E+01	2.33E-01	Yes	Po-216	7.25E-05	9.04E-07	
Am-242	7.09E-02	8.84E-04		Po-218	3.57E-05	4.45E-07	
Am-242m	7.13E-02	8.89E-04	Yes	Pu-238	1.13E+01	1.41E-01	Yes
Am-243	9.28E-01	1.16E-02	Yes	Pu-239	2.02E+01	2.51E-01	Yes
At-217	8.36E-04	1.04E-05		Pu-240	6.44E+00	8.02E-02	Yes
At-218	7.14E-09	8.90E-11		Pu-241	4.18E-02	5.21E-04	Yes
Ba-133	3.51E-13	4.38E-15		Pu-242	6.85E-03	8.54E-05	
Ba-137m	2.58E-02	3.21E-04		Pu-243	6.01E-07	7.49E-09	
Bi-210	3.01E-05	3.75E-07		Pu-244	1.18E-10	1.47E-12	
Bi-211	5.55E-06	6.92E-08		Ra-223	5.55E-06	6.92E-08	
Bi-212	7.25E-05	9.04E-07		Ra-224	7.25E-05	9.04E-07	
Bi-213	8.36E-04	1.04E-05		Ra-225	8.37E-04	1.04E-05	
Bi-214	3.57E-05	4.45E-07		Ra-226	3.57E-05	4.45E-07	
Bk-247	2.21E-03	2.75E-05		Ra-228	7.25E-05	9.04E-07	
Cd-113m	3.00E-09	3.74E-11		Rn-219	5.55E-06	6.92E-08	
Cf-249	4.02E-03	5.01E-05		Rn-220	7.25E-05	9.04E-07	
Cf-250	6.16E-14	7.68E-16		Rn-222	3.57E-05	4.45E-07	
Cf-251	1.69E-02	2.10E-04	Yes	Sb-125	7.48E-46	9.33E-48	
Cl-36	9.74E-01	1.21E-02	Yes	Sb-126	5.03E-02	6.26E-04	
Cm-242	5.88E-02	7.33E-04		Sb-126m	3.59E-01	4.47E-03	
Cm-243	1.25E-05	1.55E-07		Se-79	2.77E-03	3.45E-05	
Cm-244	8.22E-06	1.03E-07		Sm-147	3.96E-09	4.94E-11	
Cm-245	4.17E-02	5.20E-04	Yes	Sm-151	7.98E+00	9.95E-02	Yes
Cm-246	1.22E-02	1.52E-04	Yes	Sn-121	1.42E-03	1.77E-05	
Cm-247	6.01E-07	7.49E-09		Sn-121m	1.83E-03	2.28E-05	
Cm-248	3.77E-05	4.70E-07		Sn-126	3.59E-01	4.47E-03	Yes
Co-60	1.56E-23	1.95E-25		Sr-90	1.43E+00	1.78E-02	Yes
Cs-135	1.61E-03	2.01E-05		Tc-99	1.53E-01	1.90E-03	Yes
Cs-137	2.72E-02	3.40E-04	Yes	Te-125m	1.81E-46	2.26E-48	
Eu-154	3.01E-13	3.76E-15		Th-227	5.48E-06	6.83E-08	
Eu-155	5.61E-25	6.99E-27		Th-228	7.25E-05	9.04E-07	
Fr-221	8.36E-04	1.04E-05		Th-229	8.37E-04	1.04E-05	
Fr-223	7.67E-08	9.56E-10		Th-230	4.78E-04	5.96E-06	
I-129	3.77E-04	4.70E-06		Th-231	7.08E-04	8.83E-06	
Nb-93m	5.23E-01	6.52E-03	Yes	Th-232	7.25E-05	9.04E-07	
Ni-59	1.30E+00	1.61E-02	Yes	Th-234	1.67E-02	2.08E-04	
Ni-63	5.92E+00	7.38E-02	Yes	Tl-207	5.54E-06	6.90E-08	
Np-237	7.26E-02	9.05E-04	Yes	Tl-208	2.60E-05	3.25E-07	
Np-238	3.39E-04	4.23E-06		Tl-209	1.81E-05	2.25E-07	
Np-239	9.28E-01	1.16E-02		U-233	2.23E-02	2.78E-04	Yes
Np-240m	1.18E-10	1.47E-12		U-234	1.57E-01	1.95E-03	Yes
Pa-231	6.02E-06	7.51E-08		U-235	7.08E-04	8.83E-06	
Pa-233	7.26E-02	9.05E-04		U-236	1.42E-03	1.77E-05	
Pa-234	5.51E-05	6.86E-07		U-237	1.02E-06	1.28E-08	
Pa-234m	1.67E-02	2.08E-04		U-238	1.67E-02	2.08E-04	Yes
Pb-209	8.36E-04	1.04E-05		U-240	1.18E-10	1.47E-12	
Pb-210	3.01E-05	3.75E-07		Y-90	1.43E+00	1.78E-02	
Pb-211	5.55E-06	6.92E-08		Zr-93	5.23E-01	6.52E-03	Yes
Pb-212	7.25E-05	9.04E-07		TOTAL	8.02E+01	1.00E+00	
Pb-214	3.57E-05	4.45E-07					
Pd-107	1.29E-03	1.61E-05					
Pm-147	4.14E-45	5.16E-47					
Po-210	3.00E-05	3.74E-07					
Po-211	1.56E-08	1.94E-10					
Po-212	4.65E-05	5.79E-07					

APPENDIX F. ACTIVITIES OF DRIED SLUDGE IN YEAR 2515 ($\mu\text{Ci/g}$)

Nuclide	Y 2515 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2515 μCi/g	Fraction of Activity	Reportable
Ac-225	1.04E-03	1.70E-05		Po-213	1.02E-03	1.66E-05	
Ac-227	7.04E-06	1.15E-07		Po-214	5.74E-05	9.39E-07	
Ac-228	7.25E-05	1.19E-06		Po-215	7.04E-06	1.15E-07	
Am-241	1.60E+01	2.61E-01	Yes	Po-216	7.25E-05	1.19E-06	
Am-242	4.50E-02	7.35E-04		Po-218	5.74E-05	9.40E-07	
Am-242m	4.52E-02	7.39E-04	Yes	Pu-238	5.14E+00	8.41E-02	Yes
Am-243	9.19E-01	1.50E-02	Yes	Pu-239	2.01E+01	3.29E-01	Yes
At-217	1.04E-03	1.70E-05		Pu-240	6.37E+00	1.04E-01	Yes
At-218	1.15E-08	1.88E-10		Pu-241	4.15E-02	6.78E-04	Yes
Ba-133	5.54E-16	9.05E-18		Pu-242	6.85E-03	1.12E-04	Yes
Ba-137m	2.56E-03	4.18E-05		Pu-243	6.73E-07	1.10E-08	
Bi-210	5.01E-05	8.20E-07		Pu-244	1.47E-10	2.40E-12	
Bi-211	7.04E-06	1.15E-07		Ra-223	7.04E-06	1.15E-07	
Bi-212	7.25E-05	1.19E-06		Ra-224	7.25E-05	1.19E-06	
Bi-213	1.04E-03	1.70E-05		Ra-225	1.04E-03	1.70E-05	
Bi-214	5.74E-05	9.40E-07		Ra-226	5.74E-05	9.40E-07	
Bk-247	2.10E-03	3.44E-05		Ra-228	7.25E-05	1.19E-06	
Cd-113m	1.84E-11	3.01E-13		Rn-219	7.04E-06	1.15E-07	
Cf-249	3.30E-03	5.39E-05		Rn-220	7.25E-05	1.19E-06	
Cf-250	3.08E-16	5.04E-18		Rn-222	5.74E-05	9.40E-07	
Cf-251	1.56E-02	2.56E-04	Yes	Sb-125	1.02E-56	1.67E-58	
Cl-36	9.74E-01	1.59E-02	Yes	Sb-126	5.02E-02	8.21E-04	
Cm-242	3.73E-02	6.10E-04		Sb-126m	3.59E-01	5.87E-03	
Cm-243	1.10E-06	1.79E-08		Se-79	2.77E-03	4.52E-05	
Cm-244	1.79E-07	2.93E-09		Sm-147	3.96E-09	6.48E-11	
Cm-245	4.14E-02	6.77E-04	Yes	Sm-151	3.70E+00	6.04E-02	Yes
Cm-246	1.20E-02	1.96E-04	Yes	Sn-121	4.03E-04	6.58E-06	
Cm-247	6.73E-07	1.10E-08		Sn-121m	5.19E-04	8.48E-06	
Cm-248	3.77E-05	6.16E-07		Sn-126	3.59E-01	5.87E-03	Yes
Co-60	3.04E-29	4.98E-31		Sr-90	1.32E-01	2.16E-03	Yes
Cs-135	1.61E-03	2.63E-05		Tc-99	1.53E-01	2.50E-03	Yes
Cs-137	2.70E-03	4.42E-05		Te-125m	2.46E-57	4.03E-59	
Eu-154	1.14E-16	1.87E-18		Th-227	6.94E-06	1.14E-07	
Eu-155	4.79E-31	7.83E-33		Th-228	7.25E-05	1.19E-06	
Fr-221	1.04E-03	1.70E-05		Th-229	1.04E-03	1.70E-05	
Fr-223	9.72E-08	1.59E-09		Th-230	6.19E-04	1.01E-05	
I-129	3.77E-04	6.17E-06		Th-231	7.10E-04	1.16E-05	
Nb-93m	5.23E-01	8.55E-03	Yes	Th-232	7.25E-05	1.19E-06	
Ni-59	1.29E+00	2.12E-02	Yes	Th-234	1.67E-02	2.73E-04	
Ni-63	2.88E+00	4.70E-02	Yes	Tl-207	7.02E-06	1.15E-07	
Np-237	7.32E-02	1.20E-03	Yes	Tl-208	2.60E-05	4.26E-07	
Np-238	2.15E-04	3.52E-06		Tl-209	2.24E-05	3.67E-07	
Np-239	9.19E-01	1.50E-02		U-233	2.23E-02	3.65E-04	Yes
Np-240m	1.47E-10	2.40E-12		U-234	1.59E-01	2.60E-03	Yes
Pa-231	7.51E-06	1.23E-07		U-235	7.10E-04	1.16E-05	
Pa-233	7.32E-02	1.20E-03		U-236	1.44E-03	2.35E-05	
Pa-234	5.51E-05	9.01E-07		U-237	1.02E-06	1.66E-08	
Pa-234m	1.67E-02	2.73E-04		U-238	1.67E-02	2.73E-04	Yes
Pb-209	1.04E-03	1.70E-05		U-240	1.47E-10	2.40E-12	
Pb-210	5.01E-05	8.20E-07		Y-90	1.32E-01	2.16E-03	
Pb-211	7.04E-06	1.15E-07		Zr-93	5.23E-01	8.55E-03	Yes
Pb-212	7.25E-05	1.19E-06		TOTAL	6.11E+01	1.00E+00	
Pb-214	5.74E-05	9.39E-07					
Pd-107	1.29E-03	2.11E-05					
Pm-147	1.39E-56	2.28E-58					
Po-210	5.00E-05	8.18E-07					
Po-211	1.97E-08	3.22E-10					
Po-212	4.65E-05	7.60E-07					

APPENDIX G. ACTIVITIES OF DRIED SLUDGE IN YEAR 2615 ($\mu\text{Ci/g}$)

Nuclide	Y 2615 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2615 μCi/g	Fraction of Activity	Reportable
Ac-225	1.24E-03	2.38E-05		Po-213	1.21E-03	2.33E-05	
Ac-227	8.53E-06	1.64E-07		Po-214	8.43E-05	1.62E-06	
Ac-228	7.25E-05	1.39E-06		Po-215	8.53E-06	1.64E-07	
Am-241	1.36E+01	2.61E-01	Yes	Po-216	7.25E-05	1.39E-06	
Am-242	2.85E-02	5.47E-04		Po-218	8.43E-05	1.62E-06	
Am-242m	2.86E-02	5.50E-04	Yes	Pu-238	2.35E+00	4.51E-02	Yes
Am-243	9.11E-01	1.75E-02	Yes	Pu-239	2.01E+01	3.85E-01	Yes
At-217	1.24E-03	2.38E-05		Pu-240	6.30E+00	1.21E-01	Yes
At-218	1.69E-08	3.24E-10		Pu-241	4.11E-02	7.90E-04	Yes
Ba-133	8.71E-19	1.67E-20		Pu-242	6.86E-03	1.32E-04	Yes
Ba-137m	2.54E-04	4.87E-06		Pu-243	7.40E-07	1.42E-08	
Bi-210	7.54E-05	1.45E-06		Pu-244	1.76E-10	3.37E-12	
Bi-211	8.53E-06	1.64E-07		Ra-223	8.53E-06	1.64E-07	
Bi-212	7.25E-05	1.39E-06		Ra-224	7.25E-05	1.39E-06	
Bi-213	1.24E-03	2.38E-05		Ra-225	1.24E-03	2.38E-05	
Bi-214	8.43E-05	1.62E-06		Ra-226	8.43E-05	1.62E-06	
Bk-247	2.00E-03	3.84E-05		Ra-228	7.25E-05	1.39E-06	
Cd-113m	1.12E-13	2.16E-15		Rn-219	8.53E-06	1.64E-07	
Cf-249	2.70E-03	5.19E-05		Rn-220	7.25E-05	1.39E-06	
Cf-250	1.54E-18	2.95E-20		Rn-222	8.43E-05	1.62E-06	
Cf-251	1.45E-02	2.78E-04	Yes	Sb-125	1.38E-67	2.64E-69	
Cl-36	9.74E-01	1.87E-02	Yes	Sb-126	5.02E-02	9.63E-04	
Cm-242	2.36E-02	4.54E-04		Sb-126m	3.58E-01	6.88E-03	
Cm-243	9.62E-08	1.85E-09		Se-79	2.76E-03	5.30E-05	
Cm-244	3.90E-09	7.48E-11		Sm-147	3.96E-09	7.60E-11	
Cm-245	4.11E-02	7.89E-04	Yes	Sm-151	1.71E+00	3.28E-02	Yes
Cm-246	1.18E-02	2.27E-04	Yes	Sn-121	1.14E-04	2.19E-06	
Cm-247	7.40E-07	1.42E-08		Sn-121m	1.47E-04	2.82E-06	
Cm-248	3.77E-05	7.23E-07		Sn-126	3.58E-01	6.88E-03	Yes
Co-60	5.91E-35	1.13E-36		Sr-90	1.22E-02	2.35E-04	Yes
Cs-135	1.61E-03	3.09E-05		Tc-99	1.53E-01	2.93E-03	Yes
Cs-137	2.68E-04	5.15E-06		Te-125m	3.33E-68	6.39E-70	
Eu-154	4.34E-20	8.33E-22		Th-227	8.41E-06	1.61E-07	
Eu-155	4.08E-37	7.83E-39		Th-228	7.25E-05	1.39E-06	
Fr-221	1.24E-03	2.38E-05		Th-229	1.24E-03	2.38E-05	
Fr-223	1.18E-07	2.26E-09		Th-230	7.62E-04	1.46E-05	
I-129	3.77E-04	7.24E-06		Th-231	7.12E-04	1.37E-05	
Nb-93m	5.23E-01	1.00E-02	Yes	Th-232	7.25E-05	1.39E-06	
Ni-59	1.29E+00	2.48E-02	Yes	Th-234	1.67E-02	3.21E-04	
Ni-63	1.40E+00	2.68E-02	Yes	Tl-207	8.50E-06	1.63E-07	
Np-237	7.37E-02	1.41E-03	Yes	Tl-208	2.60E-05	5.00E-07	
Np-238	1.36E-04	2.62E-06		Tl-209	2.67E-05	5.13E-07	
Np-239	9.11E-01	1.75E-02		U-233	2.23E-02	4.29E-04	Yes
Np-240m	1.76E-10	3.37E-12		U-234	1.60E-01	3.07E-03	Yes
Pa-231	9.00E-06	1.73E-07		U-235	7.12E-04	1.37E-05	
Pa-233	7.37E-02	1.41E-03		U-236	1.46E-03	2.80E-05	
Pa-234	5.51E-05	1.06E-06		U-237	1.01E-06	1.94E-08	
Pa-234m	1.67E-02	3.20E-04		U-238	1.67E-02	3.21E-04	Yes
Pb-209	1.24E-03	2.38E-05		U-240	1.76E-10	3.37E-12	
Pb-210	7.54E-05	1.45E-06		Y-90	1.22E-02	2.35E-04	
Pb-211	8.53E-06	1.64E-07		Zr-93	5.23E-01	1.00E-02	Yes
Pb-212	7.25E-05	1.39E-06		TOTAL	5.21E+01	1.00E+00	
Pb-214	8.43E-05	1.62E-06					
Pd-107	1.29E-03	2.48E-05					
Pm-147	4.65E-68	8.92E-70					
Po-210	7.52E-05	1.44E-06					
Po-211	2.39E-08	4.58E-10					
Po-212	4.65E-05	8.92E-07					

APPENDIX H. ACTIVITIES OF DRIED SLUDGE IN YEAR 2715 ($\mu\text{Ci/g}$)

Nuclide	Y 2715 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 2715 μCi/g	Fraction of Activity	Reportable
Ac-225	1.44E-03	3.06E-05		Po-213	1.41E-03	2.99E-05	
Ac-227	1.00E-05	2.13E-07		Po-214	1.16E-04	2.47E-06	
Ac-228	7.25E-05	1.54E-06		Po-215	1.00E-05	2.13E-07	
Am-241	1.16E+01	2.47E-01	Yes	Po-216	7.25E-05	1.54E-06	
Am-242	1.81E-02	3.84E-04		Po-218	1.16E-04	2.47E-06	
Am-242m	1.82E-02	3.86E-04	Yes	Pu-238	1.08E+00	2.29E-02	Yes
Am-243	9.02E-01	1.92E-02	Yes	Pu-239	2.00E+01	4.26E-01	Yes
At-217	1.44E-03	3.06E-05		Pu-240	6.23E+00	1.33E-01	Yes
At-218	2.32E-08	4.94E-10		Pu-241	4.08E-02	8.69E-04	Yes
Ba-133	1.37E-21	2.92E-23		Pu-242	6.86E-03	1.46E-04	Yes
Ba-137m	2.52E-05	5.36E-07		Pu-243	8.02E-07	1.71E-08	
Bi-210	1.06E-04	2.25E-06		Pu-244	2.05E-10	4.36E-12	
Bi-211	1.00E-05	2.13E-07		Ra-223	1.00E-05	2.13E-07	
Bi-212	7.25E-05	1.54E-06		Ra-224	7.25E-05	1.54E-06	
Bi-213	1.44E-03	3.06E-05		Ra-225	1.44E-03	3.06E-05	
Bi-214	1.16E-04	2.47E-06		Ra-226	1.16E-04	2.47E-06	
Bk-247	1.90E-03	4.05E-05		Ra-228	7.25E-05	1.54E-06	
Cd-113m	6.88E-16	1.46E-17		Rn-219	1.00E-05	2.13E-07	
Cf-249	2.22E-03	4.72E-05		Rn-220	7.25E-05	1.54E-06	
Cf-250	7.69E-21	1.64E-22		Rn-222	1.16E-04	2.47E-06	
Cf-251	1.34E-02	2.85E-04	Yes	Sb-125	1.87E-78	3.99E-80	
Cl-36	9.73E-01	2.07E-02	Yes	Sb-126	5.02E-02	1.07E-03	
Cm-242	1.50E-02	3.19E-04		Sb-126m	3.58E-01	7.62E-03	
Cm-243	8.45E-09	1.80E-10		Se-79	2.76E-03	5.87E-05	
Cm-244	8.48E-11	1.81E-12		Sm-147	3.96E-09	8.43E-11	
Cm-245	4.08E-02	8.68E-04	Yes	Sm-151	7.92E-01	1.69E-02	Yes
Cm-246	1.16E-02	2.48E-04	Yes	Sn-121	3.24E-05	6.89E-07	
Cm-247	8.02E-07	1.71E-08		Sn-121m	4.17E-05	8.88E-07	
Cm-248	3.76E-05	8.01E-07		Sn-126	3.58E-01	7.62E-03	Yes
Co-60	1.15E-40	2.45E-42		Sr-90	1.13E-03	2.41E-05	
Cs-135	1.61E-03	3.43E-05		Tc-99	1.53E-01	3.25E-03	Yes
Cs-137	2.66E-05	5.67E-07		Te-125m	4.53E-79	9.65E-81	
Eu-154	1.65E-23	3.51E-25		Th-227	9.88E-06	2.10E-07	
Eu-155	3.48E-43	7.41E-45		Th-228	7.25E-05	1.54E-06	
Fr-221	1.44E-03	3.06E-05		Th-229	1.44E-03	3.06E-05	
Fr-223	1.38E-07	2.94E-09		Th-230	9.05E-04	1.93E-05	
I-129	3.77E-04	8.02E-06		Th-231	7.14E-04	1.52E-05	
Nb-93m	5.23E-01	1.11E-02	Yes	Th-232	7.25E-05	1.54E-06	
Ni-59	1.29E+00	2.75E-02	Yes	Th-234	1.67E-02	3.55E-04	
Ni-63	6.79E-01	1.44E-02	Yes	Tl-207	9.99E-06	2.13E-07	
Np-237	7.41E-02	1.58E-03	Yes	Tl-208	2.60E-05	5.54E-07	
Np-238	8.64E-05	1.84E-06		Tl-209	3.10E-05	6.60E-07	
Np-239	9.02E-01	1.92E-02		U-233	2.24E-02	4.76E-04	Yes
Np-240m	2.05E-10	4.35E-12		U-234	1.60E-01	3.41E-03	Yes
Pa-231	1.05E-05	2.23E-07		U-235	7.14E-04	1.52E-05	
Pa-233	7.41E-02	1.58E-03		U-236	1.48E-03	3.14E-05	
Pa-234	5.51E-05	1.17E-06		U-237	1.00E-06	2.13E-08	
Pa-234m	1.67E-02	3.55E-04		U-238	1.67E-02	3.55E-04	Yes
Pb-209	1.44E-03	3.06E-05		U-240	2.05E-10	4.35E-12	
Pb-210	1.06E-04	2.25E-06		Y-90	1.13E-03	2.41E-05	
Pb-211	1.00E-05	2.13E-07		Zr-93	5.23E-01	1.11E-02	Yes
Pb-212	7.25E-05	1.54E-06		TOTAL	4.70E+01	1.00E+00	
Pb-214	1.16E-04	2.47E-06					
Pd-107	1.29E-03	2.75E-05					
Pm-147	1.56E-79	3.33E-81					
Po-210	1.05E-04	2.24E-06					
Po-211	2.80E-08	5.97E-10					
Po-212	4.65E-05	9.89E-07					

APPENDIX I. ACTIVITIES OF DRIED SLUDGE IN YEAR 2815 ($\mu\text{Ci/g}$)

Nuclide	Y 2815 μCi/g	Fraction of Activity	Reportable
Ac-225	1.63E-03	3.73E-05	
Ac-227	1.15E-05	2.63E-07	
Ac-228	7.25E-05	1.66E-06	
Am-241	9.87E+00	2.26E-01	Yes
Am-242	1.14E-02	2.62E-04	
Am-242m	1.15E-02	2.63E-04	Yes
Am-243	8.94E-01	2.04E-02	Yes
At-217	1.63E-03	3.73E-05	
At-218	3.05E-08	6.98E-10	
Ba-133	2.16E-24	4.93E-26	
Ba-137m	2.50E-06	5.71E-08	
Bi-210	1.41E-04	3.21E-06	
Bi-211	1.15E-05	2.63E-07	
Bi-212	7.25E-05	1.66E-06	
Bi-213	1.63E-03	3.73E-05	
Bi-214	1.53E-04	3.49E-06	
Bk-247	1.81E-03	4.13E-05	
Cd-113m	4.20E-18	9.60E-20	
Cf-249	1.82E-03	4.16E-05	
Cf-250	3.84E-23	8.77E-25	
Cf-251	1.24E-02	2.83E-04	Yes
Cl-36	9.73E-01	2.22E-02	Yes
Cm-242	9.49E-03	2.17E-04	
Cm-243	7.42E-10	1.70E-11	
Cm-244	1.85E-12	4.22E-14	
Cm-245	4.05E-02	9.25E-04	Yes
Cm-246	1.15E-02	2.62E-04	Yes
Cm-247	8.59E-07	1.96E-08	
Cm-248	3.76E-05	8.60E-07	
Co-60	2.24E-46	5.11E-48	
Cs-135	1.61E-03	3.68E-05	
Cs-137	2.64E-06	6.03E-08	
Eu-154	6.25E-27	1.43E-28	
Eu-155	2.97E-49	6.78E-51	
Fr-221	1.63E-03	3.73E-05	
Fr-223	1.59E-07	3.63E-09	
I-129	3.77E-04	8.62E-06	
Nb-93m	5.23E-01	1.19E-02	Yes
Ni-59	1.29E+00	2.95E-02	Yes
Ni-63	3.30E-01	7.53E-03	Yes
Np-237	7.44E-02	1.70E-03	Yes
Np-238	5.48E-05	1.25E-06	
Np-239	8.94E-01	2.04E-02	
Np-240m	2.33E-10	5.34E-12	
Pa-231	1.20E-05	2.74E-07	
Pa-233	7.44E-02	1.70E-03	
Pa-234	5.51E-05	1.26E-06	
Pa-234m	1.67E-02	3.81E-04	
Pb-209	1.63E-03	3.73E-05	
Pb-210	1.41E-04	3.21E-06	
Pb-211	1.15E-05	2.63E-07	
Pb-212	7.25E-05	1.66E-06	
Pb-214	1.53E-04	3.49E-06	
Pd-107	1.29E-03	2.95E-05	
Pm-147	5.22E-91	1.19E-92	
Po-210	1.40E-04	3.21E-06	
Po-211	3.22E-08	7.36E-10	
Po-212	4.65E-05	1.06E-06	

Nuclide	Y 2815 μCi/g	Fraction of Activity	Reportable
Po-213	1.60E-03	3.65E-05	
Po-214	1.53E-04	3.49E-06	
Po-215	1.15E-05	2.63E-07	
Po-216	7.25E-05	1.66E-06	
Po-218	1.53E-04	3.49E-06	
Pu-238	4.95E-01	1.13E-02	Yes
Pu-239	2.00E+01	4.56E-01	Yes
Pu-240	6.17E+00	1.41E-01	Yes
Pu-241	4.05E-02	9.26E-04	Yes
Pu-242	6.86E-03	1.57E-04	Yes
Pu-243	8.59E-07	1.96E-08	
Pu-244	2.34E-10	5.34E-12	
Ra-223	1.15E-05	2.63E-07	
Ra-224	7.25E-05	1.66E-06	
Ra-225	1.63E-03	3.73E-05	
Ra-226	1.53E-04	3.49E-06	
Ra-228	7.25E-05	1.66E-06	
Rn-219	1.15E-05	2.63E-07	
Rn-220	7.25E-05	1.66E-06	
Rn-222	1.53E-04	3.49E-06	
Sb-125	2.53E-89	5.79E-91	
Sb-126	5.01E-02	1.15E-03	
Sb-126m	3.58E-01	8.18E-03	
Se-79	2.76E-03	6.30E-05	
Sm-147	3.96E-09	9.05E-11	
Sm-151	3.67E-01	8.38E-03	Yes
Sn-121	9.18E-06	2.10E-07	
Sn-121m	1.18E-05	2.70E-07	
Sn-126	3.58E-01	8.18E-03	Yes
Sr-90	1.05E-04	2.39E-06	
Tc-99	1.53E-01	3.49E-03	Yes
Te-125m	6.13E-90	1.40E-91	
Th-227	1.13E-05	2.59E-07	
Th-228	7.25E-05	1.66E-06	
Th-229	1.63E-03	3.73E-05	
Th-230	1.05E-03	2.40E-05	
Th-231	7.16E-04	1.64E-05	
Th-232	7.25E-05	1.66E-06	
Th-234	1.67E-02	3.82E-04	
Tl-207	1.15E-05	2.62E-07	
Tl-208	2.60E-05	5.95E-07	
Tl-209	3.53E-05	8.06E-07	
U-233	2.24E-02	5.11E-04	Yes
U-234	1.60E-01	3.66E-03	Yes
U-235	7.16E-04	1.64E-05	
U-236	1.49E-03	3.41E-05	
U-237	9.93E-07	2.27E-08	
U-238	1.67E-02	3.82E-04	Yes
U-240	2.33E-10	5.34E-12	
Y-90	1.05E-04	2.39E-06	
Zr-93	5.23E-01	1.19E-02	Yes
TOTAL	4.38E+01	1.00E+00	

APPENDIX J. ACTIVITIES OF DRIED SLUDGE IN YEAR 2915 ($\mu\text{Ci/g}$)

Nuclide	Y 2015 μCi/g	Fraction of Activity	Reportable
Ac-225	1.83E-03	4.40E-05	
Ac-227	1.30E-05	3.13E-07	
Ac-228	7.25E-05	1.75E-06	
Am-241	8.42E+00	2.03E-01	Yes
Am-242	7.25E-03	1.75E-04	
Am-242m	7.29E-03	1.76E-04	Yes
Am-243	8.86E-01	2.13E-02	Yes
At-217	1.83E-03	4.40E-05	
At-218	3.87E-08	9.33E-10	
Ba-133	3.40E-27	8.19E-29	
Ba-137m	2.48E-07	5.97E-09	
Bi-210	1.80E-04	4.34E-06	
Bi-211	1.30E-05	3.13E-07	
Bi-212	7.25E-05	1.75E-06	
Bi-213	1.83E-03	4.40E-05	
Bi-214	1.94E-04	4.66E-06	
Bk-247	1.72E-03	4.14E-05	
Cd-113m	2.57E-20	6.20E-22	
Cf-249	1.49E-03	3.60E-05	
Cf-250	1.92E-25	4.62E-27	
Cf-251	1.15E-02	2.76E-04	Yes
Cl-36	9.73E-01	2.34E-02	Yes
Cm-242	6.02E-03	1.45E-04	
Cm-243	6.52E-11	1.57E-12	
Cm-244	4.02E-14	9.68E-16	
Cm-245	4.01E-02	9.67E-04	Yes
Cm-246	1.13E-02	2.72E-04	Yes
Cm-247	9.12E-07	2.20E-08	
Cm-248	3.76E-05	9.06E-07	
Co-60	4.36E-52	1.05E-53	
Cs-135	1.61E-03	3.88E-05	
Cs-137	2.62E-07	6.31E-09	
Eu-154	2.37E-30	5.72E-32	
Eu-155	2.53E-55	6.10E-57	
Fr-221	1.83E-03	4.40E-05	
Fr-223	1.79E-07	4.32E-09	
I-129	3.77E-04	9.08E-06	
Nb-93m	5.23E-01	1.26E-02	Yes
Ni-59	1.29E+00	3.11E-02	Yes
Ni-63	1.60E-01	3.86E-03	Yes
Np-237	7.47E-02	1.80E-03	Yes
Np-238	3.47E-05	8.36E-07	
Np-239	8.86E-01	2.13E-02	
Np-240m	2.62E-10	6.32E-12	
Pa-231	1.35E-05	3.24E-07	
Pa-233	7.47E-02	1.80E-03	
Pa-234	5.51E-05	1.33E-06	
Pa-234m	1.67E-02	4.01E-04	
Pb-209	1.83E-03	4.40E-05	
Pb-210	1.80E-04	4.34E-06	
Pb-211	1.30E-05	3.13E-07	
Pb-212	7.25E-05	1.75E-06	
Pb-214	1.94E-04	4.66E-06	
Pd-107	1.29E-03	3.11E-05	
Pm-147	1.76E-102	4.23E-104	
Po-210	1.80E-04	4.33E-06	
Po-211	3.64E-08	8.76E-10	
Po-212	4.65E-05	1.12E-06	

Nuclide	Y 2015 μCi/g	Fraction of Activity	Reportable
Po-213	1.79E-03	4.31E-05	
Po-214	1.94E-04	4.66E-06	
Po-215	1.30E-05	3.13E-07	
Po-216	7.25E-05	1.75E-06	
Po-218	1.94E-04	4.66E-06	
Pu-238	2.29E-01	5.51E-03	Yes
Pu-239	1.99E+01	4.79E-01	Yes
Pu-240	6.10E+00	1.47E-01	Yes
Pu-241	4.02E-02	9.68E-04	Yes
Pu-242	6.86E-03	1.65E-04	Yes
Pu-243	9.12E-07	2.20E-08	
Pu-244	2.63E-10	6.33E-12	
Ra-223	1.30E-05	3.13E-07	
Ra-224	7.25E-05	1.75E-06	
Ra-225	1.83E-03	4.40E-05	
Ra-226	1.94E-04	4.66E-06	
Ra-228	7.25E-05	1.75E-06	
Rn-219	1.30E-05	3.13E-07	
Rn-220	7.25E-05	1.75E-06	
Rn-222	1.94E-04	4.66E-06	
Sb-125	3.45E-100	8.31E-102	
Sb-126	5.01E-02	1.21E-03	
Sb-126m	3.58E-01	8.62E-03	
Se-79	2.75E-03	6.63E-05	
Sm-147	3.96E-09	9.54E-11	
Sm-151	1.70E-01	4.09E-03	Yes
Sn-121	2.60E-06	6.27E-08	
Sn-121m	3.35E-06	8.08E-08	
Sn-126	3.58E-01	8.62E-03	Yes
Sr-90	9.69E-06	2.34E-07	
Tc-99	1.53E-01	3.67E-03	Yes
Te-125m	8.35E-101	2.01E-102	
Th-227	1.28E-05	3.09E-07	
Th-228	7.25E-05	1.75E-06	
Th-229	1.83E-03	4.40E-05	
Th-230	1.19E-03	2.87E-05	
Th-231	7.18E-04	1.73E-05	
Th-232	7.25E-05	1.75E-06	
Th-234	1.67E-02	4.02E-04	
Tl-207	1.30E-05	3.12E-07	
Tl-208	2.60E-05	6.27E-07	
Tl-209	3.95E-05	9.51E-07	
U-233	2.24E-02	5.40E-04	Yes
U-234	1.60E-01	3.86E-03	Yes
U-235	7.18E-04	1.73E-05	
U-236	1.51E-03	3.64E-05	
U-237	9.85E-07	2.37E-08	
U-238	1.67E-02	4.02E-04	Yes
U-240	2.62E-10	6.32E-12	
Y-90	9.70E-06	2.34E-07	
Zr-93	5.23E-01	1.26E-02	Yes
TOTAL	4.15E+01	1.00E+00	

APPENDIX K. ACTIVITIES OF DRIED SLUDGE IN YEAR 3015 ($\mu\text{Ci/g}$)

Nuclide	Y 3015 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 3015 μCi/g	Fraction of Activity	Reportable
Ac-225	2.02E-03	5.08E-05		Po-213	1.98E-03	4.97E-05	
Ac-227	1.45E-05	3.64E-07		Po-214	2.39E-04	6.00E-06	
Ac-228	7.25E-05	1.82E-06		Po-215	1.45E-05	3.64E-07	
Am-241	7.18E+00	1.80E-01	Yes	Po-216	7.25E-05	1.82E-06	
Am-242	4.60E-03	1.15E-04		Po-218	2.39E-04	6.00E-06	
Am-242m	4.62E-03	1.16E-04	Yes	Pu-238	1.06E-01	2.67E-03	Yes
Am-243	8.77E-01	2.20E-02	Yes	Pu-239	1.98E+01	4.98E-01	Yes
At-217	2.02E-03	5.08E-05		Pu-240	6.04E+00	1.52E-01	Yes
At-218	4.78E-08	1.20E-09		Pu-241	3.99E-02	1.00E-03	Yes
Ba-133	5.36E-30	1.35E-31		Pu-242	6.86E-03	1.72E-04	Yes
Ba-137m	2.46E-08	6.17E-10		Pu-243	9.61E-07	2.41E-08	
Bi-210	2.24E-04	5.63E-06		Pu-244	2.92E-10	7.32E-12	
Bi-211	1.45E-05	3.64E-07		Ra-223	1.45E-05	3.64E-07	
Bi-212	7.25E-05	1.82E-06		Ra-224	7.25E-05	1.82E-06	
Bi-213	2.02E-03	5.08E-05		Ra-225	2.02E-03	5.08E-05	
Bi-214	2.39E-04	6.00E-06		Ra-226	2.39E-04	6.00E-06	
Bk-247	1.64E-03	4.10E-05		Ra-228	7.25E-05	1.82E-06	
Cd-113m	1.57E-22	3.95E-24		Rn-219	1.45E-05	3.64E-07	
Cf-249	1.23E-03	3.08E-05		Rn-220	7.25E-05	1.82E-06	
Cf-250	9.59E-28	2.41E-29		Rn-222	2.39E-04	6.00E-06	
Cf-251	1.06E-02	2.67E-04	Yes	Sb-125	4.70E-111	1.18E-112	
Cl-36	9.73E-01	2.44E-02	Yes	Sb-126	5.01E-02	1.26E-03	
Cm-242	3.81E-03	9.57E-05		Sb-126m	3.58E-01	8.97E-03	
Cm-243	5.73E-12	1.44E-13		Se-79	2.75E-03	6.90E-05	
Cm-244	8.75E-16	2.20E-17		Sm-147	3.96E-09	9.94E-11	
Cm-245	3.98E-02	1.00E-03	Yes	Sm-151	7.86E-02	1.97E-03	Yes
Cm-246	1.11E-02	2.79E-04	Yes	Sn-121	7.38E-07	1.85E-08	
Cm-247	9.61E-07	2.41E-08		Sn-121m	9.51E-07	2.39E-08	
Cm-248	3.76E-05	9.44E-07		Sn-126	3.58E-01	8.97E-03	Yes
Co-60	8.49E-58	2.13E-59		Sr-90	8.97E-07	2.25E-08	
Cs-135	1.61E-03	4.04E-05		Tc-99	1.53E-01	3.83E-03	Yes
Cs-137	2.60E-08	6.53E-10		Te-125m	1.14E-111	2.85E-113	
Eu-154	9.02E-34	2.26E-35		Th-227	1.43E-05	3.59E-07	
Eu-155	2.16E-61	5.43E-63		Th-228	7.25E-05	1.82E-06	
Fr-221	2.02E-03	5.08E-05		Th-229	2.02E-03	5.08E-05	
Fr-223	2.00E-07	5.02E-09		Th-230	1.34E-03	3.35E-05	
I-129	3.77E-04	9.46E-06		Th-231	7.20E-04	1.81E-05	
Nb-93m	5.23E-01	1.31E-02	Yes	Th-232	7.25E-05	1.82E-06	
Ni-59	1.29E+00	3.23E-02	Yes	Th-234	1.67E-02	4.19E-04	
Ni-63	7.78E-02	1.95E-03	Yes	Tl-207	1.44E-05	3.63E-07	
Np-237	7.50E-02	1.88E-03	Yes	Tl-208	2.60E-05	6.54E-07	
Np-238	2.20E-05	5.52E-07		Tl-209	4.37E-05	1.10E-06	
Np-239	8.77E-01	2.20E-02		U-233	2.24E-02	5.63E-04	Yes
Np-240m	2.91E-10	7.31E-12		U-234	1.60E-01	4.03E-03	Yes
Pa-231	1.50E-05	3.75E-07		U-235	7.20E-04	1.81E-05	
Pa-233	7.50E-02	1.88E-03		U-236	1.53E-03	3.84E-05	
Pa-234	5.51E-05	1.38E-06		U-237	9.77E-07	2.45E-08	
Pa-234m	1.67E-02	4.18E-04		U-238	1.67E-02	4.19E-04	Yes
Pb-209	2.02E-03	5.08E-05		U-240	2.91E-10	7.31E-12	
Pb-210	2.24E-04	5.63E-06		Y-90	8.97E-07	2.25E-08	
Pb-211	1.45E-05	3.64E-07		Zr-93	5.23E-01	1.31E-02	Yes
Pb-212	7.25E-05	1.82E-06		TOTAL	3.98E+01	1.00E+00	
Pb-214	2.39E-04	6.00E-06					
Pd-107	1.29E-03	3.24E-05					
Pm-147	5.90E-114	1.48E-115					
Po-210	2.24E-04	5.62E-06					
Po-211	4.06E-08	1.02E-09					
Po-212	4.65E-05	1.17E-06					

APPENDIX L. ACTIVITIES OF DRIED SLUDGE IN YEAR 3115 ($\mu\text{Ci/g}$)

Nuclide	Y 3115 μCi/g	Fraction of Activity	Reportable	Nuclide	Y 3115 μCi/g	Fraction of Activity	Reportable
Ac-225	2.21E-03	5.75E-05		Po-213	2.17E-03	5.63E-05	
Ac-227	1.60E-05	4.15E-07		Po-214	2.89E-04	7.49E-06	
Ac-228	7.25E-05	1.88E-06		Po-215	1.60E-05	4.15E-07	
Am-241	6.12E+00	1.59E-01	Yes	Po-216	7.25E-05	1.88E-06	
Am-242	2.91E-03	7.57E-05		Po-218	2.89E-04	7.50E-06	
Am-242m	2.93E-03	7.61E-05		Pu-238	4.99E-02	1.30E-03	Yes
Am-243	8.69E-01	2.26E-02	Yes	Pu-239	1.98E+01	5.14E-01	Yes
At-217	2.21E-03	5.75E-05		Pu-240	5.97E+00	1.55E-01	Yes
At-218	5.77E-08	1.50E-09		Pu-241	3.96E-02	1.03E-03	Yes
Ba-133	8.43E-33	2.19E-34		Pu-242	6.86E-03	1.78E-04	Yes
Ba-137m	2.44E-09	6.34E-11		Pu-243	1.01E-06	2.61E-08	
Bi-210	2.72E-04	7.07E-06		Pu-244	3.21E-10	8.33E-12	
Bi-211	1.60E-05	4.15E-07		Ra-223	1.60E-05	4.15E-07	
Bi-212	7.25E-05	1.88E-06		Ra-224	7.25E-05	1.88E-06	
Bi-213	2.21E-03	5.75E-05		Ra-225	2.21E-03	5.75E-05	
Bi-214	2.89E-04	7.50E-06		Ra-226	2.89E-04	7.50E-06	
Bk-247	1.55E-03	4.04E-05		Ra-228	7.25E-05	1.88E-06	
Cd-113m	9.63E-25	2.50E-26		Rn-219	1.60E-05	4.15E-07	
Cf-249	1.01E-03	2.61E-05		Rn-220	7.25E-05	1.88E-06	
Cf-250	4.79E-30	1.24E-31		Rn-222	2.89E-04	7.50E-06	
Cf-251	9.84E-03	2.56E-04	Yes	Sb-125	6.35E-122	1.65E-123	
Cl-36	9.73E-01	2.53E-02	Yes	Sb-126	5.00E-02	1.30E-03	
Cm-242	2.42E-03	6.28E-05		Sb-126m	3.57E-01	9.28E-03	
Cm-243	5.03E-13	1.31E-14		Se-79	2.75E-03	7.14E-05	
Cm-244	1.90E-17	4.95E-19		Sm-147	3.96E-09	1.03E-10	
Cm-245	3.95E-02	1.03E-03	Yes	Sm-151	3.64E-02	9.45E-04	Yes
Cm-246	1.10E-02	2.85E-04	Yes	Sn-121	2.09E-07	5.44E-09	
Cm-247	1.01E-06	2.61E-08		Sn-121m	2.70E-07	7.01E-09	
Cm-248	3.76E-05	9.77E-07		Sn-126	3.57E-01	9.28E-03	Yes
Co-60	1.65E-63	4.28E-65		Sr-90	8.30E-08	2.16E-09	
Cs-135	1.61E-03	4.18E-05		Tc-99	1.52E-01	3.96E-03	Yes
Cs-137	2.58E-09	6.70E-11		Te-125m	1.54E-122	3.99E-124	
Eu-154	3.42E-37	8.88E-39		Th-227	1.58E-05	4.09E-07	
Eu-155	1.84E-67	4.79E-69		Th-228	7.25E-05	1.88E-06	
Fr-221	2.21E-03	5.75E-05		Th-229	2.21E-03	5.75E-05	
Fr-223	2.20E-07	5.73E-09		Th-230	1.48E-03	3.84E-05	
I-129	3.77E-04	9.79E-06		Th-231	7.22E-04	1.88E-05	
Nb-93m	5.23E-01	1.36E-02	Yes	Th-232	7.25E-05	1.88E-06	
Ni-59	1.29E+00	3.34E-02	Yes	Th-234	1.67E-02	4.34E-04	
Ni-63	3.78E-02	9.81E-04	Yes	Tl-207	1.59E-05	4.14E-07	
Np-237	7.52E-02	1.95E-03	Yes	Tl-208	2.60E-05	6.77E-07	
Np-238	1.39E-05	3.62E-07		Tl-209	4.78E-05	1.24E-06	
Np-239	8.69E-01	2.26E-02		U-233	2.24E-02	5.83E-04	Yes
Np-240m	3.20E-10	8.32E-12		U-234	1.60E-01	4.16E-03	Yes
Pa-231	1.64E-05	4.27E-07		U-235	7.22E-04	1.88E-05	
Pa-233	7.52E-02	1.95E-03		U-236	1.55E-03	4.02E-05	
Pa-234	5.51E-05	1.43E-06		U-237	9.70E-07	2.52E-08	
Pa-234m	1.67E-02	4.33E-04		U-238	1.67E-02	4.34E-04	Yes
Pb-209	2.21E-03	5.75E-05		U-240	3.20E-10	8.32E-12	
Pb-210	2.72E-04	7.07E-06		Y-90	8.30E-08	2.16E-09	
Pb-211	1.60E-05	4.15E-07		Zr-93	5.23E-01	1.36E-02	Yes
Pb-212	7.25E-05	1.88E-06		TOTAL	3.85E+01	1.00E+00	
Pb-214	2.89E-04	7.49E-06					
Pd-107	1.29E-03	3.35E-05					
Pm-147	1.97E-125	5.12E-127					
Po-210	2.72E-04	7.07E-06					
Po-211	4.47E-08	1.16E-09					
Po-212	4.65E-05	1.21E-06					

This page intentionally left blank.

Distribution:

S. L. Marra, 773-A
C. C. Herman, 999-W
A. B. Barnes, 999-W
B. J. Giddings, 786-5A
S. D. Fink, 773-A
L. M. Chandler, 773-A
R. H. Young, 773-A
F. M. Pennebaker, 773-42A
D. A. Crowley, 773-43A
J. P. Vaughan, 773-41A
N. E. Bibler, 773-A
J. M. Pareizs, 773-A
S. H. Reboul, 773-A
D. K. Peeler, 999-W
D. C. Koopman, 999-W
M. E. Stone, 999-W
D. P. Lambert, 999-W
B. R. Pickenheim, 999-W
T. B. Edwards, 999-W

J. E. Occhipinti, 704-S
D. C. Sherburne, 704-S
J. W. Ray, 704-S
R. N. Hinds, 704-S
J. F. Iaukea, 704-30S
M. A. Broome, 704-29S
R. T. McNew, 704-27S
A. R. Shafer, 704-27S
M. C. Clark, 704-27S
J. M. Bricker, 704-27S
T. L. Fellingner, 704-26S
H. H. Elder, 704-24S

H. B. Shah, 766-H
J. M. Gillam, 766-H
D. D. Larsen, 766-H
B. A. Hamm, 766-H