



Using SNP arrays to leverage historic data sets for improved prediction accuracy and estimation of GxE of fruit maturity in sweet cherry

S605

Craig Hardner¹, Stijn Vanderzande², Lichun Cai³, Julia Piaskowski², Nnadozie Oraguzie⁴, Jose Quero-Garcia⁵, José Antonio Campoy⁵, Teresa Barreneche⁵, Ana Wünsch⁶, Daniela Giovannini⁷, Alessandro Liverani⁷, Thomas Sotiropoulos⁸, Konstantinos Kazantzis⁸, Satish Kumar⁹, Ben Hayes², Cameron Peace²

¹ **University of Queensland, Queensland Alliance for Agriculture and Food Innovation, St Lucia 4072 QLD, Australia, 306 Carmody Road, St Lucia 4076, Australia**

² **Department of Horticulture, Washington State University, Pullman, WA 99164, USA**

³ **Michigan State Univ, Dept Hort, E Lansing, MI 48824 USA**

⁴ **Washington State Univ, Irrigated Agr Res & Extens Ctr, 24106 N Bunn Rd, Prosser, WA 99350, USA**

⁵ **INRA, UMR Biol Fruit & Pathol 1332, F-33140 Villenave Dornon, France**

⁶ **Ctr Invest & Tecnol Agroalimentaria Aragon CITA, Zaragoza, Spain**

⁷ **CRA FRF Unita Ric Frutticoltura Forli, I-47121 Forli, Italy**

⁸ **Agri & Agri Food Canada, Pacific Agri Food Res Ctr, Summerland, BC V0H 1Z0, Canada**

⁹ **New Zealand Inst Plant & Food Res Ltd, Havelock North 4157, New Zealand**

Improved prediction accuracy of the genetic potential of advanced selections and cultivars for commercially relevant traits in commercially relevant conditions will lead to increased response to breeding. While a large number of loci might influence quantitative traits, QTLs with major trait effect have been identified, and can be used for selection if the effect of alleles on commercial performance can be accurately predicted. However, selection response will be compromised if the effect of major trait loci are confounded with polygenetic effects from the background genome - if not accounted for, and external factors such as management practices or environment conditions also influence the trait. Here, we propose a novel approach to improve prediction accuracy of genetic potential under commercial conditions by using SNP arrays to characterise variation at major effect QTLs and estimate realised additive and dominance relationships among individuals that have also been evaluated for traits of interest. This approach enables data from otherwise disconnected historical data sets to be combined into a single analysis, thereby increasing prediction accuracy, without the need for replication of individuals. In addition, the different locations from which the performance data have been collected represent samples of the variability of external factors. As such, this analysis can be also used to investigate the influence of these factors on the stability of both major-effect QTLs and polygenetic effects. As part of the RosBREED project, we demonstrate this approach and results using 800 sweet cherry individuals that have were genotyped using the 6K SNP sweet cherry array and evaluated in the USA (Prosser, Washington), Canada (Summerland British Columbia), France (Boudreaux), and Italy (Forli).