

Comparative effects of the marine algal toxins azaspiracid-1, -2, and -3 on Jurkat T lymphocyte cells

Journal:	<i>Chemical Research in Toxicology</i>
Manuscript ID:	tx-2011-00553p.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Twiner, Michael; University of Michigan-Dearborn, Natural Sciences El-Ladki, Racha; University of Michigan-Dearborn, Natural Sciences Kilcoyne, Jane; Irish Marine Institute, Doucette, Gregory; NOAA/National Ocean Service, Marine Biotoxin Program, NOAA

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Comparative effects of the marine algal toxins azaspiracid-1, -2, and -3 on Jurkat T lymphocyte cells

Michael J. Twiner^{†,}, Racha El-Ladki[†], Jane Kilcoyne[‡], and Gregory J. Doucette[§]*

[†]Department of Natural Sciences, University of Michigan – Dearborn, Dearborn, MI USA

[‡]Marine Institute, Renville, Oranmore, Co. Galway, Ireland

[§]Marine Biotoxins Program, NOAA/National Ocean Service, Charleston, SC USA

Corresponding author email: mtwiner@umd.umich.edu

RECEIVED DATE:

TITLE RUNNING HEAD: Cytotoxicity of natural AZA analogs

1
2
3
4
5
6
7 * CORRESPONDING AUTHOR:
8

9
10 Dr. Michael J. Twiner

11
12 Department of Natural Sciences

13
14 University of Michigan-Dearborn

15
16 4901 Evergreen Rd
17


18
19 Dearborn, MI USA 48128
20

21
22 office: (313) 593-5298
23

24
25 fax: (313) 593-4937
26

27
28 email: mtwiner@umd.umich.edu
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TOC GRAPHIC



1
2
3
4
5
6 ABSTRACT
7
8
9

10 Azaspiracids (AZA) are polyether marine toxins of dinoflagellate origin that accumulate in
11 shellfish and represent an emerging human health risk. Although monitored and regulated in
12 many European and Asian countries, there are no monitoring programs or regulatory
13 requirements in the United States for this toxin group. This did not prove to be a problem until
14 June 2009 when AZAs were identified in US seafood for the first time resulting in human
15 intoxications and further expanding their global distribution. Efforts are now underway in several
16 laboratories to better define the effects and mechanism(s) of action for the AZAs. Our
17 investigations have employed Jurkat T lymphocyte cells as an *in vitro* model to characterize the
18 toxicological effects of AZA1, AZA2, and AZA3. Cytotoxicity experiments employing a
19 metabolically-based dye (i.e., MTS) indicated that AZA1, AZA2, and AZA3 each elicited a
20 lethal response that was both concentration- and time- dependent, with EC₅₀ values in the sub- to
21 low nanomolar range. Based on EC₅₀ comparisons, the order of potency was as follows: AZA2 >
22 AZA3 > AZA1, with toxic equivalence factors (TEFs) relative to AZA1 of 8.3-fold and 4.5-fold
23 greater for AZA2 and AZA3, respectively. Image analysis of exposed cells using Nomarski
24 differential interference contrast (DIC) imaging and fluorescent imaging of cellular actin
25 indicated that the morphological effects of AZA1 on this cell type are unique relative to the
26 effects of AZA2 and AZA3. Collectively, our data support the growing body of evidence
27 suggesting that natural analogs of AZA are highly potent and that they may have multiple
28 molecular targets.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **KEYWORDS:** azaspiracid (AZA), cytotoxicity, potency, structure-activity relationship, T
4
5 lymphocytes, toxic equivalence factor (TEF)
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Azaspiracids (AZAs) are a class of polyether marine toxins produced by the marine dinoflagellate *Azadinium spinosum* (1). AZA toxins were originally found in shellfish off the coast of Ireland and have caused multiple human intoxication events throughout Europe (reviewed by Furey et al. (2)) and most recently in the United States (3). Reports of *Azadinium* species and/or AZAs have now been documented for many other parts of the world, including Morocco (4), Chile (5), and Canada (M. Quilliam, pers. comm). AZAs were also recently detected in marine sponges (6). Human symptoms following intoxication include nausea, vomiting, and stomach cramps (7), but thus far no deaths have been attributed to AZAs.

The AZA group now includes more than 20 analogs that are produced either by phytoplankton, through biotransformation in shellfish, or as by-products formed during processing and storage of toxin-containing tissues (7-9). Although there is geographical variation, plankton samples are dominated by AZA1 and AZA2 (Figure 1), with the former often constituting over 80% of the toxin profile (1, 10). The recent isolation and culturing of *A. spinosum* has further confirmed these findings (1, 11), but interestingly, other non-toxigenic *Azadinium* species have now been described (12). Naturally contaminated shellfish often remain above the European regulatory action level for many months following exposure to AZAs (13, 14). Toxin binding proteins may be responsible, in part, for this slow rate of depuration in certain shellfish (15, 16).

Studies in mice exposed orally to AZA1 revealed severe gastrointestinal tract deterioration, adverse histopathologies of the liver, spleen, and thymus, as well as a small incidence of tumor formation (17, 18). The minimal lethal doses for AZA1 are *ca.* 500 $\mu\text{g}/\text{kg}$ (oral) (19) and *ca.* 150 $\mu\text{g}/\text{kg}$ (*i.p.*) (20). Relative analog potencies (via *i.p.*) are as follows: AZA2 > AZA3 > AZA1 > AZA4 > AZA5 (7, 21, 22). Note that AZA4 and AZA5 are the 3- and 23-hydroxyl analogs,

1
2
3 respectively, of AZA3. Although these experimental data are extremely valuable, their clear
4 interpretation has been hindered due to the use of material of unknown purity. Nonetheless, this
5 information has provided the guidance needed to propose toxic equivalence factors (TEFs) (1.0,
6 1.8, and 1.4 for AZA1, AZA2, and AZA3; respectively) that are used for determining the total
7 AZA equivalent concentrations in European shellfish (23). The EU regulatory limit is 0.16 µg/g
8 “total” AZAs in shellfish as detected by LC-MS/MS. Only AZA1-3 are currently regulated.
9

10
11
12
13
14
15
16
17
18 Multiple studies have demonstrated the *in vitro* cytotoxic potential of AZA1 wherein every cell
19 type tested to date appears to be susceptible (see Twiner et al. (7) for a review). AZAs are very
20 potent cytotoxic compounds (6, 24, 25), induce irreversible cytoskeletal rearrangements in
21 mammalian cells (24, 26, 27), stimulate secondary messenger molecules (28, 29), deplete
22 cellular ATP (30), inhibit neuronal ion flux and bioelectrical activity (31, 32), induce cell volume
23 reductions followed by c-Jun N-terminal kinase (JNK) activation (33, 34), inhibit membrane
24 protein endocytosis (35), and up-regulate low density lipoprotein receptor as well as stimulating
25 cholesterol biosynthesis (36). AZA1 has also been shown to be a potent teratogen in developing
26 fish embryos (37). In many cases the toxin concentrations necessary to observe these various
27 effects are in the low nanomolar range (7, 31, 34, 35), but for some endpoints they approach or
28 exceed micromolar concentrations (31, 38). Despite the efforts of many investigators, the
29 mechanism(s) of action of AZAs has not yet been determined.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45
46 The possible involvement of an apoptotic pathway in AZA1 cytotoxicity is complicated and
47 remains unresolved. Results of an *in vitro* study assessing mitochondrial membrane potential in
48 neuroblastoma cells (38) are suggestive of a non-apoptotic pathway; however, up-regulation of
49 caspase activity in neuroblastoma cells (39) and neocortical neurons (31) is supportive of
50 apoptosis activation. Furthermore, apoptotic lymphocyte cells have been observed in the spleens
51
52
53
54
55
56
57
58
59
60

1
2
3 of mice exposed to AZA1 (40). Nonetheless, such discrepancies may be explained, in part, by
4
5 AZA1 causing simultaneous induction of both necrotic and apoptotic mechanisms (31).
6
7

8 Due to difficulties in obtaining sufficient amounts of purified material, much less information
9
10 is available regarding the toxicological properties of the other natural AZA analogs. At relatively
11
12 high concentrations (i.e., 50-1000 nM), AZA2, AZA3, AZA4, and AZA5 have been shown to
13
14 have various effects on intracellular pH, cytosolic calcium, and cAMP in lymphocytes (29, 41,
15
16 42). In neocortical neurons, AZA1, AZA2, and AZA3 inhibited spontaneous Ca^{2+} oscillations at
17
18 moderate concentrations (EC_{50} values ranged from 138 to 445 nM), with 48 h cytotoxicity EC_{50}
19
20 values of 43, 48, and 10 nM, respectively (31). This study provided the first side-by-side
21
22 evidence for potency differences among these regulated toxins.
23
24
25
26

27 Clearly, structure-activity relationship (SAR) studies are very important from a regulatory
28
29 perspective, but to date, have been hindered by the lack of pure material. As such, many of the
30
31 previous SAR studies have been confined to the study of AZA1 and its synthetic fragments (43,
32
33 44); however, purified AZA2 and AZA3 have recently become available commercially (45).
34
35 This supply of new material enabled the present comparative study in which we characterized the
36
37 toxicological effects of the currently regulated AZAs (i.e., AZA1, AZA2, AZA3) in an *in vitro*
38
39 model system based on the Jurkat T lymphocyte cell line.
40
41
42
43
44

45 MATERIALS AND METHODS

46
47
48 **Toxin Isolation.** AZA1, AZA2 and AZA3 were isolated from whole cooked mussel tissue (*M.*
49
50 *edulis*) collected in 2005 from Bruckless, Donegal, Ireland. The toxins were purified using a 7
51
52 step isolation procedure, the details of which have recently been reported (46). Toxin purity
53
54 (>95%) was confirmed by LC-MS/MS and NMR spectroscopy and this material was used to
55
56
57
58
59
60

1
2
3 produce commercially available standards now distributed by the Certified Reference Materials
4 Program at the National Research Council of Canada (Inst. for Marine Biosciences, Halifax, NS,
5 Canada). Toxin dilutions prior to cell exposure were performed in phosphate-buffered saline
6 (PBS; pH 7.4)/10% methanol. Toxin/methanol additions never exceeded 1% vol/vol.
7
8
9
10
11
12
13
14

15 **Cell Culturing.** Human Jurkat E6-1 T lymphocyte cells (American Type Culture Collection
16 TIB-152; Manassas, VA, USA) were grown as described in Twiner et al. (24). Briefly, cells were
17 grown in RPMI medium (cat. #11875-093, Invitrogen, CA, USA) supplemented with 10% (v/v)
18 fetal bovine serum (FBS; cat. #26140, Invitrogen, CA, USA) and maintained in a humidified
19 incubator (Sanyo 18AIC-UV) with 5%:95% CO₂:air at 37 °C. Cells were subcultured with fresh
20 medium at an inoculum ratio of 1:4 every 3 to 4 days by transferring 2.5 mL of cells to 7.5 mL of
21 fresh supplemented medium in 75 cm² screw cap culture flasks.
22
23
24
25
26
27
28
29
30
31
32
33

34 **Cytotoxicity assay.** T lymphocytes were diluted in fresh medium to 3-4 x 10⁵ cells/mL and
35 seeded in 96-well plates containing 100 µL per well (3-4 x 10⁴ cells/well). Cells were allowed to
36 settle for 18-24 h prior to AZA additions to each set of three replicate wells for 24, 48, or 72 h of
37 continuous exposure. The final concentration of AZA1 ranged from 9.48 x 10⁻⁸ to 1 x 10⁻¹² M
38 and the final concentrations of AZA2 and AZA3 ranged from 1.5 x 10⁻⁸ to 5 x 10⁻¹² M. Vehicle
39 control wells with an equivalent amount of 10% methanol in PBS were used to normalize the
40 viability data. Cellular viability/cytotoxicity was assessed using the MTS assay (3-(4,5-
41 dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium; cat. no.
42 G358A, Promega Biosciences, CA, USA). Absorbance of each well was measured at 485 nm
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 using a Fluostar microplate reader (BMG Technologies, NC, USA). All data (mean \pm SE) from 3
4
5 or more separate experiments were normalized to percentage relative to vehicle controls.
6
7

8
9
10 **Morphological characteristics.** T lymphocytes were diluted in fresh medium to $3\text{-}4 \times 10^5$
11
12 cells/mL and seeded in 24-well plates containing 500 μL per well ($1.5\text{-}2 \times 10^5$ cells/well). Cells
13
14 were exposed to AZA concentrations approximately twice the EC_{50} concentration (Table 1 and
15
16 Figure 2). Final concentrations were 5×10^{-9} M AZA1, 5×10^{-10} M AZA2, and 1×10^{-9} M
17
18 AZA3. Vehicle control wells contained an equivalent amount of 10% methanol in PBS. Cells
19
20 were continuously exposed for 24, 48, or 72 h prior to microscopic examination and imaging.
21
22 Differential interference contrast (DIC) and epi-fluorescence photomicrographs were taken using
23
24 an Axioskop 50 microscope (Carl Zeiss, Inc., Thornwood, NY, USA). DIC images were taken
25
26 directly using aliquots of suspended cells. Epi-fluorescence images of intracellular actin were
27
28 obtained by preparing the cells as described in Twiner et al. (2005) and staining with Alexa Fluor
29
30 Phalloidin (cat. #12379, Invitrogen, CA, USA). Images were captured using a Canon EOS 5D
31
32 (EF 24-105 L IS USM) camera.
33
34
35
36
37
38

39
40
41 **Statistical Analysis.** Data are presented as means \pm SE of at least three separate experiments.
42
43 In addition, each cytotoxicity experiment was performed using triplicate wells. Cytotoxicity data
44
45 were blank corrected and normalized to the vehicle control (% viability). EC_{50} and 95%
46
47 confidence interval determinations were calculated using three parameter, variable slope, non-
48
49 linear regression analysis (GraphPad Prism, ver. 5.0c, San Diego, USA).
50
51
52

53 54 55 **RESULTS** 56 57 58 59 60

1
2
3 **Effect of AZA Analogs on Cell Viability.** Jurkat T lymphocyte cells were exposed to natural
4 analogs of AZA (i.e., AZA1, AZA2, AZA3) to assess cellular viability and determine relative
5 potencies. AZAs were tested across a range of concentrations and time points (24, 48, and 72 h).
6 Each AZA analog induced time- and concentration-dependent cytotoxicity to the lymphocyte
7 cells (Figure 2) but differed in their relative potencies. Cytotoxicity was minimal at 24 h with
8 cell viability ranging from 78%-86% for the three AZA analogs. Longer exposure times of 48
9 and 72 h for each AZA analog resulted in a step-wise reduction of viability causing more
10 complete cytotoxicity (20-21% viability at 48 h and 7-12% viability at 72 h) at AZA1
11 concentrations > 1 nM, AZA2 concentrations > 0.1 nM, and AZA3 concentrations > 0.5 nM.
12 EC₅₀ values (including 95% confidence intervals) are presented for each AZA analog at each
13 time point in Table 1. The mean EC₅₀ value for each AZA analog across all three time points was
14 2.5 nM for AZA1, 0.30 nM for AZA2, and 0.55 nM for AZA3. As such, AZA2 and AZA3 were
15 8.3-fold and 4.5-fold more potent than AZA1, respectively (Table 1).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 **Morphological Changes of Lymphocyte Cells Following Exposure to AZA Analogs.**
37 Photomicrographs of control T lymphocyte cells illustrated that they were clearly intact with no
38 signs of lysis and exhibited cytosolic extensions that have been identified as pseudopodia or
39 “false feet” (Figure 3A, B, and C; see arrows). However, lymphocyte cells exposed to the AZA
40 analogs suggest a variety of cytolytic effects. Cells exposed to 5 nM AZA1 displayed a lack of
41 cellular integrity (particularly at 48 and 72 h) as revealed by lysed cells and debris, organelle
42 protrusion with concurrent flattening of cells, and a distinct lack of pseudopodia (Figure 3D, E,
43 and F). Although cell diameter was significantly ($p < 0.05$) increased at 72 h, this appears to be
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 an artifact related to the flattening of the cells during cytolysis and not due to cellular swelling or
4
5
6 blebbing.

7
8 Cells treated with AZA2 and AZA3 also showed time-dependent cytolysis with some features
9
10 (i.e., lack of cellular integrity, organelle protrusion, flattening of cells) in common with AZA1-
11
12 treated cells (Figure 3G, H, I, and Figure 3J, K, L, respectively). However, despite clear
13
14 cytolysis, pseudopodia were still present in each of these treatment regimes (as indicated by
15
16 arrows). Apparent increases in cell diameter were also statistically significant ($p < 0.05$) for
17
18 AZA2 at 72 h and AZA3 at 48 and 72 h (data not shown), but similar to AZA1 these changes are
19
20 attributed to cell flattening and not actual swelling.
21
22
23
24
25
26

27 **Effects of AZA Analogs on Arrangement of F-actin.** Photomicrographs of fluorescently
28
29 stained control cells illustrate their spheroid-like shape with irregular, F-actin filled pseudopodial
30
31 extensions (Figure 4A, B, and C). Upon exposure to AZA1 for 24, 48, or 72 h, cells became
32
33 more rounded with a loss of pseudopodial extensions (Figure 4D, E, and F, respectively). Lysed
34
35 or lysing cells were not apparent, likely due to the loss of exuded material or debris during the
36
37 staining procedure. Cells exposed to AZA2 and AZA3 for 24, 48, or 72 h looked remarkably
38
39 similar to control cells with respect to the continued presence of pseudopodia (Figure 4G, H, I,
40
41 and Figure 4J, K, L, respectively). Similarly, the presence of cells undergoing lysis was not
42
43 expected due to the staining procedure.
44
45
46
47
48
49

50 **Quantitative Effects of AZA Analogs on Pseudopodia.** Jurkat T lymphocyte cells were
51
52 exposed to AZA1 across a range of concentrations for 48 h prior to enumeration of pseudopodial
53
54 extensions. AZA1 caused a concentration-dependent reduction in visible pseudopodia per cell
55
56
57
58
59
60

1
2
3 with an IC_{50} value of 0.68 nM (0.54 - 0.84 nM 95% confidence intervals) (Figure 5). Control
4
5 cells consistently had visible pseudopodia (mean = 2.1 ± 0.18 ; n = 50 cells), whereas cells treated
6
7 with ≥ 10 nM AZA1 had no discernible pseudopodia. When cells were treated for 48 h with
8
9 AZA2 or AZA3 under the same conditions outlined in Figures 3 and 4 (0.5 nM AZA2 and 1 nM
10
11 AZA3), there was an average of 2.2 ± 0.19 and 2.1 ± 0.17 (n = 50) visible pseudopodia per cell,
12
13 respectively (Figure 6).
14
15
16
17
18
19
20
21

22 DISCUSSION

23
24 Human AZA shellfish poisonings have now been reported in multiple European countries and
25
26 in the United States, caused by the import of contaminated Irish shellfish (2, 3). Although more
27
28 than 20 AZA analogs have been identified, only three (AZA1, AZA2, and AZA3) are currently
29
30 thought to be of toxicological concern and as such are regulated in European seafood products at
31
32 a maximal level of 0.16 $\mu\text{g AZA}_{\text{total}}/\text{g}$ shellfish meat (23). The determination of AZA_{total} (i.e.,
33
34 AZA1, AZA2, and AZA3) is primarily performed via LC-MS/MS quantification and the
35
36 application of toxic equivalence factors (TEFs) to account for potency differences between the
37
38 AZA analogs. Based on mouse intraperitoneal injections, TEFs of 1.0, 1.8, and 1.4 are currently
39
40 employed for AZA1, AZA2, and AZA3, respectively (21, 47), thereby implying that the *in vivo*
41
42 order of potency is $AZA2 > AZA3 > AZA1$. Our focus herein was to compare these *in vivo*
43
44 findings with *in vitro* cytotoxicity-based data, as well as assess the morphological effects of
45
46 AZAs in a T lymphocyte model cell line as a means to better understand their potential toxic
47
48 effects.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The cytotoxic potential of AZA1 towards various cell types is well established and has been
4 unambiguously demonstrated using microscopy techniques (24, 27, 34, 39), cellular protein
5 content (27), DNA content and synthesis (27, 48), release of cytosolic enzymes (31, 36), and
6 mitochondrial activity (24, 31). Within the current study, the cytotoxic potential of AZA1, as
7 well as AZA2 and AZA3, was confirmed using human Jurkat T lymphocytes as the model cell
8 line, selected based on previous work demonstrating its high level of sensitivity to AZA1 (28).
9 Each analog induced cytotoxicity in a time- and concentration-dependent manner, exhibiting
10 EC₅₀ concentrations in the sub- to low-nanomolar (0.25 – 2.7 nM) range, which are remarkably
11 lower (i.e., more potent) than the EC₅₀ values for these same AZA analogs tested against murine
12 neocortical neurons (42.7, 48.0, and 9.88 nM for AZA1, AZA2, and AZA3, respectively) (31),
13 but very similar to AZA2 studies using murine lymphoblast cells (EC₅₀ = 0.84 nM) (6).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 Consistent with previous AZA1 studies, increased exposure time to the AZA analogs did affect
30 cell viability but it did not change the calculated EC₅₀ values for time points up to 72 h (24).
31 Based on EC₅₀ values for each analog, AZA2 and AZA3 are 8.3-fold and 4.5-fold more potent
32 than AZA1, suggesting the order of *in vitro* potency as AZA2 > AZA3 > AZA1. This order is
33 consistent with the corresponding *in vivo* results reported by Satake et al. (47) and Ofuji et al.
34 (21) (based on minimal lethal concentrations) that have been implemented by the EU for
35 regulatory purposes (23). However, the TEFs estimated based on these *in vivo* studies (23)
36 suggest less drastic potency differences between these three AZA analogs as compared to the *in*
37 *vitro* data. In contrast, the relative cytotoxic potencies of these same AZA analogs differ when
38 assessed in neocortical neurons (i.e., AZA3 > AZA2 ~ AZA1) and by cytotoxicity (via LDH
39 release) and calcium ion oscillation suppression (31) (i.e., AZA3 > AZA2 > AZA1), further
40 supporting cell type differences in sensitivity. As such, methylation at C8 (AZA2) confers the
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 highest degree of cytotoxic potency in T lymphocytes, whereas de-methylation at C22 (AZA3)
4
5 confers the highest degree of potency in neocortical neurons, suggesting that structural changes
6
7
8 in analogs yield variable toxicities in different cell lines.
9

10 An interesting and rather unusual observation compared to other phycotoxin classes is that the
11 various AZA analogs showing structural differences at the C3, C8, C22, and/or the C23
12
13 positions, elicit different cellular effects and in turn, may be affecting more than one molecular
14
15 target. This was suspected early on by the Botana group when studying secondary messenger
16
17 systems (28, 29, 38, 41) and also reported more recently by the Murray group when they
18
19 observed a discrepancy between the effects of AZA1 and fragments of AZA1 on cytotoxicity and
20
21 spontaneous calcium ion oscillations (31). We have documented previously the total retraction of
22
23 pseudopodia in T lymphocyte cells caused by AZA1 in a time-dependent manner that was
24
25 specific to AZA1 but not elicited by a wide variety of other marine phycotoxins (24). In the
26
27 present study, AZA1 caused similar retraction of pseudopodia in a concentration-dependent
28
29 manner with no observed effect on ruffle formation. On the contrary, exposure of T lymphocyte
30
31 cells to AZA2 and AZA3 had no effect on morphological characteristics or number of
32
33 pseudopodia. The mechanisms involved in pseudopodia assembly are not completely understood
34
35 but they are known to involve nucleation of actin filaments by nucleation factors such as formin
36
37 proteins (49). Despite evidence that AZA1 does not directly alter actin polymerization and/or
38
39 depolymerization (7), actin nucleation factors such as formin FRL2 have been shown to be
40
41 directly involved in the formation and length of pseudopodia in Jurkat T lymphocyte cells (50),
42
43 and may be affected by AZA1. The functional relevance of pseudopodia inhibition by AZA1 has
44
45 not yet been examined but may compromise immune function in organisms exposed to AZAs.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 With the possibility of multiple targets, discerning the structure-activity relationships (SAR) of
4 AZAs becomes very complex. Although varying in toxic potency, AZA1, AZA2, AZA3, AZA4
5 and AZA5 are lethal to mice following *i.p.* injection (21, 22, 47) and AZA1, AZA2, and AZA3
6 have been shown to be highly cytotoxic *in vitro* (6, 24, 25, 31). Prior to the recent availability of
7 other natural AZA analogs, SAR studies have benefited tremendously from the *in silico* organic
8 synthesis of AZA1 (44) and the use of various fragments produced during this process. AZA1
9 fragments containing the ABCD/ABCDE rings yielded increased cytosolic calcium ion
10 concentrations without affecting viability in murine cerebellar granule cells (CGCs) (25). These
11 fragments also did not affect the growth of a lung carcinoma cell line, nor the morphology and
12 cytoskeletal features of a neuroblastoma cell line (27). The same ABCD/ABCDE fragments
13 tested against neocortical neurons did not elicit any effects on calcium ion oscillations or cellular
14 viability (31). SAR studies using the FGHI rings revealed no effects on cytosolic calcium ion
15 concentrations or viability in CGCs (25), yet inhibited calcium ion oscillations and reduced
16 cellular viability in neocortical neurons (31). The relative potencies were particularly high for the
17 FGHI rings when attached to a phenyl glycine methyl ester moiety, a functional component of
18 the natural AZA1, AZA2, and AZA3 analogs. On the contrary, the same set of fragments
19 provided by Nicolaou et al. and applied *in vivo* suggested that the full AZA structure, with its
20 correct stereochemistry, is necessary to induce lethality at levels similar to the natural analog
21 (40).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 Many of these differences in SAR observations may be due to the respective model cell
49 lines/organisms employed, coupled with the possibility of multiple molecular targets among the
50 AZA toxin class. The existence of distinct molecular targets may also help explain the
51
52
53
54
55
56
57
58
59
60

1
2
3 simultaneous appearance of cellular markers for both apoptosis and necrosis across a variety of
4
5 cell lines (7, 31, 38, 39) and following *in vivo* exposure (17, 18).
6
7
8
9

10 **CONCLUSIONS**

11
12 Our data provide evidence that AZA1, AZA2, and AZA3 are highly cytotoxic to T
13
14 lymphocytes and that there are significant differences in their relative potencies that correlate
15
16 well with *in vivo* studies. Furthermore, specific cellular effects such as pseudopodial retraction
17
18 are not conserved across all analogs within this toxin class, but thus far are only observed for
19
20 AZA1. Collectively, our data support the growing body of evidence suggesting that there may be
21
22 several molecular targets for the AZAs and it is likely that these are differentially affected by the
23
24 various AZA analogs.
25
26
27
28
29
30
31

32 **ACKNOWLEDGEMENTS**

33
34 This project has been financially supported in part by a UM-D Office of Research and
35
36 Sponsored Programs Faculty Initiation and Seed Grant and the Irish National Development Plan
37
38 (NDP) under the ASTOX2 project within the seventh European Community Framework
39
40 Programme (FP7/2007-2013) under grant agreement no. 221117. Special thanks go to Chris
41
42 Miles, Philipp Hess, Adela Keogh, Conor Duffy, Ger Clancy, Pearse McCarron, Nils Rehmann,
43
44 Daniel o'Driscoll, and Michael Quilliam.
45
46
47
48
49
50

51 **Disclaimer:** This publication does not constitute an endorsement of any commercial product or
52
53 intend to be an opinion beyond scientific or other results obtained by the National Oceanic and
54
55 Atmospheric Administration (NOAA). No reference shall be made to NOAA, or this publication
56
57
58
59
60

1
2
3 furnished by NOAA, to any advertising or sales promotion which would indicate or imply that
4
5 NOAA recommends or endorses any proprietary product mentioned herein, or which has as its
6
7 purpose an interest to cause the advertised product to be used or purchased because of this
8
9 publication.
10
11

12 13 14 15 16 ABBREVIATIONS

17
18
19 AZA, azaspiracid; CGC, cerebellar granule cells; DIC, differential interference contrast; EC₅₀,
20
21 50% maximal effective concentration; i.p., intraperitoneal; LDH, lactate dehydrogenase; SAR,
22
23 structure-activity relationship; TEF, toxic equivalence factor
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 42 REFERENCES

- 43
44
45 (1) Tillmann, U., Elbrächter, M., Krock, B., John, U., and Cembella, A. (2009) *Azadinium*
46
47 *spinosum* gen. et sp. nov. (Dinophyceae) identified as a primary producer of azaspiracid
48
49 toxins. *European J. Phycol.*, 44, 63 - 79.
50
51
52 (2) Furey, A., O'Doherty, S., O'Callaghan, K., Lehane, M., and James, K. J. (2010)
53
54 Azaspiracid poisoning (AZP) toxins in shellfish: Toxicological and health considerations.
55
56 *Toxicon*, 56, 173-190.
57
58
59
60

- 1
2
3 (3) Klontz, K. C., Abraham, A., Plakas, S. M., and Dickey, R. W. (2009) Mussel-associated
4 azaspiracid intoxication in the United States. *Annals Internal Med.*, 150, 361.
5
6
7
8 (4) Taleb, H., Vale, P., Amanhir, R., Benhadouch, A., Sagou, R., and Chafik, A. (2006) First
9 detection of azaspiracids in mussels in north west Africa. *J. Shellfish Res.*, 25, 1067-
10 1070.
11
12
13
14 (5) Álvarez, G., Uribe, E., Ávalos, P., Mariño, C., and Blanco, J. (2010) First identification
15 of azaspiracid and spirolides in *Mesodesma donacium* and *Mulinia edulis* from Northern
16 Chile. *Toxicon*, 55, 638-641.
17
18
19
20
21
22 (6) Ueoka, R., Ito, A., Izumikawa, M., Maeda, S., Takagi, M., Shin-ya, K., Yoshida, M., van
23 Soest, R. W. M., and Matsunaga, S. (2009) Isolation of azaspiracid-2 from a marine
24 sponge *Echinoclathria* sp. as a potent cytotoxin. *Toxicon*, 53, 680-684.
25
26
27
28
29 (7) Twiner, M. J., Rehmann, N., Hess, P., and Doucette, G. J. (2008) Azaspiracid shellfish
30 poisoning: A review on the chemistry, ecology, and toxicology with an emphasis on
31 human health impacts. *Mar. Drugs*, 6, 39-72.
32
33
34
35
36 (8) Rehmann, N., Hess, P., and Quilliam, M. (2008) Discovery of new analogs of the marine
37 biotoxin azaspiracid in blue mussels (*Mytilus edulis*) by ultra-performance liquid
38 chromatography/tandem mass spectrometry. *Rapid Comm. Mass Spectrom.*
39
40
41
42
43 (9) James, K. J., Sierra, M. D., Lehane, M., Braña Magdalena, A., and Furey, A. (2003)
44 Detection of five new hydroxyl analogues of azaspiracids in shellfish using multiple
45 tandem mass spectrometry. *Toxicon*, 41, 277-283.
46
47
48
49
50 (10) Rundberget, T., Sandvik, M., Hovgaard, P., Nguyen, L., Aasen, J. A. B., Castberg, T.,
51 Gustad, E., and Miles, C. O. (2006) Use of SPATT disks in Norway: detection of AZAs
52
53
54
55
56
57
58
59
60

- 1
2
3 & DTXs and comparison with algal cell counts and toxin profiles in shellfish. , In *Marine*
4 *Biotoxin Science Workshop No. 23* pp 35-38, NZFSA, Wellington, New Zealand.
5
6
7
8 (11) Krock, B., Tillmann, U., John, U., and Cembella, A. D. (2009) Characterization of
9 azaspiracids in plankton size-fractions and isolation of an azaspiracid-producing
10 dinoflagellate from the North Sea. *Harmful Algae*, 8, 254-263.
11
12
13 (12) Tillmann, U., Elbrächter, M., John, U., Krock, B., and Cembella, A. (2010) *Azadinium*
14 *obesum* (Dinophyceae), a new nontoxic species in the genus that can produce azaspiracid
15 toxins. *Phycologia*, 49, 169-182.
16
17
18 (13) James, K. J., Fidalgo Saez, M. J., Furey, A., and Lehane, M. (2004) Azaspiracid
19 poisoning, the food-borne illness associated with shellfish consumption. *Food Addit.*
20 *Contam.*, 21, 879-892.
21
22
23 (14) James, K. J., Furey, A., Satake, M., and Yasumoto, T. (2001) Azaspiracid poisoning
24 (AZP): A new shellfish toxic syndrome in Europe, In *Harmful Algal Blooms 2000*
25 (Hallegraeff, G. M., Blackburn, S. I., Bolch, C. J., and Lewis, R. J., Eds.) pp 250-253,
26 Intergovernmental Oceanographic Commission of UNESCO.
27
28
29 (15) Nzoughet, J. K., Hamilton, J. T. G., Botting, C. H., Douglas, A., Devine, L., Nelson, J.,
30 and Elliott, C. T. (2009) Proteomics identification of azaspiracid toxin biomarkers in blue
31 mussels, *Mytilus edulis*. *Mol. Cell. Proteomics*, 8.8, 1811-1822.
32
33
34 (16) Nzoughet, K. J., Hamilton, J. T. G., Floyd, S. D., Douglas, A., Nelson, J., Devine, L., and
35 Elliott, C. T. (2008) Azaspiracid: First evidence of protein binding in shellfish. *Toxicon*,
36 51, 1255-1263.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (17) Ito, E., Satake, M., Ofuji, K., Higashi, M., Harigaya, K., McMahon, T., and Yasumoto, T.
4
5 (2002) Chronic effects in mice caused by oral administration of sublethal doses of
6
7 azaspiracid, a new marine toxin isolated from mussels. *Toxicon*, *40*, 193-203.
8
9
10 (18) Ito, E., Satake, M., Ofuji, K., Kurita, N., McMahon, T., James, K., and Yasumoto, T.
11
12 (2000) Multiple organ damage caused by a new toxin azaspiracid, isolated from mussels
13
14 produced in Ireland. *Toxicon*, *38*, 917-930.
15
16
17 (19) Ito, E., Terao, K., McMahon, T., Silke, J., and Yasumoto, T. (1998) Acute pathological
18
19 changes in mice caused by crude extracts of novel toxins isolated from Irish mussels, In
20
21 *Harmful Algae* (Reguera, B., Blanco, J., Fernandez, M. L., and Wyatt, T., Eds.) pp 588-
22
23 589, Santiago de Compostela: Xunta de Galicia and Intergovernmental Oceanographic
24
25 Commission of UNESCO.
26
27
28 (20) Satake, M., Ofuji, K., James, K. J., Furey, A., and Yasumoto, T. (1998) New toxic event
29
30 caused by Irish mussels, In *Harmful Algae* (Reguera, B., Blanco, J., Fernandez, M. L.,
31
32 and Wyatt, T., Eds.) pp 468-469, Santiago de Compostela: Xunta de Galicia and
33
34 Intergovernmental Oceanographic Commission of UNESCO.
35
36
37 (21) Ofuji, K., Satake, M., McMahon, T., Silke, J., James, K. J., Naoki, H., Oshima, Y., and
38
39 Yasumoto, T. (1999) Two analogs of azaspiracid isolated from mussels, *Mytilus edulis*,
40
41 involved in human intoxication in Ireland. *Nat. Toxins*, *7*, 99-102.
42
43
44 (22) Ofuji, K., Satake, M., McMahon, T., James, K. J., Naoki, H., Oshima, Y., and Yasumoto,
45
46 T. (2001) Structures of azaspiracid analogs, azaspiracid-4 and azaspiracid-5, causative
47
48 toxins of azaspiracid poisoning in Europe. *Biosci. Biotechnol. Biochem.*, *65*, 740-742.
49
50
51 (23) EFSA. (2008) Marine biotoxins in shellfish – Azaspiracid group: Scientific Opinion of
52
53 the Panel on Contaminants in the Food chain. *The EFSA Journal*, *723*, 1-52.
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- (24) Twiner, M. J., Hess, P., Bottein Dechraoui, M.-Y., McMahon, T., Samons, M. S., Satake, M., Yasumoto, T., Ramsdell, J. S., and Doucette, G. J. (2005) Cytotoxic and cytoskeletal effects of azaspiracid-1 on mammalian cell lines. *Toxicol.*, *45*, 891-900.
- (25) Vale, C., Nicolaou, K. C., Frederick, M. O., Gomez-Limia, B., Alfonso, A., Vieytes, M. R., and Botana, L. M. (2007) Effects of azaspiracid-1, a potent cytotoxic agent, on primary neuronal cultures. A structure-activity relationship study. *J. Med. Chem.*, *50*, 356-363.
- (26) Vilariño, N. (2008) Marine toxins and the cytoskeleton: azaspiracids. *FEBS Journal*, *275*, 6075-6081.
- (27) Vilariño, N., Nicolaou, K. C., Frederick, M. O., Cagide, E., Ares, I. R., Louzao, M. C., Vieytes, M. R., and Botana, L. M. (2006) Cell growth inhibition and actin cytoskeleton disorganization induced by azaspiracid-1 structure-activity studies. *Chem. Res. Toxicol.*, *19*, 1459-1466.
- (28) Alfonso, A., Roman, Y., Vieytes, M. R., Ofuji, K., Satake, M., Yasumoto, T., and Botana, L. M. (2005) Azaspiracid-4 inhibits Ca^{2+} entry by stored operated channels in human T lymphocytes. *Biochem. Pharmacol.*, *69*, 1627-1636.
- (29) Roman, Y., Alfonso, A., Vieytes, M. R., Ofuji, K., Satake, M., Yasumoto, T., and Botana, L. M. (2004) Effects of azaspiracids 2 and 3 on intracellular cAMP, $[Ca^{2+}]$, and pH. *Chem. Res. Toxicol.*, *17*, 1338-1349.
- (30) Kellmann, R., Schaffner, C. A. M., Grønset, T. A., Satake, M., Ziegler, M., and Fladmark, K. E. (2009) Proteomic response of human neuroblastoma cells to azaspiracid-1. *J. Proteomics*, *72*, 695-707.

- 1
2
3 (31) Cao, Z., LePage, K. T., Frederick, M. O., Nicolaou, K. C., and Murray, T. F. (2010)
4
5 Involvement of caspase activation in azaspiracid-induced neurotoxicity in neocortical
6
7 neurons. *Toxicol. Sci.*, *114*, 323-334.
8
9
10 (32) Kulagina, K. V., Twiner, M. J., Hess, P., McMahon, T., Satake, M., Yasumoto, T.,
11
12 Ramsdell, J. S., Doucette, G. J., Ma, W., and O'Shaughnessy, T. J. (2006) Azaspiracid-1
13
14 inhibits bioelectrical activity of spinal cord neuronal networks. *Toxicon*, *47*, 766-773.
15
16
17 (33) Vale, C., Gomez-Limia, B., Nicolaou, K. C., Frederick, M. O., Vieytes, M. R., and
18
19 Botana, L. M. (2007) The c-Jun-N-terminal kinase is involved in the neurotoxic effect of
20
21 azaspiracid-1. *Cell Physiol Biochem*, *20*, 957-966.
22
23
24 (34) Vale, C., Nicolaou, K. C., Frederick, M. O., Vieytes, M. R., and Botana, L. M. (2010)
25
26 Cell volume decrease as a link between azaspiracid-induced cytotoxicity and c-Jun-N-
27
28 terminal kinase activation in cultured neurons. *Toxicol. Sci.*, *113*, 158-168.
29
30
31 (35) Bellocci, M., Sala, G. L., Callegari, F., and Rossini, G. P. (2010) Azaspiracid-1 inhibits
32
33 endocytosis of plasma membrane proteins in epithelial cells. *Toxicol. Sci.*, *117*, 109-121.
34
35
36 (36) Twiner, M. J., Ryan, J. C., Morey, J. S., Smith, K. J., Hammad, S. M., Van Dolah, F. M.,
37
38 Hess, P., McMahon, T., Satake, M., Yasumoto, T., and Doucette, G. J. (2008)
39
40 Transcriptional profiling and inhibition of cholesterol biosynthesis in human lymphocyte
41
42 T cells by the marine toxin azaspiracid. *Genomics*, *91*, 289-300.
43
44
45 (37) Colman, J. R., Twiner, M. J., Hess, P., McMahon, T., Satake, M., Yasumoto, T.,
46
47 Doucette, G. J., and Ramsdell, J. S. (2005) Teratogenic effects of azaspiracid-1 identified
48
49 by microinjection of Japanese medaka (*Oryzias latipes*) embryos. *Toxicon*, *45*, 881-890.
50
51
52 (38) Roman, Y., Alfonso, A., Louzao, M. C., de la Rosa, L. A., Leira, F., Vieites, J. M.,
53
54 Vieytes, M. R., Ofuji, K., Satake, M., Yasumoto, T., and Botana, L. M. (2002)
55
56
57
58
59
60

- 1
2
3 Azaspiracid-1, a potent, nonapoptotic new phycotoxin with several cell targets. *Cell.*
4
5
6 *Signal.*, 14, 703-716.
- 7
8 (39) Vilariño, N., Nicolaou, K. C., Frederick, M. O., Vieytes, M. R., and Botana, L. M. (2007)
9
10 Irreversible cytoskeletal disarrangement is independent of caspase activation during *in*
11
12 *vitro* azaspiracid toxicity in human neuroblastoma cells. *Biochem. Pharm.*, 74, 327-335.
- 13
14 (40) Ito, E., Frederick, M. O., Koftis, T. V., Tang, W., Petrovic, G., Ling, T., and Nicolaou, K.
15
16 C. (2006) Structure toxicity relationships of synthetic azaspiracid-1 and analogs in mice.
17
18 *Harmful Algae*, 5, 586-591.
- 19
20 (41) Alfonso, A., Vieytes, M. R., Ofuji, K., Satake, M., Nicolaou, K. C., Frederick, M. O., and
21
22 Botana, L. M. (2006) Azaspiracids modulate intracellular pH levels in human
23
24 lymphocytes. *Biochem. Biophys. Res. Comm.*, 346, 1091-1099.
- 25
26 (42) Alfonso, C., Alfonso, A., Otero, P., Rodriguez, P., Vieytes, M. R., Elliott, C. T., Higgins,
27
28 C., and Botana, L. M. (2008) Purification of five azaspiracids from mussel samples
29
30 contaminated with DSP toxins and azaspiracids. *J. Chromatogr. B*, 865, 133-140.
- 31
32 (43) Nicolaou, K. C., Koftis, T. V., Vyskocil, S., Petrovic, G., Ling, T., Yamada, T. M. A.,
33
34 Tang, W., and Frederick, M. O. (2004) Structural revision and total synthesis of
35
36 azaspiracid-1, Part 2: Definition of the ABCD domain and total synthesis. *Angew. Chem.*
37
38 *Int. Ed.*, 43, 4318 - 4324.
- 39
40 (44) Nicolaou, K. C., Koftis, T. V., Vyskocil, S., Petrovic, G., Tang, W., Frederick, M. O.,
41
42 Chen, D. Y. K., Yiwei, L., Ling, T., and Yamada, T. M. A. (2004) Total synthesis and
43
44 structural elucidation of azaspiracid-1. Final assignment and total synthesis of the correct
45
46 structure of azaspiracid-1. *J. Am. Chem. Soc.*, 128, 2859-2872.
- 47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (45) Perez, R., Rehmann, N., Crain, S., LeBlanc, P., Craft, C., MacKinnon, S., Reeves, K.,
4
5
6 Burton, I., Walter, J., Hess, P., Quilliam, M., and Melanson, J. (2010) The preparation of
7
8 certified calibration solutions for azaspiracid-1, -2, and -3, potent marine biotoxins found
9
10 in shellfish. *Anal. Bioanal. Chem.*, 398, 2243-2252.
11
12 (46) Kilcoyne, J., Keogh, A., Clancy, G., Le Blanc, P., Burton, I., Quilliam, M., Hess, P., and
13
14 Miles, C. O. (in press) Improved isolation procedure for azaspiracids from shellfish,
15
16 structural elucidation of azaspiracid-6 and stability studies. *J. Agric. Food Chem.*
17
18
19 (47) Satake, M., Ofuji, K., Naoki, H., James, K. J., Furey, A., McMahon, T., Silke, J., and
20
21 Yasumoto, T. (1998) Azaspiracid, a new marine toxin having unique spiro ring assemblies,
22
23 isolated from Irish mussels, *Mytilus edulis*. *J. Amer. Chem. Soc.*, 120, 9967-9968.
24
25
26 (48) Ronzitti, G., Hess, P., Rehmann, N., and Rossini, G. P. (2007) Azaspiracid-1 alters the E-
27
28 cadherin pool in epithelial cells. *Toxicol. Sci.*, 95, 427-435.
29
30
31 (49) Harris, E. S., Rouiller, I., Hanein, D., and Higgs, H. N. (2006) Mechanistic differences in
32
33 actin bundling activity of two mammalian formins, FRL1 and mDia2. *J. Biol. Chem.*,
34
35 281, 14383-14392.
36
37
38 (50) Harris, E. S., Gauvin, T. J., Heimsath, E. G., and Higgs, H. N. (2010) Assembly of
39
40 filopodia by the formin FRL2 (FMNL3). *Cytoskeleton*, 67, 755-772.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables and Figures Legends

Table 1. Cytotoxicity EC₅₀ values (nM) and 95% confidence intervals for AZA1, AZA2, and AZA3.

AZA analog	24 h		48 h		72 h		Mean EC ₅₀	Rel. Pot.
	EC ₅₀	95% CI	EC ₅₀	95% CI	EC ₅₀	95% CI		
AZA-1	2.1	0.56 - 8.3	2.7	1.2 - 6.2	2.6	1.4 - 4.9	2.5	1.0
AZA-2	0.38	0.10 - 1.4	0.28	0.15 - 0.52	0.25	0.14 - 0.43	0.30	8.3
AZA-3	0.44	0.15 - 1.3	0.58	0.29 - 1.2	0.63	0.32 - 1.2	0.55	4.5

Figure Legends

1
2
3 Figure 1. Chemical structures of AZA1, AZA2, and AZA3.
4
5
6
7
8

9
10 Figure 2. Effect of various AZA analogs on cell viability. Lymphocyte T cells were exposed to
11 AZA1, AZA2, or AZA3 for 24, 48, or 72 h and viability was assessed using the MTS assay. All
12 data (mean \pm SE; $n \geq 3$) were normalized to the control (methanol vehicle) and analyzed using
13 non-linear sigmoidal dose-response (variable slope). Calculated EC₅₀ values are shown in Table
14
15
16
17
18
19 1.
20
21
22
23
24


25
26 Figure 3. Effect of various AZA analogs on cell morphology. Jurkat T lymphocyte cells were
27 exposed to AZA analogs at *ca.* twice EC₅₀ concentrations for 24, 48, and 72 h (from left to right)
28 prior to photographs being taken. Panels A, B, and C illustrate control cells exposed to
29 equivalent volumes of methanol, panels D, E, and F illustrate cells exposed to 5 nM AZA1,
30 panels G, H, and I illustrate cells exposed to 0.5 nM AZA2, and panels J, K, and L illustrate cells
31 exposed to 1 nM AZA3. Arrows indicate the location of pseudopodia. Note: Each panel is a
32 composite of various individual DIC images/cells and not a single field of view, thus illustrating
33 a representative range of cell morphologies.
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48
49 Figure 4. Fluorescence staining and visualization of F-actin in Jurkat T lymphocyte cells
50 following exposure to various AZA analogs. Jurkat T lymphocyte cells were exposed to AZA
51 analogs at *ca.* twice EC₅₀ concentrations for 24, 48, and 72 h (from left to right) prior to
52 fluorescent staining and visualizing cellular F-actin using Alexa Fluor 488 phalloidin. Panels A,
53
54
55
56
57
58
59
60

1
2
3 B, and C illustrate control cells exposed to equivalent volumes of methanol, panels D, E, and F
4
5 illustrate cells exposed to 5 nM AZA1, panels G, H, and I illustrate cells exposed to 0.5 nM
6
7 AZA2, and panels J, K, and L illustrate cells exposed to 1 nM AZA3. Arrows indicate the
8
9 location of pseudopodia. Note: Each panel is a composite of various individual epi-fluorescence
10
11 images/cells and not a single field of view, thus illustrating a representative range of cell
12
13 morphologies.
14
15
16
17
18
19
20




21
22 Figure 5. Effect of AZA1 on the abundance of pseudopodia. Lymphocyte T cells were exposed
23
24 to various concentrations of AZA1 for 48 h and the number of pseudopodia per cell were
25
26 counted using DIC microscopy. All data (mean \pm SE; n = 50 individual cells) were analyzed
27
28 using non-linear sigmoidal dose-response (variable slope).
29
30
31
32
33
34

35
36 Figure 6. Effects of various AZA analogs on the abundance of pseudopodia. Lymphocyte T cells
37
38 were exposed to AZA1 (5 nM), AZA2 (0.5 nM), or AZA3 (1 nM) for 48 h and the number of
39
40 pseudopodia per cell were counted using DIC microscopy. Data are illustrated as mean \pm SE (n =
41
42 50 individual cells) and an asterisk (*) indicates a significant difference from vehicle control (p <
43
44 0.05).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60







	R_1	R_2	R_3	R_4
AZA1	H	H	CH ₃	H
AZA2	H	CH ₃	CH ₃	H
AZA3	H	H	H	H

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60








116x224mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60





110x76mm (300 x 300 DPI)



115x77mm (300 x 300 DPI)