

Painful faces-induced attentional blink modulated by top-down and bottom-up mechanisms

Chun Zheng^{1,2,3}, Jin-Yan Wang^{1*} and Fei Luo¹

¹ Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, China, ² University of Chinese Academy of Sciences, Beijing, China, ³ Department of Psychology, Southwest University for Nationalities, Chengdu, China

OPEN ACCESS

Edited by:

Narayanan Srinivasan,
University of Allahabad, India

Reviewed by:

Adrian Von Muhlenen,
University of Warwick, UK
Sonia Baloni Ray,
Centre of Behavioural and Cognitive
Sciences, India
Priyanka Srivastava,
International Institute of Information
Technology, Hyderabad, India

*Correspondence:

Jin-Yan Wang,
Key Laboratory of Mental Health,
Institute of Psychology, Chinese
Academy of Sciences,
16 Lincui Road, Chaoyang District,
Beijing 100101, China
wangjy@psych.ac.cn

Specialty section:

This article was submitted to
Cognitive Science,
a section of the journal
Frontiers in Psychology

Received: 17 December 2014

Accepted: 11 May 2015

Published: 01 June 2015

Citation:

Zheng C, Wang J-Y and Luo F (2015)
Painful faces-induced attentional blink
modulated by top-down
and bottom-up mechanisms.
Front. Psychol. 6:695.
doi: 10.3389/fpsyg.2015.00695

Pain-related stimuli can capture attention in an automatic (bottom-up) or intentional (top-down) fashion. Previous studies have examined attentional capture by pain-related information using spatial attention paradigms that involve mainly a bottom-up mechanism. In the current study, we investigated the pain information-induced attentional blink (AB) using a rapid serial visual presentation (RSVP) task, and compared the effects of task-irrelevant and task-relevant pain distractors. Relationships between accuracy of target identification and individual traits (i.e., empathy and catastrophizing thinking about pain) were also examined. The results demonstrated that task-relevant painful faces had a significant pain information-induced AB effect, whereas task-irrelevant faces showed a near-significant trend of this effect, supporting the notion that pain-related stimuli can influence the temporal dynamics of attention. Furthermore, we found a significant negative correlation between response accuracy and pain catastrophizing score in task-relevant trials. These findings suggest that active scanning of environmental information related to pain produces greater deficits in cognition than does unintentional attention toward pain, which may represent the different ways in which healthy individuals and patients with chronic pain process pain-relevant information. These results may provide insight into the understanding of maladaptive attentional processing in patients with chronic pain.

Keywords: attentional blink, bottom-up, pain, RSVP, top-down

Introduction

As an alarm signal of bodily threat, pain has an inherently stronger ability than other stimuli to draw attention and interrupt ongoing goals (Eccleston and Crombez, 1999; Legrain et al., 2009; Van Damme et al., 2010; Schoth et al., 2012). Similarly, pain-related information also has the advantage of capturing attention preferentially, consuming attentional resources, and making attention switching difficult. Using the dot-probe task (Roelofs et al., 2005; Liossi et al., 2009; Haggman et al., 2010; Baum et al., 2013) and spatial cueing paradigm (Van Damme et al., 2004a,b, 2006b), researchers have found significant attentional bias toward painful words/faces and visual cues predictive of pain compared with neutral words/faces and non-pain cues, respectively.

Abbreviations: AB, attentional blink; ANOVA, analysis of variance; IRI, Interpersonal Reactivity Index; PCS, Pain Catastrophizing Scale; RSVP, rapid serial visual presentation; SOA, stimulus onset asynchrony.

These studies have examined attentional bias toward pain-related stimuli based on the spatial distribution of attention. Few studies have addressed these issues in terms of the temporal dynamics of attention. Moreover, available evidence supporting pain-related attentional bias comes largely from patients with chronic pain; this bias is rarely found in healthy subjects (Crombez et al., 2013).

The AB paradigm is used widely to study the temporal deployment of attention (Martens and Wyble, 2010). In a typical AB paradigm, two targets are embedded in a stream of items in a RSVP. The AB effect is characterized by a failure to detect the second target, which follows the first target after a very short interval (Raymond et al., 1992; Maciokas and Crognale, 2003). This paradigm has been used to explore the effect of emotional information on attention capture (de Oca et al., 2012; Schonenberg and Abdelrahman, 2013; Yerys et al., 2013; de Jong et al., 2014). Recent studies using a single-target RSVP task have found that task-irrelevant emotional words (Barnard et al., 2005; Arnell et al., 2007) or pictures (Kennedy and Most, 2012; Quaedflieg et al., 2013) are capable of inducing an AB. Most et al. (2005) defined the difference in task performance when viewing emotional and neutral pictures as emotion-induced AB (Smith et al., 2006).

Pain-related information can capture attention in a bottom-up or top-down fashion (Crombez et al., 2005; Van Damme et al., 2010). Most previous studies investigating pain-related attention capture have used task-irrelevant stimuli (i.e., stimulus-driven or bottom-up mode; Van Damme et al., 2010; Schoth et al., 2012; Sharpe, 2012; Crombez et al., 2013). No study other than that conducted by Grynberg et al. (2014) has employed pain-related material as a task-relevant stimulus (i.e., goal-driven or top-down mode). Distinct patterns of attentional processing of pain-related stimuli have been thought to represent different ways in which healthy individuals and patients with chronic pain process pain-relevant information, with the former using mainly a stimulus-driven, bottom-up pattern and the latter using mainly a top-down process (Van Damme et al., 2010; Crombez et al., 2013). Selective attention to and difficulty in disengaging from pain-related information lead to cognitive impairment, as well as the maintenance or exacerbation of pain, in patients with chronic pain (Vlaeyen and Linton, 2000; Leeuw et al., 2007). No study to date has compared the effects of bottom-up and top-down attention engagement using pain-related information. Thus, the present study aimed to examine the effects of these two attention modes using task-irrelevant and task-relevant AB paradigms, to provide a possible basis for the understanding of maladaptive attentional processing in patients with chronic pain.

This study involved the administration of a single-task RSVP, in which only a target image was identified, and a dual-task RSVP, in which subjects were asked to respond to a distractor and a target, to healthy subjects. Photographs depicting painful and neutral facial expressions were used as distractors. Subjects were asked to complete the self-reported PCS and IRI. Pain catastrophizing is believed to play an important role in shaping the experiences of acute and chronic pain, and has been shown to be associated with attentional bias for pain-relevant stimuli (Van Damme et al., 2002, 2004a). The IRI measures trait empathy;

human elements such as faces and voices have been suggested to provide important cues triggering empathy (Jabbi et al., 2007; Choi and Watanuki, 2014). The following two hypotheses were tested: (1) pain information-induced AB effects (i.e., worse target performance in the painful condition than in the neutral condition) would be observed for both tasks, and would be more significant in the top-down (dual task) model than in the bottom-up (single task) model; and (2) pain catastrophizing and/or empathy trait would be related to the target accuracy in pain condition.

Materials and Methods

Participants

Fifty-two healthy, right-handed undergraduate and graduate students, with normal or corrected-to-normal vision, participated in this study (mean age 22.0 ± 2.6 years, 35 females). All participants had no history of psychiatric or neurological disorders, and none of them reported current pain or a past history of chronic pain. The Institutional Review Board of the Institute of Psychology, Chinese Academy of Sciences, approved the experimental procedures. Informed consent was obtained from each subject before the experiment. All individuals were paid for participation.


Experimental Procedure

Experiments were performed in a sound-attenuated room with the temperature maintained at 26°C , relative humidity of 48–53%, and luminance level of 361–374 lux. Participants were seated in a comfortable chair at a viewing distance of 60 ± 5 cm from a computer monitor. An RSVP paradigm was delivered to measure the attention-capturing effect of pain-related information. Participants were randomly assigned to a single-task RSVP group ($n = 26$, 18 females), in which only one target was identified, or a dual-task RSVP group ($n = 26$, 17 females), in which participants were instructed to respond to a distractor and a target. Each subject was allowed to practice before the formal experiment. After terminating the task, participants were asked to complete the IRI and PCS. **Figure 1A** illustrates the experimental procedure.

Visual Stimuli

The stimuli were presented in the center of a gray background on a 17-inch cathode ray tube monitor with a refresh rate of 100 Hz using E-prime 1.1 software (Psychology Software Tools, Inc., Pittsburgh, PA, USA). The screen resolution was set to 800×600 pixels, resulting in a photograph display size of about $13.8 \text{ cm} \times 10.3 \text{ cm}$.

Visual stimuli were 320×240 -pixel color photographs gathered from the Internet (www.gettyimages.cn). The stimuli consisted of 180 photographs of adult male and female faces (distractors; 60 painful, 60 neutral, and 60 scrambled), 180 photographs of tree/architectural elements (targets; 90 rotated 90° clockwise and 90 rotated 90° counterclockwise), and 240 upright tree/architectural images (fillers). The painful distractors depicted men and women with painful expressions and hands



on the forehead, indicating headache. The painful and neutral faces were matched by gender. The scrambled distractors were created by segmenting each painful image using an 8×6 grid and randomly reordering the segments using Adobe Photoshop CS4 (Adobe Systems, Inc., San Jose, CA, USA). The scrambled pictures served as controls to ensure that the behavioral differences elicited by painful and neutral conditions were not due to low-level visual properties, such as color and luminance. **Figure 1B** depicts an example of an RSVP image stream. Each target and distractor image appeared only once during the task.

Prior to the experiment, subjects not involved in the main study rated the degrees of pain and arousal depicted in facial photographs on a seven-point scale ranging from 0 (“no feeling” and “calm” respectively) to 6 (“intolerably painful” and “excited” respectively). An initial sample of 240 face photographs was divided into two sets, each containing 120 photographs and rated

by an independent group of subjects ($n = 20$, 10 females) to prevent fatigue. The ages of subjects in the two groups were matched (22.1 ± 3.3 and 22.6 ± 1.4 years; $t_{(38)} = 0.57$, $p = 0.57$). A total of 120 face photographs with the highest and lowest pain scores ($n = 60$ each) were used as painful and neutral distractors, respectively, in the present study. Subjective ratings of these painful and neutral distractors were shown to differ significantly before the formal experiment was conducted (pain score: 3.67 ± 0.53 vs. 0.47 ± 0.12 , $t_{(118)} = 45.52$, $p < 0.001$; arousal score: 3.20 ± 0.32 vs. 2.47 ± 0.36 , $t_{(118)} = 11.68$, $p < 0.001$).

Rapid Serial Visual Presentation Paradigm

The experiment comprised 180 RSVP trials. Participants started trials by pressing the space bar on the computer keyboard. Each trial began with the presentation of a fixation cross (random

duration, 100–300 ms) in the center of the monitor. The RSVP trial consisted of a stream of 17 images, including a distractor image, a target image, and 15 filler images, each presented for 100 ms. Distractors were positioned as the fourth, sixth, or eighth image in each trial, similar to the method used by Most et al. (2005). Distractor position was randomized across trials. The target appeared two (lag 2, SOA = 200 ms) or eight (lag 8, SOA = 800 ms) places after the distractor. **Figure 1B** illustrates the sequence of events for one RSVP trial. Participants were required to provide responses after presentation of the stream, at their own pace. One second after responding, participants could press the space bar again to start the next trial.

The single-task RSVP was preceded by a brief practice session (18 trials) that included no distractor. Targets were similar to those used in the experiment, and filler images were selected from the set used in the experiment. Each item in the RSVP stream was presented for 200 ms to ensure that participants could identify the target and learn the rules of the experiment. In the dual-task RSVP, participants followed the same procedure as in the single-task experiment, but they were given different instructions (see Instructions). The initial 18-trial practice session was similar to that preceding the single-task RSVP, but it contained distractor images (similar to those used in the experiment). No feedback was provided to participants.

Instructions

For the single-task RSVP, participants were given the following instructions: “You will see a series of rapidly presented photographs of landscapes and faces. You should pay attention to a rotated photo and indicate its orientation by pressing the corresponding (left or right) arrow key. Note that only one landscape photo is rotated.”

Instructions for the dual-task RSVP were: “You will see a series of rapidly presented photographs of landscapes and faces. You should pay attention to both a face photo and a rotated landscape photo, and identify the type of face by pressing the corresponding key on the keyboard (“1” = painful, “2” = neutral, “0” = scrambled). Then, indicate the orientation of rotated photo by pressing the corresponding (left or right) arrow key. Note that only one landscape photo is rotated.”

Interpersonal Reactivity Index

The IRI was developed by Davis (1980). It comprises 28 items rated on a five-point scale, and measures trait empathy using four subscales. The perspective-taking subscale assesses the tendency to spontaneously adopt others’ psychological points of view; the fantasy subscale measures respondents’ tendency to transpose themselves imaginatively into the feelings and actions of fictitious characters in books, movies, and plays; the empathic concern subscale assesses “other-oriented” feelings of sympathy and concern for unfortunate others; and the personal distress subscale measures “self-oriented” feelings of personal anxiety and unease in tense interpersonal settings. The IRI total score is the sum of all subscale scores. The IRI subscales have been shown to have good internal consistency (ranging between 0.71 and 0.77; Davis, 1980, 1983). The Chinese version

of the IRI, used in the present study, has been demonstrated to have satisfactory construct reliability (0.59, 0.75, 0.60, and 0.69 for perspective-taking, fantasy, empathic concern, and personal distress subscales, respectively) and retest reliability (0.64, 0.75, 0.59, and 0.78 for perspective-taking, fantasy, empathic concern, and personal distress subscales, respectively; Rong et al., 2010).

Pain Catastrophizing Scale

The PCS was designed to measure catastrophic thinking about pain (Sullivan et al., 1995). It contains 13 items in three subscales: rumination (e.g., “I can’t stop thinking about how much it hurts”), magnification (e.g., “I worry that something serious may happen”), and helplessness (e.g., “It’s awful and I feel that it overwhelms me”). Each item is rated on a five-point scale. The PCS total score is the sum of all subscale scores. The PCS has been shown to have good internal consistency (Cronbach’s $\alpha = 0.87$ for the total PCS, 0.87 for rumination, 0.60 for magnification, 0.79 for helplessness; Sullivan et al., 1995). The Chinese version of the PCS (HK-PCS), used in the present study, has been demonstrated to have satisfactory internal consistency (Cronbach’s alpha = 0.927, 0.809, 0.768, and 0.839 for the total scale and rumination, magnification, and helplessness subscales, respectively; Yap et al., 2008).

Data Analyses

For both tasks, the accuracy of target direction judgment was recorded for analysis. For the dual-task group, only trials in which distractors were identified correctly were included in the analysis. The pain information-induced AB was defined as the difference in target accuracy between the neutral and painful conditions at lag 2.

All statistical analyses were performed using SPSS 13.0 (SPSS, Inc., Chicago, IL, USA). The accuracy of face identification in the dual task was assessed using 3×2 (distractor type \times lag) repeated-measures ANOVA. Task performance (i.e., target accuracy) data were entered into a $2 \times 3 \times 2$ [task (single, dual) \times distractor type (painful, neutral, scrambled) \times lag (lag 2, lag 8)] mixed-design ANOVA, with repeated measures on the last two factors. Split-up ANOVAs were performed in cases of three-way and two-way interaction. Separate 3×2 (distractor type \times task) ANOVAs were conducted for lag 2 and lag 8 data. When the ANOVA of lag 2 data showed significant interaction, one-way ANOVA was conducted for distractor type. *Post hoc* comparisons were performed using Fisher’s least significant difference method. Effect sizes for ANOVAs were reported using partial eta squared (η^2) values. Pearson’s correlations were computed to explore relationships between target accuracy and IRI and PCS scores. The significance level was set at $p < 0.05$.

Results

Performance of Face Identification in the Dual Task

For the dual task, the accuracy of identification of painful, neutral, and scrambled faces was 93 ± 7 , 95 ± 6 , and $95 \pm 4\%$,

respectively, at lag 2 and 95 ± 8 , 96 ± 6 , and $97 \pm 4\%$, respectively at lag 8. Two-way ANOVA revealed no significant main effect (lag: $F_{(1,25)} = 1.57$, $p = 0.22$, $\eta^2 = 0.06$; distractor type: $F_{(2,50)} = 1.26$, $p = 0.29$, $\eta^2 = 0.05$) or interaction (distractor type \times lag: $F_{(2,50)} = 0.17$, $p = 0.79$, $\eta^2 = 0.01$).

Pain Information-Induced Attentional Blink Effect

Three-way ANOVA revealed significant interaction effects of task \times lag ($F_{(1,50)} = 82.29$, $p < 0.001$, $\eta^2 = 0.62$), lag \times distractor type ($F_{(2,100)} = 13.32$, $p < 0.001$, $\eta^2 = 0.21$),

and task \times lag \times distractor type ($F_{(2,100)} = 3.79$, $p < 0.05$, $\eta^2 = 0.07$). Significant main effects of task ($F_{(1,50)} = 47.29$, $p < 0.001$, $\eta^2 = 0.49$), lag ($F_{(1,50)} = 94.67$, $p < 0.001$, $\eta^2 = 0.65$), and distractor type ($F_{(2,100)} = 11.58$, $p < 0.001$, $\eta^2 = 0.19$) were observed. **Figure 2A** shows the effect of distractor type on target identification accuracy according to target position (i.e., lag) for both tasks. Overall, mean accuracy was significantly lower at lag 2 than at lag 8 (78 ± 14 vs. $89 \pm 5\%$, $p < 0.001$), indicating the presence of an AB effect (**Figure 2B**).

Two-way ANOVA (task \times distractor type) of lag 2 data revealed significant task \times distractor type interaction




FIGURE 2 | Response accuracy in the single and dual tasks. (A) Temporal distribution of target identification accuracy. **(B)** Overall effect of lag on response accuracy. **(C)** Effect of task relevance on response accuracy at lag 2. **(D)** Effects of distractor type on response accuracy at lag 2 in the single and dual tasks. Error bars indicate standard errors. * $p < 0.05$, *** $p < 0.001$.

($F_{(2,100)} = 3.93, p < 0.05, \eta^2 = 0.07$), and main effects of task ($F_{(1,50)} = 74.97, p < 0.001, \eta^2 = 0.60$) and distractor type ($F_{(2,100)} = 16.21, p < 0.001, \eta^2 = 0.25$). The mean response accuracy was significantly higher in the single task than in the dual task, suggesting that the effect of interruption on target identification was more prominent for task-relevant than for task-irrelevant distractors (Figure 2C). *Post hoc* comparisons revealed significant differences between the single and dual tasks for all three distractor types (painful: 86 ± 10 vs. $61 \pm 12\%$, $p < 0.001$; neutral: 90 ± 9 vs. $66 \pm 14\%$, $p < 0.001$; scrambled: 91 ± 7 vs. $75 \pm 14\%$, $p < 0.001$).

For both tasks, one-way ANOVA revealed significant main effects of distractor type at lag 2 (single task: $F_{(2,50)} = 3.61, p < 0.05, \eta^2 = 0.13$; dual task: $F_{(2,50)} = 13.76, p < 0.001, \eta^2 = 0.36$). *Post hoc* comparisons revealed a significant difference in target accuracy between painful and neutral conditions in the dual task (61 ± 12 vs. $66 \pm 14\%$, $p < 0.05$), and a marginally significant difference in the single task (86 ± 10 vs. $90 \pm 9\%$, $p = 0.068$; Figure 2D). Two-way ANOVA (task \times distractor type) of lag 8 data showed a main effect of distractor type ($F_{(2,100)} = 3.97, p < 0.05, \eta^2 = 0.07$), but no interaction and no effect of task.

Correlations between Accuracy in Painful Distractor Trials and Questionnaire Scores

Results of correlation analysis between response accuracy and individual variables of self-reported measures are presented in Table 1. In general, significant correlations were found only in painful trials, not in neutral trials. For the single task, target accuracy and IRI perspective-taking subscale score were positively correlated ($r = 0.45, p < 0.05$). The positive correlation between target accuracy and IRI total score ($r = 0.47, p < 0.05$) was due mainly to the perspective-taking subscale score. By contrast, for the dual task, significant negative correlations were observed between response accuracy and PCS total ($r = -0.46, p < 0.05$) and rumination subscale ($r = -0.43, p < 0.05$) scores. No other significant correlation was identified. Scatter plots of significant correlations between accuracy and questionnaire scores are presented in Figure 3.

Discussion

The current study demonstrated that task-relevant painful faces had a significantly stronger AB effect (i.e., temporary visual processing impairment for targets appearing soon after faces) than did neutral faces, and task-irrelevant faces showed a near-significant trend of this effect. The attention-capturing effect was greater in the dual task than in the single task, consistent with our hypothesis. Importantly, we found that response accuracy for targets following painful faces was correlated positively with IRI perspective-taking subscale score in the single-task condition, and negatively correlated with PCS score in the dual-task condition.


Previous research has demonstrated that pain-related stimuli attract attention. Most evidence has come from studies of attentional bias, which have used paradigms such as the modified Stroop task (Muris et al., 1995; Roelofs et al., 2002), the dot-probe task (Keogh et al., 2001; Vervoort et al., 2011), and the spatial cueing task (Van Damme et al., 2004a, 2006a). In these studies, pain-related stimuli were always task-irrelevant cues, rather than task-relevant targets. Participants were not required to pay attention to the pain-related stimuli, and attention was captured by painful materials in a bottom-up fashion. In the present study, we used single-target and dual-target tasks to investigate the existence of a pain information-induced AB, and compared the bottom-up and top-down modulation of attention using task-irrelevant and task-relevant paradigms. Our results showed a significant pain information-induced AB effect (i.e., worse target performance in the painful condition than in the neutral condition) in the task-relevant condition, consistent with our hypothesis and with the findings of previous studies that emotional expressions or words can capture attention and interfere with an ongoing task (Mathewson et al., 2008; Srivastava and Srinivasan, 2010). As painful (negative emotional) expressions are more salient than neutral expressions, they obviously capture and hold attention more effectively, thereby causing more disruption of subsequent target identification.

Unexpectedly, we failed to find a significant pain information-induced AB effect in the task-irrelevant condition, although the

TABLE 1 | Mean scores of the questionnaires (Interpersonal Reactivity Index and Pain Catastrophizing Scale) and correlation coefficient values between the questionnaire and target accuracy (at lag 2).

	Single task			Dual task		
	Mean (SD)	Painful	Neutral	Mean (SD)	Painful	Neutral
IRI total (0–112)	67.04 (10.57)	0.47*	–0.01	69.92 (12.26)	0.16	0.22
<i>Perspective-taking</i>	18.12 (4.23)	0.45*	0.12	18.38 (4.32)	0.13	–0.18
<i>Fantasy</i>	16.27 (4.68)	0.29	0.10	16.50 (4.93)	0.20	0.22
<i>Empathic concern</i>	17.77 (3.71)	0.27	0.05	19.85 (3.94)	0.07	0.18
<i>Personal distress</i>	14.88 (4.68)	0.15	–0.28	15.19 (5.23)	0.02	0.32
PCS total (0–52)	16.77 (10.37)	0.10	0.01	21.85 (8.56)	–0.46*	–0.29
<i>Rumination</i>	8.65 (4.37)	0.18	0.10	8.77 (3.72)	–0.43*	–0.29
<i>Magnification</i>	3.27 (2.62)	0.02	–0.16	4.92 (2.61)	–0.25	–0.00
<i>Helplessness</i>	4.85 (4.58)	0.04	0.03	8.15 (4.72)	–0.36	–0.29

* $p < 0.05$; IRI, Interpersonal Reactivity Index; PCS, Pain Catastrophizing Scale.



results showed a near-significant trend of this effect. These results are not in accord with those of Most et al. (2005), who employed a single-task RSVP and found that negative images produced greater impairment of target processing than did neutral images. This inconsistency may be due to differences between studies in the intensity of negative emotional images. Most et al. (2005) used negative photographs of people and animals and included graphic images of violence, distress, and medical trauma. These bloody, violent scenes may produce a stronger emotional effect than did those depicting merely painful facial expressions in our study. Additionally, the high level of target accuracy (86%) at lag 2 in the painful condition indicates that the painful faces used in our study were less threatening than the images used by Most et al. (2005; who reported 71% accuracy). Furthermore, previous research has failed to demonstrate attentional bias toward pain-related words or images in healthy volunteers (Crombez et al., 2013). Most evidence supporting pain-related attentional bias has come from patients with chronic pain (Snider et al., 2000) or studies using conditioned visual signals of pain as cues (Crombez et al., 2013). Thus, the occurrence of attentional bias may depend on prior pain experience. In the present study, we used healthy subjects and found a marginally significant pain information-induced AB effect in the task-irrelevant painful condition,

suggesting that the temporal attention paradigm is more sensitive than spatial attention paradigms for the measurement of pain-related attention-capturing effects.

As expected, we found that task-relevant painful faces interfered more with target identification than did task-irrelevant faces, suggesting that painful faces that were processed in a top-down manner engaged more attentional resources. The different effects of bottom-up and top-down attention to painful faces may be explained by the distinct temporal characteristics of these two systems. In the bottom-up pattern, attention is captured transiently by a pain distractor; in the top-down pattern, a painful face holds attention for a longer time and leaves limited resources available for target identification (Nakayama and Mackeben, 1989; Dux and Marois, 2009; Martens and Wyble, 2010; Pinto et al., 2013). Given that attention capturing was guided in the task-relevant condition compared with the task-irrelevant condition, target impairment in the dual task can be better explained by the holding, rather than capturing, of attention (Mathewson et al., 2008; Srivastava and Srinivasan, 2010). Additionally, when a subject is prepared to identify a target or has knowledge about the target while scanning an environment, relevant stimuli capture his/her attention more rapidly and accurately than if he/she was not prepared. This case may

also reflect interplay between top-down and bottom-up processing. In real life, pain-related information is often not a currently pursued goal, but a signal of physical threat, for healthy individuals. It automatically attracts attention in a bottom-up fashion (Eccleston and Crombez, 1999). By contrast, physical symptoms have a persistent influence on the lives of patients with chronic pain, who must direct their attention toward physical pain or potential pain indicators in an attempt to control pain in a timely and efficient manner (Eccleston and Crombez, 1999, 2007; Van Damme et al., 2008). In this case, pain-related information is selectively attended to, and top-down processing may result in hypervigilance, i.e., increased arousal to environmental change.

In the present study, target performance was modulated by different individual traits in the two tasks. In the single-target task, response accuracy was positively correlated with the IRI perspective-taking subscale; in the dual-target task, a negative correlation was found between response accuracy and catastrophic thinking about pain (particularly the PCS rumination subscale). The positive correlation between IRI perspective-taking subscale score and target performance is difficult to interpret, and may not be very meaningful. The painful faces used in this study did not likely evoke empathic responses. First, the photographs had a low degree of emotional strength. Second, the duration of stimulus presentation (100 ms) was much shorter than that typically used in empathy studies (≥ 200 ms; Fan and Han, 2008; Sessa et al., 2014). Finally, in the task-irrelevant condition, subjects were asked to ignore the distractor and the visual stimulus was processed only modestly. A possible explanation for this is that the perspective-taking subscale, which measures only the cognitive dimension of empathy, probably reflects general cognitive ability. That is, individuals with better ability to identify emotional faces had better task performance.

The negative correlation between target accuracy and catastrophizing thinking about pain was not surprising. In the dual task in which pain was task relevant, participants were required to focus on pain distractors, which may result in a greater degree of attention holding than in the task-irrelevant condition. Moreover, subjects who engage in more pain catastrophizing may have more difficulty disengaging their attention from pain-related stimuli. Our results fit well with the findings of Van Damme et al. (2002, 2004a) who demonstrated that catastrophizing thinking about pain predicted the retardation of disengagement from pain-related information

using a cueing task (Van Damme et al., 2002) and a spatial cueing paradigm (Van Damme et al., 2004a). Previous studies have demonstrated that personality traits can modulate the AB by showing that greater extraversion and openness predicted smaller AB while greater neuroticism predicted larger AB (Maclean and Arnell, 2010). Our results also found the relationship between personality characteristics (i.e., catastrophizing thoughts) and AB in pain-face-as-target trials rather than neutral-face-as-target trials, suggesting that such a relation was specific to pain. Highly catastrophizing individuals may be inclined to extensive processing of pain, which may impede the disengagement of attentional resources from pain-related thoughts and feelings, thereby impairing the processing of other environmental stimuli (Vervoort et al., 2011).

Conclusion

This study examined the attention-capturing effects of painful facial expressions using task-relevant and task-irrelevant AB paradigms. In the task-relevant condition, painful faces had a significant pain information-induced AB effect; task-irrelevant faces showed a near-significant trend of this effect. This difference was mediated by the distinct patterns of attentional processing (i.e., top-down vs. bottom-up), which may represent the different ways in which healthy individuals and patients with chronic pain process pain-relevant information. Moreover, in the dual-task condition, a higher level of pain catastrophizing predicted poorer task performance, suggesting that catastrophizing thoughts play a critical role in cognitive impairment in patients with chronic pain. These results extend previous findings by showing different effects of top-down and bottom-up processing of pain-related stimuli on cognitive function, and provide insight into maladaptive cognition in patients with chronic pain.

Acknowledgments

This work was funded by an NSFC Grant (31271092), a Youth Innovation Promotion Association CAS grant to J-YW, and NSFC grants (61033011, 31171067, and 31471061) and a grant from the Key Laboratory of Mental Health, Institute of Psychology – Chinese Academy of Sciences (KLMH2014ZK01) to FL, and KLMH2014G01 to Feng Yu.

References

- Arnell, K. M., Killman, K. V., and Fijavz, D. (2007). Blinded by emotion: target misses follow attention capture by arousing distractors in RSVP. *Emotion* 7, 465–477. doi: 10.1037/1528-3542.7.3.465
- Barnard, P. J., Ramponi, C., Battye, G., and Mackintosh, B. (2005). Anxiety and the deployment of visual attention over time. *Vis. Cogn.* 12, 181–211. doi: 10.1080/13506280444000139
- Baum, C., Schneider, R., Keogh, E., and Lautenbacher, S. (2013). Different stages in attentional processing of facial expressions of pain: a dot-probe task modification. *J. Pain* 14, 223–232. doi: 10.1016/j.jpain.2012.11.001
- Choi, D., and Watanuki, S. (2014). Effect of empathy trait on attention to faces: an event-related potential (ERP) study. *J. Physiol. Anthropol.* 33:4. doi: 10.1186/1880-6805-33-4
- Crombez, G., Van Damme, S., and Eccleston, C. (2005). Hypervigilance to pain: an experimental and clinical analysis. *Pain* 116, 4–7. doi: 10.1016/j.pain.2005.03.035
- Crombez, G., Van Ryckeghem, D. M. L., Eccleston, C., and Van Damme, S. (2013). Attentional bias to pain-related information: a meta-analysis. *Pain* 154, 497–510. doi: 10.1016/j.pain.2012.11.013
- Davis, M. H. (1980). A multidimensional approach to individual differences in empathy. *JSAS Catal. Select. Doc. Psychol.* 10, 85.

- Davis, M. H. (1983). Measuring individual differences in empathy: evidence for a multidimensional approach. *J. Pers. Soc. Psychol.* 44, 113–126. doi: 10.1037/0022-3514.44.1.113
- de Jong, P. J., Koster, E. H. W., Wessel, I., and Martens, S. (2014). Distinct temporal processing of task-irrelevant emotional facial expressions. *Emotion* 14, 12–16. doi: 10.1037/A0034630
- de Oca, B. M., Villa, M., Cervantes, M., and Welbourne, T. (2012). Emotional modulation of the attentional blink by pleasant and unpleasant pictures. *J. Gen. Psychol.* 139, 289–314. doi: 10.1080/00221309.2012.708681
- Dux, P. E., and Marois, R. (2009). The attentional blink: a review of data and theory. *Atten. Percept. Psychophys.* 71, 1683–1700. doi: 10.3758/app.71.8.1683
- Eccleston, C., and Crombez, G. (1999). Pain demands attention: a cognitive-affective model of the interruptive function of pain. *Psychol. Bull.* 125, 356–366. doi: 10.1037//0033-2909.125.3.356
- Eccleston, C., and Crombez, G. (2007). Worry and chronic pain: a misdirected problem solving model. *Pain* 132, 233–236. doi: 10.1016/j.pain.2007.09.014
- Fan, Y., and Han, S. (2008). Temporal dynamic of neural mechanisms involved in empathy for pain: an event-related brain potential study. *Neuropsychologia* 46, 160–173. doi: 10.1016/j.neuropsychologia.2007.07.023
- Grynberg, D., Vermeulen, N., and Luminet, O. (2014). Amplification of attentional blink by distress-related facial expressions: relationships with alexithymia and affectivity. *Int. J. Psychol.* 49, 371–380. doi: 10.1002/ijop.12006
- Haggman, S. P., Sharpe, L. A., Nicholas, M. K., and Refshauge, K. M. (2010). Attentional biases toward sensory pain words in acute and chronic pain patients. *J. Pain* 11, 1136–1145. doi: 10.1016/j.jpain.2010.02.017
- Jabbi, M., Swart, M., and Keysers, C. (2007). Empathy for positive and negative emotions in the gustatory cortex. *Neuroimage* 34, 1744–1753. doi: 10.1016/j.neuroimage.2006.10.032
- Kennedy, B. L., and Most, S. B. (2012). Perceptual, not memorial, disruption underlies emotion-induced blindness. *Emotion* 12, 199–202. doi: 10.1037/a0026380
- Keogh, E., Ellery, D., Hunt, C., and Hannent, I. (2001). Selective attentional bias for pain-related stimuli amongst pain fearful individuals. *Pain* 91, 91–100. doi: 10.1016/S0304-3959(00)00422-X
- Leeuw, M., Goossens, M., Linton, S. J., Crombez, G., Boersma, K., and Vlaeyen, J. W. S. (2007). The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J. Behav. Med.* 30, 77–94. doi: 10.1007/s10865-006-9085-0
- Legrain, V., Damme, S. V., Eccleston, C., Davis, K. D., Seminowicz, D. A., and Crombez, G. (2009). A neurocognitive model of attention to pain: behavioral and neuroimaging evidence. *Pain* 144, 230–232. doi: 10.1016/j.pain.2009.03.020
- Lioffi, C., Schoth, D. E., Bradley, B. P., and Mogg, K. (2009). Time-course of attentional bias for pain-related cues in chronic daily headache sufferers. *Eur. J. Pain* 13, 963–969. doi: 10.1016/j.ejpain.2008.11.007
- Maciokas, J. B., and Crognale, M. A. (2003). Cognitive and attentional changes with age: evidence from attentional blink deficits. *Exp. Aging Res.* 29, 137–153. doi: 10.1080/03610730390176095
- Maclean, M. H., and Arnell, K. M. (2010). Personality predicts temporal attention costs in the attentional blink paradigm. *Psychon. Bull. Rev.* 17, 556–562. doi: 10.3758/Pbr.17.4.556
- Martens, S., and Wyble, B. (2010). The attentional blink: past, present, and future of a blind spot in perceptual awareness. *Neurosci. Biobehav. Rev.* 34, 947–957. doi: 10.1016/j.neubiorev.2009.12.005
- Mathewson, K. J., Arnell, K. M., and Mansfield, C. A. (2008). Capturing and holding attention: the impact of emotional words in rapid serial visual presentation. *Mem. Cogn.* 36, 182–200. doi: 10.3758/Mc.36.1.182
- Most, S. B., Chun, M. M., Widders, D. M., and Zald, D. H. (2005). Attentional rubbernecking: cognitive control and personality in emotion-induced blindness. *Psychon. Bull. Rev.* 12, 654–661. doi: 10.3758/BF03196754
- Muris, P., Merckelbach, H., and De Jongh, A. (1995). Colour-naming of dentist-related words: role of coping style, dental anxiety, and trait anxiety. *Pers. Individ. Diff.* 18, 685–688. doi: 10.1016/0191-8869(94)00185-U
- Nakayama, K., and Mackeben, M. (1989). Sustained and transient components of focal visual-attention. *Vision Res.* 29, 1631–1647. doi: 10.1016/0042-6989(89)90144-2
- Pinto, Y., Van Der Leij, A. R., Slight, I. G., Lamme, V. A. F., and Scholte, H. S. (2013). Bottom-up and top-down attention are independent. *J. Vis.* 13:16. doi: 10.1167/13.3.16
- Quaedflieg, C. W., Giesbrecht, T., Meijer, E., Merckelbach, H., De Jong, P. J., Thorsteinsson, H., et al. (2013). Early emotional processing deficits in depersonalization: an exploration with event-related potentials in an undergraduate sample. *Psychiatry Res.* 212, 223–229. doi: 10.1016/j.psychres.2012.08.001
- Raymond, J. E., Shapiro, K. L., and Arnell, K. M. (1992). Temporary suppression of visual processing in an RSVP task: an attentional blink? *J. Exp. Psychol. Hum. Percept. Perform.* 18, 849–860. doi: 10.1037/0096-1523.18.3.849
- Roelofs, J., Peters, M. L., Fassaert, T., and Vlaeyen, J. W. S. (2005). The role of fear of movement and injury in selective attentional processing in patients with chronic low back pain: a dot-probe evaluation. *J. Pain* 6, 294–300. doi: 10.1016/j.jpain.2004.12.011
- Roelofs, J., Peters, M. L., and Vlaeyen, J. W. S. (2002). Selective attention for pain-related information in healthy individuals: the role of pain and fear. *Eur. J. Pain* 6, 331–339. doi: 10.1016/S1090-3801(02)00021-6
- Rong, X., Sun, B. H., Huang, X. Z., Cai, M. Y., and Li, W. J. (2010). Reliabilities and validities of Chinese version of Interpersonal Reactivity Index. *Chin. J. Clin. Psychol.* 18, 158–160.
- Schonenberg, M., and Abdelrahman, T. (2013). In the face of danger: exploring the attentional blink to emotional facial expressions in PTSD. *Psychiatry Res.* 209, 180–185. doi: 10.1016/j.psychres.2012.11.011
- Schoth, D. E., Nunes, V. D., and Lioffi, C. (2012). Attentional bias towards pain-related information in chronic pain: a meta-analysis of visual-probe investigations. *Clin. Psychol. Rev.* 32, 13–25. doi: 10.1016/j.cpr.2011.09.004
- Sessa, P., Meconi, F., Castelli, L., and Dell'acqua, R. (2014). Taking one's time in feeling other-race pain: an event-related potential investigation on the time-course of cross-racial empathy. *Soc. Cogn. Affect. Neurosci.* 9, 454–463. doi: 10.1093/scan/nst003
- Sharpe, L. (2012). Attentional biases and their role in the management of pain. *Pain* 153, 2307–2308. doi: 10.1016/j.pain.2012.09.004
- Smith, S. D., Most, S. B., Newsome, L. A., and Zald, D. H. (2006). An emotion-induced attentional blink elicited by aversively conditioned stimuli. *Emotion* 6, 523–527. doi: 10.1037/1528-3542.6.3.523
- Snider, B. S., Asmundson, G. J. G., and Wiese, K. C. (2000). Automatic and strategic processing of threat cues in patients with chronic pain: a modified Stroop evaluation. *Clin. J. Pain* 16, 144–154. doi: 10.1097/00002508-200006000-00008
- Srivastava, P., and Srinivasan, N. (2010). Time course of visual attention with emotional faces. *Atten. Percept. Psychophys.* 72, 369–377. doi: 10.3758/App.72.2.369
- Sullivan, M. J., Bishop, S. R., and Pivik, J. (1995). The pain catastrophizing scale: development and validation. *Psychol. Assess.* 7, 524–532. doi: 10.1037/1040-3590.7.4.524
- Van Damme, S., Crombez, G., and Eccleston, C. (2002). Retarded disengagement from pain cues: the effects of pain catastrophizing and pain expectancy. *Pain* 100, 111–118. doi: 10.1016/S0304-3959(02)00290-7
- Van Damme, S., Crombez, G., and Eccleston, C. (2004a). The anticipation of pain modulates spatial attention: evidence for pain-specificity in high-pain catastrophizers. *Pain* 111, 392–399. doi: 10.1016/j.pain.2004.07.022
- Van Damme, S., Lorenz, J., Eccleston, C., Koster, E. H. W., De Clercq, A., and Crombez, G. (2004b). Fear-conditioned cues of impending pain facilitate attentional engagement. *Clin. Neurophysiol.* 34, 33–39. doi: 10.1016/j.neucli.2003.11.001
- Van Damme, S., Crombez, G., and Eccleston, C. (2008). Coping with pain: a motivational perspective. *Pain* 139, 1–4. doi: 10.1016/j.pain.2008.07.022
- Van Damme, S., Crombez, G., Eccleston, C., and Koster, E. H. W. (2006a). Hypervigilance to learned pain signals: a componential analysis. *J. Pain* 7, 346–357. doi: 10.1016/j.jpain.2005.12.006
- Van Damme, S., Crombez, G., Hermans, D., Koster, E. H. W., and Eccleston, C. (2006b). The role of extinction and reinstatement in attentional bias to threat: a conditioning approach. *Behav. Res. Ther.* 44, 1555–1563. doi: 10.1016/j.brat.2005.11.008
- Van Damme, S., Legrain, V., Vogt, J., and Crombez, G. (2010). Keeping pain in mind: a motivational account of attention to pain. *Neurosci. Biobehav. Rev.* 34, 204–213. doi: 10.1016/j.neubiorev.2009.01.005
- Vervoort, T., Caes, L., Crombez, G., Koster, E., Van Damme, S., Dewitte, M., et al. (2011). Parental catastrophizing about children's pain and selective attention to varying levels of facial expression of pain in children: a dot-probe study. *Pain* 152, 1751–1757. doi: 10.1016/j.pain.2011.03.015

- Vlaeyen, J. W., and Linton, S. J. (2000). Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 85, 317–332. doi: 10.1016/S0304-3959(99)00242-0
- Yap, J. C., Lau, J., Chen, P. P., Gin, T., Wong, T., Chan, I., et al. (2008). Validation of the Chinese Pain Catastrophizing Scale (HK-PCS) in patients with chronic pain. *Pain. Med.* 9, 186–195. doi: 10.1111/j.1526-4637.2007.00307.x
- Yerys, B. E., Ruiz, E., Strang, J., Sokoloff, J., Kenworthy, L., and Vaidya, C. J. (2013). Modulation of attentional blink with emotional faces in typical development and in autism spectrum disorders. *J. Child Psychol. Psychiatry* 54, 636–643. doi: 10.1111/jcpp.12013

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Zheng, Wang and Luo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.