

# Spontaneous haemorrhage in a healthy newborn

Eulália Viveiros,<sup>1</sup> Marcos Sanches,<sup>2</sup> Alexandra Couto,<sup>3</sup> Helena Ramos<sup>3</sup>

<sup>1</sup>Pediatric Department, Hospital do Divino Espírito Santo de Ponta Delgada, EPE, Ponta Delgada, Azores, Portugal

<sup>2</sup>Pediatric Department, Hospital Distrital de Santarém, EPE, Santarém, Portugal

<sup>3</sup>Department of Neonatal Intensive Care, Maternidade Dr Alfredo da Costa, Lisboa, Portugal

## Correspondence to

Dr Eulália Viveiros,  
eulaliaviveiros@gmail.com

Accepted 3 May 2014

## DESCRIPTION

A full-term male neonate, who was born by vaginal delivery without trauma, presented a small purple tumefaction on his back at 8 h of life, with a tender consistency. It was well defined and interpreted as a haemangioma. At 24 h of life he became lethargic, pale, tachycardic and the tumefaction worsened (figures 1 and 2).<sup>1 2</sup> A careful review of his family history revealed haemophilia A in first-degree to third-degree relatives on the maternal side.<sup>1 2</sup> Haemoglobin dropped from 14.6 to 9.5 g/dL in 4 h. His activated thromboplastin time was 85.2 s with normal prothrombin time and number of platelets. The coagulation factor dosage revealed a severe factor VIII deficiency (FVIII <0.1%). He received red blood cells, fresh frozen plasma and intravenous recombinant factor VIII transfusions, and started phototherapy for jaundice.<sup>2</sup> Cranial, abdominal ultrasounds and thorax radiographies were normal. In the following days, his general condition and haematoma significantly improved.

He was discharged at 7 days of life with FVIII dosage in normal range (85.2%) and negative inhibitors for FVIII.

## Learning points

- ▶ Haemophilia A is an X-linked coagulation disorder, so mothers are usually asymptomatic carriers and manifestations are almost exclusively in male neonates.
- ▶ A review of the family history in an otherwise healthy newborn, with unexplained bleeding, can lead to the diagnosis in two-thirds of the population.
- ▶ The severe form of haemophilia should be part of the differential diagnosis of haemodynamically unstable newborns with spontaneous superficial haematoma.



**Figure 1** Diffuse haematoma on the neonate's back and the presence of some central yellowish papule surrounded by a halo of erythema (erythema toxicum).



CrossMark

**To cite:** Viveiros E, Sanches M, Couto A, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-202253

**Figure 2** Extensive haematoma on the back of the neonate.



**Acknowledgements** The authors would like to thank all in the neonatal intensive care and haematology department.

**Competing interests** None.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

#### REFERENCES

- 1 Chalmers EA. Neonatal coagulation problems. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F475–8.
- 2 Dyson H. Neonatal haemophilia—a guide for recognition and management. *Haemophilia* 2006;2:156–9.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow