



# Clinical phenotypes of Cerebral Amyloid Angiopathy

Luís F. Maia<sup>a</sup>, Ian R.A. Mackenzie<sup>b</sup>, Howard H. Feldman<sup>c,d,\*</sup>

<sup>a</sup> Department of Neurology, Hospital Geral Santo António, Porto, Portugal

<sup>b</sup> Division of Neuropathology, Department of Pathology, University of British Columbia, Vancouver, British Columbia, Canada

<sup>c</sup> Division of Neurology, Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

<sup>d</sup> Clinic for Alzheimer's Disease and Related Disorders, University of British Columbia Hospital and Vancouver Coastal Health, Vancouver, British Columbia, Canada

Available online 6 March 2007

## Abstract

The term Cerebral Amyloid Angiopathy (CAA) is used to describe the pathological changes occurring in cerebral blood vessels, both leptomeningeal and cortical that result from the deposition of amyloid proteins. This CNS vasculopathy is associated with a spectrum of clinical phenotypes that include both ischemic and hemorrhagic presentations. Dementia, cognitive impairment and transient neurological symptoms or signs are also being increasingly recognized as part of the CAA clinical spectrum.

This review covers the clinical, pathological and neuroimaging aspects of CAA.

© 2007 Elsevier B.V. All rights reserved.

**Keywords:** Cerebral Amyloid Angiopathy; Clinical presentations; AD; Dementia; MRI; Stroke

## 1. Introduction

Cerebral Amyloid Angiopathy is defined by the deposition of amyloid proteins within leptomeningeal and cortical arteries, arterioles, capillaries, and rarely veins. The term amyloid refers to any amorphous, eosinophilic, extracellular protein deposit with specific staining characteristics, such as Congo red binding and birefringence with polarized light (Fig. 1) [1,2]. There are a number of different proteins that can deposit as amyloid in human CAA (Table 1).

In sporadic and most of the hereditary forms of CAA the amyloid includes  $\beta$ -amyloid peptides 1–40 and 1–42 which aggregate within vessel walls, occasionally spreading and depositing in the surrounding neuropil or within the glia limitans when capillaries are involved (Fig. 2) [1–3]. The composition of the vascular amyloid deposits differs from senile plaques (SP) of Alzheimer's Disease (AD) by having an

increased A $\beta$  40/42 ratio, which seems to shift the amyloid deposition from the parenchyma to the vascular wall [4,5].

CAA occurs within a heterogeneous group of hereditary and sporadic diseases. Hereditary forms account for the minority of the CAA cases but have more pleomorphic phenotypes (Table 1). Sporadic CAA increases dramatically with age, rarely being present before the age of 50, and increasing in prevalence to more than 50% in people over age 90 [3,6]. Data from autopsy series have confirmed age as being a significant risk factor in a variety of different populations [7–9]. Though CAA is frequently presumed to be asymptomatic, it is now increasingly recognized to present with a spectrum of clinical presentations.

## 2. Historical overview

The histological vascular abnormalities recognizable as vascular amyloid deposition were first described by Gustav Oppenheim, in the brain parenchyma adjacent to hyalinized capillary walls in 6 of 14 autopsied brains of individuals with dementia and the pathology of AD [10]. In 1938, Scholz published the first article in which the primary focus was the

\* Corresponding author. Division of Neurology, University of British Columbia Hospital, S192-2211 Wesbrook Mall Vancouver, BC, Canada V6T 2B5. Tel.: +1 604 822 7979; fax: +1 604 822 7703.

E-mail address: [hfeldman@interchange.ubc.ca](mailto:hfeldman@interchange.ubc.ca) (H.H. Feldman).

vascular abnormality now recognizable as CAA. He detected this abnormality in the brains of 15 of 104 unselected autopsies, and he suggested it to be a disease related to aging [10].

Although it has been nearly a century since the first pathological descriptions of CAA, knowledge of its link with intracerebral hemorrhage was recognized only in 1960, when Neumann [11] reported on a 45-year old woman with severe Cerebral Amyloid Angiopathy who had 2 symptomatic lobar hemorrhages and multiple small asymptomatic lobar petechial hemorrhages (PH). After this description further interest followed when it was appreciated that there was CAA underlying a significant proportion of cerebral hemorrhages occurring in non-hypertensive individuals [11,12]. Following these seminal descriptions, there has been an emerging clinical and neuroimaging interest in both sporadic and hereditary cases in the correlations with this pathology.

### 3. The clinical spectrum of sporadic CAA

The different clinical presentations that are recognized to be associated with CAA may be divided syndromically into stroke; transient neurological events including TIAs, seizures, and migrainous phenomena; cognitive impairment and dementia [1,3].

#### 3.1. Stroke

##### 3.1.1. Lobar cerebral hemorrhage

The most recognized clinical presentation of CAA and the basis for the clinical criteria set by the Boston group [13,14], is lobar cerebral hemorrhage which is frequently recurrent [2,3,15]. Indeed, the antemortem diagnosis of CAA is most often established around this presentation [13,15,16]. CAA-related lobar hemorrhages usually affect normotensive individuals over age 55. They account for 5–20% of all spontaneous (non-traumatic) cerebral hemorrhages in elderly subjects [17,18]. The CAA lobar hematoma typically involves the cortical–subcortical regions and can extend from the cortex to the subarachnoid space or less commonly, to the ventricles. Recurrent and multiple hemorrhages are a feature of CAA-related lobar hemorrhage in patients who survive the initial bleed [1,3]. The localization of CAA-related hemorrhage follows the localization of CAA in the cerebral cortex and cortico–subcortical or lobar regions [1–3,15]. Despite the high prevalence of CAA in the occipital cortex, CAA-related hemorrhages have been shown to be more evenly distributed [1,3], with a slight predominance of occipital and frontal cortices [15]. Patients present with cortical dysfunction that correlates to the extent and location of lesions, typically preceded or coincident with a headache, vomiting and nuchal rigidity [19]. CAA is typically absent in regions characteristic of hypertensive hemorrhages (deep brain regions and cerebellum) (Fig. 3). The presence of petechial hemorrhages in a cortical/subcortical location detected by low signal on gradient-recalled-echo (GRE) MRI sequences with T2\*-weighting

further supports the diagnosis of CAA (Fig. 2) [13]. This low signal is caused by deposited iron products (Fig. 4) [20]. These small cortical–subcortical petechial hemorrhages correlate well with the pathological diagnosis of CAA [13,20]. From the diagnostic standpoint gradient echo MRI has emerged as the most important diagnostic aid for CAA identification during life. The lack of other accurate and accessible biomarkers has otherwise limited knowledge of early detection, characterization of the different clinical presentations of CAA and the potential impact on the patients' neurological status.

Fibrinoid necrosis and aneurysm formation are both implicated in the in the pathogenesis of vessel rupture [1–3]. Possession of the apoE  $\epsilon$ 2 and 4 alleles has been associated with both fibrinoid necrosis and concentric splitting (“double-barrel” appearance) of the blood-vessel walls (Fig. 5) [21]. It is, therefore, not unexpected that patients with CAA-related hemorrhages and the 2/4 genotype (i.e. with both “risk” alleles) have early and recurrent lobar hemorrhages [21–23]. Nevertheless, at present apoE genotyping does not have a recommended place in the clinical setting for this problem.

There is an emerging question regarding the risks of hemorrhagic complications of thrombolytic agents (e.g. tissue plasminogen activator — rtPA) in patients with CAA. Thrombolysis-related intracerebral hemorrhage in patients with myocardial infarction has similar risk factors and the hematoma features as sporadic CAA related hemorrhages [24]. On the other hand, in a small series of patients




Fig. 1. CAA in sporadic case of Alzheimer's Disease. (A) Congo-red positive material in vessel wall, (B) confirmed as amyloid by showing apple-green birefringence when viewed with polarized light.

Table 1  
Genetic, clinical, radiological and pathological data from hereditary forms of CAA

Hereditary CAA	Gene/ mutation	Amyloid Protein	Age of onset	Clinical Phenotype	Neuroradiological phenotype	Pathological phenotype
Hereditary cerebral hemorrhage with amyloidosis-Dutch type [30,54–58]	APP E693Q mutation	A $\beta$	39–76	Lobar brain hemorrhages; dementia in survivors; epilepsy in survivors.	Lobar hemorrhages; PHs in the gray-white matter junction in T2*MRI; infarcts; diffuse white matter damage.	CAA affects cerebral and cerebellar meningeal arteries and cerebro-cortical arterioles. CAA is more severe occipitally; Diffuse plaques; Hemorrhages and infarcts.
Hereditary CAA - Iowa type [31,59]	APP D694N mutation	A $\beta$	60–70	Progressive aphasia/dementia; Lobar brain hemorrhage (in Spanish kindred).	Occipital gyriform calcification and l eukoencephalopathy. Cortical PHs in T2*MRI.	Extensive CAA; Calcified deposits in vessel walls; Diffuse plaques; NFTs; Cortical infarcts and hemorrhages.
Hereditary CAA - Flemish type [60–62]	APP A692G mutation	A $\beta$	41–61	Amnesic syndrome/dementia; Lobar brain hemorrhage.	Cortical central atrophy; white matter lesions; Lobar brain hemorrhages.	Extensive CAA large amyloid core SP cerebral and cerebellar and NFTs.
Hereditary CAA - Italian kindred [63,64]	APP E693K mutation	A $\beta$	50s	Lobar brain hemorrhages.	NA	Extensive CAA.
Hereditary cerebral hemorrhage with amyloidosis-Icelandic type [2,65]	CystC Leu68Gln	Acys	20–50	Lobar brain hemorrhages; cognitive impairment or dementia in survivors.	NA	Extensive CAA within small arteries and arterioles of leptomeninges, cerebral cortex, basal ganglia, brainstem, and cerebellum.
Familial amyloid polyneuropathy/meningo-vascular amyloidoses [66–71]	TTR Asp18Gly or Val30Gly or Tyr69His	ATTR	36–53	Dementia, ataxia, and spasticity. Dementia, seizures, ataxia, hemiparesis, and decreased vision. Subarachnoid hemorrhage.	Symmetric calcification of the cortex; MRI contrast enhancement at the surface of the cortex. Superficial siderosis.	TTR-amyloid deposits predominated in cerebrospinal leptomeningeal vessels and in subpial and subependymal regions. Brain parenchyma is not involved.
Familial amyloidoses Finnish type [72]	GEL G654A or G654T	AGel	30	Dementia, ataxia, neuropsychiatric features; corneal lattice dystrophy, cranial and peripheral neuropathy	White matter lesions in postmortem MRI.	Minor cortical CAA, white matter AGel angiopathy and myelin loss; major spinal angiopathy.
Prion disease with cerebral amyloid angiopathy [73]	PRNP Y145Stop	APrP	38	Dementia, extrapyramidal signs, ataxia and spastic paraparesis.	Late cerebral atrophy and ventricular dilatation.	Extensive CAA in leptomeningeal and parenchymal vessels, NFTs in cortex. Severe gliosis and neuronal loss.
Familial British dementia [2,53]	BRI2 Mutation in stop codon	ABRi	40–60	Dementia; spasticity; ataxia; rare Lobar brain hemorrhages.	Frontal and occipital white matter lesions; lesion in the corpus callosum. No evidence of hemorrhages.	Extensive CAA affecting vessels in the leptomeninges and both gray and white matter; abundant amyloid plaques and severe neurofibrillary degeneration.
Familial Danish dementia [74,75]	BRI2 decamer duplication in codon 265–266	ADan	30	Cataracts; deafness; dementia; spasticity; ataxia; rare strokes.	White matter lesions.	Extensive CAA in cerebral vessels, choroid plexus, cerebellum, spinal cord, and retina; plaques and neurofibrillary tangles; ischemic lesions in white matter. Uniform diffuse atrophy .
Familial Alzheimer's Disease and Down Syndrome [76–81]	PS1 PS2; APP	A $\beta$	Variable	Dementia; Lobar brain hemorrhages and spastic paraparesis	Variable	Variable CAA.

the presence of petechial hemorrhages on gradient echo imaging was reported to not substantially increase the risk of either symptomatic or asymptomatic brain hemorrhage

following IV rtPA administered between 3 and 6 h after stroke onset [25]. More prospective data is required in order to clarify the risk of hemorrhagic complications with the use




Fig. 2. Sporadic case of Alzheimer's Disease. A-beta Immunohistochemistry demonstrating A-beta protein deposition in the walls of leptomeningeal and parenchymal vessels (CAA) and in senile plaques.

of thrombolysis in patients with CAA. There are currently no recommendations to guide antithrombotic or fibrinolytic treatment based on the detection of cerebral petechial hemorrhages on MRI [24,26].

### 3.1.2. Ischemic infarctions

Manifestations of cerebral ischemia should be considered within the clinical spectrum of CAA. CAA related ischemic infarcts are more often located in the cerebral cortex, with lesions that have been described as 'cortical lacunes'. These cortically based lesions may present as TIAs or minor strokes in elderly CAA patients [12,27]. CAA with ischemic infarctions are observed in patients, with both hereditary and, less frequently, sporadic forms of CAA [12,28–31].

In tissue biopsies of patients with a history of recent cerebral or cerebellar infarctions, CAA was found in 13% of cases, compared to 3.7% of controls, OR 3.8 (95% confidence-interval 1.3–10.9) [32].




Fig. 3. CT scan of a biopsy proven CAA patient that presented with a stroke syndrome. Acute left frontal lobe hemorrhage. Peri-ventricular white matter hyperdensity.




Fig. 4. Axial T2\*GRE MR with T2\*-weighting image of sporadic CAA case. Multiple petechial cortical hemorrhages (black arrowheads) and a major lobar brain hemorrhage (white arrowhead).

The pathogenic mechanisms underlying ischemic lesions related to CAA remains uncertain. Whether it is through vascular stenosis or obstruction, or whether it results from a dynamic change in the vessel physiology or both, is presently debated [33,34].

### 3.1.3. Subarachnoid hemorrhage (SAH)

Over age 60, CAA is a rare cause of primary SAH yet it is the most frequent cause of ICH accompanying secondary SAH [35]. The SAH source may be intraparenchymal with hemorrhage extending into the subarachnoid space or from leptomeningeal vessels leaking into the subarachnoid space. Further studies are still needed to clarify this issue [36].

There have been descriptions of localized SAH or superficial siderosis over cortical *sulci* in patients that presented with transient neurological symptoms, in patients




Fig. 5. Double barrel appearance of CAA of an amyloid-laden vessel in a case of sporadic CAA.




Fig. 6. CT Scan of a probable CAA patient that presented with left sensitive seizures and post-ictal left hemiparesis, showing a right perirolandic subarachnoid hemorrhage.

demonstrated to have had CAA (Figs. 6 and 7) [37,38]. The extensive involvement of the meningo-cortical vessels by the amyloid deposits and its rupture underlies the SAH and related clinical manifestations that can be recognized in CAA.

### 3.2. Transient focal neurological symptoms and signs (TFNSS)

TFNSS can include different phenomena such as focal seizures with or without Todd's paralysis, focal deficits (TIAs) or positive visual symptoms similar to migrainous auras. Most of the published cases describe brief stereotypic phenomena that sometimes preceded major hemorrhagic events. A favorable response to antiepileptic drugs supports an epileptic mechanism for some TFNSS [38].

There have been no systematic studies elucidating these phenomena in patients with CAA with only case reports in the literature [27,37,38]. The transitory nature of the clinical symptoms and signs, coupled with the absence of a clear CAA biomarker make clinical–pathological correlation difficult to establish. Underlying the TFNSS different pathologies have been described including: petechial cortical hemorrhages [38], subpial–subarachnoid hemorrhages [37] or ischemic lesions (“cortical lacunes”) (Figs. 4, 6 and 7) [27].

### 3.3. Cognitive impairment and dementia

The relationship of CAA with cognitive impairment and dementia has been well recognized. There is an inverse correlation between CAA presence and cognitive function that has been reported in both case series [39] as well as in the population based studies. In the MRC study, severe CAA was identified in 36.5% of individuals with dementia compared to 7% without dementia, OR 7.7 (95% CI, 3.3 to 20.4) [7]. CAA is present in >80% of AD case-series [8,17]. In the Honolulu-Asia Aging Study (HAAS), the comorbid presence of CAA with AD was associated with significantly

worse cognitive test performance [40]. CAA can cause both mild cognitive impairment and full blown dementia, in sporadic and hereditary cases [41].

CAA impairs cognitive function through a number of different mechanisms that include: 1. — Ischemic or hemorrhagic lesions which disrupt neuronal circuits or cause strategic lesions [42]; 2. — Interference with cerebrovascular auto-regulation in response to blood pressure with resultant ischemia to white matter supplied by A $\beta$ -laden meningocortical arteries [43]; 3. — Lowering the threshold of dementia in a predisposed patient (e.g. AD patient) [40,44] either by neuronal loss associated with the severity of CAA [45] or by functional disturbance of the transport of essential nutrients across the blood–brain barrier as a result of the deposition of A $\beta$  in capillaries [44]. The observed patterns of cognitive impairment are related to the underlying mechanisms and location of vascular injury in CAA. Stepwise decline can occur both with recurrent lobar hemorrhage as well as with recurrent ischemia [1–3]. Though CAA related hemorrhages are known to involve cortical–subcortical regions and spare the structures most usually associated with strategic infarcts (e.g. thalamus and caudate), affected cortical regions (e.g. right parietal) can be, solely responsible for the cognitive decline of patients with CAA [46]. As well, the severity of white matter disease (WMD) and the presence of silent infarction on MRI scan are identified as important risk factors for subsequent cognitive decline and development of dementia [47].

Several studies support the hypothesis that there are links between white matter damage and CAA. There is diffuse white matter damage in hereditary forms of CAA (e.g. Dutch and Iowa) while in sporadic CAA there is often periventricular




Fig. 7. Axial T2\* GRE MR image of autopsy proven sporadic CAA case that experimented transient neurological symptoms (aphasia and right motor deficit). This image shows low signal in a gyriform pattern over the left parietal lobe, attributable to deposited iron products in a superficial meningeal or subpial location.

leukoencephalopathy featuring gliosis, myelin loss, and hyalinization of the blood vessel walls [30,48]. White matter disease is associated with increased cognitive decline in about 20% of the patients, particularly those where it is severe. Patients with WMD have a high incidence of concomitant petechial hemorrhages that can negatively affect their cognitive performance [39]. Though not specifically addressed, it is presumed that the involvement of the subcortical and particularly periventricular white matter, and the concomitant circuit disruption leads to a pattern of cognitive decline where apathy, impaired executive function and psychomotor slowing predominating over impaired episodic memory. On the other hand, when the memory deficit is the dominant presenting feature, over speed of processing, gait and other cognitive domains it is plausible that both parenchymal and vascular amyloid are playing synergistic roles in the process.

### 3.4. Subacute to rapidly progressive cognitive decline

There are some clinical circumstances, where CAA becomes associated with a more rapidly progressive rate of cognitive decline and dementia. In such instances the A $\beta$  vascular deposits or CAA seem to become associated with a full blown angitis, where there is a profuse inflammatory response including multinucleated giant cells in vessel walls and where macrophages can be seen to have internalized amyloid [49,50]. There have been a total of 41 cases with such a CAA associated angitis wherein a rapidly progressive dementia has been the most frequent presentation. Language and praxis are amongst the most frequently affected cognitive domains within these cases while headaches, seizures, hallucinations and confusion have also been reported concomitantly. These case series are likely biased towards the most serious presentations that progressed to autopsy, leaving open to speculation whether there might be more minor clinical symptoms might be seen in CAA patients with lesser inflammatory responses.

The aggressive clinical and radiographic picture observed in these patients, and the responses to immunosuppressive treatment has suggested that the inflammation is playing a direct role.

In isolated case-reports CAA has been described with a reversible leukoencephalopathy, sometimes associated with documented vessel inflammation, and other times with hypertension though these are not frequent presentations [51,52].

### 3.5. Hereditary forms of CAA

Hereditary forms of CAA are rare and account for a minority of the CAA cases. [2] Most of these forms of CAA are associated with amyloid precursor protein mutations while they also occur with mutations of cystatin C, presenilin, prion protein, transthyretin, gelsolin and in other amyloidoses [30,31,53–80]. There is some recognized phenotypic variability where the clinical characteristics within the hereditary group include a broader spectrum of presentations than

within sporadic disease, usually with a younger age of onset (Table 1). Clinical signs in the hereditary forms can include spastic paraparesis, extrapyramidal signs, progressive ataxia or ocular disturbances, features that have not been part of the spectrum of sporadic cases [66–75].

These already characterized families (Table 1) demonstrate the pleomorphism of CAA clinical presentations and the need for further characterization of sporadic CAA to elaborate whether these are systematic differences between the hereditary and sporadic forms.

### 3.6. Aging and CAA

CAA is a frequent finding in autopsy series of individuals without dementia (3–50%), particularly over the age of 65. It is not clear if in this context CAA is part of the normal aging process [3,7,18] or whether individuals were completely asymptomatic as these relationships are most often addressed retrospectively. The majority of these individuals have mild or moderate CAA, with symptoms and signs that may be very subtle or remain unnoticed by clinicians [2,3].

## 4. Conclusion

CAA describes pathological changes that occur in cerebral blood vessels, both leptomeningeal and cortical resulting from the deposition of amyloid proteins. The spectrum of clinical manifestations of CAA is evolving with recognition of hemorrhagic complications that range from subtle siderosis to large lobar hemorrhages. CAA has important relationships with cerebral white matter disease, subarachnoid hemorrhage and intraparenchymal lesions. Vascular function may be impeded in a variety of ways in CAA which allows for a broad clinical spectrum to be appreciated. In rare instances it may act as a trigger to a more aggressive dementia and inflammatory vasculopathy. Intracerebral hemorrhage in a cortical–subcortical location remains the most reliable method for diagnosing CAA during life aided through the use of gradient echo MRI.

## Acknowledgements

The authors gratefully acknowledge the technical support of Mr. Jacob Grand in the preparation of the figures in this paper.

## References

- [1] Vinters HV. Cerebral amyloid angiopathy. A critical review. Stroke Mar–Apr 1987;18(2):311–24.
- [2] Revesz T, Ghiso J, Lashley T, Plant G, Rostagno A, Frangione B, et al. Cerebral amyloid angiopathies: a pathologic, biochemical, and genetic view. J Neuropathol Exp Neurol Sep 2003;62(9):885–98.
- [3] Greenberg SM. Cerebral amyloid angiopathy: prospects for clinical diagnosis and treatment. Neurology 1998;51:690–4.
- [4] Herzig MC, Winkler DT, Burgermeister P, Pfeifer M, Kohler E, Schmidt SD, et al. Abeta is targeted to the vasculature in a mouse model of hereditary cerebral hemorrhage with amyloidosis. Nat Neurosci Sep 2004;7(9):954–60.

- [5] Fryer JD, Simmons K, Parsadanian M, Bales KR, Paul SM, Sullivan PM, et al. Human apolipoprotein E4 alters the amyloid-beta 40:42 ratio and promotes the formation of cerebral amyloid angiopathy in an amyloid precursor protein transgenic model. *J Neurosci* Mar 16 2005;25(11):2803–10.
- [6] Vonsattel JP, Myers RH, Hedley-Whyte ET, Ropper AH, Bird ED, Richardson Jr EP. Cerebral amyloid angiopathy without and with cerebral hemorrhages: a comparative histological study. *Ann Neurol* 1991;30:637–49.
- [7] Neuropathology Group. Medical Research Council Cognitive Function and Aging Study. Pathological correlates of late-onset dementia in a multicentre, community-based population in England and Wales. Neuropathology Group of the Medical Research Council Cognitive Function and Ageing Study (MRC CFAS). *Lancet* Jan 20 2001;357(9251):169–75.
- [8] Jellinger KA, Attems J. Prevalence and pathogenic role of cerebrovascular lesions in Alzheimer disease. *J Neurol Sci* Mar 15 2005;229–230:37–41.
- [9] Mastaglia FL, Byrnes ML, Johnsen RD, Kakulas BA. Prevalence of cerebral vascular amyloid-beta deposition and stroke in an aging Australian population: a postmortem study. *J Clin Neurosci* Mar 2003;10(2):186–9.
- [10] Smith EE, Eichler F. Cerebral amyloid angiopathy and lobar intracerebral hemorrhage. *Arch Neurol* Jan 2006;63(1):148–51.
- [11] Neumann MA. Combined amyloid vascular changes and argyrophilic plaques in the central nervous system. *J Neuropathol Exp Neurol* 1960;19:370–82.
- [12] Okazaki H, Reagan TJ, Campbell RJ. Clinicopathologic studies of primary cerebral amyloid angiopathy. *Mayo Clin Proc* 1979;54:22–31.
- [13] Knudsen KA, Rosand J, Karluk D, Greenberg SM. Clinical diagnosis of cerebral amyloid angiopathy: validation of the Boston criteria. *Neurology* Feb 27 2001;56(4):537–9.
- [14] Smith EE, Greenberg SM. Clinical diagnosis of cerebral amyloid angiopathy: validation of the Boston criteria. *Curr Atheroscler Rep* Jul 2003;5(4):260–6.
- [15] Yamada M. Cerebral amyloid angiopathy: an overview. *Neuropathology* Mar 2000;20(1):8–22.
- [16] Haan J, Maat-Schieman ML, Roos RA. Clinical aspects of cerebral amyloid angiopathy. *Dementia* May–Aug 1994;5(3–4):210–3.
- [17] Ellis RJ, Olichney JM, Thal LJ, Mirra SS, Morris JC, Beekly D, et al. Cerebral amyloid angiopathy in the brains of patients with Alzheimer's disease: the CERAD experience, Part XV. *Neurology* Jun 1996;46(6):1592–6.
- [18] Jellinger KA. Alzheimer disease and cerebrovascular pathology: an update. *J Neural Transm* 2002;109:813–36.
- [19] Warlow CP, Dennis MS, van Gijn J, Hankey GJ, Sandercock PAG, Bamford JM, et al. What pathological type of stroke is it? In: Warlow CP, editor. *Stroke, a practical guide to management*, 2nd ed. Oxford: Blackwell Science; 2001. p. 151–222.
- [20] Greenberg SM, Finklestein SP, Schaefer PW. Petechial hemorrhages accompanying lobar hemorrhage: detection by gradient-echo MRI. *Neurology* 1996;46:1751–4.
- [21] O'Donnell HC, Rosand J, Knudsen KA, Furie KL, Segal AZ, Chiu RI, et al. Apolipoprotein E genotype and the risk of recurrent lobar intracerebral hemorrhage. *N Engl J Med* 2000;342:240–5.
- [22] Greenberg SM, Vonsattel JP, Segal AZ, Chiu RI, Clatworthy AE, Liao A, et al. Association of apolipoprotein E epsilon2 and vasculopathy in cerebral amyloid angiopathy. *Neurology* Apr 1998;50(4):961–5.
- [23] Nicoll JA, Burnett C, Love S, Graham DI, Ironside JW, Vinters HV. High frequency of apolipoprotein E epsilon 2 in patients with cerebral hemorrhage due to cerebral amyloid angiopathy. *Ann Neurol* May 1996;39(5):682–3.
- [24] McCarron MO, Nicoll JA. Cerebral amyloid angiopathy and thrombolysis-related intracerebral haemorrhage. *Lancet Neurol* Aug 2004;3(8):484–92.
- [25] Kakuda W, Thijs VN, Lansberg MG, Bammer R, Wechsler L, Kemp S, et al. DEFUSE Investigators. Clinical importance of microbleeds in patients receiving IV thrombolysis. *Neurology* Oct 25 2005;65(8):1175–8.
- [26] Koennecke HC. Cerebral microbleeds on MRI: prevalence, associations, and potential clinical implications. *Neurology* Jan 24 2006;66(2):165–71.
- [27] Greenberg SM, Vonsattel JP, Stakes JW, Gruber M, Finklestein SP. The clinical spectrum of cerebral amyloid angiopathy: presentations without lobar hemorrhage. *Neurology* Oct 1993;43(10):2073–9.
- [28] Olichney JM, Hansen LA, Hofstetter CR, Grundman M, Katzman R, Thal LJ. Cerebral infarction in Alzheimer's disease is associated with severe amyloid angiopathy and hypertension. *Arch Neurol* 1995;52:702–8.
- [29] Olichney JM, Ellis RJ, Katzman R, Sabbagh MN, Hansen L. Types of cerebrovascular lesions associated with severe cerebral amyloid angiopathy in Alzheimer's disease. *Ann N Y Acad Sci* 1997;826:493–7.
- [30] Bornebroek M, Haan J, Maat-Schieman ML, Van Duinen SG, Roos RA. Hereditary cerebral hemorrhage with amyloidosis-Dutch type (HCHWA-D): I — A review of clinical, radiologic and genetic aspects. *Brain Pathol* Apr 1996;6(2):111–4.
- [31] Grabowski TJ, Cho HS, Vonsattel JP, Rebeck GW, Greenberg SM. Novel amyloid precursor protein mutation in an Iowa family with dementia and severe cerebral amyloid angiopathy. *Ann Neurol* Jun 2001;49(6):697–705.
- [32] Cadavid D, Mena H, Koeller K, Frommelt RA. Cerebral beta amyloid angiopathy is a risk factor for cerebral ischemic infarction. A case control study in human brain biopsies. *J Neuropathol Exp Neurol* Sep 2000;59(9):768–73.
- [33] Greenberg SM. Cerebral amyloid angiopathy and vessel dysfunction. *Cerebrovasc Dis* 2002;13(suppl 2):42–7.
- [34] Kalaria RN. The role of cerebral ischemia in Alzheimer's disease. *Neurobiol Aging* Mar–Apr 2000;21(2):321–30.
- [35] Yamada M, Itoh Y, Otomo E, Hayakawa M, Miyatake T. Subarachnoid hemorrhage in the elderly: a necropsy study of the association with cerebral amyloid angiopathy. *J Neurol Neurosurg Psychiatry* 1993;56:543–7.
- [36] Takeda S, Yamazaki K, Miyakawa T, Onda K, Hinokuma K, Ikuta F, et al. Subcortical hematoma caused by cerebral amyloid angiopathy: does the first evidence of hemorrhage occur in the subarachnoid space? *Neuropathology* 2003;23:254–61.
- [37] Maia LF, Botelho L, Correia MM. Commentary on 'Subcortical hematoma caused by cerebral amyloid angiopathy: does the first evidence of hemorrhage occur in the subarachnoid space?' (*Neuropathology* 2003; 23, 254–261). *Neuropathology* Dec 2004;24(4):354–5.
- [38] Roch JA, Nighoghossian N, Hermier M, Cakmak S, Picot M, Honnorat J, et al. Transient neurologic symptoms related to cerebral amyloid angiopathy: usefulness of T2\*-weighted imaging. *Cerebrovasc Dis* 2005;20(5):412–4.
- [39] Smith EE, Gurol ME, Eng JA, Engel CR, Nguyen TN, Rosand J, et al. White matter lesions, cognition, and recurrent hemorrhage in lobar intracerebral hemorrhage. *Neurology* Nov 9 2004;63(9):1606–12.
- [40] Pfeifer LA, White LR, Ross GW, Petrovitch H, Launer LJ. Cerebral amyloid angiopathy and cognitive function: the HAAS autopsy study. *Neurology* 2002;58:1629–34.
- [41] Greenberg SM, Gurol ME, Rosand J, Smith EE. Amyloid angiopathy-related vascular cognitive impairment. *Stroke* Nov 2004;35(11 Suppl 1):2616–9.
- [42] Wallin A, Milos V, Sjogren M, Pantoni L, Erkinjuntti T. Classification and subtypes of vascular dementia. In: Erkinjuntti T, Gauthier S, editors. *Vascular cognitive impairment*, 1st ed. Martin Dunitz; 2002. p. 27–42.
- [43] Imaoka K, Kobayashi S, Fujihara S, Shimode K, Nagasaki M. Leukoencephalopathy with cerebral amyloid angiopathy: a semiquantitative and morphometric study. *J Neurol* Aug 1999;246(8):661–6.
- [44] Attems J. Sporadic cerebral amyloid angiopathy: pathology, clinical implications, and possible pathomechanisms. *Acta Neuropathol (Berl)* Oct 2005;110(4):345–59.
- [45] Zarow C, Zaias B, Lyness SA, Chui H. Cerebral amyloid angiopathy in Alzheimer disease is associated with apolipoprotein E4 and cortical neuron loss. *Alzheimer Dis Assoc Disord* Jan 1999;13(1):1–8.

- [46] Knopman DS. Vascular dementia. *Contin Neurol* Feb 2004(1):113–34.
- [47] Vermeer SE, Hollander M, van Dijk EJ, Hofman A, Koudstaal PJ, Breteler MM. Rotterdam Scan Study. Silent brain infarcts and white matter lesions increase stroke risk in the general population: the Rotterdam Scan Study. *Stroke* May 2003;34(5):1126–9.
- [48] Pantoni L, Garcia JH. Cognitive impairment and cellular/vascular changes in the cerebral white matter. *Ann N Y Acad Sci* Sep 26 1997;826:92–102.
- [49] Eng JA, Frosch MP, Choi K, Rebeck GW, Greenberg SM. Clinical manifestations of cerebral amyloid angiopathy-related inflammation. *Ann Neurol* Feb 2004;55(2):250–6.
- [50] Scolding NJ, Joseph F, Kirby PA, Mazanti I, Gray F, Mikol J, et al. Abeta-related angiitis: primary angiitis of the central nervous system associated with cerebral amyloid angiopathy. *Brain* Mar 2005;128(Pt 3):500–15.
- [51] Caulo M, Tampieri D, Brassard R, Christine Guiot M, Melanson D. Cerebral amyloid angiopathy presenting as nonhemorrhagic diffuse encephalopathy: neuropathologic and neuroradiologic manifestations in one case. *AJNR Am J Neuroradiol* Jun–Jul 2001;22(6):1072–6.
- [52] Oh U, Gupta R, Krakauer JW, Khandji AG, Chin SS, Elkind MS. Reversible leukoencephalopathy associated with cerebral amyloid angiopathy. *Neurology* Feb 10 2004;62(3):494–7.
- [53] Mead S, James-Galton M, Revesz T, Doshi RB, Harwood G, Pan EL, et al. Familial British dementia with amyloid angiopathy: early clinical, neuropsychological and imaging findings. *Brain* 2000;123(Pt 5):975–91.
- [54] Wattendorff AR, Bots GT, Went LN, Endtz LJ. Familial cerebral amyloid angiopathy presenting as recurrent cerebral hemorrhage. *J Neurol Sci* Aug 1982;55(2):121–35.
- [55] Levy E, Carman MD, Fernandez-Madrid IJ, Power MD, Lieberburg I, van Duinen SG, et al. Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage, Dutch type. *Science* Jun 1 1990;248(4959):1124–6.
- [56] Natte R, Maat-Schieman ML, Haan J, Bornebroek M, Roos RA, van Duinen SG. Dementia in hereditary cerebral hemorrhage with amyloidosis-Dutch type is associated with cerebral amyloid angiopathy but is independent of plaques and neurofibrillary tangles. *Ann Neurol* Dec 2001;50(6):765–72.
- [57] Maat-Schieman M, Roos R, van Duinen S. Hereditary cerebral hemorrhage with amyloidosis-Dutch type. *Neuropathology* Dec 2005;25(4):288–97.
- [58] van den Boom R, Bornebroek M, Behloul F, van den Berg-Huysmans AA, Haan J, van Buchem MA. Microbleeds in hereditary cerebral hemorrhage with amyloidosis-Dutch type. *Neurology* Apr 12 2005;64(7):1288–9.
- [59] Greenberg SM, Shin Y, Grabowski TJ, Cooper GE, Rebeck GW, Iglesias S, et al. Hemorrhagic stroke associated with the Iowa amyloid precursor protein mutation. *Neurology* Mar 25 2003;60(6):1020–2.
- [60] Roks G, Van Harskamp F, De Koning I, Cruts M, De Jonghe C, Kumar-Singh S, et al. Presentation of amyloidosis in carriers of the codon 692 mutation in the amyloid precursor protein gene (APP692). *Brain* Oct 2000;123(Pt 10):2130–40.
- [61] Cras P, van Harskamp F, Hendriks L, Ceuterick C, van Duijn CM, Stefanko SZ, et al. Presenile Alzheimer dementia characterized by amyloid angiopathy and large amyloid core type senile plaques in the APP 692Ala→Gly mutation. *Acta Neuropathol (Berl)* Sep 1998;96(3):253–60.
- [62] Hendriks L, van Duijn CM, Cras P, Cruts M, Van Hul W, van Harskamp F, et al. Presenile dementia and cerebral haemorrhage linked to a mutation at codon 692 of the beta-amyloid precursor protein gene. *Nat Genet* Jun 1992;1(3):218–21.
- [63] Tagliavini F, Rossi G, Padovani A, Magoni M, Andora G, Sgarzi M, et al. A new bPP mutation related to hereditary cerebral haemorrhage. *Alzheimer's Rep* 1999;2:S28 [Suppl.].
- [64] Melchor JP, McVoy L, Van Nostrand WE. Charge alterations of E22 enhance the pathogenic properties of the amyloid beta-protein. *J Neurochem* May 2000;74(5):2209–12.
- [65] Palsdottir A, Abrahamson M, Thorsteinsson L, Arnason A, Olafsson I, Grubb A, et al. Mutation in cystatin C gene causes hereditary brain haemorrhage. *Lancet* Sep 10 1988;2(8611):603–4.
- [66] Vidal R, Garzuly F, Budka H, Lalowski M, Linke RP, Brittig F, et al. Meningocerebrovascular amyloidosis associated with a novel transthyretin mis-sense mutation at codon 18 (TTR D18G). *Am J Pathol* Feb 1996;148(2):361–6.
- [67] Garzuly F, Vidal R, Wisniewski T, Brittig F, Budka H. Familial meningocerebrovascular amyloidosis, Hungarian type, with mutant transthyretin (TTR Asp18Gly). *Neurology* Dec 1996;47(6):1562–7.
- [68] Petersen RB, Goren H, Cohen M, Richardson SL, Tresser N, Lynn A, et al. Transthyretin amyloidosis: a new mutation associated with dementia. *Ann Neurol* Mar 1997;41(3):307–13.
- [69] Blevins G, Macaulay R, Harder S, Fladeland D, Yamashita T, Yazaki M, et al. Oculoleptomeningeal amyloidosis in a large kindred with a new transthyretin variant Tyr69His. *Neurology* May 27 2003;60(10):1625–30.
- [70] Jin K, Sato S, Takahashi T, Nakazaki H, Date Y, Nakazato M, et al. Familial leptomeningeal amyloidosis with a transthyretin variant Asp18Gly representing repeated subarachnoid haemorrhages with superficial siderosis. *J Neurol Neurosurg Psychiatry* Oct 2004;75(10):1463–6.
- [71] Ellie E, Camou F, Vital A, Rummens C, Grateau G, Delpech M, et al. Recurrent subarachnoid hemorrhage associated with a new transthyretin variant (Gly53Glu). *Neurology* Jul 10 2001;57(1):135–7.
- [72] Kiuru S, Salonen O, Haltia M. Gelsolin-related spinal and cerebral amyloid angiopathy. *Ann Neurol* Mar 1999;45(3):305–11.
- [73] Ghetti B, Piccardo P, Spillantini MG, Ichimiya Y, Porro M, Perini F, et al. Vascular variant of prion protein cerebral amyloidosis with tau-positive neurofibrillary tangles: the phenotype of the stop codon 145 mutation in PRNP. *Proc Natl Acad Sci U S A* Jan 23 1996;93(2): 744–8.
- [74] Vidal R, Revesz T, Rostagno A, Kim E, Holton JL, Bek T, et al. A decamer duplication in the 3' region of the BRI gene originates an amyloid peptide that is associated with dementia in a Danish kindred. *Proc Natl Acad Sci U S A* Apr 25 2000;97(9):4920–5.
- [75] Holton JL, Lashley T, Ghiso J, Braendgaard H, Vidal R, Guerin CJ, et al. Familial Danish dementia: a novel form of cerebral amyloidosis associated with deposition of both amyloid-Dan and amyloid-beta. *J Neuropathol Exp Neurol* Mar 2002;61(3):254–67.
- [76] Houlden H, Baker M, McGowan E, Lewis P, Hutton M, Crook R, et al. Variant Alzheimer's disease with spastic paraparesis and cotton wool plaques is caused by PS-1 mutations that lead to exceptionally high amyloid-beta concentrations. *Ann Neurol* Nov 2000;48(5):806–8.
- [77] Dermaut B, Kumar-Singh S, De Jonghe C, Cruts M, Lofgren A, Lubke U, et al. Cerebral amyloid angiopathy is a pathogenic lesion in Alzheimer's disease due to a novel presenilin 1 mutation. *Brain* Dec 2001;124(Pt 12):2383–92.
- [78] Nochlin D, Bird TD, Nemens EJ, Ball MJ, Sumi SM. Amyloid angiopathy in a Volga German family with Alzheimer's disease and a presenilin-2 mutation (N141I). *Ann Neurol* Jan 1998;43(1):131–5.
- [79] Rossor MN, Newman S, Frackowiak RS, Lantos P, Kennedy AM. Alzheimer's disease families with amyloid precursor protein mutations. *Ann N Y Acad Sci* Sep 24 1993;695:198–202.
- [80] Kalaria RN, Cohen DL, Greenberg BD, Savage MJ, Bogdanovic NE, Winblad B, et al. Abundance of the longer A beta 42 in neocortical and cerebrovascular amyloid beta deposits in Swedish familial Alzheimer's disease and Down's syndrome. *Neuroreport* May 31 1996;7(8):1377–81.
- [81] Donahue JE, Khurana JS, Adelman LS. Intracerebral hemorrhage in two patients with Down's syndrome and cerebral amyloid angiopathy. *Acta Neuropathol (Berl)* Feb 1998;95(2):213–6.