

NOT TO BE CITED WITHOUT PRIOR
REFERENCE TO THE AUTHOR(S)

Northwest Atlantic



Fisheries Organization

Serial No. N5758

NAFO SCR Doc. 10/09

SCIENTIFIC COUNCIL MEETING – JUNE 2010

Atlantic cod and Yellowtail flounder indices from the Spanish Survey conducted in Divisions 3NO
of the NAFO Regulatory Area

by

D. González-Troncoso, C. Gonzalez and X. Paz

Instituto Español de Oceanografía,
P.O. Box 1552. Vigo, Spain.
e-mail: diana.gonzalez@vi.ieo.es

Abstract

Since 1995, Spain carries out a stratified random spring bottom trawl survey in Div. 3NO of the NAFO Regulatory Area. The entire series of mean catches, biomass and length distribution for Atlantic cod (*Gadus morhua*) are presented for the period 1997-2009 and for Yellowtail flounder (*Limanda ferruginea*) for the period 1995-2009. For Atlantic cod we can see a general decreasing in the biomass between 2002 and 2005, and an increasing since then, especially in 2006 and, higher, in 2009, although this values is mainly due to two hauls with high catches. In 2007 the biomass decreased, but the level is over the value in the period 2002-2005. For this species, an increase in the recruitment can be seen in 2004 and 2005, and from 2007 the youngest length classes are much over the rest of the length classes. For Yellowtail flounder, there is no a clear trend since 1998; its indices are almost constant along this period.

Material and methods

The survey in Div. 3NO of NAFO Regulatory Area was initiated by Spain in 1995. Until 2001, the survey was carried out in Spring (May), on board the Spanish vessel C/V *Playa de Menduiña* (338 GT and 800 HP) using bottom trawl net type *Pedreira*. Since 2001, the R/V *Vizconde de Eza* replaced the C/V *Playa de Menduiña* as the research vessel for the survey, and *Campelen* net replaced *Pedreira* net as survey gear. The main specifications and geometry of these gears, as the rigging profile and the net plan, and a sheet with the resume of the main technical data of the survey are described in a previous paper (Walsh *et. al.*, 2001). Table 1 presents the number of valid tows, the depth strata covered and the dates of the survey series. The survey area was stratified following the standard stratification schemes (Bishop, 1994). Set number was allocated to strata proportionally to their size, with a minimum of two planned hauls per stratum and the trawl positions were chosen at random (Doubleday, 1981). Biomass indices were calculated by the swept area method (Cochran, 1997), assuming catchability factor of 1.

For Atlantic cod, the series are presented since 1997 because in years 1995 and 1996 the surveyed depth strata were only until 1000 meters, so they are not representative. As the strata where the Yellowtail flounder is presented were well surveyed, the series for this species are presented since 1995.

The catch from each haul was sorted by species and weighted. Random samples of each species catches were measured to total length to the nearest lower cm. Length distribution scaled from catches was estimated for the period 1997-2009 (Atlantic cod) and 1995-2009 (Yellowtail flounder) in two cm range. Data were grouping beginning with the pair number.

For each species, the haul mean catch, with its variance, and the stratified mean catches by stratum and year, with the annual variance, are presented, transformed until 2000 and no-transformed in the period 2002-2009. In the year 2001, there are data transformed from the former vessel with original data from the new vessel. Besides this, the biomass per stratum and year, with the annual variance, are presented, as the stratified mean catches per haul length distribution. To more information about the calculation of these indices, see González Troncoso *et al.*, 2004 and Paz *et al.*, 2004.

Due to technical problems in the vessel, this year two strata were not surveyed and six more have only one haul, so there are no standard deviations for these strata. As only one of these strata usually have catch of Atlantic cod and Yellowtail flounder, this fact is not significant for the calculation of the final standard deviation for the mean weight per tow and biomass.

Results

Atlantic cod

Atlantic cod in Divisions 3NO has been under moratorium to directed fishing since 1994. According to the NAFO Scientific Council, the stock of Atlantic cod in Divisions 3NO declined dramatically during the mid-1980s, and the total biomass and the spawning biomass are currently estimated to be at an extremely low levels (NAFO, 2009).

Mean Catches and Biomass

The Atlantic cod haul mean catches by stratum are presented in Table 2, included swept area, number of hauls and SD. Atlantic cod stratified mean catches per tow by stratum and year and their SD are presented in Table 3.

The entire time series (1997-2009) of biomass and their SD estimates for Atlantic cod are presented in Table 4. Estimated parameters a and b values of length-weight relationship are presented in Table 5.

We can see a great variation in the cod indices since 1997, but this is due to a few hauls in which the presence of cod was very high. For example, in 1998 and 2001, the C/V *Playa de Menduiña* made a more than seven tons cod catch in a single haul. Besides this, in 2001, the R/V *Vizconde de Eza* made two hauls with more than a ton of cod catches. But before year 2006, and apart from those hauls, the catches of cod were very poor. Between 2002 and 2005 there was a decreasing in the biomass. Since 2006, we can see an increasing trend in the biomass of this species. Although the 2006 increase is above all for a single catch of almost 2 tons, in general the catches of Atlantic cod in the survey of 2006 were over the mean. In 2007 we can see a decrease in the biomass over the 2006 biomass, but still remains greater than in the 2002-2005 period. In 2008 a new high increase is shown, reaching the second highest value in the time series, and in this case there is no haul with very high catches (the maximum was 585.5 kg). And in 2009 the biomass reaches a new maximum, well above the rest of the values of the series, due most of all to two hauls with 2.5 and 3.2 tons of catch, respectively. If these two hauls are not taking into account, we obtain a level more or less as the 2006 value. The great value of the variance in some years is due to the tows with a large catch (Fig. 1 and 2).

Length Distribution

Table 6 and Figures 3 and 4 show the stratified mean catches per haul length distribution by year, besides the sampled size and its catch, for the period 1997-2009. The data have been grouped two by two, so we present the data every two cm. The modal values used to be very low before the year 2006, except in 2001, and in general all lengths presence is very low, even it is very difficult to follow the modal values. In 2001 we have a good presence of individuals between 36 and 58 cm, probably due to the three hauls with great catches of this year. From 2006 it can be seen a series of great modal values along the length distribution. In 2006 there is two modes in the length distribution, one around 30 cm and another one around 40 cm. There is no good recruitment until 2004, in which the individuals between 12 and 16 cm correspond to the greatest presence in the series, and in 2005 between 24 and 32, with a new mode between 12 and 16 cm, as in last year. In 2007 the youngest lengths dominate the length range, with the highest mode in the lengths 12-16, that are between 2 and 4 times the abundance of the 48 cm length class, the following mode. In 2008 and in 2009 we can follow the evolution of these lengths, being in 2007 the dominant lengths the ones between 20 and 26 and in 2009 between 30 and 36 cm.

Yellowtail flounder

After a moratorium between 1994 and 1997, the Yellowtail flounder fishery is under TAC. According to the Report of NAFO Scientific Council Meeting, the stock size had a minimum in the mid 1990's, but since 1994 has steadily increased and now it is estimated to be at a level well above that of the mid-1980s (NAFO, 2009).

Mean Catches and Biomass

In Table 7 we present the haul mean catches by stratum for Yellowtail flounder, included swept area, number of hauls and SD. The stratified mean catches per tow by stratum and year and their SD are presented in Table 8 for this species.

The entire time series (1995-2009) of biomass by the swept area method and their SD estimates of Yellowtail flounder are presented in Table 9. The parameters a and b for the calculation of the length-weight relationship are presented in Table 10.

The Yellowtail flounder indices show no clear trend along the time (in the entire series). There was an increasing between 1995 and 1999 and since 2001 the indices are stabilised at a high level (Figures 5 and 6).

Length Distribution

The stratified mean catches per haul length distribution by sex and year, besides the sampled size and its catch, are presented in Table 11 and Figure 7 the period 1995-2009. The data have been grouped two by two, so we present the data every two cm. There is no presence of good recruitment in last years. In Figure 8, we can see the evolution of a modal value since the beginning of the series, but, although there is a presence of juveniles in the lengths, this presence is very low. In the length distribution it can be seen a small movement of the adult segment for several years, with a mode for males of 33 cm in this year, only 2 cm more than in the five last years, and a mode for females of 36 cm, only 1 cm more than in the two last years. Despite that, there is a small proportion of individuals with lengths lower than 20 cm. This situation is possibly due to a high exploitation rate that compensates the growth.

References

- Bishop, C A.. 1994. Revisions and additions to stratification schemes used during research vessel surveys in NAFO subareas 2 and 3. NAFO SCR Doc. 94/43, Serial nº N2413, 23 pp.
- Cochran, W. G.. 1997. Sampling techniques. J. Wiley and Sons, N.Y., 428 pp.
- Doubleday, W. G.. 1981. Manual on groundfish surveys in the Northwest Atlantic. NAFO Sci. Coun. Studies, 2, 55.
- González Troncoso, D., X. Paz and C. González. 2004. Atlantic cod population indices obtained from the Spring surveys conducted by Spain in the NAFO Regulatory Area of Divisions 3NO, 1995-2003. NAFO SCR Doc. 04/12, Serial Number N4957, 21 pp.
- NAFO, 2006. Report of Scientific Council Meeting, 1-15 June 2006
- NAFO, 2009. Report of Scientific Council Meeting, 4-18 June 2009
- Paz, X., D. González Troncoso and E. Román. 2004. New time series for Yellowtail flounder from the comparative experience between the C/V *Playa de Mendumá* and the R/V *Vizconde de Eza* in the NAFO Regulatory Area of Divisions 3NO, 1995-2003. NAFO SCR Doc. 04/10, Serial Number N4955, 19 pp.
- Walsh, J.S., X. Paz and P. Durán. 2001. A preliminary investigation of the efficiency of Canadian and Spanish Survey bottom trawls on the Southern Bank. NAFO SCR Doc., 01/74, Serial nº N4453, 18 pp.

TABLE 1.- Spanish spring bottom trawl surveys on NAFO Div. 3NO: 1995-2009

Year	Vessel	Valid tows	Depth strata covered (m)	Dates
1995	C/V <i>Playa de Menduíña</i>	77	42-684	May 18-May 29
1996	C/V <i>Playa de Menduíña</i>	112	41-1135	May 07-May 24
1997	C/V <i>Playa de Menduíña</i>	128	42-1263	April 26-May 18
1998	C/V <i>Playa de Menduíña</i>	124	42-1390	May 06-May 26
1999	C/V <i>Playa de Menduíña</i>	114	41-1381	May 07-May 26
2000	C/V <i>Playa de Menduíña</i>	118	42-1401	May 07-May 28
2001 ^(*)	R/V <i>Vizconde de Eza</i>	83	36-1156	May 03-May 24
	C/V <i>Playa de Menduíña</i>	121	40-1500	May 05-May 23
2002	R/V <i>Vizconde de Eza</i>	125	38-1540	April 29-May 19
2003	R/V <i>Vizconde de Eza</i>	118	38-1666	May 11-June 02
2004	R/V <i>Vizconde de Eza</i>	120	43-1539	June 06-June 24
2005	R/V <i>Vizconde de Eza</i>	119	47-1485	June 10-June 29
2005	R/V <i>Vizconde de Eza</i>	119	47-1485	June 10-June 29
2006	R/V <i>Vizconde de Eza</i>	120	45-1480	June 7-June 27
2007	R/V <i>Vizconde de Eza</i>	110	45-1374	May 29-June 19
2008	R/V <i>Vizconde de Eza</i>	122	45-1374	May 27-June 16
2009	R/V <i>Vizconde de Eza</i>	109	45-1374	May 31-June 18

(*) We took, for the calculation of the series, 83 hauls from the R/V *Vizconde de Eza* and 40 hauls from the C/V *Playa de Menduíña* (123 hauls in total)

TABLE 2.- Swept area, number of hauls and Atlantic cod mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1997-2009. Swept area in square miles. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Mendiña* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1997				1998				1999				2000				2001			
	Swept area	Tow number	A. cod Mean	A. cod SD	Swept area	Tow number	A. cod Mean	A. cod SD	Swept area	Tow number	A. cod Mean	A. cod SD	Swept area	Tow number	A. cod Mean	A. cod SD	Swept area	Tow number	A. cod Mean	A. cod SD
353	0.0480	4	0.00	0.000	0.0465	4	0.00	0.004	0.0360	3	6.26	8.593	0.0356	3	8.59	9.984	0.0341	3	7.33	8.145
354	0.0233	2	0.00	0.000	0.0356	3	17.67	29.046	0.0218	2	4.92	3.192	0.0356	3	18.44	27.099	0.0338	3	16.07	3.315
355	0.0233	2	4.29	5.711	0.0221	2	27.05	3.662	0.0229	2	6.39	2.549	0.0233	2	94.83	76.209	0.0240	2	56.11	64.898
356	0.0225	2	7.80	0.495	0.0221	2	6.23	0.247	0.0229	2	41.19	0.346	0.0225	2	16.34	17.172	0.0240	2	149.60	76.650
357	0.0443	4	91.55	174.202	0.0240	2	7.45	0.742	0.0236	2	10.12	11.461	0.0124	1	9.15	-	0.0244	2	27.20	36.062
358	0.0563	5	1.77	1.655	0.0236	3	4.46	4.030	0.0349	3	9.98	4.006	0.0341	3	184.88	194.829	0.0345	3	3.42	2.592
359	0.0690	6	1.13	2.385	0.0698	6	0.39	0.858	0.0364	3	7.25	11.394	0.0469	4	18.26	17.367	0.0803	7	176.35	433.935
360	0.3754	32	0.11	0.226	0.2561	25	0.22	0.700	0.2325	19	2.33	3.801	0.2396	20	2.16	3.561	0.2423	20	11.36	27.470
374	0.0353	3	0.06	0.099	0.0353	3	0.00	0.000	0.0244	2	0.58	0.594	0.0240	2	0.00	0.000	0.0240	2	0.00	0.000
375	0.0116	1	0.00	-	0.0345	3	0.78	0.403	0.0236	2	0.97	0.579	0.0244	2	0.00	0.000	0.0338	3	0.00	0.000
376	0.1583	14	0.00	0.000	0.0930	10	0.20	0.187	0.1219	10	0.62	0.545	0.1200	10	0.90	1.852	0.1155	10	0.04	0.119
377	0.0116	1	0.27	-	0.0229	2	1.89	2.375	0.0240	2	0.21	0.302	0.0229	2	0.02	0.027	0.0229	2	0.00	0.000
378	0.0210	2	2.34	3.316	0.0120	2	3.46	0.940	0.0229	2	7.76	5.951	0.0233	2	10.65	11.169	0.0236	2	11.98	15.726
379	0.0206	2	3.68	0.307	0.0356	3	8.30	5.847	0.0236	2	5.22	4.147	0.0225	2	41.12	54.683	0.0229	2	9.54	9.001
380	0.0210	2	0.36	0.515	0.0113	2	2.33	1.361	0.0236	2	38.58	48.720	0.0236	2	8.21	3.236	0.0206	2	6.00	2.895
381	0.0221	2	0.07	0.099	0.0229	2	0.21	0.187	0.0229	2	0.87	0.388	0.0236	2	1.74	0.730	0.0236	2	0.66	0.891
382	0.0461	4	0.00	0.000	0.0229	3	0.32	0.336	0.0484	4	0.05	0.036	0.0499	4	0.71	0.561	0.0469	4	0.12	0.145
721	0.0221	2	20.98	7.052	0.0203	2	0.61	0.866	0.0244	2	88.29	106.743	0.0236	2	28.34	17.122	0.0248	2	4.85	6.859
722	0.0214	2	0.31	0.139	0.0101	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.90	1.277	0.0233	2	0.00	0.000
723	0.0210	2	9.90	2.425	0.0233	2	4.39	3.736	0.0229	2	16.87	20.735	0.0248	2	22.02	12.010	0.0240	2	676.15	932.179
724	0.0225	2	1.30	1.269	0.0206	2	1488.84	2101.820	0.0225	2	0.02	0.032	0.0233	2	0.70	0.341	0.0353	3	6.16	10.254
725	0.0206	2	23.50	17.734	0.0086	1	30.86	-	0.0229	2	13.65	19.102	0.0210	2	4.34	3.857	0.0116	2	1367.61	1856.733
726	n.s.	n.s.	n.s.	n.s.	0.0094	2	4.74	5.617	0.0225	2	0.81	0.492	0.0221	2	8.85	12.221	0.0116	2	1.83	2.593
727	0.0094	1	0.12	-	0.0233	2	2.66	2.821	0.0236	2	9.20	4.701	0.0210	2	9.16	10.803	0.0225	2	10.40	4.810
728	0.0214	2	1.17	0.569	0.0206	2	1.54	2.177	0.0233	2	0.00	0.000	0.0210	2	0.90	1.267	0.0229	2	0.00	0.000
752	0.0218	2	0.00	0.000	0.0229	2	0.00	0.000	0.0233	2	0.00	0.000	0.0206	2	0.00	0.000	0.0210	2	0.00	0.000
753	0.0214	2	0.00	0.000	0.0218	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000
754	0.0330	3	0.00	0.000	0.0210	2	0.00	0.000	0.0206	2	0.00	0.000	0.0195	2	0.00	0.000	0.0195	2	0.00	0.000
755	n.s.	n.s.	n.s.	n.s.	0.0206	2	0.00	0.000	0.0311	3	0.00	0.000	0.0431	4	0.00	0.000	0.0416	4	0.00	0.000
756	0.0109	1	0.00	-	0.0225	2	0.32	0.449	0.0225	2	0.24	0.334	0.0203	2	0.36	0.257	0.0113	2	0.04	0.057
757	0.0304	3	0.00	0.000	0.0206	2	0.00	0.000	0.0233	2	0.00	0.000	0.0214	2	0.00	0.000	0.0233	2	0.00	0.000
758	0.0214	2	0.00	0.000	0.0105	2	0.00	0.000	0.0214	2	0.00	0.000	0.0210	2	0.00	0.000	0.0218	2	0.00	0.000
759	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.000	0.0218	2	0.00	0.000	0.0210	2	0.00	0.000	0.0221	2	0.00	0.000
760	0.0105	1	0.00	-	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000	0.0210	2	0.00	0.000	0.0229	2	0.00	0.000
761	0.0315	3	0.00	0.000	0.0206	2	0.00	0.000	0.0210	2	0.00	0.000	0.0221	2	0.00	0.000	0.0225	2	0.00	0.000
762	0.0308	3	0.00	0.000	0.0094	2	0.00	0.000	0.0210	2	0.00	0.000	0.0203	2	0.00	0.000	0.0116	2	0.00	0.000
763	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.000	0.0311	3	0.00	0.000	0.0416	4	1.08	2.170	0.0330	3	0.00	0.000
764	0.0206	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0240	2	0.00	0.000
765	0.0206	2	0.00	0.000	0.0098	2	0.00	0.000	0.0221	2	0.00	0.000	0.0203	2	0.00	0.000	0.0113	2	0.00	0.000
766	0.0308	3	0.00	0.000	0.0191	2	0.00	0.000	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000	0.0203	2	0.00	0.000
767	n.s.	n.s.	n.s.	n.s.	0.0109	2	0.00	0.000	0.0214	2	0.00	0.000	0.0210	2	0.00	0.000	0.0225	2	0.00	0.000

TABLE 2 (cont.).- Swept area, number of hauls and Atlantic cod mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1997-2009. Swept area in square miles. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Menduiña* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2002				2003				2004				2005			
	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD
353	0.0476	4	0.00	0.003	0.0334	3	0.00	0.000	0.0338	3	10.21	8.691	0.0353	3	4.20	3.962
354	0.0356	3	0.01	0.012	0.0338	3	7.63	13.221	0.0345	3	4.76	3.335	0.0353	3	6.76	8.311
355	0.0236	2	0.96	0.370	0.0229	2	3.02	2.390	0.0229	2	5.09	3.267	0.0225	2	1.97	0.255
356	0.0233	2	15.20	10.889	0.0225	2	15.61	1.605	0.0221	2	2.97	0.714	0.0233	2	1.43	1.478
357	0.0240	2	6.65	1.909	0.0229	2	5.28	7.460	0.0229	2	13.30	17.727	0.0233	2	3.98	4.603
358	0.0345	3	2.63	1.429	0.0338	3	207.22	260.186	0.0330	3	14.41	12.455	0.0349	3	22.75	17.967
359	0.0686	6	2.72	3.436	0.0791	7	1.03	1.522	0.0791	7	29.83	54.712	0.0814	7	57.31	134.609
360	0.2865	25	0.82	2.887	0.2254	20	1.14	2.952	0.2310	20	3.55	4.484	0.2325	20	2.47	4.698
374	0.0345	3	0.00	0.000	0.0225	2	0.00	0.000	0.0232	2	0.00	0.000	0.0229	2	0.11	0.148
375	0.0353	3	0.47	0.503	0.0330	3	0.48	0.826	0.0338	3	0.05	0.081	0.0349	3	0.00	0.000
376	0.1140	10	0.00	0.000	0.1125	10	0.65	1.987	0.1166	10	0.60	0.733	0.1174	10	0.76	0.963
377	0.0229	2	0.00	0.000	0.0225	2	1.25	1.768	0.0218	2	19.60	24.020	0.0233	2	61.19	64.955
378	0.0233	2	1.45	2.051	0.0225	2	19.18	19.141	0.0225	2	17.75	3.989	0.0225	2	8.59	10.087
379	0.0229	2	24.83	32.492	0.0229	2	4.35	0.481	0.0124	1	23.95	-	0.0236	2	5.70	7.078
380	0.0225	2	0.31	0.035	0.0229	2	1.09	0.976	0.0221	2	7.77	2.305	0.0229	2	27.53	24.784
381	0.0229	2	0.04	0.057	0.0229	2	0.00	0.000	0.0225	2	5.47	4.150	0.0233	2	3.63	3.765
382	0.0341	3	0.04	0.076	0.0454	4	0.00	0.000	0.0461	4	0.47	0.888	0.0458	4	0.97	0.639
721	0.0233	2	1.01	1.430	0.0225	2	9.40	13.287	0.0221	2	2.20	3.111	0.0229	2	0.00	0.000
722	0.0236	2	0.00	0.000	0.0221	2	1.73	2.447	0.0218	2	0.00	0.000	0.0233	2	0.00	0.000
723	0.0233	2	55.60	69.155	0.0229	2	0.65	0.919	0.0229	2	1.94	2.744	0.0233	2	0.00	0.000
724	0.0225	2	49.80	70.428	0.0225	2	10.46	14.786	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
725	0.0225	2	9.25	7.849	0.0229	2	2.17	3.062	0.0225	2	0.29	0.403	0.0236	2	1.47	2.073
726	0.0214	2	1122.95	1569.289	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0113	1	0.00	-
727	0.0233	2	2.80	3.960	0.0218	2	7.45	9.405	0.0232	2	0.00	0.000	0.0229	2	0.00	0.000
728	0.0229	2	21.40	30.264	0.0225	2	0.00	0.000	0.0180	2	0.00	0.000	0.0109	1	0.00	-
752	0.0116	1	0.00	0.000	0.0229	2	0.00	0.000	0.0214	2	0.00	0.000	0.0236	2	0.00	0.000
753	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
754	0.0341	3	0.00	0.000	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
755	0.0338	3	0.00	0.000	0.0221	2	0.00	0.000	0.0319	3	0.00	0.000	0.0450	4	0.00	0.000
756	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000	0.0233	2	0.00	0.000
757	0.0225	2	64.40	91.075	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
758	0.0225	2	2.80	3.960	0.0221	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
759	0.0225	2	0.00	0.000	0.0113	1	0.00	-	0.0214	2	0.00	0.000	0.0229	2	0.00	0.000
760	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
761	0.0225	2	0.17	0.236	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0221	2	0.00	0.000
762	0.0225	2	0.15	0.212	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000
763	0.0225	2	0.00	0.000	0.0311	3	0.00	0.000	0.0326	3	0.00	0.000	0.0334	3	0.00	0.000
764	0.0236	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000	0.0233	2	0.00	0.000
765	0.0236	2	0.00	0.000	0.0113	1	0.00	-	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
766	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
767	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0113	1	0.00	-

TABLE 2 (cont.).- Swept area, number of hauls and Atlantic cod mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1997-2009. Swept area in square miles. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Menduiña* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2006				2007				2008				2009			
	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD	Swept area	Tow number	A. cod Mean catch	A. cod SD
353	0.0371	3	11.53	7.341	0.0364	3	0.14	0.138	0.0341	3	0.04	0.070	0.0345	3	0.00	0.000
354	0.0364	3	10.98	14.032	0.0364	3	16.81	14.624	0.0345	3	64.76	69.913	0.0338	3	25.17	21.163
355	0.0248	2	3.04	0.078	0.0240	2	41.34	12.820	0.0221	2	2.30	3.253	0.0233	2	3.63	4.448
356	0.0240	2	3.88	3.247	0.0240	2	0.96	1.351	0.0236	2	13.45	13.011	0.0229	2	2.94	3.585
357	0.0244	2	12.75	8.400	0.0360	3	1.42	1.323	0.0233	2	6.31	8.917	0.0116	2	14.29	15.293
358	0.0349	3	82.54	80.442	0.0368	3	113.84	43.776	0.0345	3	249.58	302.829	0.0341	3	50.33	41.797
359	0.0975	8	372.36	643.214	0.0855	7	3.17	4.658	0.0799	7	224.94	196.538	0.0795	7	520.11	821.106
360	0.2340	19	7.35	8.119	0.2378	20	2.42	4.606	0.2340	20	10.10	14.465	0.2273	20	162.21	719.968
374	0.0236	2	0.00	0.000	0.0240	2	0.00	0.000	0.0233	2	0.57	0.812	0.0225	2	0.00	0.000
375	0.0364	3	13.53	15.862	0.0364	3	1.71	1.646	0.0334	3	18.64	29.958	0.0341	3	0.00	0.000
376	0.1219	10	6.84	11.380	0.1185	10	0.68	1.167	0.1129	10	11.60	9.917	0.1133	10	0.67	0.921
377	0.0236	2	90.62	69.919	0.0240	2	698.56	987.885	0.0233	2	234.80	189.646	0.0225	2	11.89	16.568
378	0.0240	2	90.32	85.680	0.0233	2	85.98	23.723	0.0240	2	213.40	239.992	0.0229	2	709.31	0.269
379	0.0236	2	6.30	8.627	0.0240	2	3.13	0.394	0.0229	2	2.26	1.965	0.0229	2	54.61	74.091
380	0.0229	2	8.70	1.697	0.0240	2	4.20	5.945	0.0225	2	21.80	11.738	0.0229	2	4.11	1.727
381	0.0229	2	8.43	1.167	0.0240	2	2.19	0.354	0.0229	2	4.49	6.242	0.0229	2	0.02	0.028
382	0.0469	4	0.75	1.033	0.0484	4	0.00	0.000	0.0458	4	0.13	0.167	0.0450	4	0.00	0.000
721	0.0236	2	0.00	0.000	0.0116	1	0.00	-	0.0225	2	1.24	1.747	0.0229	2	1.00	1.414
722	0.0240	2	0.00	0.000	0.0225	2	0.00	0.000	0.0206	2	0.00	0.000	0.0225	2	0.00	0.000
723	0.0236	2	0.00	0.000	0.0240	2	3.15	4.455	0.0225	2	5.54	7.835	0.0225	2	17.29	8.641
724	0.0233	2	0.00	0.000	0.0233	2	0.00	0.000	0.0221	2	0.00	0.000	0.0233	2	0.00	0.000
725	0.0233	2	0.00	0.000	0.0225	2	11.89	11.823	0.0229	2	0.61	0.863	0.0229	2	2.19	3.090
726	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	1.38	1.945
727	0.0225	2	0.00	0.000	0.0240	2	0.00	0.000	0.0221	2	0.39	0.554	0.0113	1	5.01	-
728	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
752	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0229	2	0.25	0.354
753	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0116	1	0.00	-
754	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0113	1	0.00	-
755	0.0338	3	0.00	0.000	0.0338	3	0.00	0.000	0.0431	4	0.00	0.000	0.0116	1	0.00	-
756	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
757	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
758	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
759	0.0225	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	0.0221	2	0.00	0.000	0.0113	1	0.00	-
760	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
761	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
762	0.0233	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.000	0.0225	2	0.00	0
763	0.0225	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	0.0311	3	0.00	0.000	n.s.	n.s.	n.s.	n.s.
764	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0116	1	0.00	-
765	0.0236	2	0.00	0.000	0.0225	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
766	0.0229	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
767	0.0233	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.

TABLE 3.- Stratified mean catches (Kg) by stratum and year and SD by year of Atlantic cod (1997-2009). n.s. means stratum not surveyed. 1997-2000 data are transformed C/V Playa de Mendumá data. 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
353	0.00	0.59	1684.29	2310.56	1972.67	0.40	0.00	2746.49	1129.80	3100.67	36.76	10.85	0.00
354	0.00	4347.10	1209.44	4536.47	3954.04	1.64	1877.80	1172.11	1662.39	2700.42	4134.28	15931.78	6190.92
355	317.46	2001.63	472.57	7017.36	4152.14	71.15	223.48	376.66	145.78	224.59	3058.79	170.20	268.25
356	366.75	292.75	1935.74	768.05	7031.20	714.40	733.44	139.36	66.98	182.17	44.89	632.15	137.95
357	15014.55	1222.35	1659.07	1500.68	4460.80	1090.60	865.10	2180.38	651.90	2091.00	232.33	1034.02	2344.22
358	397.76	1002.53	2246.51	41597.12	768.75	592.50	46625.25	3241.50	5119.50	18570.75	25614.00	56156.25	11325.00
359	473.87	164.50	3052.91	7687.04	74245.15	1146.52	435.31	12557.95	24128.71	156764.14	1334.99	94698.54	218966.47
360	301.58	616.24	6478.57	6017.33	31605.14	2283.17	3169.28	9886.61	6869.14	20449.63	6724.01	28119.71	451440.17
374	12.23	0.00	124.31	0.00	0.00	0.00	0.00	0.00	22.47	0.00	0.00	122.84	0.00
375	0.00	211.79	261.73	0.00	0.00	126.47	129.18	12.65	0.00	3665.73	464.22	5050.09	0.00
376	0.00	263.27	822.50	1202.94	50.03	0.00	864.70	801.87	1010.91	9129.90	911.39	15474.27	898.32
377	26.59	188.96	21.35	1.92	0.00	0.00	125.00	1959.50	6119.00	9062.00	69855.95	23480.00	1188.50
378	325.88	481.53	1078.58	1480.09	1665.22	201.55	2665.33	2466.56	1194.36	12553.79	11950.53	29662.60	98594.09
379	390.21	880.31	553.41	4358.29	1010.71	2631.45	461.10	2538.70	603.67	667.80	331.94	239.51	5788.66
380	34.94	223.39	3703.59	788.08	576.11	30.19	104.64	745.92	2642.40	835.20	403.58	2092.80	394.32
381	10.08	30.36	125.22	250.68	95.74	5.76	0.00	787.90	523.08	1213.20	315.36	646.06	2.88
382	0.00	108.42	18.00	243.65	41.41	14.98	0.00	160.78	332.28	255.54	0.00	45.96	0.00
721	1363.56	39.80	5738.57	1842.35	315.25	65.75	610.68	143.00	0.00	0.00	0.00	80.28	65.00
722	26.16	0.00	0.00	75.84	0.00	0.00	145.32	0.00	0.00	0.00	0.00	0.00	0.00
723	1534.94	680.69	2614.28	3413.20	104803.25	8618.00	100.75	300.70	0.00	0.00	488.25	858.70	2679.95
724	161.20	184615.64	2.82	87.21	764.25	6175.20	1296.42	0.00	0.00	0.00	0.00	0.00	0.00
725	2467.77	3240.64	1432.94	455.78	143598.88	971.25	227.33	29.93	153.93	0.00	1248.45	64.05	229.43
726	n.s.	341.39	58.07	637.55	132.02	80852.04	0.00	0.00	0.00	0.00	0.00	0.00	99.00
727	11.42	255.30	883.49	879.12	998.37	268.80	715.20	0.00	0.00	0.00	0.00	37.58	480.96
728	91.43	120.09	0.00	69.87	0.00	1669.20	0.00	0.00	0.00	0.00	0.00	0.00	0.00
752	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	32.75
753	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
754	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
755	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
756	0.00	32.07	23.86	36.40	4.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
757	0.00	0.00	0.00	0.00	0.00	6568.80	0.00	0.00	0.00	0.00	0.00	0.00	0.00
758	0.00	0.00	0.00	0.00	0.00	277.20	0.00	0.00	0.00	0.00	0.00	0.00	0.00
759	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.	0.00	0.00
760	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
761	0.00	0.00	0.00	0.00	0.00	28.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00
762	0.00	0.00	0.00	0.00	0.00	31.80	0.00	0.00	0.00	0.00	n.s.	0.00	0.00
763	n.s.	0.00	0.00	283.12	0.00	0.00	0.00	0.00	0.00	0.00	n.s.	0.00	n.s.
764	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
765	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
766	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.	0.00	0.00
767	n.s.	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	n.s.	0.00	n.s.
TOTAL	23328	201361	36202	87541	382245	114437	61375	42249	52376	241467	127150	274608	801127
Ȳ	2.50	19.47	3.50	8.46	36.96	11.07	5.93	4.09	5.06	23.35	13.47	26.55	80.73
S.D.	1.54	17.82	0.75	2.58	17.97	7.82	3.29	0.95	2.16	9.39	7.44	5.71	46.81

TABLE 4.- Survey estimates (by the swept area method) of Atlantic cod biomass (t) and SD by stratum and year on NAFO Div. 3NO. n.s. means stratum not surveyed. 1997-2000 data are transformed C/V *Playa de Menduiña* data. 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
353	0	0	140	195	173	0	0	244	96	251	3	1	0
354	0	366	111	382	351	0	167	102	141	223	341	1385	550
355	27	181	41	604	346	6	20	33	13	18	255	15	23
356	33	26	169	68	586	61	65	13	6	15	4	54	12
357	1357	102	140	121	366	91	76	191	56	172	19	89	403
358	35	86	194	3657	67	52	4144	295	440	1597	2091	4883	996
359	41	14	252	656	6476	100	39	1111	2076	12863	109	8299	21377
360	26	53	529	502	2609	199	281	856	591	1660	566	2403	39731
374	1	0	10	0	0	0	0	0	2	0	0	11	0
375	0	18	22	0	0	11	12	1	0	302	38	454	0
376	0	23	67	100	4	0	77	69	86	749	77	1371	79
377	2	17	2	0	0	0	11	180	526	767	5821	2020	106
378	31	41	95	127	141	17	237	219	106	1046	1028	2472	8620
379	38	74	47	387	88	230	40	205	51	57	28	21	506
380	3	20	314	67	56	3	9	67	231	73	34	186	34
381	1	3	11	21	8	1	0	70	45	106	26	56	0
382	0	10	1	20	4	1	0	14	29	22	0	4	0
721	123	4	471	156	25	6	54	13	0	0	0	7	6
722	2	0	0	7	0	0	13	0	0	0	0	0	0
723	146	59	229	276	8734	741	9	26	0	0	41	76	238
724	14	17902	0	8	65	549	115	0	0	0	0	0	0
725	239	376	125	43	12347	86	20	3	13	0	111	6	20
726	n.s.	33	5	58	11	7565	0	0	0	0	0	0	9
727	1	22	75	84	89	23	66	0	0	0	0	3	43
728	9	12	0	7	0	146	0	0	0	0	0	0	0
752	0	0	0	0	0	0	0	0	0	0	0	0	3
753	0	0	0	0	0	0	0	0	0	0	0	0	0
754	0	0	0	0	0	0	0	0	0	0	0	0	0
755	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
756	0	3	2	4	0	0	0	0	0	0	0	0	0
757	0	0	0	0	0	584	0	0	0	0	0	0	0
758	0	0	0	0	0	25	0	0	0	0	0	0	0
759	n.s.	0	0	0	0	0	0	0	0	0	n.s.	0	0
760	0	0	0	0	0	0	0	0	0	0	0	0	0
761	0	0	0	0	0	3	0	0	0	0	0	0	0
762	0	0	0	0	0	3	0	0	0	0	n.s.	0	0
763	n.s.	0	0	27	0	0	0	0	0	0	n.s.	0	n.s.
764	0	0	0	0	0	0	0	0	0	0	0	0	0
765	0	0	0	0	0	0	0	0	0	0	0	0	0
766	0	0	0	0	0	0	0	0	0	0	n.s.	0	0
767	n.s.	0	0	0	0	0	0	0	0	0	n.s.	0	n.s.
TOTAL	2131	19444	3054	7576	32548	10502	5455	3712	4509	19921	10592	23817	72757
S.D.	1322	18206	655	2566	15903	7971	3016	848	1984	8109	5853	5221	40466

TABLE 5.- Length weight relationships in the calculation of Atlantic cod biomass. The equation is $Weight = a(l + 0.5)^b$
 Spanish Spring Surveys on NAFO Div. 3NO: 1997-2009.

	1997	1998	1999	2000	2001	2002	2003
a	0.0102 Error = 0.2480	0.0061 Error = 0.0748	0.0048 Error = 0.0788	0.0060 Error = 0.0706	0.0048 Error = 0.0893	0.0057 Error = 0.1025	0.0046 Error = 0.0581
b	2.9387 Error = 0.0629	3.0671 Error = 0.0197	3.1313 Error = 0.0203	3.0822 Error = 0.0179	3.1198 Error = 0.0228	3.0783 Error = 0.0274	3.1370 Error = 0.0153
	R ² = 0.975 N = 431	R2 = 0.997 N = 687	R ² = 0.997 N = 430	R ² = 0.997 N = 877	R ² = 0.996 N = 488	R ² = 0.995 N = 678	R ² = 0.998 N = 516

	2004	2005	2006	2007	2008	2009
a	0.0052 Error = 0.0698	0.0052 Error = 0.0715	0.0058 Error = 0.0678	0.0059 Error = 0.0570	0.0047 Error = 0.0858	0.0052 Error = 0.0833
b	3.1107 Error = 0.0185	3.1238 Error = 0.0189	3.0965 Error = 0.0174	3.0762 Error = 0.0153	3.1341 Error = 0.0217	3.0937 Error = 0.0220
	R ² = 0.997 N = 656	R ² = 0.997 N = 612	R ² = 0.999 N = 1129	R2 = 0.998 N= 1011	R2 = 0.998 N= 1266	R2 = 0.996 N= 795

TABLE 6.- Atlantic cod length distribution per haul mean catches by sex and year. Number per stratified mean catches. Spanish Spring Survey on NAFO 3NO; 1997-2009. Indet. means indeterminate. 1997-2000 data are transformed C/V *Playa de Mendoña* data. 2002-2009 data are original R/V *Vizconde de Eza* data. (*) indicates untransformed data.

Length (cm.)	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
6	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.014	0.000	0.000	0.000
8	0.000	0.000	0.013	0.000	0.009	0.011	0.000	0.000	0.000	0.000	0.015	0.000	0.000
10	0.000	0.000	0.011	0.000	0.000	0.039	0.000	0.408	0.088	0.014	0.416	0.000	0.009
12	0.000	0.010	0.010	0.006	0.000	0.030	0.054	2.141	0.722	0.008	4.160	0.000	0.111
14	0.000	0.025	0.205	0.020	0.000	0.000	0.065	1.542	1.129	0.014	8.177	0.013	0.132
16	0.000	0.011	0.741	0.124	0.021	0.040	0.028	0.828	0.644	0.013	4.890	0.057	0.190
18	0.000	0.007	0.774	0.115	0.019	0.007	0.037	0.186	0.245	0.007	1.020	0.910	0.055
20	0.000	0.004	0.287	0.211	0.103	0.000	0.061	0.063	0.036	0.041	0.070	5.006	0.225
22	0.000	0.002	0.478	0.285	0.184	0.083	0.062	0.076	0.094	0.367	0.024	8.175	0.448
24	0.002	0.005	1.173	0.241	0.175	0.096	0.075	0.119	0.224	0.948	0.008	8.532	0.847
26	0.002	0.009	1.740	0.319	0.194	0.139	0.150	0.219	0.515	3.595	0.000	6.053	2.145
28	0.013	0.028	1.251	0.385	0.216	0.153	0.189	0.141	1.044	5.313	0.041	1.953	5.742
30	0.013	0.037	0.525	0.417	0.132	0.284	0.243	0.162	1.151	3.854	0.086	1.112	12.908
32	0.051	0.028	0.172	0.327	0.162	0.312	0.087	0.149	0.551	1.710	0.317	0.301	19.091
34	0.087	0.030	0.096	0.438	0.380	0.524	0.067	0.330	0.393	1.102	0.442	0.152	15.295
36	0.142	0.022	0.070	0.559	0.923	0.596	0.071	0.421	0.189	2.258	0.729	0.176	16.078
38	0.184	0.026	0.090	1.038	1.787	0.572	0.121	0.420	0.129	5.496	0.925	0.539	8.529
40	0.108	0.105	0.086	1.030	3.363	0.689	0.081	0.217	0.135	5.305	0.881	0.962	4.828
42	0.066	0.075	0.031	0.897	3.463	1.005	0.078	0.248	0.113	4.004	0.885	1.337	2.320
44	0.106	0.365	0.047	0.473	4.234	1.141	0.117	0.101	0.097	2.317	0.788	1.617	2.777
46	0.073	0.603	0.025	0.307	5.028	1.483	0.111	0.110	0.136	1.054	1.632	1.683	4.296
48	0.091	0.931	0.045	0.183	5.686	1.090	0.175	0.077	0.173	0.487	2.035	1.327	3.909
50	0.043	0.963	0.044	0.137	4.959	1.058	0.225	0.060	0.101	0.279	1.748	1.465	7.314
52	0.074	0.924	0.063	0.099	4.098	1.111	0.298	0.088	0.128	0.276	1.412	1.556	3.843
54	0.087	1.499	0.106	0.109	3.195	0.895	0.390	0.072	0.026	0.227	0.651	1.750	3.711
56	0.142	1.537	0.081	0.069	1.224	0.691	0.428	0.065	0.028	0.231	0.401	1.537	5.611
58	0.124	1.764	0.113	0.136	0.693	0.223	0.322	0.110	0.012	0.256	0.262	1.104	3.879
60	0.195	1.026	0.130	0.101	0.532	0.370	0.306	0.074	0.055	0.229	0.094	0.624	2.342
62	0.114	0.540	0.098	0.065	0.181	0.126	0.183	0.093	0.078	0.204	0.054	0.348	2.164
64	0.088	0.505	0.072	0.152	0.032	0.005	0.227	0.104	0.092	0.114	0.079	0.280	0.701
66	0.111	0.163	0.049	0.134	0.047	0.057	0.098	0.063	0.089	0.098	0.056	0.241	0.459
68	0.014	0.271	0.067	0.101	0.014	0.000	0.093	0.071	0.077	0.092	0.096	0.075	0.867
70	0.029	0.157	0.019	0.137	0.015	0.061	0.085	0.042	0.093	0.074	0.037	0.075	0.123
72	0.004	0.193	0.013	0.104	0.028	0.007	0.027	0.031	0.083	0.096	0.029	0.121	0.129
74	0.013	0.136	0.018	0.142	0.012	0.000	0.011	0.033	0.078	0.071	0.012	0.087	0.129
76	0.002	0.086	0.011	0.066	0.017	0.002	0.015	0.030	0.079	0.121	0.042	0.056	0.060
78	0.003	0.080	0.008	0.034	0.022	0.000	0.010	0.017	0.056	0.051	0.029	0.031	0.011
80	0.006	0.079	0.015	0.073	0.039	0.000	0.027	0.036	0.047	0.103	0.008	0.038	0.029
82	0.001	0.038	0.005	0.032	0.013	0.000	0.000	0.009	0.018	0.057	0.036	0.051	0.077
84	0.003	0.000	0.004	0.044	0.000	0.011	0.025	0.003	0.006	0.041	0.000	0.086	0.015
86	0.001	0.048	0.012	0.026	0.021	0.000	0.008	0.000	0.022	0.041	0.000	0.057	0.019
88	0.000	0.042	0.010	0.021	0.003	0.007	0.002	0.022	0.014	0.013	0.000	0.030	0.013
90	0.001	0.000	0.000	0.016	0.011	0.000	0.000	0.008	0.014	0.039	0.015	0.024	0.008
92	0.000	0.003	0.019	0.020	0.000	0.000	0.000	0.009	0.000	0.005	0.000	0.000	0.000
94	0.000	0.000	0.000	0.005	0.003	0.000	0.000	0.013	0.000	0.000	0.000	0.026	0.000
96	0.000	0.000	0.005	0.003	0.012	0.000	0.008	0.000	0.000	0.026	0.000	0.057	0.000
98	0.000	0.000	0.005	0.003	0.008	0.000	0.000	0.000	0.000	0.025	0.000	0.000	0.000
100	0.000	0.000	0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.020	0.008
102	0.000	0.000	0.000	0.010	0.000	0.000	0.008	0.000	0.000	0.014	0.000	0.019	0.000
104	0.000	0.001	0.000	0.000	0.000	0.011	0.000	0.027	0.000	0.014	0.000	0.026	0.000
106	0.000	0.000	0.000	0.005	0.000	0.000	0.014	0.000	0.000	0.000	0.000	0.013	0.000
108	0.000	0.000	0.000	0.000	0.011	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
110	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
112	0.000	0.000	0.000	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
114	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.004	0.000	0.000
116	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
118	0.002	0.000	0.003	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.016	0.000
120	0.000	0.000	0.006	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
122	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
124	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
126	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
128	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
130	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
132	0.000	0.000	0.000	0.000	0.009	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	1.997	12.378	8.847	9.220	41.290	12.930	4.684	9.035	9.005	40.718	32.605	49.717	131.444
Nº samples (*):	40	55	72	70	32	41	42	58	59	64	58	66	55
Nº Ind. (*):	742	967	2770	2753	1591	1030	539	939	1126	2909	2301	4404	2746
Sampled catch:	248	410	527	752	1107	776	654	554	778	2026	1115	3394	1417
Range (*):	24-118	12-104	9-121	13-118	8-132	9-104	12-106	10-105	11-91	7-104	9-114	14-118	11-100
Total catch:	572	3873	613	1274	3487	2806	846	554	794	3994	2182	3907	9165
Total hauls (*):	128	124	114	118	123	125	118	120	119	120	110	122	109

TABLE 7.- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2009. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduiña* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1995				1996				1997				1998			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0353	3	5.82	4.105	0.0371	3	74.88	94.62	0.0480	4	12.55	14.26	0.0465	4	12.22	20.16
354	0.0353	3	1.78	3.089	0.0319	3	1.11	0.84	0.0233	2	1.41	1.56	0.0356	3	1.22	0.24
355	n.s.	n.s.	n.s.	n.s.	0.0221	2	0.25	0.35	0.0233	2	2.20	0.31	0.0221	2	0.13	0.18
356	n.s.	n.s.	n.s.	n.s.	0.0203	2	0.00	0.00	0.0225	2	0.32	0.46	0.0221	2	0.00	0.00
357	0.0109	1	0.00	-	0.0218	2	0.00	0.00	0.0443	4	0.00	0.00	0.0240	2	0.00	0.00
358	0.0319	3	0.00	0.000	0.0319	3	0.13	0.23	0.0563	5	0.02	0.04	0.0236	3	0.00	0.00
359	0.0345	3	1.35	2.336	0.0548	5	0.92	0.83	0.0690	6	0.08	0.14	0.0698	6	0.17	0.22
360	0.3563	31	20.44	40.707	0.3761	31	142.09	128.86	0.3754	32	80.92	155.59	0.2561	25	373.90	629.84
374	0.0225	2	0.00	0.000	0.0233	2	0.00	0.00	0.0353	3	0.00	0.00	0.0353	3	0.04	0.02
375	0.0225	2	1.48	1.875	0.0229	2	41.40	58.54	0.0116	1	0.20	-	0.0345	3	12.37	21.37
376	0.1729	15	35.06	58.691	0.1650	14	71.40	86.94	0.1583	14	162.35	179.83	0.0930	10	279.27	181.29
377	0.0221	2	0.00	0.000	0.0229	2	0.00	0.00	0.0116	1	0.00	-	0.0229	2	0.00	0.00
378	0.0435	4	0.00	0.000	0.0330	3	0.06	0.10	0.0210	2	0.00	0.00	0.0120	2	0.00	0.00
379	0.0221	2	0.00	0.000	0.0113	1	0.00	-	0.0206	2	0.00	0.00	0.0356	3	0.00	0.00
380	n.s.	n.s.	n.s.	n.s.	0.0221	2	0.00	0.00	0.0210	2	0.00	0.00	0.0113	2	0.00	0.00
381	n.s.	n.s.	n.s.	n.s.	0.0229	2	0.00	0.00	0.0221	2	0.00	0.00	0.0229	2	0.00	0.00
382	n.s.	n.s.	n.s.	n.s.	0.0338	3	0.00	0.00	0.0461	4	0.00	0.00	0.0229	3	0.00	0.00
721	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.03	0.05	0.0221	2	0.75	1.06	0.0203	2	0.00	0.00
722	n.s.	n.s.	n.s.	n.s.	0.0206	2	0.00	0.00	0.0214	2	0.00	0.00	0.0101	2	0.00	0.00
723	n.s.	n.s.	n.s.	n.s.	0.0109	1	0.00	-	0.0210	2	0.00	0.00	0.0233	2	0.00	0.00
724	0.0105	1	0.00	-	0.0203	2	0.00	0.00	0.0225	2	0.00	0.00	0.0206	2	0.00	0.00
725	0.0334	3	0.00	0.000	0.0225	2	0.00	0.00	0.0206	2	0.00	0.00	0.0086	1	0.00	-
726	0.0214	2	0.00	0.000	0.0218	2	0.00	0.00	n.s.	n.s.	n.s.	n.s.	0.0094	2	0.00	0.00
727	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0094	1	0.00	-	0.0233	2	0.00	0.00
728	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.00	0.0214	2	0.00	0.00	0.0206	2	0.00	0.00
752	n.s.	n.s.	n.s.	n.s.	0.0109	1	0.00	-	0.0218	2	0.00	0.00	0.0229	2	0.00	0.00
753	n.s.	n.s.	n.s.	n.s.	0.0199	2	0.00	0.00	0.0214	2	0.00	0.00	0.0218	2	0.00	0.00
754	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0330	3	0.00	0.00	0.0210	2	0.00	0.00
755	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0206	2	0.00	0.00
756	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0109	1	0.00	-	0.0225	2	0.00	0.00
757	n.s.	n.s.	n.s.	n.s.	0.0188	2	0.00	0.00	0.0304	3	0.00	0.00	0.0206	2	0.00	0.00
758	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.00	0.0105	2	0.00	0.00
759	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.00
760	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0105	1	0.00	-	0.0214	2	0.00	0.00
761	n.s.	n.s.	n.s.	n.s.	0.0199	2	0.00	0.00	0.0315	3	0.00	0.00	0.0206	2	0.00	0.00
762	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0308	3	0.00	0.00	0.0094	2	0.00	0.00
763	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.00
764	n.s.	n.s.	n.s.	n.s.	0.0210	2	0.00	0.00	0.0206	2	0.00	0.00	0.0218	2	0.00	0.00
765	n.s.	n.s.	n.s.	n.s.	0.0199	2	0.00	0.00	0.0206	2	0.00	0.00	0.0098	2	0.00	0.00
766	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0308	3	0.00	0.00	0.0191	2	0.00	0.00
767	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.0109	2	0.00	0.00

TABLE 7 (cont.).- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2009. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Mendoña* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1999				2000				2001				2002			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0360	3	150.18	182.44	0.0356	3	67.87	91.37	0.0341	3	61.42	102.797	0.0476	4	75.13	88.259
354	0.0218	2	0.08	0.12	0.0356	3	1.79	1.93	0.0338	3	0.34	0.322	0.0356	3	0.17	0.289
355	0.0229	2	0.00	0.00	0.0233	2	0.00	0.00	0.0240	2	0.00	0.000	0.0236	2	0.00	0.000
356	0.0229	2	0.00	0.00	0.0225	2	0.00	0.00	0.0240	2	0.01	0.007	0.0233	2	0.00	0.000
357	0.0236	2	0.00	0.00	0.0124	1	0.00	-	0.0244	2	0.00	0.000	0.0240	2	0.00	0.000
358	0.0349	3	0.00	0.00	0.0341	3	0.00	0.00	0.0345	3	0.00	0.000	0.0345	3	0.00	0.000
359	0.0364	3	0.34	0.47	0.0469	4	2.36	2.93	0.0803	7	1.42	2.836	0.0686	6	0.11	0.261
360	0.2325	19	545.18	424.37	0.2396	20	391.18	331.64	0.2423	20	536.80	488.657	0.2865	25	340.23	356.687
374	0.0244	2	74.16	103.18	0.0240	2	20.47	23.55	0.0240	2	238.75	111.369	0.0345	3	32.04	52.542
375	0.0236	2	347.15	168.25	0.0244	2	153.36	2.06	0.0338	3	100.33	68.319	0.0353	3	48.61	68.927
376	0.1219	10	551.60	165.61	0.1200	10	435.27	236.60	0.1155	10	443.12	196.619	0.1140	10	533.62	416.745
377	0.0240	2	0.00	0.00	0.0229	2	0.05	0.06	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
378	0.0229	2	0.00	0.00	0.0233	2	0.00	0.00	0.0236	2	0.00	0.000	0.0233	2	0.00	0.000
379	0.0236	2	0.00	0.00	0.0225	2	0.00	0.00	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
380	0.0236	2	0.00	0.00	0.0236	2	0.00	0.00	0.0206	2	0.00	0.000	0.0225	2	0.00	0.000
381	0.0229	2	0.00	0.00	0.0236	2	0.00	0.00	0.0236	2	0.00	0.000	0.0229	2	0.00	0.000
382	0.0484	4	0.00	0.00	0.0499	4	0.00	0.00	0.0469	4	0.02	0.030	0.0341	3	0.00	0.000
721	0.0244	2	0.00	0.00	0.0236	2	0.00	0.00	0.0248	2	0.00	0.000	0.0233	2	0.00	0.000
722	0.0229	2	0.00	0.00	0.0218	2	0.00	0.00	0.0233	2	0.00	0.000	0.0236	2	0.00	0.000
723	0.0229	2	0.00	0.00	0.0248	2	0.00	0.00	0.0240	2	0.00	0.000	0.0233	2	0.00	0.000
724	0.0225	2	0.00	0.00	0.0233	2	0.00	0.00	0.0353	3	0.00	0.000	0.0225	2	0.00	0.000
725	0.0229	2	0.00	0.00	0.0210	2	0.00	0.00	0.0116	2	0.00	0.000	0.0225	2	0.00	0.000
726	0.0225	2	0.00	0.00	0.0221	2	0.00	0.00	0.0116	2	0.00	0.000	0.0214	2	0.00	0.000
727	0.0236	2	0.00	0.00	0.0210	2	0.00	0.00	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000
728	0.0233	2	0.00	0.00	0.0210	2	0.00	0.00	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
752	0.0233	2	0.00	0.00	0.0206	2	0.00	0.00	0.0210	2	0.06	0.083	0.0116	1	0.00	-
753	0.0229	2	0.00	0.00	0.0218	2	0.00	0.00	0.0214	2	0.00	0.000	0.0229	2	0.00	0.000
754	0.0206	2	0.00	0.00	0.0195	2	0.00	0.00	0.0195	2	0.00	0.000	0.0341	3	0.00	0.000
755	0.0311	3	0.00	0.00	0.0431	4	0.00	0.00	0.0416	4	0.00	0.000	0.0338	3	0.00	0.000
756	0.0225	2	0.00	0.00	0.0203	2	0.00	0.00	0.0113	2	0.00	0.000	0.0229	2	0.00	0.000
757	0.0233	2	0.00	0.00	0.0214	2	0.00	0.00	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000
758	0.0214	2	0.00	0.00	0.0210	2	0.00	0.00	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
759	0.0218	2	0.00	0.00	0.0210	2	0.00	0.00	0.0221	2	0.00	0.000	0.0225	2	0.00	0.000
760	0.0225	2	0.00	0.00	0.0210	2	0.00	0.00	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
761	0.0210	2	0.00	0.00	0.0221	2	0.00	0.00	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
762	0.0210	2	0.00	0.00	0.0203	2	0.00	0.00	0.0116	2	0.00	0.000	0.0225	2	0.00	0.000
763	0.0311	3	0.00	0.00	0.0416	4	0.00	0.00	0.0330	3	0.00	0.000	0.0225	2	0.00	0.000
764	0.0225	2	0.00	0.00	0.0218	2	0.00	0.00	0.0240	2	0.00	0.000	0.0236	2	0.00	0.000
765	0.0221	2	0.00	0.00	0.0203	2	0.00	0.00	0.0113	2	0.00	0.000	0.0236	2	0.00	0.000
766	0.0218	2	0.00	0.00	0.0214	2	0.00	0.00	0.0203	2	0.00	0.000	0.0233	2	0.00	0.000
767	0.0214	2	0.00	0.00	0.0210	2	0.00	0.00	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000

TABLE 7 (cont.).- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2009. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Mendumá* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2003				2004				2005				2006			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0334	3	11.15	19.307	0.0338	3	8.79	14.005	0.0353	3	58.83	99.610	0.0371	3	71.98	122.954
354	0.0338	3	0.00	0.000	0.0345	3	0.62	1.065	0.0353	3	0.21	0.188	0.0364	3	0.21	0.371
355	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0248	2	0.00	0.000
356	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0233	2	0.00	0.000	0.0240	2	0.00	0.000
357	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0233	2	0.00	0.000	0.0244	2	0.00	0.000
358	0.0338	3	0.00	0.000	0.0330	3	0.26	0.442	0.0349	3	0.00	0.000	0.0349	3	0.00	0.000
359	0.0791	7	0.00	0.000	0.0791	7	25.01	38.371	0.0814	7	99.52	142.727	0.0975	8	169.33	359.779
360	0.2254	20	360.55	298.992	0.2310	20	403.19	333.463	0.2325	20	342.14	223.566	0.2340	19	361.02	266.205
374	0.0225	2	16.13	8.238	0.0233	2	193.46	225.058	0.0229	2	300.46	128.092	0.0236	2	610.03	73.518
375	0.0330	3	28.45	35.557	0.0338	3	543.04	155.015	0.0349	3	288.64	138.290	0.0364	3	287.65	109.715
376	0.1125	10	391.60	257.289	0.1166	10	481.06	140.810	0.1174	10	500.53	238.908	0.1219	10	489.81	231.495
377	0.0225	2	0.70	0.990	0.0218	2	0.00	0.000	0.0233	2	42.84	60.518	0.0236	2	6.09	8.605
378	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0240	2	0.00	0.000
379	0.0229	2	0.00	0.000	0.0124	1	0.00	-	0.0236	2	0.00	0.000	0.0236	2	0.00	0.000
380	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
381	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000	0.0229	2	0.00	0.000
382	0.0454	4	0.00	0.000	0.0461	4	0.00	0.000	0.0458	4	0.00	0.000	0.0469	4	0.00	0.000
721	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000	0.0236	2	0.00	0.000
722	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000	0.0233	2	0.00	0.000	0.0240	2	0.00	0.000
723	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000	0.0233	2	0.00	0.000	0.0236	2	0.18	0.247
724	0.0225	2	0.52	0.735	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000
725	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0236	2	0.00	0.000	0.0233	2	0.00	0.000
726	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0113	1	0.00	-	0.0225	2	0.00	0.000
727	0.0218	2	0.00	0.000	0.0233	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000
728	0.0225	2	0.00	0.000	0.0180	2	0.00	0.000	0.0109	1	0.00	-	0.0225	2	0.00	0.000
752	0.0229	2	0.00	0.000	0.0214	2	0.00	0.000	0.0236	2	0.00	0.000	0.0225	2	0.00	0.000
753	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
754	0.0218	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
755	0.0221	2	0.00	0.000	0.0319	3	0.00	0.000	0.0450	4	0.00	0.000	0.0338	3	0.00	0.000
756	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000	0.0233	2	0.00	0.000	0.0229	2	0.00	0.000
757	0.0221	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
758	0.0221	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
759	0.0113	1	0.00	-	0.0214	2	0.00	0.000	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000
760	0.0218	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.35	0.488	0.0225	2	0.00	0.000
761	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0221	2	0.00	0.000	0.0233	2	0.00	0.000
762	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0233	2	0.00	0.000
763	0.0311	3	0.00	0.000	0.0326	3	0.00	0.000	0.0334	3	0.00	0.000	0.0225	2	0.00	0.000
764	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000	0.0233	2	0.00	0.000	0.0233	2	0.00	0.000
765	0.0113	1	0.00	-	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0236	2	0.00	0.000
766	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
767	0.0229	2	0.00	0.000	0.0218	2	0.00	0.000	0.0113	1	0.00	-	0.0233	2	0.00	0.000

TABLE 7 (cont.).- Swept area, number of hauls and Yellowtail flounder mean catch (kg) and SD by stratum. Spanish Spring Surveys on NAFO Div. 3NO: 1995-2009. Swept area in square miles. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Mendumá* data, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	2007				2008				2009			
	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD	Swept area	Tow number	Y. flounder Mean catch	Y. flounder SD
353	0.0364	3	0.64	0.172	0.0341	3	18.63	30.202	0.0345	3	0.15	0.259
354	0.0364	3	0.16	0.283	0.0345	3	1.03	0.775	0.0338	3	0.00	0.000
355	0.0240	2	0.00	0.000	0.0221	2	0.00	0.000	0.0233	2	0.00	0.000
356	0.0240	2	0.00	0.000	0.0236	2	0.00	0.000	0.0229	2	0.00	0.000
357	0.0360	3	0.00	0.000	0.0233	2	0.00	0.000	0.0116	2	0.00	0.000
358	0.0368	3	0.00	0.000	0.0345	3	0.00	0.000	0.0341	3	0.00	0.000
359	0.0855	7	102.63	116.690	0.0799	7	26.40	38.865	0.0795	7	11.16	31.077
360	0.2378	20	349.70	307.902	0.2340	20	339.09	220.066	0.2273	20	358.38	377.704
374	0.0240	2	1057.60	455.094	0.0233	2	696.25	157.331	0.0225	2	1392.90	938.048
375	0.0364	3	145.73	86.977	0.0334	3	574.00	461.113	0.0341	3	335.84	149.894
376	0.1185	10	460.24	203.990	0.1129	10	421.05	280.644	0.1133	10	514.96	250.661
377	0.0240	2	165.35	233.840	0.0233	2	173.40	8.202	0.0225	2	0.12	0.163
378	0.0233	2	0.00	0.000	0.0240	2	0.00	0.000	0.0229	2	0.00	0.000
379	0.0240	2	0.00	0.000	0.0229	2	0.05	0.067	0.0229	2	0.00	0.000
380	0.0240	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
381	0.0240	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
382	0.0484	4	0.00	0.000	0.0458	4	0.00	0.000	0.0450	4	0.00	0.000
721	0.0116	1	0.00	-	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
722	0.0225	2	0.00	0.000	0.0206	2	0.00	0.000	0.0225	2	0.00	0.000
723	0.0240	2	0.00	0.000	0.0225	2	0.00	0.000	0.0225	2	0.00	0.000
724	0.0233	2	0.00	0.000	0.0221	2	0.00	0.000	0.0233	2	0.00	0.000
725	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000	0.0229	2	0.00	0.000
726	0.0229	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
727	0.0240	2	0.00	0.000	0.0221	2	0.00	0.000	0.0113	1	0.00	-
728	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
752	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0229	2	0.00	0.000
753	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0116	1	0.00	-
754	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0113	1	0.00	-
755	0.0338	3	0.00	0.000	0.0431	4	0.00	0.000	0.0116	1	0.00	-
756	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
757	0.0229	2	0.00	0.000	0.0221	2	0.00	0.000	0.0229	2	0.00	0.000
758	0.0225	2	0.00	0.000	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
759	n.s.	n.s.	n.s.	n.s.	0.0221	2	0.00	0.000	0.0113	1	0.00	-
760	0.0233	2	0.00	0.000	0.0225	2	0.00	0.000	0.0229	2	0.00	0.000
761	0.0225	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
762	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.000	0.0225	2	0.00	0.0000
763	n.s.	n.s.	n.s.	n.s.	0.0311	3	0.00	0.000	n.s.	n.s.	n.s.	n.s.
764	0.0225	2	0.00	0.000	0.0221	2	0.00	0.000	0.0116	1	0.00	-
765	0.0225	2	0.00	0.000	0.0214	2	0.00	0.000	0.0225	2	0.00	0.000
766	n.s.	n.s.	n.s.	n.s.	0.0218	2	0.00	0.000	0.0225	2	0.00	0.000
767	n.s.	n.s.	n.s.	n.s.	0.0214	2	0.00	0.000	n.s.	n.s.	n.s.	n.s.

TABLE 8.- Stratified mean catches (Kg) by stratum and year and SD by year of Yellowtail flounder (1995-2009). n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
353	1565	20142	3377	3288	40399	18256	16521	20209	2998	2365	15825	19364	173	5011	40
354	439	0	346	299	21	440	84	41	0	151	52	53	40	253	0
355	n.s.	0	163	9	0	0	0	0	0	0	0	0	0	0	0
356	n.s.	0	15	0	0	0	0	0	0	0	0	0	0	0	0
357	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
358	0	30	4	0	0	0	0	0	0	57	0	0	0	0	0
359	568	386	34	73	143	995	598	45	0	10528	41896	71290	43209	11113	4700
360	56885	395449	225203	1040562	1517233	1088648	1493909	946848	1003413	1122078	952164	1004708	973222	943689	997385
374	0	0	0	10	15871	4380	51093	6857	3451	41400	64297	130545	226326	148998	298081
375	402	11218	54	3353	94077	41561	27190	13173	7710	147165	78221	77953	39494	155554	91013
376	46775	95247	216576	372549	735836	580654	591126	711849	522389	641737	667712	653413	613960	561677	686958
377	0	0	0	0	0	5	0	0	70	0	4284	609	16535	17340	12
378	0	8	0	0	0	0	0	0	0	0	0	0	0	0	0
379	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0
380	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
381	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
382	n.s.	0	0	0	0	0	5	0	0	0	0	0	0	0	0
721	n.s.	2	49	0	0	0	0	0	0	0	0	0	0	0	0
722	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
723	n.s.	0	0	0	0	0	0	0	0	0	0	0	27	0	0
724	0	0	0	0	0	0	0	0	64	0	0	0	0	0	0
725	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
726	0	0	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
727	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
728	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
752	n.s.	0	0	0	0	0	8	0	0	0	0	0	0	0	0
753	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
754	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0
755	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
756	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
757	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
758	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0
759	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.	0
760	n.s.	0	0	0	0	0	0	0	0	0	0	53	0	0	0
761	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
762	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	n.s.	0
763	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.	n.s.
764	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
765	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
766	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	n.s.	0
767	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.	0
TOTAL	106633	522481	445822	1420143	2403580	1734937	2180533	1699022	1540096	1965481	1824505	1957961	1912960	1843639	2078188
\bar{Y}	16.22	59.54	47.74	137.32	232.41	167.76	210.84	164.28	148.92	190.05	176.42	189.32	202.64	178.27	209.43
S.D.	4.37	8.41	10.69	34.70	27.41	22.21	30.58	24.92	20.84	21.27	17.06	19.83	23.61	19.00	29.75

TABLE 9.- Survey estimates (by the swept area method) of Yellowtail flounder biomass (t) and SD by stratum and year on NAFO Div. 3NO. n.s. means stratum not surveyed. 1995-2000 data are transformed C/V *Playa de Menduña* data. 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.

Stratum	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
353	133	1628	281	282	3367	1537	1452	1697	270	210	1347	1565	14	440	4
354	37	26	30	25	2	37	7	3	0	13	4	4	3	22	0
355	n.s.	2	14	0	0	0	0	0	0	0	0	0	0	0	0
356	n.s.	0	1	0	0	0	0	0	0	0	0	0	0	0	0
357	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
358	0	3	0	0	0	0	0	0	0	5	0	0	0	0	0
359	49	35	3	6	12	85	52	4	0	931	3604	5849	3538	974	473
360	4950	32593	19198	89742	123989	90863	123341	82622	89057	97150	81907	81579	81869	80657	87779
374	0	0	0	0	1302	365	4258	596	307	3561	5622	11051	18861	12817	26496
375	36	981	5	291	7964	3410	2417	1121	701	13081	6729	6429	3257	13982	8001
376	4059	8082	19160	32255	60376	48388	51175	62443	46435	55026	56887	53613	51811	49761	60659
377	0	0	0	0	0	0	0	0	6	0	368	52	1378	1492	1
378	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
379	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
380	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
381	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
382	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
721	n.s.	0	4	0	0	0	0	0	0	0	0	0	0	0	0
722	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
723	n.s.	0	0	0	0	0	0	0	0	0	0	0	2	0	0
724	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
725	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
726	0	0	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
727	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
728	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
752	n.s.	0	0	0	0	0	1	0	0	0	0	0	0	0	0
753	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
754	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0
755	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	0
756	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
757	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
758	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0
759	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.	0
760	n.s.	0	0	0	0	0	0	0	0	0	0	5	0	0	0
761	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
762	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	n.s.	0
763	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.	0
764	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
765	n.s.	0	0	0	0	0	0	0	0	0	0	0	0	0	0
766	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	0	n.s.	0
767	n.s.	n.s.	n.s.	0	0	0	0	0	0	0	0	0	0	n.s.	0
TOTAL	9264	43349	38697	122601	197012	144685	182704	148487	136775	169978	156472	160145	160731	160146	183412
S.D.	2484	6032	8527	31359	22938	19097	25847	23368	19287	18869	15271	16458	18852	17297	25736

TABLE 10.- Length weight relationships in the calculation of Yellowtail flounder biomass. The equation is $Weight = a(l + 0.5)^b$
 Spanish Spring Surveys on NAFO Div. 3NO: 1995-2009. To calculate the parameters for the indeterminate individuals,
 we used the total data (males + females + indeterminate individuals). *E* means Error.

		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Males	a	0.0079 E = 0.2653	0.0080 E = 0.0907	0.0081 E = 0.0936	0.0075 E = 0.1034	0.0084 E = 0.2119	0.0036 E = 0.0994	0.0081 E = 0.1248	0.0075 E = 0.0729	0.0121 E = 0.1109	0.0053 E = 0.1352	0.0027 E = 0.0882	0.0096 E = 0.0825	0.0074 E = 0.0655	0.0085 E = 0.1149	0.0051 E = 0.1710
	b	3.0416 E = 0.0799	3.0342 E = 0.0269	3.0197 E = 0.0281	3.0376 E = 0.0313	3.0098 E = 0.0610	3.2403 E = 0.0300	3.0176 E = 0.0374	3.0271 E = 0.0226	2.8978 E = 0.0348	3.1236 E = 0.0419	3.3274 E = 0.0274	2.9463 E = 0.0263	3.0190 E = 0.0201	2.9716 E = 0.0353	3.1109 E = 0.0519
		R2 = 0.984 N=137	R2 = 0.998 N=430	R2 = 0.997 N=556	R2 = 0.997 N=523	R2 = 0.994 N=56	R2 = 0.997 N=270	R2 = 0.995 N=271	R2 = 0.998 N=274	R2 = 0.995 N=316	R2 = 0.995 N=411	R2 = 0.997 N=311	R2 = 0.999 N=371	R2 = 0.999 N=578	R2 = 0.998 N=479	R2 = 0.993 N= 270
Females	a	0.0063 E = 0.1251	0.0056 E = 0.0632	0.0056 E = 0.0517	0.0067 E = 0.1290	0.0073 E = 0.2607	0.0026 E = 0.0914	0.006 E = 0.0841	0.0051 E = 0.0901	0.0061 E = 0.0995	0.0047 E = 0.0630	0.0027 E = 0.0634	0.0069 E = 0.1137	0.0043 E = 0.1973	0.0060 E = 0.0801	0.0066 E = 0.1594
	b	3.1083 E = 0.0367	3.1496 E = 0.0179	3.1382 E = 0.0152	3.0788 E = 0.0384	3.0577 E = 0.0739	3.3504 E = 0.0267	3.1122 E = 0.0249	3.1448 E = 0.0274	3.1079 E = 0.0307	3.1768 E = 0.0191	3.329 E = 0.0177	3.0584 E = 0.0347	3.1915 E = 0.0582	3.0850 E = 0.0237	3.0549 E = 0.0464
		R2 = 0.995 N=246	R2 = 0.999 N=735	R2 = 0.999 N=910	R2 = 0.994 N=682	R2 = 0.989 N=62	R2 = 0.998 N=344	R2 = 0.997 N=378	R2 = 0.997 N=343	R2 = 0.996 N=513	R2 = 0.999 N=547	R2 = 0.998 N=569	R2 = 0.997 N=507	R2 = 0.987 N= 731	R2 = 0.999 N= 594	R2 = 0.991 N= 378
Indet.	a	0.0088 E = 0.1109	0.006 E = 0.0656	0.006 E = 0.0580	0.0071 E = 0.0652	0.0078 E = 0.1656	0.0026 E = 0.0835	0.0092 E = 0.1075	0.006 E = 0.0402	0.0069 E = 0.1095	0.004 E = 0.0608	0.0025 E = 0.0523	0.0102 E = 0.1453	0.0068 E = 0.1078	0.0065 E = 0.0785	0.0067 E = 0.1293
	b	3.0144 E = 0.0330	3.1285 E = 0.0188	3.1166 E = 0.0171	3.0614 E = 0.0195	3.0406 E = 0.0477	3.3423 E = 0.0245	2.9883 E = 0.0329	3.0977 E = 0.0123	3.0737 E = 0.0337	3.2137 E = 0.0186	3.3552 E = 0.0148	2.9471 E = 0.0448	3.0606 E = 0.0327	3.0642 E = 0.0233	3.0502 E = 0.0379
		R2 = 0.996 N=391	R2 = 0.999 N=1181	R2 = 0.999 N=1466	R2 = 0.994 N=1211	R2 = 0.995 N=118	R2 = 0.999 N=614	R2 = 0.994 N=703	R2 = 0.999 N=620	R2 = 0.995 N=833	R2 = 0.999 N=969	R2 = 0.999 N=884	R2 = 0.995 N=887	R2 = 0.995 N=1312	R2 = 0.999 N= 1074	R2 = 0.994 N= 648

TABLE 11.- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2009. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Menduíña* data. 2002-2009 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	1995				1996				1997				1998				
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total	
4	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
6	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
8	0.000	0.000	0.185	0.185	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
10	0.000	0.000	0.456	0.456	0.000	0.000	0.498	0.498	0.000	0.000	0.000	0.000	0.000	0.000	0.071	0.071	
12	0.103	0.870	2.350	3.323	0.000	0.000	0.877	0.877	1.356	0.560	0.000	1.916	0.000	0.000	1.538	1.538	
14	1.557	1.441	2.842	5.840	0.000	0.048	2.711	2.759	0.155	0.819	0.000	0.974	0.121	0.157	0.000	0.278	
16	2.045	3.581	0.277	5.903	0.288	3.152	5.167	8.607	2.947	1.811	0.000	4.758	1.500	1.535	0.000	3.034	
18	2.649	3.358	0.031	6.038	2.334	15.279	3.167	20.780	5.076	4.415	0.000	9.491	8.365	5.129	0.000	13.495	
20	2.984	3.212	0.000	6.196	5.319	26.981	0.750	33.050	13.857	15.055	0.000	28.912	8.974	10.166	0.000	19.140	
22	4.807	6.015	0.000	10.823	8.522	32.231	0.065	40.818	28.296	23.048	0.000	51.345	25.957	20.452	0.000	46.409	
24	4.810	6.082	0.000	10.892	10.962	32.203	0.000	43.165	31.348	27.786	0.000	59.134	44.950	37.421	0.000	82.371	
26	2.340	2.446	0.000	4.786	9.552	16.875	0.000	26.427	24.015	26.970	0.000	50.985	72.376	60.520	0.000	132.896	
28	2.704	2.544	0.000	5.248	9.151	11.591	0.000	20.742	13.921	21.248	0.000	35.169	57.459	62.401	0.000	119.861	
30	2.588	4.738	0.000	7.325	7.206	9.915	0.000	17.122	6.159	10.349	0.000	16.508	32.472	56.275	0.000	88.747	
32	1.664	4.451	0.000	6.115	6.379	6.166	0.000	12.545	3.761	5.090	0.000	8.851	15.566	32.294	0.000	47.859	
34	1.290	3.070	0.000	4.361	5.565	6.928	0.000	12.493	1.894	2.803	0.000	4.698	5.840	22.613	0.000	28.453	
36	0.661	1.797	0.000	2.459	4.143	9.508	0.000	13.651	1.195	2.683	0.000	3.878	2.638	12.385	0.000	15.023	
38	0.475	1.395	0.000	1.870	2.083	6.687	0.000	8.771	0.485	2.407	0.000	2.892	2.475	8.439	0.000	10.914	
40	0.373	0.937	0.000	1.310	0.724	5.018	0.000	5.742	0.245	1.723	0.000	1.968	1.060	7.705	0.000	8.765	
42	0.059	0.588	0.000	0.647	0.694	3.305	0.000	4.000	0.099	0.801	0.000	0.899	0.065	3.260	0.000	3.324	
44	0.004	0.471	0.000	0.475	0.087	1.550	0.000	1.637	0.031	0.281	0.000	0.311	0.008	1.729	0.000	1.737	
46	0.004	0.081	0.000	0.085	0.081	0.969	0.000	1.050	0.006	0.044	0.000	0.049	0.000	0.600	0.000	0.600	
48	0.000	0.191	0.000	0.191	0.018	0.286	0.000	0.304	0.000	0.052	0.000	0.052	0.004	0.273	0.000	0.277	
50	0.000	0.027	0.000	0.027	0.000	0.045	0.000	0.045	0.000	0.018	0.000	0.018	0.000	0.000	0.000	0.000	
52	0.000	0.052	0.000	0.052	0.000	0.053	0.000	0.053	0.000	0.018	0.000	0.018	0.000	0.000	0.000	0.000	
54	0.000	0.005	0.000	0.005	0.000	0.039	0.000	0.039	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
56	0.000	0.005	0.000	0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
Total	31.117	47.358	6.141	84.616	73.109	188.829	13.235	275.173	134.845	147.982	0.000	282.827	279.828	343.354	1.609	624.791	
Nº samples (*):					43				33				54				48
Nº Ind. (*):	1876	3003	81	4960	1837	4584	249	6670	3635	4469	0	8104	2848	3693	3	6544	
Sampled catch:					375				532				585				536
Range (*):					9-56				10-55				12-53				11-49
Total catch:					2731				5721				4956				12231
Total hauls (*):					77				112				128				124

TABLE 11 (cont.).- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2009. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Mendoña* data. 2002-2009 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	1999				2000				2001				2002			
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total
4	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
6	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.325	0.325	0.000	0.141	0.475	0.616
8	0.000	0.000	1.516	1.516	0.000	0.000	0.000	0.000	0.000	0.000	1.937	1.937	0.349	0.639	0.332	1.321
10	5.154	3.352	2.960	11.465	0.000	0.793	0.000	0.793	0.104	0.356	1.850	2.310	1.315	0.712	0.000	2.027
12	12.807	8.911	0.000	21.718	3.716	1.266	0.000	4.982	0.320	1.239	1.187	2.746	0.620	0.675	0.000	1.295
14	19.227	16.710	0.000	35.938	7.773	11.915	0.000	19.687	0.952	1.477	1.114	3.543	1.544	1.064	0.000	2.608
16	13.999	15.356	0.000	29.355	10.311	10.506	0.000	20.817	3.575	4.509	0.412	8.497	1.889	2.134	0.000	4.023
18	8.893	10.757	0.000	19.650	14.266	16.475	0.000	30.741	10.107	10.530	0.149	20.786	3.180	2.479	0.000	5.660
20	14.809	10.199	0.000	25.008	16.177	19.576	0.000	35.753	17.815	24.898	0.000	42.713	7.908	6.122	0.000	14.030
22	33.285	22.789	0.000	56.073	17.231	18.660	0.000	35.891	21.299	29.178	0.000	50.477	16.552	12.664	0.000	29.217
24	61.756	39.009	0.000	100.765	21.395	20.983	0.000	42.378	24.254	23.840	0.000	48.094	21.724	22.245	0.000	43.968
26	98.561	59.521	0.000	158.083	48.000	33.100	0.000	81.100	28.911	24.809	0.000	53.720	27.246	24.307	0.000	51.553
28	107.816	84.193	0.000	192.009	67.229	39.182	0.000	106.412	58.237	33.305	0.000	91.542	40.151	22.443	0.000	62.594
30	72.947	92.236	0.000	165.183	64.336	44.684	0.000	109.020	72.412	45.107	0.000	117.519	57.549	34.445	0.000	91.994
32	28.850	75.169	0.000	104.018	36.450	53.416	0.000	89.865	49.179	59.052	0.000	108.232	46.938	50.680	0.000	97.618
34	15.810	43.595	0.000	59.405	12.695	39.970	0.000	52.665	22.267	64.772	0.000	87.039	18.047	57.599	0.000	75.646
36	9.185	24.775	0.000	33.960	6.653	25.712	0.000	32.365	8.702	46.598	0.000	55.300	7.014	45.699	0.000	52.713
38	3.658	14.964	0.000	18.623	3.526	15.747	0.000	19.274	6.293	30.315	0.000	36.608	2.651	25.514	0.000	28.165
40	1.466	8.582	0.000	10.049	1.996	10.642	0.000	12.638	2.145	12.925	0.000	15.070	1.183	12.427	0.000	13.610
42	0.262	5.318	0.000	5.580	0.286	6.803	0.000	7.089	0.857	7.788	0.000	8.645	0.616	6.257	0.000	6.873
44	0.111	2.620	0.000	2.731	0.013	4.005	0.000	4.018	0.614	4.596	0.000	5.210	0.042	2.690	0.000	2.732
46	0.028	0.988	0.000	1.016	0.000	1.806	0.000	1.806	0.221	1.968	0.000	2.190	0.024	1.150	0.000	1.174
48	0.096	0.486	0.000	0.582	0.003	0.845	0.000	0.848	0.000	0.775	0.000	0.775	0.000	0.818	0.000	0.818
50	0.000	0.140	0.000	0.140	0.000	0.246	0.000	0.246	0.000	0.242	0.000	0.242	0.020	0.149	0.000	0.169
52	0.000	0.032	0.000	0.032	0.000	0.000	0.000	0.000	0.000	0.051	0.000	0.051	0.000	0.038	0.000	0.038
54	0.000	0.000	0.000	0.000	0.000	0.033	0.000	0.033	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
56	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	508.721	539.702	4.475	1052.898	332.057	376.364	0.000	708.421	328.265	428.326	6.975	763.567	256.565	333.090	0.807	590.462
Nº samples (*):				39					42			43				43
Nº Ind. (*):	4616	5076	6	9698	3323	4100	0	7423	3358	4684	80	8122	3419	4576	7	8002
Sampled catch:				796					717			2298				2269
Range (*):				8-52					11-54			6-53				6-52
Total catch:				17169					12742			16141				14385
Total hauls (*):				114					118			123				125

TABLE 11 (cont.).- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2009. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Mendoña* data. 2002-2009 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	2003				2004				2005				2006			
	Males	Females	Indet.	Total												
4	0.000	0.000	0.009	0.009	0.000	0.000	0.116	0.116	0.000	0.000	0.000	0.000	0.060	0.000	0.000	0.060
6	0.000	0.107	0.297	0.404	0.000	0.000	0.337	0.337	0.000	0.013	0.192	0.205	0.000	0.000	0.079	0.079
8	0.036	0.121	0.274	0.431	0.109	0.049	0.741	0.899	0.269	0.018	0.054	0.341	0.187	0.162	0.245	0.594
10	0.847	0.572	0.140	1.559	0.528	0.637	0.000	1.165	1.725	0.467	0.051	2.243	0.686	0.384	0.276	1.346
12	0.969	1.205	0.000	2.174	2.005	1.577	0.000	3.582	2.353	1.877	0.000	4.229	2.026	1.734	0.000	3.760
14	0.977	0.869	0.000	1.846	3.503	2.632	0.000	6.135	4.728	3.053	0.000	7.780	3.645	3.862	0.000	7.507
16	0.946	0.289	0.000	1.234	4.580	3.608	0.000	8.188	4.674	3.630	0.000	8.304	5.776	6.009	0.000	11.785
18	1.665	1.689	0.000	3.355	4.649	3.543	0.000	8.192	3.334	3.348	0.000	6.682	5.989	5.547	0.000	11.536
20	1.695	2.233	0.000	3.928	5.414	6.205	0.000	11.619	4.905	4.847	0.000	9.752	9.721	8.196	0.000	17.917
22	4.214	4.602	0.000	8.817	5.563	5.757	0.000	11.321	8.934	6.836	0.000	15.770	10.735	10.545	0.000	21.280
24	11.364	8.741	0.000	20.105	8.232	7.732	0.000	15.964	8.930	7.162	0.000	16.092	11.073	12.977	0.000	24.050
26	27.765	19.581	0.000	47.347	25.572	16.572	0.000	42.145	15.997	8.451	0.000	24.447	13.117	13.439	0.000	26.556
28	37.413	29.153	0.000	66.566	57.974	27.637	0.000	85.611	34.840	17.504	0.000	52.344	26.251	15.412	0.000	41.663
30	52.296	29.328	0.000	81.624	87.376	52.285	0.000	139.661	75.001	34.103	0.000	109.105	64.180	25.059	0.000	89.238
32	45.761	40.076	0.000	85.836	74.712	58.683	0.000	133.396	70.556	58.866	0.000	129.423	74.126	52.415	0.000	126.541
34	19.769	52.100	0.000	71.869	30.847	58.596	0.000	89.443	28.072	62.961	0.000	91.032	38.379	67.737	0.000	106.116
36	6.757	39.555	0.000	46.312	7.531	46.290	0.000	53.820	8.105	48.672	0.000	56.777	11.021	63.706	0.000	74.727
38	2.130	23.649	0.000	25.779	2.056	26.594	0.000	28.650	1.965	26.547	0.000	28.512	3.046	39.877	0.000	42.923
40	0.832	9.444	0.000	10.276	1.716	10.932	0.000	12.648	0.908	11.697	0.000	12.606	0.981	17.493	0.000	18.474
42	0.256	3.895	0.000	4.151	0.514	3.725	0.000	4.240	0.172	4.746	0.000	4.918	0.081	5.709	0.000	5.789
44	0.268	2.432	0.000	2.700	0.028	2.033	0.000	2.061	0.050	2.020	0.000	2.070	0.072	2.190	0.000	2.262
46	0.000	1.113	0.000	1.113	0.000	0.575	0.000	0.575	0.000	1.128	0.000	1.128	0.000	1.341	0.000	1.341
48	0.000	0.525	0.000	0.525	0.000	0.303	0.000	0.303	0.000	0.200	0.000	0.200	0.000	0.560	0.000	0.560
50	0.000	0.202	0.000	0.202	0.000	0.009	0.000	0.009	0.000	0.030	0.000	0.030	0.000	0.231	0.000	0.231
52	0.000	0.009	0.000	0.009	0.000	0.055	0.000	0.055	0.000	0.000	0.000	0.000	0.000	0.012	0.000	0.012
54	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.079	0.000	0.079	0.000	0.091	0.000	0.091
56	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	215.959	271.489	0.721	488.169	322.910	336.032	1.193	660.136	275.518	308.254	0.297	584.069	281.150	354.688	0.601	636.440
Nº samples (*):				37				45				48				45
Nº Ind. (*):	2424	3254	12	5690	3703	4234	16	7953	4790	6556	6	11352	4404	6012	10	10426
Sampled catch:				1864				2587				3784				3407
Range (*):				5-52				5-53				6-55				5-54
Total catch:				11280				15117				14275				15424
Total hauls (*):				118				120				119				120

TABLE 11 (cont.).- Yellowtail flounder length distribution. Estimated numbers per haul mean catches. Spanish Spring Survey on NAFO 3NO: 1995-2009. Indet. means indeterminate. 1995-2000 data are transformed C/V *Playa de Mendoña* data. 2002-2009 data are original R/V *Vizconde de Eza* data. In 2001, there are data from the two vessels. (*) indicates untransformed data.

Length (cm.)	2007				2008				2009			
	Males	Females	Indet.	Total	Males	Females	Indet.	Total	Males	Females	Indet.	Total
4	0.000	0.000	0.000	0.000	0.000	0.000	0.054	0.054	0.000	0.000	0.000	0.000
6	0.000	0.000	0.103	0.103	0.000	0.000	0.000	0.000	0.000	0.000	0.050	0.050
8	0.000	0.000	0.000	0.000	0.013	0.000	0.000	0.013	0.000	0.000	0.057	0.057
10	0.041	0.059	0.000	0.101	0.039	0.000	0.037	0.076	0.000	0.155	0.000	0.155
12	0.536	0.449	0.000	0.985	0.184	0.183	0.000	0.367	0.000	0.063	0.370	0.433
14	1.148	0.578	0.000	1.725	0.238	0.331	0.054	0.624	0.000	0.096	0.000	0.096
16	2.222	2.551	0.000	4.773	0.741	0.964	0.000	1.705	0.920	0.498	0.000	1.418
18	5.728	4.614	0.000	10.342	2.364	2.973	0.000	5.337	2.260	1.452	0.000	3.712
20	9.024	7.293	0.000	16.317	7.593	6.160	0.000	13.753	4.032	3.251	0.000	7.283
22	13.286	14.190	0.000	27.476	11.867	13.532	0.000	25.399	11.271	7.825	0.000	19.096
24	17.380	19.046	0.000	36.426	18.209	18.285	0.000	36.495	15.826	15.693	0.000	31.518
26	20.689	18.113	0.000	38.802	23.627	25.866	0.000	49.493	28.577	26.217	0.000	54.793
28	35.157	19.170	0.000	54.327	37.293	23.056	0.000	60.349	38.271	24.052	0.000	62.323
30	75.144	25.235	0.000	100.379	67.815	22.281	0.000	90.096	59.751	26.094	0.000	85.844
32	76.329	50.253	0.000	126.582	73.491	42.910	0.000	116.401	73.655	42.701	0.000	116.356
34	42.232	68.548	0.000	110.780	38.260	59.348	0.000	97.609	44.085	74.201	0.000	118.285
36	12.733	61.691	0.000	74.424	9.789	54.190	0.000	63.979	13.976	81.708	0.000	95.684
38	3.973	41.839	0.000	45.812	2.389	37.201	0.000	39.590	4.267	54.934	0.000	59.200
40	1.430	20.920	0.000	22.350	0.914	16.185	0.000	17.099	0.983	22.221	0.000	23.203
42	0.213	6.891	0.000	7.104	0.288	6.719	0.000	7.007	0.103	11.373	0.000	11.476
44	0.000	2.454	0.000	2.454	0.000	3.120	0.000	3.120	0.039	4.532	0.000	4.571
46	0.071	1.043	0.000	1.114	0.000	1.097	0.000	1.097	0.000	1.183	0.000	1.183
48	0.000	0.367	0.000	0.367	0.000	0.616	0.000	0.616	0.000	0.173	0.000	0.173
50	0.000	0.107	0.000	0.107	0.000	0.077	0.000	0.077	0.000	0.460	0.000	0.460
52	0.000	0.120	0.000	0.120	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
54	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
56	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total	317.336	365.532	0.103	682.971	295.113	335.096	0.145	630.355	298.014	398.879	0.477	697.369
Nº samples (*):				47				50				38
Nº Ind. (*):	5083	5533	1	10617	4795	5147	3	9945	3969	4682	5	8656
Sampled catch:				2761				2759				2604
Range (*):				7-52				5-51				7-50
Total catch:				15200				14697				16201
Total hauls (*):				110				122				109




FIGURE 1.- Atlantic cod stratified mean catches in Kg and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1997-2009 (1997-2000 transformed data from C/V *Playa de Menduña*; 2002-2009 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 2.- Atlantic cod biomass calculated by the swept area method in tons and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1997-2009 (1997-2000 transformed data from C/V *Playa de Menduña*; 2002-2009 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 3.- Atlantic cod length distribution (cm) on NAFO 3NO: 1997-2009 . Mean catches per tow numbers. 1997-2000 data are transformed data from C/V *Playa de Menduña*, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels.




FIGURE 4.- Atlantic cod mean catches per tow length distribution (cm) on NAFO 3NO: 1997-2009.




FIGURE 5.- Yellowtail flounder stratified mean catches in Kg and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1995-2009 (1995-2000 transformed data from C/V *Playa de Menduña*; 2002-2009 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 6.- Yellowtail flounder biomass calculated by the swept area method in tons and \pm SD by year. Spanish Spring surveys on NAFO Div. 3NO: 1995-2009 (1995-2000 transformed data from C/V *Playa de Menduña*; 2002-2009 original data from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels).




FIGURE 7.- Yellowtail flounder length distribution (cm) on NAFO 3NO: 1995-2009. Mean catches per tow numbers. 1995-2000 data are transformed data from C/V *Playa de Menduña*, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels




FIGURE 7 (Cont.).- Yellowtail flounder length distribution (cm) on NAFO 3NO: 1995-2009. Mean catches per tow numbers. 1995-2000 data are transformed data from C/V *Playa de Menduña*, and 2002-2009 data are original from R/V *Vizconde de Eza*. In 2001, there are data from the two vessels




FIGURE 8.- Yellowtail flounder mean catches per tow length distribution (cm) on NAFO 3NO: 1995-2009.